

Adverse effect of Maternal Anemia on Fetal Doppler Indices and estimated fetal weight during the Last trimester of pregnancy in rural and suburban area

Abdel Samie Abdel Monaïem Abdel Samie, Al Mandouh Hussein Bosilah, Rana Ahmed Mohammed Farouk

Department of Obstetrics and Gynecology, Faculty of Medicine, Fayoum University, Fayoum, Egypt.

ABSTRACT

Background: Anemia is an important global health problem affecting nearly 529 million women of reproductive age, including 38% of all pregnant women. Anemia during pregnancy is a key contributor to maternal mortality and poor birth outcomes in both low- and high-income countries. Reducing maternal anemia is widely recognized as central to the health of women and children. Current WHO global targets call for a 50% reduction of anemia in women of reproductive age by 2025. Maternal anemia is a hypoxic condition that could be responsible for the redistribution of fetal blood flow, however, no evidence of placental insufficiency has been documented currently. In the presence of fetal hypoxemia, vasodilatation of the fetal middle cerebral artery occurs resulting in the so-called "brain sparing effect" to preserve cerebral oxygenation this plays a major role in fetal adaptations to oxygen deprivation (compensatory flow or adaptation changes).

Objective: The aim of this study was to evaluate the effect of maternal anemia on fetal Doppler Indices namely umbilical artery and middle cerebral artery and subsequently fetal growth rate.

Patients and Methods: This study was designed as a prospective comparative study was carried out in the Departments of Obstetrics & Gynecology, Faculty of Medicine, Fayoum University including 200 pregnant women. The pregnant women were divided into three groups: Group A: Patients with mild anemia (Hb concentration: 9.0-10.9g/dl). Group B: Patients with moderate anemia (Hb concentration: 7.0- 8.9g/dl). Group C: Patients with severe anemia (Hb concentration: >7.0g/dl). A "Mild" is a misnomer: iron deficiency is already advanced by the time anemia is detected. The duration of the study ranged from 6 to 12 months. Patients aged 20-35 years with singleton living fetus of >_32 weeks gestation at time of first visit with no history of chronic illness or medical disorder other than iron deficiency anemia such as chronic hypertension, D.M. or

chronic blood loss e.g.; peptic ulcer and no history of recurrent perinatal deaths, recent blood transfusion or other vitamin deficiency anemia

Results: Our results reported that there was no significant difference between the studied groups regarding age and BMI. As regard fetal Doppler cerebral artery Indices of the three studied groups, the current study shows that there is a significant difference between the three studied groups regarding middle cerebral artery resistance index, middle cerebral artery pulsatility index and middle cerebral artery systolic/diastolic ratio. In the study in our hands, there is a significant difference between the three studied groups regarding umbilical artery resistance index, umbilical artery pulsatility index, umbilical artery systolic/diastolic ratio and middle cerebral/ umbilical artery resistance index. The present study shows that there is a significant difference between the three studied groups regarding biparietal diameter, abdominal circumference and femur length. There is a significant difference between the three studied groups regarding amniotic fluid index at 10 days later and at delivery. Our results reported that as for the AFI, no significant effect was noticed in the 3 groups. In the study in our hands, there is a significant difference between the three studied groups regarding middle cerebral artery resistance index at 10 days later and at delivery, umbilical artery resistance index at 10 days later, and middle cerebral/ umbilical artery resistance index at 10 days later and at delivery. There is a significant difference between the three studied groups regarding Apgar at 5 min and birth weight.

Conclusion: There was a significant effect of maternal anemia on fetal Doppler indices in both umbilical artery and middle cerebral artery in the last trimester of pregnancy and this effect is directly proportionate to the severity of anemia.

Keywords: Hemoglobin concentration, middle cerebral artery resistance index, umbilical artery resistance index, middle cerebral/ umbilical artery resistance index, birth weight.

INTRODUCTION

Anemia is defined as a condition in which the body has a decreased number of circulating erythrocytes, or red blood cells (RBCs). It can also be defined as a decrease hemoglobin concentration or RBC mass compared with age-matched controls. As with almost all human laboratory assays, normal value is a statistical term used to define a range within which 95% of the population's values fall. The World Health Organization (WHO) defines anemia as a hemoglobin less than 13 g/dL in adult men and less than 12 g/dL in non-pregnant adult women. However, these values were chosen somewhat arbitrarily; most laboratories define anemia as the lowest 2.5% of the distribution of hemoglobin values from a normal, healthy population (1).

Erythrocytes originate in the bone marrow as hematopoietic progenitor and precursor cells. After several cell divisions, mature RBCs emerge as discoid, pliable anucleate cells, each containing 4 hemoglobin molecules. An erythrocyte typically survives for 100 to 120 days before undergoing apoptosis (programmed cell death). Erythropoiesis, or the process of RBC production, occurs in a regulated fashion under the control of the hormone erythropoietin (EPO).

Patients and Methods

Study Design: This study was designed as a prospective a prospective comparative cross-sectional study include 200 pregnant women who attended.

Study Site: at The Departments of Obstetrics & Gynecology. Faculty of Medicine, Fayoum University

Duration of the study: The study was done during the period between July 2019 and January 2019.

Sample Size: Total sample size in the study was calculated to be two hundred patients (200). The pregnant women were divided into three groups:
- Group A: Patients with mild anemia (Hb concentration: 9.0-10.9g/dl). Group B: Patients

In pregnancy, more than 50% of women in underdeveloped or developing nations will develop anemia. In developed nations, this rate decreases to 20%. In the United States, the biggest risk factor for developing anemia in pregnancy is low socioeconomic status; nutritional deficiencies and chronic disease also contribute (2).

Folate deficiencies since 1960s and B12 vitamin deficiencies since 1990s defined as contributing causes of nutritional anemia. Folate deficiency is common as a result of dietary deficiency or increased demand as in pregnancy. The prevalence of folate deficiency in pregnant patients varies from

1 to 50%. The prevalence is higher among economically deprived patients. B12 vitamin is mostly found in foods of animal origin and deficiency was not reported as a frequent cause of anemia. Anemia due to zinc, copper, vitamin A or other vitamin and minerals are also considered in the literature (3).

Aim of the Study

This study aims to evaluate the effect of maternal anemia on fetal Doppler Indices namely umbilical artery and middle cerebral artery and subsequently fetal growth rate.

with moderate anemia (Hb concentration: 7.0-8.9g/dl). Group C: Patients with severe anemia (Hb concentration: >7.0g/dl). A "Mild" is a misnomer: iron deficiency is already advanced by the time anemia is detected. The deficiency has consequences even when no anemia is clinically apparent. The hemoglobin cut-off of 11 g/dl for pregnant women was first presented in the 1968.

With the following criteria for three groups:

Inclusion criteria: Patients aged 20-35 years with singleton living fetus of >_32 weeks' gestation at time of first visit with no history of chronic illness or medical disorder other than iron deficiency anemia such as chronic hypertension, D.M. or chronic blood loss e.g.; peptic ulcer and

no history of recurrent perinatal deaths, recent blood transfusion or other vitamin deficiency anemia

Exclusion criteria: Patients with pregnancy associated disorders as: Preeclampsia, gestational diabetes or placenta previa (recurrent vaginal bleeding) were excluded from the study.

Data management and Statistical Analysis: Data collected throughout history, basic clinical examination, laboratory investigations and outcome measures coded, entered and analyzed using Microsoft Excel software. Data were then

imported into Statistical Package for the Social Sciences (SPSS version 20.0) (Statistical Package for the Social Sciences) software for analysis. According to the type of data qualitative represent as number and percentage, quantitative continues group represent by mean \pm SD, the following tests were used to test differences for significance; correlation by Pearson's correlation or Spearman's. P value was set at <0.05 for significant results & <0.001 for high significant result. Data were collected and submitted to statistical analysis. The following statistical tests and parameters were used.

RESULTS

Table 1 shows that There is no significant difference between the three studied groups regarding age and BMI. Table 2 shows There is a

significant difference between the three studied groups regarding ferritin.

Table 1: Demographic data of the three studied groups

	<i>Group A</i> (N=67)	<i>Group B</i> (N=67)	<i>Group C</i> (N=66)	<i>F</i>	<i>P</i>
<i>Age (years) Mean \pm SD</i>	29.56 \pm 5.31	30.7 \pm 4.91	31.1 \pm 4.12	1.84	.161
<i>BMI (kg/m²) Mean \pm SD</i>	27.29 \pm 2.69	26.12 \pm 3.39	26.61 \pm 3.84	2.08	.128

Table 2: Specific Laboratory parameters of the three studied groups.

	<i>Group A</i> (N=67)	<i>Group B</i> (N=67)	<i>Group C</i> (N=66)	<i>F</i>	<i>P</i>
<i>Ferritin (ng/mL) Mean \pm SD</i>	35.2 \pm 7.64	28.44 \pm 8.55	22.71 \pm 9.65	34.8	.000
<i>RBS (mg/dL) Mean \pm SD</i>	110.75 \pm 14.5	141.96 \pm 17.65	111.8 \pm 14.52	1.32	.271

Table 3 shows There is a significant difference between the three studied groups regarding biparietal diameter, abdominal circumference and femur length. Table 4 shows

that There is a significant difference between the three studied groups regarding Apgar at 5 min and birth weight.

Table 3: Fetal biometry of the three studied groups.

	Group A (N=67)	Group B (N=67)	Group C (N=66)	F	P
Biparietal diameter <i>Mean ± SD</i>	77.15 ± 3.86	75.42 ± 4.53	72.87 ± 5.18	8.14	.000
Abdominal circumference <i>Mean ± SD</i>	25.25 ± 3.68	24.11 ± 3.92	22.34 ± 4.22	9.18	.000
Femur length <i>Mean ± SD</i>	56.9 ± 3.49	55.18 ± 3.98	52.97 ± 4.17	11.1	.000

Table 4: Fetal characteristics of the three studied groups

	Group A (N=67)	Group B (N=67)	Group C (N=66)	F	P
Apgar at 5 min <i>Mean ± SD</i>	8.62 ± 1.46	8.81 ± 1.56	7.46 ± 2.55	14.2	.000
Birth weight (kg) <i>Mean ± SD</i>	3.33 ± 0.98	2.91 ± 1.06	2.64 ± 1.21	5.81	.002

Table 5 shows that There is a significant difference between the three studied groups regarding middle cerebral artery resistance index,

middle cerebral artery pulsatility index and middle cerebral artery systolic/diastolic ratio.

Table 5: Fetal Doppler cerebral artery Indices of the three studied groups

	Group A (N=67)	Group B (N=67)	Group C (N=66)	F	P
Middle cerebral artery resistance index. Mean ± SD	.963 ± .066	.939 ± .041	1.08 ± .079	21.6	.000
Middle cerebral artery pulsatility index. Mean ± SD	1.41 ± .225	1.47 ± .165	1.62 ± .155	7.91	.001
Middle cerebral artery systolic/diastolic ratio index. Mean ± SD	3.82 ± .744	3.92 ± .701	3.41 ± .548	3.56	.034
Middle cerebral artery peak systolic velocity index. Mean ± SD	40.74 ± 6.33	40.78 ± 6.53	42.43 ± 7.12	0.460	.634

Table 6 shows that There is a significant difference between the three studied groups regarding amniotic fluid index at 10 days later and at delivery. Table 7 shows that There is a significant difference between the three studied

groups regarding umbilical artery resistance index, umbilical artery pulsatility index, umbilical artery systolic/diastolic ratio and middle cerebral/ umbilical artery resistance index.

Table 6: Amniotic fluid index of the three studied groups

	Group A (N=67)	Group B (N=67)	Group C (N=66)	F	P
AFI on admission (cm) Mean \pm SD	13.44 \pm 0.75	13.61 \pm 0.56	13.39 \pm 0.81	2.74	.087
AFI 10-days later (cm) Mean \pm SD	12.12 \pm 1.23	12.43 \pm 1.31	12.78 \pm 1.09	4.92	.007
AFI at delivery (cm) Mean \pm SD	10.89 \pm 1.65	11.24 \pm 2.03	11.95 \pm 2.16	5.05	.005

Table 7: Fetal Doppler Umbilical artery Indices of the three studied groups.

	Group A (N=67)	Group B (N=67)	Group C (N=66)	F	P
Umbilical artery resistance index Mean \pm SD	.872 \pm .067	.818 \pm .056	.987 \pm .062	8.63	.000
Umbilical artery pulsatility index Mean \pm SD	.729 \pm .081	.777 \pm .093	1.11 \pm .275	13.8	.000
Umbilical artery systolic/diastolic ratio Mean \pm SD	2.18 \pm .043	2.42 \pm .345	2.75 \pm .705	8.72	.000
Middle cerebral/ Umbilical artery resistance index Mean \pm SD	.876 \pm .093	.868 \pm .081	.978 \pm .084	18.7	.000

DISCUSSION

Anemia in general is characterized by a decrease in number of red blood cells or less than the normal quantity of hemoglobin. The condition is determined by the expected normal range of hemoglobin in a population, and is defined as existing in an individual whose hemoglobin concentration (Hb) has fallen below a threshold lying at two standard deviations below the median for a healthy population of the same demographic characteristics, including age, sex and pregnancy status (4).

Most of the work of erythropoiesis occurs in the bone marrow. It takes about 25 days. In children younger than 5 years old, the marrow of all the bones of the body is enlisted for. According to United Nation declaration 1997, anemia is a major public health problem that needs total elimination. It is estimated that

globally two billion people suffer from anemia or iron deficiency (6).

Maternal anemia is frequently associated with premature delivery, reduced neonatal weight, infant iron deficiency, neonatal death, and low APGAR scores at 1 min. It is also suspected to reduce the oxygen supply to the growing fetus, leading to the redistribution of fetal blood flow (7).

There is no significant difference between the three studied groups regarding age and BMI. Our results agree with study of Khalafallah et al., (8) as they reported that there was no significant difference between the studied groups regarding age and BMI. Furthermore, Sharbaf et al., (9) demonstrated that independent t-test analysis did not show any significant

differences between both groups regarding maternal age ($p=0.51$).

Anemia is an important global health problem affecting nearly 529 million women of reproductive age, including 38% of all pregnant women. Anemia during pregnancy is a key contributor to maternal mortality and poor birth outcomes in both low- and high- income countries. Reducing maternal anemia is widely recognized as central to the health of women and children. Current WHO global targets call for a 50% reduction of anemia in women of reproductive age by 2025 (10).

The present study shows that there is a significant difference between the three studied producing red blood cells. As a person ages, red blood cells are eventually produced only in the marrow of the spine, ribs, and pelvis (5). The groups regarding Hb, MCV, MCH and MCHC. There is a significant difference between the three studied groups regarding ferritin. Our results are supported by study of Abdel- Megeed et al., (11) as they reported that in this study the hemoglobin level in anemic patients ranged from 5.0-10.9 g/dl while in non-anemic was 11-13.8 g/dl, there was a significant decrease in hemoglobin level in anemic more than non-anemic, these results obeying the inclusion and exclusion criteria.

Furthermore, Kumar et al., (12) found that there is a significant difference between the three studied groups regarding Hb, MCV, MCH. Prevalence of anemia in pregnancy is quite high. Their figures showed more than half of the mothers were anemic to start with. Similar figures are quoted in other studies as well. The trend in Hb level is quite interesting. The mid-trimester drops seen in nonanemic mothers was not seen in anemic mothers. This indicates that mid-trimester drop in Hb is a very essential physiological arrangement. However, in anemic mothers, as the mean Hb in the mid-trimester was below that in nonanemic mothers, no drop phenomenon was seen (13).

Furthermore, Lasheen et al., (14) revealed that there was a statistically significant difference among Hb levels on admission among the three studied groups.

The middle cerebral artery (MCA) may also predict fetal outcomes from alterations in cerebral blood flow and its direction. It was confirmed that during the prenatal period, the resistance in the cerebral artery of the fetus is high. However, this parameter can change in threatening conditions, such as placental

insufficiency and hypoxemia, due to stimulation of chemoreceptors and alteration in vasodilator or vasoconstrictor agents. MCA Doppler measurement has been advocated as an efficient modality for detection of fetal hypoxia, which leads to perinatal adverse outcome and fetal compromise (15).

As regard fetal Doppler cerebral artery Indices of the three studied groups, the current study shows that there is a significant difference between the three studied groups regarding middle cerebral artery resistance index, middle cerebral artery pulsatility index and middle cerebral artery systolic/diastolic ratio.

Our results are in line with study of Abdel-Megeed et al., (11) as they showed comparison between the different studied groups regarding Middle cerebral artery resistance index, pulsatility index, systolic/diastolic ratio and peak systolic velocity. Middle cerebral artery resistance index in control group ranged from 0.792- 0.923 with mean value 0.821 ± 0.035 , in mild anemia group ranged from 0.78-0.98 with mean value 7.34 ± 0.06 , in moderate anemia group ranged from 0.79-0.91 with mean value 0.81 ± 0.06 , in severe anemia group ranged from 0.56-0.80 with mean value 0.70 ± 0.08 .

There was statistically significant relation between the different studied groups regarding Middle cerebral artery resistance index ($P< 0.05$). Ali et al., (16) stated that fetal Doppler examination at admission found that severe maternal anemia induced significant hemodynamic changes at the cerebral level. The mean middle cerebral artery resistance index was significantly lower in the anemia group than in the control group. This could be due to hypoxia-induced cerebral vasodilatation affecting the fetus during severe anemia. Furthermore, Sharbaf et al., (9) demonstrated that with regard to Doppler examination, MCA PI between the studied groups

was significantly different. Some studies have demonstrated that diagnosis and management of FGR rely on umbilical artery Doppler (UAD) and others have shown the benefits of uterine artery Doppler assessment in the first-trimester development of FGR. The value of combining pulsatility indices (PI) of both the umbilical artery (UA) and uterine artery for improving prediction of FGR has also been demonstrated (18).

In the study in our hands, there is a significant difference between the three studied groups regarding umbilical artery resistance index, umbilical artery pulsatility index, umbilical artery systolic/diastolic ratio and middle cerebral/ umbilical artery resistance index.

Our results are supported by study of Abdel-Megeed et al., (11) as they reported that the umbilical artery resistance index in control group ranged from 0.50-0.68, in mild anemia group ranged from 0.54-0.71, in moderate anemia group ranged from 0.59-0.74, in severe anemia group ranged from 0.60-0.77. There was statistically significant relation between the different studied groups regarding Umbilical artery resistance index ($P < 0.05$).

Umbilical artery pulsatility index in control group ranged from 0.77-0.85, in mild anemia group ranged from 0.60-0.88, in moderate anemia group ranged from 0.67-0.90, in severe anemia group ranged from 0.77-1.80. There was statistically significant relation between the different studied groups regarding Umbilical artery pulsatility index ($P < 0.05$). Umbilical artery systolic/diastolic ratio in control group ranged from 2.09-2.2, in mild anemia group ranged from 2.13-2.23 with mean value, in moderate anemia group ranged from 2.10-3.40, in severe anemia group ranged from 2.15-4.30. There was statistically significant relation between the different studied groups regarding Umbilical artery systolic/diastolic ratio ($P < 0.05$).

In Ali et al. (16) study, the middle cerebral/umbilical artery resistance ratio was significantly lower at admission in the anemia group; suggesting that the fetus had adapted to

anemic conditions by redistributing blood flow towards brain.

The middle cerebral/umbilical artery resistance ratio in the anemia group increased significantly between admission and 4weeks, and between admission and 6weeks. Severe maternal anemia results in the fetus experiencing a hypoxic state and this triggers a marked fetal cerebral vasodilatation to maintain oxygenation of brain tissue, which was evident in the present study by the observed decrease in the middle cerebral artery resistance index. Long-standing cerebral vasodilatation can result in a fetus experiencing cerebral edema and brain damage.

In moiety & Ahmed (21) study, as for the C/U ratio, in groups B and C [moderate and severe anemia], the C/U values were below the normal range (<1.1), which confirms that the fetus had to adapt by increasing its blood flow redistribution towards the brain. Such adaptation was confirmed by the increase of the C/U ratio after maternal red blood cell transfusion and parenteral iron supplementation. The increase in cerebral resistance after the transfusion without significant change in umbilical or uterine resistance confirms that maternal anemia does not create placental dysfunction and that the situation can be restored quickly by two units of red blood transfusion to the patient or intravenous iron.

In Group (A) [mild anemia], the C/U values were within the normal range, which means that the blood flow distribution between the brain and placenta was normal, despite the maternal hemoglobin content being significantly lower compared to normal

The timing of the measurement of maternal anemia is also variable across studies. This is problematic for meta- analyses, as existing literature suggests a differential impact of maternal Hb concentration depending on the timing during pregnancy.

For low Hb concentration, the link with adverse birth outcomes is more evident when Hb concentration is measured in early pregnancy. As summarized in a recent review, several studies have observed a U- shaped curve for the risk of adverse birth outcomes with maternal Hb, that is, there is a higher risk of adverse outcomes among

women with both low and high Hb concentrations. However, the consistency of this relationship varies based on the trimester of Hb assessment (19).

The present study shows that there is a significant difference between the three studied groups regarding biparietal diameter, abdominal circumference and femur length. There is a significant difference between the three studied groups regarding amniotic fluid index at 10 days later and at delivery.

Findings of our results are not coincided with study of Lasheen et al., (14) as they reported that as for the AFI, no significant effect was noticed in the 3 groups but a potential benefit might rise which is the prevention of development of oligohydramnios as pregnancy advances especially in group (C) [severe anemia]. This is similar to what have been reported by Milan et al., (20) in 2005 that there was no statistically significant increase in AFI after treatment.

Anemia in pregnant women has been regarded as detrimental to the fetal growth and pregnancy outcome. Low birth weight and preterm delivery have been persistently linked to anemia in pregnancy. The role of anemia in pregnancy and iron on the growing fetus has been studied in the last few decades.

The outcome of these studies is either inconclusive or at the most supportive of the popular notions held so far regarding pregnancy outcome and anemia. Hence, most of the countries have adopted the policy of supplementing pregnant women with iron and folic acid with a view that increasing the Hb levels has some beneficial effect (17).

In the study in our hands, there is a significant difference between the three studied groups regarding middle cerebral artery resistance index at 10 days later and at delivery, umbilical artery resistance index at 10 days later, and middle cerebral/ umbilical artery resistance index at 10 days later and at delivery. There is a significant difference between the three studied groups regarding Apgar at 5 min and birth weight.

However, Swain et al., (22) have reported an inverse relation between the second trimester Hb and birth weight. The analysis of qualitative variables in our data has revealed noteworthy clinically relevant results.

The incidence of low birth weight babies in the two groups remained similar with minimal difference and there was no statistical significance in the first two trimesters. The difference was significant only if the mothers were anemic in the third trimester. Many studies conducted by Levy et al., (23) have reported significant association between low birth weight and maternal anemia.

A retrospective study has shown no association between the first trimester anemia and low birth weight (24). Most of these studies have considered the Hb levels in the third trimester or at delivery for comparison (23). This could well suggest that third trimester Hb is an important factor in determining birth weight.

It is well known that rapid growth of fetus occurs in the third trimester. Iron and other micronutrient accretion rates are the highest in the same trimester as well. This physiology explains the association of third trimester Hb and low birth weight. In contrast to the above, association of preterm delivery with maternal anemia is quite striking. Except for the first trimester, anemia in other trimesters has shown significantly increased incidence of preterm delivery. This association appears strongest in the third trimester. There are many studies conducted by Lone et al., (25) showing similar association.

CONCLUSION

There was a significant effect of maternal anemia on fetal Doppler indices in both umbilical artery and middle cerebral artery in the last trimester of pregnancy and this effect is directly proportionate to the severity of anemia. Close monitoring of the fetal umbilical and cerebral circulation by Doppler examination in anemia pregnant females allows the measurement of the amplitude of the fetal vascular response as in severe maternal anemia strongly associated with marked fetal hemodynamic adaptation and trigger fetal blood flow redistribution.

REFERENCES

- 1) Calis, J. C., Phiri, K. S., Faragher, E. B., Brabin, B. J., Bates, I., Cuevas, L. E., ... & Hulshof, P. J. (2016). Research Article (New England Journal of Medicine) Severe anemia in Malawian children. *Malawi Medical Journal*, 28(3), 99-107.
- 2) Sawyer, S. L., Tian, L., Kähkönen, M., Schwartzentruber, J., Kircher, M., Majewski, J., ... & Moilanen, J. S. (2015). Biallelic mutations in BRCA1 cause a new Fanconi anemia subtype. *Cancer discovery*, 5(2), 135-142.
- 3) Karaoglu L, Pehlivan E, Egri M (2010): The prevalence of nutritional anemia in pregnancy in an east Anatolian province, Turkey. *BMC Public Health*, 10: 329.
- 4) McLean E, Cogswell M, Egli I (2009): Worldwide prevalence of anaemia, WHO Vitamin and Mineral Nutrition Information System, 1993-2005. *Public Health Nutr.*, 12: 444-54.
- 5) Kalantar-Zadeh K, Block G, McAllister CJ, Humphreys MH, Kopple JD (2004): The American Journal of Clinical Nutrition, 80(2): 299-307.
- 6) RCOG guidelines (2014): New S., Wirth M. Anemia, pregnancy, and maternal mortality: The problem with globally standardized hemoglobin cutoffs. *B.J.O.G.*, 122: 166-9, 2015.
- 7) CHRISTOPHER V. CHARLES: 2012. Iron Deficiency Anemia: A Public Health Problem of Global Proportions, *Public Health-Methodology, Environmental and Systems Issues*, Prof. Jay Maddock (Ed.), ISBN: 978-953-51-0641-8.
- 8) Khalafallah, A., Dennis, A., Bates, J., Bates, G., Robertson, I. K., Smith, L., ... & Rasko, J. E. J. (2010). A prospective randomized, controlled trial of intravenous versus oral iron for moderate iron deficiency anaemia of pregnancy. *Journal of internal medicine*, 268(3), 286-295.
- 9) Sharbaf, F. R., Movahed, F., Pirjani, R., Teimoory, N., Shariat, M., & Farahani, Z. (2018). Comparison of fetal middle cerebral artery versus umbilical artery color Doppler ultrasound for predicting neonatal outcome in complicated pregnancies with fetal growth restriction. *BIOMEDICAL RESEARCH AND THERAPY*, 5(5), 2296-2304.
- 10) WHO . 2014. Global nutrition targets 2025: anaemia policy brief (WHO/NMH/NHD/14.4). Geneva: World Health Organization.
- 11) Abdel-Megeed, A. M. I., Riad, A. A. M., & Elsherif, S. M. M. M. (2018). Effect of Maternal Anemia on Fetal Doppler Indices during the Last Trimester of Pregnancy. *The Egyptian Journal of Hospital Medicine*, 73(2), 6082-6089.
- 12) Kumar, K. J., Asha, N., Murthy, D. S., Sujatha, M. S., & Manjunath, V. G. (2013). Maternal anemia in various trimesters and its effect on newborn weight and maturity: an observational study. *International journal of preventive medicine*, 4(2), 193.
- 13) Kalaivani K. 2009. Prevalence and consequences of anemia in pregnancy. *Indian J Med Res.*;130:627-33.
- 14) Lasheen, Y. S. A. (2011). Effect of maternal iron deficiency anemia on fetal cerebral hemodynamic response by doppler, biophysical profile and APGAR score.
- 15) Nomura, JI Niigaki, FT Horigome, RPV Francisco, M Zugaib. 2013. Doppler velocimetry of the fetal middle cerebral artery and other parameters of fetal well-being in neonatal survival during pregnancies with placental insufficiency. *Revista da Associação Médica Brasileira.*; 59 : 392-399.
- 16) Ali E, Kumar M, Naqvi SE et al. (2016): Fetal vascular adaptation before and after treatment of severe maternal anemia in pregnancy. *International Journal of Gynecology and Obstetrics*, 133(3): 284-286.
- 17) Kalaivani K. 2009. Prevalence & consequences of anaemia in pregnancy. *Indian J Med Res.* 2009 Nov; 130(5):627- 33.
- 18) Levytska, M Higgins, S Keating, N Melamed, M Walker, NJ Sehire. 2017. Placental pathology in relation to uterine artery Doppler findings in pregnancies with severe intrauterine growth restriction and abnormal umbilical artery Doppler changes. *American journal of perinatology.*; 34 : 451-457 .

-
- 19) Dewey K.G. & Oaks B.M.. 2017. U- shaped curve for risk associated with maternal hemoglobin, iron status, or iron supplementation. *Am. J. Clin. Nutr.* 106(Suppl. 6): 1694s–1702s.
 - 20) MILAN STEFANOVIC, MILOSAVLJEVIC M., RADOVIC-JANOFEVIC V., KUTLESIC R. and VUKOMANOVIC P2005..: Maternal anemia and fetal cerebral hemodynamic response- Doppler assessment. *Medison and Biology*, (1.12) No. 2 : 93-96,
 - 21) 21) Moiety G and Ahmed Y (2012): Health care professionals, teachers, students and information professionals Arab World rely on Medical journal of Cairo university as a trusted source.The Medical Journal of Cairo University, 80(2): 15-35.
 - 22) Rasmussen S, Oian P. 1993. First- and second-trimester hemoglobin levels. Relation to birth weight and gestational age. *Acta Obstet Gynecol Scand.* May; 72(4):246-51.
 - 23) Levy A, Fraser D, Katz M, Mazor M, Sheiner E.2005. Maternal anemia during pregnancy is an independent risk factor for low birthweight and preterm delivery. *Eur J Obstet Gynecol Reprod Biol.* 2005 Oct 1; 122(2):182-6.
 - 24) Phaloprakarn C, Tangjitgamol S.2008. Impact of high maternal hemoglobin at first antenatal visit on pregnancy outcomes: a cohort study. *J Perinat Med.* 2008; 36(2):115-9.
 - 25) Lone FW, Qureshi RN, Emanuel F.2004. Maternal anaemia and its impact on perinatal outcome. *Trop Med Int Health.* 2004 Apr; 9(4):486-490.