



STUDY OF SOME BIOLOGICAL AND DYNAMIC CHARACTERISTICS OF THE MEDEIRAN SARDINELLA FISH *SARDINELLA MADERENSIS* OF EASTERN MEDITERRANEAN (NORTH SINAI COAST EGYPT)

Esraa A. El-Sayed^{1*}; G.D.I. Hassanan¹; Rasha M. Faisal¹ and A.A.O. El-Aiatt²

1. Dept. Fisheries and Aquaculture, Fac. Environ. Agric. Sci., Arish Univ., Egypt.

2. Nat. Inst. Oceanography and Fisheries, Egypt.

ARTICLE INFO

Article history:

Received: 17/04/2023

Revised: 04/05/2023

Accepted: 10/05/2023

Keywords:

Sardinella maderensis,
Growth parameters,
Mortality,
Gonadosomatic index.

ABSTRACT

Monthly collected samples of *S. maderensis* from January to December 2021 in Mediterranean coast of Sinai for determination the growth and mortality. The lengths ranged from 8.5 - 26.3 cm and the weights ranged from 5g to 171g. Value b of the relationship between length-weight was 3.0524 thus, the growth of *S. maderensis* categorized as displaying isometric growth. (VBGP) of L_{∞} , (k) and (t₀) were 30.39cm, 0.2937 and -0.6049/year, respectively. The growth performance index (Φ') was 2.334. GSI ranged from 0.11 to 2.6 of males and ranged from 0.19 to 3.4 of females, also the peak spawning periods of *S. maderensis* were the period from April to September. The lengths at first maturity were 15.5 cm (males), 16.7 cm (females) and 16.0 cm for combined sexes. Values of total mortality (Z), natural mortality (M), fishing mortality (F) and exploitation rate (E) of combined sexes of *S. maderensis* were 0.7124, 0.3229, 0.3895 year⁻¹ and 0.547, respectively. Length at first capture L₅₀ was 13.5 cm. Should be not increase the fishing effort but should be increasing the size of the nets so that the length at first capture increases from 13.0 cm to 16.0 cm in order to equal the length at first maturity to preserve the stock of these fish.



INTRODUCTION

Sardine fish in the Port of El Arish represents the first product of the Sinai coast. There are several types of sardines, including round sardinella *S. aurita* and maderian sardinella *S. maderensis*, which the maderian sardinella the second part of the production of sardines in North Sinai and there are a little previous studies in this region on *Sardinella maderensis*. Production of sardines on the Mediterranean coast in 1998 was more than 5,000 ton, reaching 11700 ton in 1999, then it decreased to 4,000 in year 2000 (El-Aiatt, 2004), and it decreased further until it reached 210 ton in year 2010 (Gaber, 2012), and this may

be due to change in environmental conditions and this is one of the reasons for this study. *S. maderensis* is from the family Clupeidae, and genus sardinella. It is classified as a small ray-finned fish and is usually found in the Eastern Atlantic and southern Mediterranean (Ba et al., 2016). It is a silvery fish like other sardinella species (Fig. 1). The growth parameters of *S. maderensis* were estimated in studies carried out in several regions of the African Atlantic coast, based on scale readings. Ghéno and Le Guen (1968) reported that this species can reach six years of age with accelerated growth up to three years of age (Camarena, 1986).

* Corresponding author: E-mail address: rokaaamin73@gmail.com

<https://doi.org/10.21608/sinjas.2023.205104.1198>

2023 SINAI Journal of Applied Sciences. Published by Fac. Environ. Agric. Sci., Arish Univ. All rights reserved.



Fig. 1. *Sardinella maderensis*

Several studies have been carried out in various regions of the West African coast on the determination of the growth of this species, using readings of scales, and frequency distributions of lengths (Shcherbich, 1981; Showers, 1996). These studies indicate that in general *S. aurita* is a fast-growing species, especially in the first two years of life. According to Ghéno and Le Guen (1968), longevity in this species is generally 6 to 7 years of life. In Angola, only one study on growth is known, based on the reading of scales carried out by Shcherbich (1981). Konoyima *et al* (2020) reported that the Mean lengths (18.5 cm) and mean weights (62.4 g) for *S. maderensis* also growth exponent gave negative allometric, the correlation between length and weight ($r=0.7282$) and the (K) was low ($K<1$). This research aims to study some of the biological and dynamic characteristics of this important specie to evaluate the status and provide some information to inventory management.

MATERIALS AND METHODS

4047 samples were collected of *S. maderensis* monthly from January to December 2021. Lengths were measured to the nearest 0.1 cm and weights to the nearest 0.1 g. Through simple observation, is determined the development of gonads in fish. Scales were taken from 2333 fish to calculate the age and also the dissection

of these fish to know the sex and weight of the gonad to know the development of the gonads (The weights of gonads are rounded to 0.1 grams) and calculate the gonadosomatic index and to know the spawning season for these fish.

The Relationship between Length-Scale Radius

The relationship between average length and Scale radius was calculated. Lee (1920) reported that the back calculation length of the length is done through the following equation: $(L_n = (S_n/S) (L-a) + a)$. Where L_n = length at the end of year n ; S_n = radius of scale to ring n ; S = overall radius of the scale; L = total length at capture and a = constant of the relationship between length and Scale radius.

Le Cren (1951) reported that the relationship between length and weight is measured by the following equation: $W = aL^b$ where W and L total length and weight a and b are constants the length – weight relationship. Several mathematical models are used to describe the theoretical growth of fishes; the most widely used model is that of Von Bertalanffy equation (1949) for calculating theoretical growth in length and weight $L_t = L_\infty (1 - e^{-k(t-t_0)})$ and $wt = w_\infty [(1 - e^{-k(t-t_0)})]^b$ where L_t = length at age t and wt = weight at age t , L_∞ = the asymptotic length, $W = aL_\infty^b$, k = growth coefficient and t_0 = the age at which the length is nil.

Pauly and Munro (1984) reported that the equation Growth performance index (@) is calculated by the following equation ($\Phi = \log k + 2 \log L_{\infty}$). Monthly condition factor was calculated by two methods, namely:

1. Condition factor "Kc" $Kc = (W \times 100) / L^3$ (**Hile, 1936**) Where: Kc = composite coefficient of condition, W = weight in g, L = length in cm
2. Relative condition "Kn" $Kn = W / W^*$ Where: Kn = relative coefficient of condition, W = observed weight in g, W* = calculated weight in g. The gonads removed and weighed to the nearest 0.1 g; calculated (GSI) Gonado somatic index every month by equation of **Bariche et al. (2003)**. Gonadosomatic Index parameter was calculated using the formula: $GSI = [(\text{gonad weight}) / (\text{body weight})] \times 100$. The total mortality coefficients were obtained by using the following method: The Powell-Wetherall plot based on data (**Powell, 1979**) discussed in **Wetherall et al. (1987)** $Z = 1 - k$. Natural mortality coefficient was estimated by using the equations of (**Ursin, 1967**) formula $M = W'^{-1/3}$ where $W' =$ Average weight of the total samples. $F = Z - M$ and $(E) E = F / (F + M)$ Where E: is exploiting rate, F: is fishing mortality, M: is natural mortality.

RESULTS

The lengths of *S. maderensis* ranged from 8.5 to 26.3 (mean 14.63 cm) and the weights ranged from 5.0 to 171 g (mean 29.1g). The relationship between length and weight in Fig. 2 and the length – weight relationship was: $W = 0.007 \times L^{3.0524}$.

Determine the Age

From reading the scales of *S. maderensis*, there were five age stages from 0 to age 4, and through this equation ($Ln = (L - 7.6046) / (Sn / S + 7.6046)$) the length was calculated at the end of each year. The percentage of attendance for each age group was as follows 9.82, 42.31, 27.73, 14.1 and 6.04% as a percent for 0, 1st, 2nd, 3rd and 4th age

groups, respectively and the lengths at the end year were 11.45, 16.1, 20.1, and 22.52 cm for age 1, 2, 3, and 4 years, respectively and the annual increment of length is given in Fig. 3 and the weight at the end year and the annual increment in Fig. 4.

Growth

The growth parameters were $L_{\infty} = 30.39$, $K = 0.2937$ and $t_0 = 0.3094$, respectively. To obtain the theoretical growth of height and weight through the two von Bertalanffy equations for growth in height and weight

Growth of length: $L_t = 30.39 (1 - e^{-0.2937(t + 0.3094)})$

Growth of weight: $W_t = 235.0 [(1 - e^{-0.2937(t + 0.3094)})]^{3.0524}$

Growth performance index (ϕ') for *S. maderensis* in Eastern Mediterranean (North Sinai Coast) were 2.4334 for length and 1.0486 for weight.

The length at first capture (L_c) which 50% of fishes retained by the gear of *S. maderensis* in Eastern Mediterranean (North Sinai Coast) were estimated at 13.5 cm (Fig. 5). Total mortality, natural mortality, fishing mortality and Exploitation rate for *S. maderensis* were 0.7124; 0.3229; 0.3895 and 0.547, respectively.

Monthly average gonads somatic index (GSI) for both males and females of *S. maderensis* are given in Fig. 6. In females and males, the gonads somatic index increases progressively from April and the top in July these results indicate that the spawning season for these fish is between March to September. The first sexual maturity for of *S. maderensis* season, 2021 was determined by examination of gonads to determine the sex and the stage of maturity. The length at first maturity was estimated at 15.5, 16.7 cm and 16.0 cm for males, females and combined sexes respectively (Fig. 7). The size at first maturity of fish may be important to assess the optimum size of first capture of a fish and the age at first maturity $T_{m50} = 2.57$ years for combined sexes of *S. maderensis* during 2021.




Fig. 2. Length-weight relationship (♀♂) of *S. maderensis* during 2021




Fig. 3. Growth and annual increment in length (♀♂) of *S. maderensis* 2021




Fig. 4. Back-calculation weight at the end of different years of life (♂♀) of *S. maderensis* during 2021




Fig. 5. Length at first capture of combined sexes (♀♂) of *S. maderensis* season, 2021




Fig. 6. GSI for males and females of *S. maderensis* during, 2021




Fig. 7. Length at first maturity of combined sexes (♀♂) of *S. maderensis*, 2021

DISCUSSION

Length weight relationships are important for comparative growth studies (**Moutopoulos and Stergiou, 2002**). Length–weight relationships are very important in order to (i) calculate the weight of a given individual fish of known length or the total weight of a fish from the length and frequency distribution; (ii) estimation of weight growth rate and several other aspects of fish populations; (iii) converting the growth equations in length to the growth equations in weight for use in inventory valuation models; (iv) Estimating the status of fish in a particular geographical area. (5) making interregional comparisons of the life history and morphology of specific species (**Kohler *et al.*, 1995; Stergiou and Moutopoulos, 2001**).

In this study the percentage of small fish is high. The length –weight relationship values (a and b are constants vary with sex, age, seasons, growth, health, habitat, feeding, and differences in how long a specimen is caught, stage of maturity, and sampling techniques from fishing gear (**Tesch, 1968; Begenal and Tesch, 1978**)).

Table 1 shows the differences in the value of b in different regions. **Richter *et al.* (2000)** reported that the growth and the relationship between length and weight very important in fishery management.

In this study, the back-calculated length at the end of life year and the highest annual increment occurred during the first year of life, The increase of length decreased gradually with age. In this study the increment of length in the first year was 50.8% then decreased gradually with age. These results agreement with **Sossoukpe *et al* (2016)** .The increment of *S. maderensis* in the nearshore waters of Benin (West Africa) 42.7%, also with **Mahfoudh *et al* (2018)** 65.7% in Mauritania. In this study, there are four age groups, the first age

group dominated hunting (42.31%), followed by the second, third, zero and finally the fourth age group (27.73%), (14.1%), (9.82%). (6.04%) respectively, average length at different ages of *S. maderensis* in different localities (Table 2).

In present study the growth parameters $L_{\infty} = 30.39\text{cm}$, $W_{\infty} = 234.9\text{g}$, $K = 0.29$, $T_0 = -0.6094$ and the growth performance index (ϕ') for *S. maderensis* was about 2.433. for combined sexes These results are higher, lower or equal to some of the results presented in Table 3 for some scientists in different regions of the world.

Estimating the mortality factor of fish stocks is an essential step for calculating the potential yield, the optimal yield per recruit, and the optimal fishing effort. In this study, although estimates of the total mortality coefficient (Z) could be affected by the absence of large individuals due to the qualitative selectivity of the size of the small study fish. In the present study the annual rates of total mortality coefficient "Z", the natural mortality "M" and the fishing mortality "F" for combined sexes of *S. maderensis* were estimated to be 0.7124, 0.3229 and 0.3895 respectively. **Olopade, *et al.* (2019)** found that the values of Z, M and F in the Sombreiro River, Nigeria for *S. maderensis* was "Z" =2.74, "M"= 1.32and "F"= 1.42, while **Wehye (2017)** in the Sombreiro River, Nigeria reported that "Z" =1.24, "M"= 0.81 and "F"= 0.43 and **Sossoukpe, *et al* (2016)** in the nearshore waters of Benin found that "Z" =3.93, "M"= 1.3 and "F"= 2.63 these results are high compared to our results.

Gulland (1971) reported that the exploitation rate is the fraction of an age class that is caught during the life span of a population exposed to fishing pressure, *i.e.*, the number caught versus the total number of individuals dying due to fishing and other reasons. which allows one to (roughly) assess if a stock is overfished or not, on the assumption that the optimal

Table 1. Constants of relationship between Length and weight in different localities

| Area of Study | Author | a | b |
|---------------------------------|-------------------------------------|---------|--------|
| Mediterranean Sea, North Sinai | Present study | 0.07 | 3.0524 |
| Nigeria | Olopade <i>et al.</i> (2019) | 0.0225 | 2.58 |
| Mauritanie (Artisanal fishery) | Mahfoudh <i>et al.</i> (2018) | 0.0071 | 3.09 |
| Nigeria | Abdul <i>et al.</i> (2016) | 0.011 | 2.9 |
| Benin | Sossoukpe <i>et al.</i> (2016) | 0.023 | 2.86 |
| Senegal | Ba <i>et al.</i> (2016) | 0.01560 | 3.00 |
| Senegal | Samba (2011) | 0.0006 | 3.01 |
| Mauritania (industrial fishery) | Pascual-Alayón <i>et al.</i> (2008) | 0.0006 | 3.12 |
| Gambia | Ecoutin <i>et al.</i> (2005) | 0.00007 | 3.15 |

Table 2. Average length at different ages of *S. maderensis* in different localities

| Area of Study | Author | Age | | | |
|--------------------------------|--------------------------------|-------|------|------|-------|
| | | 1 | 2 | 3 | 4 |
| Mediterranean Sea, North Sinai | Present study | 11.45 | 16.1 | 20.1 | 22.52 |
| West Africa | Sossoukpe <i>et al.</i> (2016) | 13.01 | 22.9 | 28.1 | 30.68 |
| Mauritania | Mahfoudh <i>et al.</i> (2018) | 23.00 | 28.5 | 32.3 | 35.0 |

Table 3. Growth parameters of *S. maderensis* in different places

| Area of Study | Author | Parameter | | | |
|--------------------------------|--------------------------------|--------------|------|-------|--------|
| | | L_{∞} | K | T0 | ϕ |
| Mediterranean Sea, North Sinai | Present study | 30.39 | 0.29 | -0.61 | 2.433 |
| Nigeria | Olopade <i>et al.</i> (2019) | 23.21 | 0.54 | -0.03 | 2.464 |
| Ghana | Arizi (2019) | 32.21 | 0.58 | -0.49 | 2.776 |
| Ghana | Samuel <i>et al.</i> (2019) | 23.36 | 0.61 | -0.28 | 2.532 |
| Mauritania | Mahfoudh <i>et al.</i> (2018) | 38.44 | 0.53 | -0.6 | 2.894 |
| West Africa | Sossoukpe <i>et al.</i> (2016) | 33.6 | 0.65 | -0.24 | 2.866 |
| Nigerian waters | Marcus (1989) | 37.5 | 0.34 | -0.25 | 2.68 |

value of E is equal to 0.5, the use of $E \approx 0.5$ as optimal value for the exploitation ratio itself resting on the assumption that sustainable yield is optimized when $F \approx M$.

Pauly (1987) suggested a less optimal E value of 0.4, thus the values of fishing mortality rate and exploitation rate were relatively high indicating a high level of exploitation. In present study the exploitation rate was 0.5468 for combined sexes. Sardines are exposed to many predators and to many fish of recreational and commercial importance such as *Scomber japonicus* (**Rizkalla and Faltas, 1997**). Feeding habits of chub mackerel (*Scomber japonicus*) in Egyptian Mediterranean waters.

In this study, GSI values differed between males and females, with common values occurring in the period from January to March. The GSI value in this study ranged from 0.11 to 2.6 for males and from 0.19 to 3.4 for females, and the peak spawning season was from April to September. These results differ with studies on *S. maderensis* conducted by **Youmbi *et al.* (1991)** where the peak spawning periods were from April to July. Also **Anonymous (1988)** showed that the spawning periods vary with season in Sardinella species, and this corresponds to the breeding season (August to December) in Ivory Coast and Ghana. **Boely *et al.* (1979)** showed that there is one major spawning period (April to October) for *S. maderensis* with two breeding peaks for *S. aurita*, which are in the dry and cool seasons (February to June). **Longhurst and Pauly (1987)** also noted that reproductive peaking is a feature of commercial fish and invertebrates in tropical waters.

Length at first maturity (L_m) is very important for fisheries management, **Osman *et al.* (2020)** reported that according to length at first maturity we can set a minimum legal size (MLS) for not overfishing. **Woodhead (1978)** found that the length at first maturity is approximately two-thirds of the maximum total length. This assumption is consistent

with our results where the maximum length was 26.3 cm and lengths at first maturity were 16.7 cm for *S. maderensis* females. These results show that the *S. maderensis* stock in the Mediterranean is overexploited, with the length at first capture in our results being 13.5 cm, which is significantly less than the length at first maturity.

Double recruitment annually is for tropical fish species and short-lived species (**Pauly 1982**). Recruitment is year-round for tropical fish and shrimp species (**Weber, 1976**). Results obtained by **Diouf *et al.* (2010)** show continuous breeding of *S. maderensis* throughout the year.

Conclusion

The fishing effort during the spawning season should be reduce especially from June to August and should be not increase the fishing effort but should be increasing the size of the nets so that the length at first capture increases from 13.0 cm to 16.0 cm in order to equal the length at first maturity to preserve the stock of these fish.

REFERENCES

- Abdul, W.O.; Omoniyi, I.T.; Adekoya, E.O.; Adeosun, F.I; Odulate, O.O.; Idowu, A.A.; Olajide, A.E. and Olowe, O.S. (2016)**. Length-weight relationship and condition factor of some commercial fish species in Ogun State coastal Estuary, Nigeria. *Ife J. Agric.*, 28: 1-10.
- Anonymous (1988)**. The Fisheries Management and Development Act, 1988 (Act No. 4 of 1988). The Fisheries Regulations. *Sierra Leone Gazette*, 119 : 35.
- Arizi, E.K. (2019)**. Towards sustainable management of the sardinella fishery in Ghana. Doctor Philosophy in Biological and Environ. Sci., Univ. Rhode Island.
- Ba, K.; Thiaw, M.; Lazar, N.; Sarr, A.; Brochier, T. and Ndiaye, I. (2016)**. Resilience of Key Biological Parameters

- of the Senegalese Flat Sardinellato Overfishing and Climate Change. PLoS ONE11(6):e0156143.doi:10.1371/journal.pone.0156143.
- Begenal, T.B. and Tesch, F.W. (1978).** Age and Growth. In: Methods for Assessment of Fish Production in Fresh Waters, third edit. (Bagenal, T. Ed.), IBP Handbook No. 3. Oxford: Blackwell Scientific Publications, 101-36.
- Bariche, M.; V.M. Harmelin and J.P. Quignard (2003).** Reproductive cycles and spawning periods of two Lessepsian siganid fishes on the Lebanese coast. J. Fish Biol., 62:129–142.
- Boely, T.J.; Chabanne, F.P. and Stequert, B. (1979).** Cycle sexuel et migration de *Sardinella aurita* sur le plateau continental Ouest Africain, des Iles Bissagos a la Mauritanie. Rapp. P.-V. Reun. Cons. Int. Explor. Mer., 180: 350-355.
- Camarena, L.T. (1986).** Lacroissance de *Sardinella maderensis* (Lowe, 1841) au Senegal. Oceanography tropical, 21:143-151. ISSN0245-9418.
- Diouf, K.; Samb, B. and Sylla, M. (2010).** Contribution à la connaissance de la biologie des sardinelles (*Sardinella aurita* et *Sardinella maderensis*) du littoral sénégalais. In: Garcia, D.S., Tandstad, M., Caramelo, A.M. (ed.) Science et aménagement des petits pélagiques. Symposium sur la science et le défi de l'aménagement des pêcheries de petits pélagiques sur les stocks partagés en Afrique nord-occidentale. Casablanca, Maroc, Rome, FAO, 39-56.
- Écoutin, J.M.; Albaret, J.J. and Trape, S. (2005).** Length-weight relationships for fish populations of a relatively undistributed tropical estuary: the Gambia. Fisheries Res., 72:347–351.doi: 10.1016/j.fishres.2004.10.007.
- El-Aiatt, A.A.O. (2004).** study on fish production of the Mediterranean coast of Sinai, Theses, Ph.D., Fac. Environ. Agric. Sci., Suez Canal Univ.
- Gaber M. (2012).** Studies on fish production of Mediterranean coast in North Sinai Governorate, Ph.D., Fac. Agric., Cairo, El-Azhar Univ.
- Gheno, Y. and Le Guen, J.C. (1968).** Determination de l'age et croissance de Sardine and Eba (Val.) dam la region de Pointe-Noire. Cah. ORSTOMSer. Oceanogr., 6: 69-82.
- Gulland, J.A. (1971).** The fish resources of the Oceans. Fishing News Books Ltd., England., 255.
- Hile, R. (1936).** Age and growth of the Cisco, Leu cichthys artedi (Le Sueur) in the lakes of the Northeastern high-lands, Wisconsin. Bull. Bur. Fish. US, 48 (19): 211-317.
- Kohler, N.E.; Casey, J.G. and Turner, P.A. (1995).** Length weight relationships for 13 species of sharks from the western North Atlantic. Fishery Bulletin., 93: 412 - 418.
- Konoyima, K.J.; Mansaray, M. and Coker, I.C.R. (2020).** Some aspects of catch, gonad maturation and growth pattern of *sardinella maderensis* and *sardinella aurita* in Sierra Leone. Int. J. Basic, Appl. and Innovative Res., IJBAIR, 9 (3):82-98 www.arpjournals.com; www.antrescentpub.com.
- Le Cren, E.D. (1951).** The length-weight relationship and seasonal cycle in gonad weight and condition in the Perch (*Perca fluviatilis*). J. Anim. Ecol., 20: 201-219.
- Lee, R. (1920).** A review of the methods of age and growth determination in fishes by means of scales. Fishery investigations, Series 2, Marine fisheries, Greet Britain Minist. Agric., Fisheries and Food, 4: 2.

- Longhurst, A.R. and Pauly, D. (1987).** Ecology of Tropical Oceans. Academic Press, San Diego, 407 pp.
- Mahfoudh, A.; Zongo, B. and Kabre, T.J. A. (2018).** Growth parameters estimate of Madeiran sardenella (*Sardinella maderensis*) exploited from artisanal fisheries and landed at the Nouakchott fishermen's beach in Mauritania. International J. Agric. Policy and Res., 6 (11): 207-219.
- Marcus, O. (1989).** Breeding, age and growth in *Sardinella maderensis* (Lowe 1839) Pisces: Clupeidae from coastal waters around Lagos, Nigeria. Nigerian J. Sci., 23 (1-2): 1-5.
- Moutopoulos, D.K. and Stergiou, K.I. (2002).** Length-weight and length-length relationships of fish species from Aegean Sea (Greece). Appl. Ichthyol., 18: 200-203.
- Olopade, O.A.; Dienye, H.E. and Bamidele, N.A. (2019).** Some Population Parameters of the *Sardinella maderensis* (Lowe, 1838) in the Sombreiro River of Niger Delta, Nigeria. Acta Aquatica Turcica, 15 (3): 354-364. <https://doi.org/10.22392/actaquatr.532284> .
- Osman, H.M.; Saber, M.A.; El-Ganainy, A.A. and Shaaban, A.M. (2020).** Fisheries biology of the haffara bream *Rhabdosaragus haffara* (Family: Sparidae) in Suez Bay, Egypt. Egypt. J. Aquatic Biol. and Fisheries, 24(4): 361-372.
- Pascual-Alayón, P.; Santamaría, M.T.G.; Balguerías, E.; Hernández, E.; Bravo de Laguna, L.; Sancho, A. and Duque, V. (2008).** Activity of European pelagic trawlers fishing in Mauritania and landing in the port of Las Palmas de Gran Canaria (Canary Islands, Spain) from 2004-2007 - Relationship between catches and SST/SSTA). Science and the challenge of managing small pelagic fisheries on shared stocks in North West Africa, Casablanca, Morocco, 11–14.
- Pauly, D. (1987).** A review of the ELEFAN system for analysis of length frequency data in fish and aquatic invertebrates. ICLARM Conf. Proa., 13: 7-34.
- Pauly, D. (1982).** Studying single-species dynamics in a tropical multi-species context. In: Pauly D, Murphy GI (Eds). Theory and management of tropical fisheries. ICLARM Fishbyte, 3: 13-14.
- Pauly, D. and Munro, J.L. (1984).** Once more on the comparison of growth in fish and invertebrates. ICLARM Fishbyte, 2 (1): 21.
- Powell, D.G. (1979).** Estimation of mortality and growth parameters from the length frequency of a catch. Rapp.P.-v. Réunion. CIEM, 175:167-169.
- Richter, H.; Lukstadt, C.; Focken, U. and Becker, K. (2000).** An improved procedure to assess fish condition on the basis of length-weight relationships. Archive of Fishery and Marine Res., 48 (3): 255-264.
- Rizkalla, S.I. and Faltas, S.N. (1997).** Feeding habits of chub mackerel (*Scomber japonicus*) in Egyptian Mediterranean waters. J. King Abdulaziz Marine Sci., 8: 127-136.
- Samba, O. (2011).** Nouvelle évaluation des caractéristiques biologiques de *Sardinella aurita* (Valenciennes, 1847) et *Sardinella maderensis* (Lowe, 1841). Mémoire de DEA,IUPA/UCAD, Dakar (Sénégal), 45.
- Samuel, K.K.A.; Patrick, K.O.; Francis, K.E.N. and Godfred, A.A. (2019).** Estimates of Population parameters for *Sardinella maderensis* (Lowe, 1838) in the coastal waters of Ghana. Greener J. Agric. Sci., 9(1): 23-31.
- Shcherbich, L.V. (1981).** Natural Mortality and optimum age of exploitation of *Sardinella aurita* (Walbaum 1847) in the southeast Atlantic. ICSEAF 1981 (Parte 11) Colln scient. Pa. Int. Comm., LSE, Atl, Fish. 8:253054.

- Showers, P.A.T. (1996).** Comparative growth performance for species of the Family Clupeidae of Sierra Leone. Croissance compare des espèces de la famille des Clupéidés en Sierra Leone. Naga ICLARM Q. 19 (4): 42-44.
- Sossoukpe, E.; Djidohokpin, G. and Fiogbe, E.D. (2016).** Demographic parameters and exploitation rate of *S. maderensis* (Pisces: Lowe 1838) in the nearshore waters of Benin (West Africa) and their implication for management and conservation. Int. J. Fisheries and Aquatic Studies, 4(1): 165-171.
- Stergiou, K.I. and Moutopoulos, D.K. (2001).** A review of length weight relationships of fishes from Greek Marine Waters. Naga, the ICLARM Quarterly, 24: 23-39.
- Tesch, F.W. (1968).** Age and Growth in Methods for Assessment of Fishes Production. In: W.E. Ricer (Ed.), in methods for assessment of fish production in Freshwater, IBP Handbook, Blackwell Sci. Publication, London, 93-123.
- Ursin, E. (1967).** A mathematical model of some aspects of fish growth, respiration and mortality. J. Fish. Res. Bd. Can, 24: 2355-2453.
- Wehye, A. (2017).** Growth, mortality and exploitation of *Sardinella maderensis* (Lowe, 1838) in the Fisheries and Aquacul. J., 8: 2150-3508.
- Von Bertalanffy, L. (1949).** Problems of organic growth. Nature, 163: 156-158.
- Weber, W. (1976).** The influence of hydrographic factors on the spawning time of tropical fish. In: Tiews K (ed.), Fisheries resources and their management in Southeast Asia: 269-281.
- Wetherall, J.A.; J.J. Polovina and S. Ralston (1987).** Estimating growth and mortality in steady-state fish stocks from length-frequency data. ICLARM Conf. Proc., (13): 53-74.
- Woodhead, A.D. (1978).** Fish in studies of aging. Exp. Geront., 13 (3-4): 125-140.
- Youmbi, J.T., Djama, T. and Gabche, C. (1991).** Reproductive patterns of *Sardinella maderensis* (Lowe, 1841) off Cameroon. J. Appl. Ichthyology, 7: 60-63.

المخلص العربي

دراسة بعض الخصائص البيولوجية والديناميكية لاسماك السردين المفطر (*Sardinella maderensis*) بشرق البحر المتوسط (ساحل شمال سيناء مصر)إسراء أمين السيد^{1*}، جابر دسوقي إبراهيم حساتين¹، رشا محمد فيصل¹، عطية على عمر العياط²

١- قسم الثروة السمكية والأحياء المائية، كلية العلوم الزراعية البيئية، جامعة العريش، مصر.

٢- المعهد القومي لعلوم البحار والمصايد، مصر.

تم جمع عينات شهرية من *S. maderensis* من يناير إلى ديسمبر ٢٠٢١ في ساحل البحر الأبيض المتوسط في سيناء لتحديد النمو والوفيات. تراوحت الأطوال من ٨,٥ - ٢٦,٣ سم وتراوحت الأوزان من ٥ جرام إلى ١٧١ جرام، وبلغت القيمة b للعلاقة بين الطول والوزن ٣,٠٥٢٤ وبالتالي فإن نمو *S. maderensis* يصنف على أنه يظهر نمو متساوي القياس. (VBGP) من L_{∞} و (k) و (t_0) كانت ٣٠,٣٩ سم و ٠,٢٩٣٧ و -٠,٦٠٤٩/سنة على التوالي. بلغ مؤشر أداء النمو (\emptyset') ٢.٣٣٤. تراوحت GSI من ٠,١١ إلى ٢,٦ للذكور وتراوحت بين ٠,١٩ و ٣,٤ للإناث، كما كانت فترات ذروة التبويض في *S. maderensis* من أبريل إلى سبتمبر. كانت الأطوال عند بداية النضج الجنسي ١٥,٥ سم (ذكور)، ١٦,٧ سم (إناث) و ١٦,٠ سم للجنس المختلط. تم تقدير قيم الوفيات (Z) و (M) و (F) ومعدل الاستغلال (E) للجنسين معا لاسماك السردين المفطر (*S. maderensis*) كانت ٠.٧١٢٤ و ٠,٣٢٢٩ و ٠,٣٨٩٥ لكل سنة و ٠,٥٤٧ على التوالي. كان الطول عند بداية الصيد ١٣,٥ سم. يجب عدم زيادة جهد الصيد ولكن يجب زيادة حجم الشباك بحيث يزيد الطول عند بداية الصيد من ١٣,٥ سم إلى ١٦,٠ سم لكي يتساوى مع الطول عند بداية النضج للحفاظ على مخزون هذه الأسماك.

الكلمات الاسترشادية: السردين المفطر، دليل المناسل، معايير النمو، النفق.

REVIEWERS:

Dr. Mohamed M. El-Sayed Al-Murr

Dept., Marine Sciences, Fac. Fish and Fisheries Biology, Suez Canal Univ., Egypt.

Dr. Ahmed Kamel

National Inst. Oceanography and Fisheries, Egypt.

| melmor88@yahoo.com

| akelhammad@yahoo.com