



Phytochemistry and some biological activities of the Genus Hypericum

Ruba Joujeh^{1*}, and Dima Joujeh²

¹ Ph.D in Plant Sciences, lecturer at Agriculture faculty, University of Aleppo, Syria

² Ph.D in Biotechnology engineering, Faculty of Technical engineering, University of Aleppo, Syria

ARTICLE INFO

Article History:

Received: 3 July 2023

Accepted: 17 Sept 2023

Published online: 25 Dec 2023

Key words:

Hypericum,
Hypericaceae,
chemical composition,
antibacterial,
antioxidant

ABSTRACT

Medicinal plants are the best source of medicine. Therefore, studies on these plants have increased because of the search for new active substances that can be used in the herbal pharmaceutical industries. Hypericum is a large genus of flowering plants, comprising many species, and presents an almost worldwide distribution. The importance of its species depends mainly on the presence of specialized secondary metabolites that display pharmaceutical and cosmetic properties. Hypericum species are known worldwide for their ethno-medicinal uses including treating infections and diseases. They are used as diuretics, cholagogues, antispasmodics, antiepileptics, and also for treating rheumatism, neuralgia, parasites, dyspepsia, diarrhea, etc. There are only few reports about the chemical composition of the genus Hypericum. The aim of the present study is to provide an overview of the importance of the genus Hypericum. A summary of the chemical composition, as well as the antibacterial and antioxidant activities of different Hypericum species.

1. Introduction

In developing countries, 70-95% of people depend on plants as their main method of treatment these days [1]. According to the WHO, medicinal plants are the best source of medicines [2]. Some researchers believed that two-thirds of the world's plants are medically beneficial [3]. Therefore, research on medicinal plants has increased rapidly in order to find new active substances that can be used in herbal medicine industries [2].

On the other hand, the problem of food preservation is becoming more complex, requiring a longer shelf

life and greater protection from microbial spoilage and oxidative damage. Some statistics indicated that about 30% of people in industrialized countries suffer annually from foodborne [4].

Medicinal plants are considered as powerful and easily available sources of antioxidants, due to the chemical compounds they contain that act individually or synergistically to treat diseases [5]. It has been reported that many phenolic compounds such as procyanidins, coumarins, flavonoids and tannins can scavenge radicals -in a dose-dependent manner- and are thus

considered as therapeutic medicines for free radical diseases [6].

Medicinal plants also contain compounds that may inhibit the growth of viruses, protozoa, bacteria and fungi by various mechanisms and this can be of a significant clinical value in treating of resistant microbial strains [7]. These bioactive compounds are used as a starting point for the synthesis of antibiotics in order to treat infectious diseases [8]. In general, the extent of the antioxidant and antimicrobial effects of the extracts could be attributed to their phenolic compounds [9].

Hypericum is a large genus of flowering plants, including many species [10], and presents an almost worldwide distribution. The importance of its species depends mainly on the presence of specialized secondary metabolites that display pharmaceutical and cosmetic properties [11]. Over the years, there has been a great interest in studying the different biological activities of *Hypericum* species [10]. Therefore, the purpose of this study is to provide an overview of the importance of *Hypericum*. A summary of the typical essential oil constituents, as well as the antibacterial and antioxidant activities of different species belonging to the genus *Hypericum*.

2. The genus *Hypericum*

Hypericaceae family contains shrubs or herbaceous plants. They have glandular hairs that contains essential oils (EOs) [12]. This family includes nine genera, one of which is *Hypericum* [13]. *Hypericum* contains nearly 500 species [14], perennial, herbaceous, or scrubby plants [15], widely distributed in mountainous, tropical and temperate regions [16]. The genus received a great as it is a source of a variety of compounds with various biological effects [15].

Hypericum species are globally recognised for their significant ethno-medicinal properties, which are

utilised in the treatment of various infections and diseases [17]. These species exhibit a wide range of therapeutic effects, including diuretic, cholagogue, antispasmodic, antiepileptic, and antimigraine properties. Additionally, they are employed in the management of conditions such as rheumatism, neuralgia, parasitic infections, dyspepsia, sciatica, and diarrhea [13]". *Hypericum* plants have been included in Pharmacopoeias of many countries such as Germany, Czechoslovakia, Russia, Romania, Poland and France [18]. Various products containing *Hypericum* plants or their extracts have been developed as additives and many brands such as beverages and yogurts include these plants [18].

The biological effects of the plants depend on their contents of secondary metabolites [19]. The chemical content of *Hypericum* plants is diverse with xanthenes, glycosides, pyrones, flavonoids, tannins, anthraquinones, lactones, lipids, phloroglucinols, coumarins and EOs [20].

The major phytochemical compounds of *Hypericum* species are naphthodianthrones (hypericin and pseudohypericin), phloroglucinol derivatives (adhyperforin and hyperforin), flavonoids (rutin, biapigenin, hyperoside, quercetin, quercitrin), chlorogenic acid and caffeic acid which possess many biological properties [14].

Hypericum perforatum is one of the most widely used medicinal plants -among all species of the genus *Hypericum*- by the publics of both more industrialized and less developed countries [21]. It is used as a mild antidepressant [22]. This effect of *H. perforatum* is due to its main component hyperforin. The combined effect of the other components within the plant extract also contributes to its overall impact synergistically [19].

3. Chemical Composition of the Essential Oils of Hypericum Species

The volatile substances in plants are generally analyzed through extraction steps, concentration, chromatography and detection steps [23].

Hypericum plants are generally known to be poor in essential oil (generally oil yield <1%, w/w) [18].

Table 1. Chemical composition of *Hypericum* essential oils

Species	Plant part	Main compounds	Ref.
<i>H. elegans</i>	AE	g-gurjunene, aromadendrene, and undecane	[24]
<i>H. tetrapterum</i>	AE	Copaene, α -longipinene, cadinene	
<i>H. hirsutum</i>	AE	caryophyllene oxide, phytol, α -caryophyllene and undecane	
<i>H. perforatum</i>	AE	β -caryophyllene, caryophyllene oxide, α -pinene, β -cadinene, and β -pinene	
<i>H. scabrum</i>	AE	α -pinene, β -caryophyllene, myrcene, cadalene and β -pinene	[25]
<i>H. scabroides</i>	AE	hexadecanoic acid, spathulenol, nonacosane, dodecanoic acid, baeckeol and γ -muurolene	[26]
<i>H. kotschyannum</i>	AE	α -pinene, nonacosane, hexadecanoic acid, β -pinene, spathulenol and limonene	
<i>H. salsugineum</i>	AE	nonacosane, hexadecanoic acid and baeckeol	
<i>H. thymopsis</i>	AE	α -pinene, baeckeol, spathulenol, limonene and camphene	
<i>H. uniglandulosum</i>	AE	2,6-Dimethyl-3,5-heptadien-2-one, nonacosane, hexadecanoic acid and α -pinene	[27]
<i>H. uniglandulosum</i>	AE	α -pinene, undecane, benzoic acid, cyclohexasiloxane	
<i>H. lydiium</i>	AE	α -pinene, β -pinene, β -myrcene	
<i>H. thymopsis</i>	AE	α -pinene, spathulenol, limonene	[28]
<i>H. perforatum</i>	AE	(germacrene D); ((E)-caryophyllene); (2-methyloctane); (α -pinene) and (bicyclogermacrene)	[29]
<i>H. perforatum</i>	FL	(E- β -farnesene); (n-hexadecanal); (E-nerolidol)	[30]
<i>H. dogonbadanicum</i>	FL	(phenyl ethyl octanoate); (terpin-4-ol); (α -phellandrene)	
<i>H. dogonbadanicum</i>	LE	β -pinene, α -pinene and p-cymene	
<i>H. helianthemoides</i>	FL	α -pinene, Z- β -ocimene and β -pinene	
<i>H. hyssopifolium</i>	FL	α -pinene, β -pinene and n-tetradecan	
<i>H. hyssopifolium</i>	LE	E-nerolidol, n-tetradecane and α -himachalene	
<i>H. lysimachioides</i>	FL	α -pinene, Z- β -ocimene and n-tetradecane	
<i>H. lysimachioides</i>	FL	n-tetradecane, α -himachalene and α -pinene	
<i>H. triquetrifolium</i>	LE	α -himachalene, n-tetradecane and n-pentadecane	
<i>H. triquetrifolium</i>	AE	Germacrene-D, β -caryophyllene, δ -cadinene, trans- β -farnesene, α -humulene, β -selinene, γ -cadinene and trans-phytol	[31]
<i>H. richeri</i>	AE	germacrene D, bicyclogermacrene, α -pinene, β -pinene, decanoic acid, β -caryophyllene, δ -cadinene, spathulenol and tetracosane	[32]
<i>H. patulum</i>	AE	β -pinene, α -pinene, limonene and α -humulene	[33]
<i>H. laricifolium</i>	AE	and nonane α -pinene, verticilol, 3-methyl-nonane, 2-methyl-octane	[13]
<i>H. rumeliacum</i>	AE	undecane, dodecanal, and germacrene D	[34]
<i>H. rumeliacum</i>	AE	(E)- β -ocimene, β -pinene, (Z)- β -ocimene, dodecanal, germacrene D, myrcene	[35]
<i>H. rumeliacum</i>	AE	α -pinene, β -pinene, dehydro-aromadendrene, α -copaene	[36]
<i>H. lydiium</i>	AE	α -pinene, β -pinene and β -myrcene	[27]

<i>H. ericoides</i>	AE	n-nonane, n-undecane, α -cubebene, α -pinene	[18]
<i>H. androsaemum</i>	AE	longifolene, β -gurjunene, and γ -gurjunene	[22]
<i>H. apricum</i>	AE	cis-piperitol acetate, p-cymenene, α -pinene	
<i>H. armenum</i>	AE	γ -cadinene, longifolene, E-nerolidol	
<i>H. asperulum</i>	AE	α -muurolol, cis-sesquisabienene hydrate, germacrene B	
<i>H. hirsutum</i>	AE	germacrene B, citronellyl propanoate, γ -gurjunene	
<i>H. linarioides</i>	AE	(E, E)-farnesyl acetate, cis-cadinene ether, 1-tridecene	
<i>H. tetrapterum</i>	AE	trans-linalool oxide, p-cymenene, (E, E)-farnesyl acetate	
<i>H. vermiculare</i>	AE	α -pinene, myrcene, E- β -farnesene	
<i>H. asperulum</i>	AE	(α -pinene); (caryophyllene oxide); (E-caryophyllene) and (spathulenol)	[37]
<i>H. scabrum</i>	AE	(α -pinene); (β -pinene); (limonene) and (E-caryophyllene)	
<i>H. vermiculare</i>	AE	(α -pinene); (caryophyllene oxide); (E-caryophyllene) and (spathulenol)	
<i>H. pseudolaeva</i>	AE	(trans-caryophyllene); (δ -limonene); (α -cadinol); (caryophyllene oxide); (α -pinene); (spathulenol) and (β -selinene)	[38]
<i>H. thymbrifolium</i>	AE	α -pinene, undecane, germacrene D, β -pinene, β -myrcene, spathuleneol, naphthalene	[18]
<i>H. humifusum</i>	AE	(n-undecane); (α -pinene); (β -pinene); (limonene); (myrcene)	
<i>H. bupleuroides</i>	AE	(β -sesquiphellandrene); (β -caryophyllene); (selina-3,7(11)-diene); (γ -elemene); (undecane); (germacrene-B)	[20]
<i>H. gaitii</i>	AE	α -pinene, allo-aromadendrene, δ -cadinene, n-nonane, β -caryophyllene, α -selinene	[39]
<i>H. mexicanum</i>	AE	n-nonane, α -pinene	[40]
<i>H. myricariifolium</i>	AE	α -pinene, β -caryophyllene	[41]
<i>H. juniperinum</i>	AE	n-nonane, α -pinene, geranyl acetate, and β -caryophyllene	
<i>H. heterophyllum</i>	AE	germacrene-D, bicyclogermacrene, d-cadinene, spathulenol, a-guaiaene, and valencene	[41]
<i>H. amblysepalum</i>	AE	δ -3-carene, caryophyllene-oxide, cis-ocimene, β -caryophyllene, α -pinene	[42]
<i>H. spectabile</i>	AE	β -caryophyllene, germacrene D, α -cadinol, caryophyllene oxide	
<i>H. helianthemoides</i>	AE	α -pinene, δ -3-carene, d-limonene, cis-ocimene, undecane	[43]
<i>H. acmosepalum</i>	AE	(ar-curcumene); (β -selinene)	
<i>H. beanii</i>	AE	(γ -muurolene); (β -selinene); (caryophyllene oxide)	
<i>H. calycinum</i>	AE	α -terpineol, P-pinene	
<i>H. choisyianum</i>	AE	cis-eudesma-6,11-diene	
<i>H. forrestii</i>	AE	α -pinene, caryophyllene oxide	
<i>H. kouytchense</i>	AE	cis- β -guaiaene, γ -muurolene	
<i>H. lancasteri</i>	AE	β -selinene, eudesmadienone	
<i>H. leschenaultii</i>	AE	(Cuparene) and (γ -muurolene)	
<i>H. monogynum</i>	AE	Tricosane, myrcene	
<i>H. patulum</i>	AE	β -selinene	
<i>H. pseudohenryi</i>	AE	β -selinene	
<i>H. X moserianum</i>	AE	γ -muurolene, δ -cadinene	
<i>H. salsolifolium</i>	AE	α -pinene, limonene, spathulenol, β -pinene, germacrene D	[44]
<i>H. retusum</i>	AE	α -pinene, limonene, spathulenol, β -pinene, germacrene D	[45]
<i>H. maculatum</i>	-	β -farnesene, n-undecane, β -caryophyllene, δ -cadinene, muurolene	
<i>H. foliosum</i>	-	n-nonane, limonene, terpinolene, β -caryophyllene, β -pinene	[46]
<i>H. brasiliense</i>	WP	β -Caryophyllene, α -Humulene, Caryophyllene oxide, Cubenol, aromadendrene	[47]

AE: Aerial parts; FL: flower; LE: leaf; WP: whole plant

The essential oil (EO) composition of about 50 different *Hypericum* species have so far been identified [26]. The main components of some *Hypericum* species EOs are presented in Table 1. Some components were detected in more than one species, such as α -pinene was identified in the EO of *H. vermiculare*, *H. pseudolaeva*, *H. thymbrifolium*, *H. humifusum*, *H. ericoides*, *H. apricum*, *H. laricifolium*, *H. lydium*, *H. richeri*, *H. patulum*, *H. lysimachioides*, *H. triquetrifolium*, *H. dogonbadanicum*, *H. helianthemoides*, *H. hyssopifolium*, *H. uniglandulosum*, *H. kotschyianum*, *H. thymopsis*, *H. perforatum*, *H. scabrum*. Whereas β -Pinene was identified in the EO of *H. humifusum*, *H. lydium*, *H. richeri*, *H. patulum*, *H. thymbrifolium*, *H. dogonbadanicum*, *H. helianthemoides*, *H. hyssopifolium*, *H. kotschyianum*, *H. perforatum*, *H. scabrum*. Undecane was also introduced as one of the main compounds of *H. bupleuroides*, *H. thymbrifolium*, *H. humifusum*, *H. ericoides*, *H. hirsutum*, *H. rumeliacum*, *H. elegans* EOs. Spathuleneol was also identified in the EOs of *H. pseudolaeva*, *H. thymbrifolium*, *H. kotschyianum*, *H. richeri*, *H. thymopsis*, *H. scabroides*. hexadecanoic acid was identified in the EOs of *H. scabroides*, *H. uniglandulosum*, *H. kotschyianum*, *H. salsugineum*.

In contrast, some compounds were detected as main constituents in the unique species, such as copaene in the EO of *H. tetrapterum*, α -muurolol in the EO of *H. asperulum*, cis-piperitol acetate in the EO of *H. apricum* and verticilol in the EO of *H. laricifolium*.

Previous investigations on the EO composition of *Hypericum* plants did not give homogenous results. These studies have shown that this variance depends on genetics and environment, ontogeny, season, analytical method [48], geographical distribution, type of glands and phenological cycle [32].

For instance, the *H. perforatum* EO extracted from aerial parts obtained from the Republic of Moldova contained β -caryophyllene, caryophyllene oxide, α -pinene, β -cadinene, and β -pinene in the following percentages (12.175%, 12.119%, 8.574%, 4.155%,

3.216%), respectively [24], while another *H. perforatum* EO extracted from aerial parts obtained from southeastern Serbia showed germacrene D, (E)-caryophyllene, 2-methyloctane, α -pinene, bicyclogermacrene and (E)- β -ocimene as main volatile constituents in the following percentages (18.6%, 11.2%, 9.5%, 6.5%, 5.0%, 4.6%), respectively [29].

The variability of the EOs composition depend also on the part analyzed [48]. For instance, the flowers oil of *H. dogonbadanicum* was dominated by phenyl ethyl octanoate, terpin-4-ol, and α -phellandrene, whereas the oil from the leaves of the same plant had α -pinene, β -pinene and p-cymene as its main compounds.

The flowers oil of *H. hyssopifolium* was quite different from the leaves oil. α -pinene, β -pinene and n-tetradecan were the main compounds of the flowers oil, where the leaves contained E-nerolidol, n-tetradecane and α -himachalene.

4. Chemical Composition

This review summarizes the chemical constituents from the genus *Hypericum* which are mainly responsible for its pharmacological benefits. We documented about 100 different natural products and 34 different plant species as sources for these natural products (Table 2). Most of the chemical constituents were mainly isolated from the aerial parts and a few from roots and fruits. The isolated compounds are in the class of phenolic compounds, flavonoids and its glycosides, phloroglucinols, triterpenoid, benzophenones, naphthodianthrone. Flavonoids were the most abundant among the isolated compounds. *Hypericum* species were also characterized by containing several compounds of the xanthenes class.

According to this survey, quercetin, rutin, hypericin, isoquercitrin, quercetrin, chlorogenic acid, pseudohypericin and I3-II8 Biapigenin are the most abundant among secondary metabolites isolated from *Hypericum* species. The pharmacological researches mainly focus on flavonoids such as rutin, hyperoside, and quercetin, and naphthodianthrone such as hypericin.

Table 2. Compounds isolated from *Hypericum* species

Compound	Plant species	Plant part	Ref.
Phenolic compounds			
3-geranyl-1-(2'-methylbutanoyl) phloroglucinol	<i>H. scruglii</i>	AE	[49]
	<i>H. empetrifolium</i>	FR	[50]
3-geranyl-1-(2'-methylpropanoyl) phloroglucinol	<i>H. scruglii</i>	AE	[49]
	<i>H. empetrifolium</i>	FR	[50]
3-(13-hydroxygeranyl)-1-(2'-methylbutanoyl) phloroglucinol	<i>H. scruglii</i>	AE	[49]
adhyperfoliatin	<i>H. trichocaulon</i>	AE	[51]
(E)-chlorogenic acid	<i>H. trichocaulon</i>	AE	
procyanidin A2	<i>H. trichocaulon</i>	AE	
Uliginosin B	<i>H. caprifoliatum</i>	AE	[52]
	<i>H. denudatum</i>	AE	
	<i>H. polyanthemum</i>	AE	
Uliginosin A	<i>H. brasiliense</i>	LE - FL	[53]
Isouliginosin B	<i>H. denudatum</i>	AE	[52]
	<i>H. polyanthemum</i>	AE	
	<i>H. brasiliense</i>	LE - FL	[53]
Hyperbrasilol B	<i>H. austrobrasiliense</i>	AE	[52]
	<i>H. caprifoliatum</i>	AE	
	<i>H. pedersenii</i>	AE	
Hyperbrasilol A	<i>H. brasiliense</i>	LE - FL	[53]
	<i>H. denudatum</i>	AE	
Isohyperbrasilol B	<i>H. caprifoliatum</i>	AE	
	<i>H. pedersenii</i>	AE	
Austrobrasilol B	<i>H. austrobrasiliense</i>	AE	
	<i>H. caprifoliatum</i>	AE	
	<i>H. pedersenii</i>	AE	
Isoaustrobrasilol B	<i>H. austrobrasiliense</i>	AE	
Austrobrasilol A	<i>H. austrobrasiliense</i>	AE	[52]
	<i>H. caprifoliatum</i>	AE	
	<i>H. pedersenii</i>	AE	
	<i>H. denudatum</i>	AE	
Denudatin A	<i>H. denudatum</i>	AE	
Japonicine A	<i>H. caprifoliatum</i>	AE	
	<i>H. pedersenii</i>	AE	
	<i>H. brasiliense</i>	LE - FL	[53]

hengshanols A	<i>H. hengshanense</i>	AE	[54]
Hyperforin	<i>H. perforatum</i>	-	[55]
	<i>H. perforatum</i>	FLS	[56]
	<i>H. humifusum</i>	AE	[57]
	<i>H. confertum</i>	FL	[58]
	<i>H. pruinatum</i>	FL	
	<i>H. lydium</i>	FL	
adhyperforin	<i>H. confertum</i>	FL	
	<i>H. pruinatum</i>	FL	
	<i>H. lydium</i>	FL	
1-(6-hydroxy-2,4-dimethoxyphenyl)-2-methyl-1-propanone	<i>H. cistifolium</i>	AE	[59]
hydroquinone	<i>H. lanceolatum</i>	STB	[60]
3-O-Caffeoylquinic acid (chlorogenic acid)	<i>H. richeri</i>	AE	[61]
	<i>H. perforatum</i>	AE	
	<i>H. hirsutum</i>	AE	
	<i>H. androsaemum</i>	AE	
	<i>H. acutum</i>	AE	
	<i>H. maculatum</i>	AE	
	<i>H. confertum</i>	ST- LE- FL	[58]
	<i>H. thymifolium</i>	ST- LE- FL	
	<i>H. linarioides</i>	ST- LE- FL	
	<i>H. pruinatum</i>	ST- LE- FL	
	<i>H. olivieri</i>	LE- FL	
	<i>H. scabrum</i>	ST- LE- FL	
	<i>H. lydium</i>	LE- FL	[56]
	<i>H. perforatum</i>	FLS	
4-O-Caffeoylquinic acid	<i>H. barbatum</i>	AE	[61]
5-O-Caffeoylquinic acid	<i>H. richeri</i>	AE	
	<i>H. barbatum</i>	AE	
	<i>H. perforatum</i>	AE	
	<i>H. hirsutum</i>	AE	
	<i>H. androsaemum</i>	AE	
	<i>H. acutum</i>	AE	

Neochlorogenic acid	<i>H. confertum</i>	ST- LE- FL	[58]
	<i>H. thymifolium</i>	ST- LE- FL	
	<i>H. linarioides</i>	ST- LE- FL	
	<i>H. pruinatum</i>	ST- LE- FL	
	<i>H. olivieri</i>	ST- LE- FL	
	<i>H. scabrum</i>	LE- FL	
	<i>H. lydium</i>	ST- LE- FL	
caffeic acid	<i>H. confertum</i>	LE- FL	
	<i>H. thymifolium</i>	FL	
	<i>H. linarioides</i>	FL	
	<i>H. pruinatum</i>	LE- FL	
2,4-dihydroxybenzoic acid	<i>H. confertum</i>	ST- LE- FL	
	<i>H. thymifolium</i>	LE- FL	
	<i>H. linarioides</i>	LE- FL	
	<i>H. pruinatum</i>	ST- LE- FL	
	<i>H. olivieri</i>	FL	
	<i>H. scabrum</i>	LE- FL	
	<i>H. lydium</i>	FL	
3,4-dihydroxybenzoic acid	<i>H. scruglii</i>	AE	[49]
Quinic acid	<i>H. perforatum</i>	FLS	[56]
3-p-Coumaroylquinic acid	<i>H. perforatum</i>	FLS	
3-Feruloylquinic acid	<i>H. perforatum</i>	FLS	
Flavonoids			
amentoflavone	<i>H. confertum</i>	LE- FL	[58]
	<i>H. thymifolium</i>	FL	
	<i>H. linarioides</i>	LE- FL	
	<i>H. pruinatum</i>	ST- LE- FL	
	<i>H. olivieri</i>	LE- FL	
	<i>H. scabrum</i>	FL	
	<i>H. lydium</i>	LE- FL	
Rutin (quercetin 3-O-rutinoside)	<i>H. perforatum</i>	FLS	[56]
	<i>H. triquetrifolium</i>	AE	[62]

	<i>H. richeri</i>	AE	[61]
	<i>H. perforatum</i>		
	<i>H. hirsutum</i>		
	<i>H. confertum</i>	ST- LE- FL	[58]
	<i>H. thymifolium</i>	ST- LE- FL	
	<i>H. linarioides</i>	ST- LE- FL	
	<i>H. pruinatum</i>	ST- LE- FL	
	<i>H. olivieri</i>	ST- LE- FL	
	<i>H. scabrum</i>	ST- LE- FL	
	<i>H. lydium</i>	ST- LE- FL	
	<i>H. trichocaulon</i>	AE	[51]
	<i>H. calycinum</i>	-	[63]
Hyperoside (quercetin 3-O-galactoside)	<i>H. perforatum</i>	FLS	[56]
	<i>H. trichocaulon</i>	AE	[51]
	<i>H. richeri</i>	AE	[61]
	<i>H. perforatum</i>	AE	
	<i>H. barbatum</i>	AE	
	<i>H. hirsutum</i>	AE	
	<i>H. androsaemum</i>	AE	
	<i>H. acutum</i>	AE	
	<i>H. maculatum</i>	AE	
	<i>H. brasiliense</i>	LE - FL	[53]
	<i>H. ternum</i>	-	[64]
	<i>H. confertum</i>	ST- LE- FL	[58]
	<i>H. thymifolium</i>	ST- LE- FL	
	<i>H. linarioides</i>	ST- LE- FL	
	<i>H. pruinatum</i>	ST- LE- FL	
	<i>H. olivieri</i>	ST- LE- FL	
	<i>H. scabrum</i>	ST- LE- FL	
<i>H. lydium</i>	ST- LE- FL		
<i>H. calycinum</i>	-	[63]	

Quercitrin (quercetin 3-O-rhamnoside)	<i>H. perforatum</i>	FLS	[56]
	<i>H. confertum</i>	ST- LE- FL	[58]
	<i>H. thymifolium</i>	ST- LE- FL	
	<i>H. linarioides</i>	FL	
	<i>H. pruinatum</i>	ST- LE- FL	
	<i>H. olivieri</i>	LE- FL	
	<i>H. scabrum</i>	ST- LE- FL	
	<i>H. lydium</i>	LE	
	<i>H. richeri</i>	AE	[61]
	<i>H. perforatum</i>	AE	
	<i>H. barbatum</i>	AE	
	<i>H. acutum</i>	AE	
	<i>H. maculatum</i>	AE	
	<i>H. trichocaulon</i>	AE	[51]
	<i>H. brasiliense</i>	LE - FL	[53]
<i>H. calycinum</i>	-	[63]	
Isoquercitrin	<i>H. richeri</i>	AE	[61]
	<i>H. perforatum</i>	AE	
	<i>H. barbatum</i>	AE	
	<i>H. hirsutum</i>	AE	
	<i>H. androsaemum</i>	AE	
	<i>H. acutum</i>	AE	
	<i>H. maculatum</i>	AE	
	<i>H. ternum</i>	-	[64]
	<i>H. brasiliense</i>	LE - FL	[53]
	<i>H. confertum</i>	ST- LE- FL	[58]
	<i>H. thymifolium</i>	ST- LE- FL	
	<i>H. linarioides</i>	ST- LE- FL	
	<i>H. pruinatum</i>	ST- LE- FL	
<i>H. olivieri</i>	ST- LE- FL		
<i>H. scabrum</i>	ST- LE- FL		

	<i>H. lydium</i>	ST- LE- FL	
	<i>H. calycinum</i>	-	[63]
Quercetin	<i>H. perforatum</i>	FLS	[56]
	<i>H. hyssopifolium</i>	AE	[65]
	<i>H. confertum</i>	ST- LE- FL	[58]
	<i>H. thymifolium</i>	ST- LE- FL	
	<i>H. linarioides</i>	ST- LE- FL	
	<i>H. pruinatum</i>	ST- LE- FL	
	<i>H. olivieri</i>	ST- LE- FL	
	<i>H. scabrum</i>	ST- LE- FL	
	<i>H. lydium</i>	ST- LE- FL	
	<i>H. richeri</i>	AE	[61]
	<i>H. perforatum</i>	AE	
	<i>H. barbatum</i>	AE	
	<i>H. hirsutum</i>	AE	
	<i>H. androsaemum</i>	AE	
	<i>H. acutum</i>	AE	
	<i>H. maculatum</i>	AE	
	<i>H. brasiliense</i>	LE - FL	[53]
	<i>H. laricifolium</i>	AE	[66]
	<i>H. lanceolatum</i>	FL	[67]
	<i>H. coadunatum</i>	AE	[68]
<i>H. calycinum</i>	-	[63]	
Quercetin 3-methyl ether	<i>H. ternum</i>	-	[64]
3,7-dimethyl ether Quercetin	<i>H. ternum</i>	-	
arabinofuranoside Quercetin-3-O- α -l	<i>H. hyssopifolium</i>	AE	[65]
Quercetin-3-O- β -d-galactopyranoside	<i>H. hyssopifolium</i>	AE	
Quercetin-3-O- β -d-galactopyranoside-7-O- β -d-glucopyranoside	<i>H. hyssopifolium</i>	AE	

Avicularin	<i>H. confertum</i>	ST- LE- FL	[58]
	<i>H. thymifolium</i>	ST- LE	
	<i>H. linarioides</i>	ST- LE- FL	
	<i>H. pruinatum</i>	LE- FL	
	<i>H. olivieri</i>	FL	
	<i>H. scabrum</i>	FL	
	<i>H. lydium</i>	FL	
Catechin	<i>H. confertum</i>	ST- LE- FL	[56]
	<i>H. thymifolium</i>	ST- LE- FL	
	<i>H. linarioides</i>	ST- LE- FL	
	<i>H. pruinatum</i>	ST- LE- FL	
	<i>H. olivieri</i>	ST- LE- FL	
	<i>H. scabrum</i>	ST- LE- FL	
	<i>H. lydium</i>	ST- LE- FL	
	<i>H. perforatum</i>	FLS	[63]
Epicatechin	<i>H. confertum</i>	ST- LE- FL	[58]
	<i>H. thymifolium</i>	ST- LE- FL	
	<i>H. linarioides</i>	ST- LE- FL	
	<i>H. pruinatum</i>	ST- LE- FL	
	<i>H. olivieri</i>	ST- LE- FL	
	<i>H. scabrum</i>	ST- LE- FL	
	<i>H. lydium</i>	ST- LE- FL	
	<i>H. calycinum</i>	-	[63]
Amentoflavone	<i>H. perforatum</i>	FLS	[56]
	<i>H. confertum</i>	LE- FL	[58]
Cyanidin 3-O-glycoside	<i>H. perforatum</i>	FLS	[56]
Cyanidin 3-O-rhamnoside	<i>H. perforatum</i>	FLS	
miquelianin	<i>H. calycinum</i>	-	[63]
myricitrin	<i>H. trichocaulon</i>	AE	[51]

myricetin	<i>H. lanceolatum</i>	FL	[67]
orientin	<i>H. coadunatum</i>	AE	[68]
	<i>H. hirsutum</i>	AE	[61]
2"-O-Acetyl-orientin	<i>H. hirsutum</i>	AE	
I3-II8 Biapigenin	<i>H. perforatum</i>	FLS	[56]
	<i>H. richeri</i>	AE	[61]
	<i>H. perforatum</i>	AE	
	<i>H. barbatum</i>	AE	
	<i>H. hirsutum</i>	AE	
	<i>H. acutum</i>	AE	
	<i>H. maculatum</i>	AE	
	<i>H. lanceolatum</i>	FL	[67]
	<i>H. hyssopifolium</i>	AE	[65]
	<i>H. trichocaulon</i>	AE	[51]
	<i>H. ternum</i>	-	[64]
	<i>H. calycinum</i>	-	[63]
Myricetin 3-O-rutinoside	<i>H. richeri</i>	AE	[61]
Myricetin 3-O-galactoside	<i>H. richeri</i>	AE	
Myricetin 3-O-glucoside	<i>H. richeri</i>	AE	
luteolin	<i>H. brasiliense</i>	LE - FL	[53]
Kaempferol 3-O-rutinoside	<i>H. perforatum</i>	FLS	[56]
kaempferol-3-O-β-Dglucopyranoside	<i>H. lanceolatum</i>	FL	[67]
kaempferol	<i>H. brasiliense</i>	LE - FL	[53]
friedelanone	<i>H. lanceolatum</i>	FL	[67]
friedelan-3-β-ol	<i>H. lanceolatum</i>	FL	
B-type procyanidin dimer	<i>H. perforatum</i>	FLS	[56]
	<i>H. perforatum</i>	FLS	
	<i>H. perforatum</i>	FLS	
Procyanidin trimer	<i>H. perforatum</i>	FLS	

8-C- β -L- arabinopiranosylapigenin	<i>H. coadunatum</i>	AE	[68]
guaijaverin	<i>H. coadunatum</i>	AE	
	<i>H. ternum</i>	-	[64]
	<i>H. brasiliense</i>	LE - FL	[53]
Hovetrichoside C	<i>H. coadunatum</i>	AE	[68]
Naphthodianthrone			
Hypericin	<i>H. triquetrifolium</i>	AE	[62]
	<i>H. perforatum</i>	FLS	[56]
	<i>H. perforatum</i>	-	[55]
	<i>H. humifusum</i>	AE	[57]
	<i>H. confertum</i>	LE- FL	[58]
	<i>H. thymifolium</i>	LE- FL	
	<i>H. linarioides</i>	LE- FL	
	<i>H. pruinatum</i>	LE- FL	
	<i>H. olivieri</i>	LE- FL	
	<i>H. scabrum</i>	FL	
	<i>H. lydium</i>	LE- FL	[61]
	<i>H. richeri</i>	AE	
	<i>H. perforatum</i>	AE	
	<i>H. barbatum</i>	AE	
	<i>H. hirsutum</i>	AE	
	<i>H. acutum</i>	AE	
	<i>H. maculatum</i>	AE	
	<i>H. hyssopifolium</i>	AE	[65]
Protopseudohypericin	<i>H. perforatum</i>	FLS	[56]
Pseudohypericin	<i>H. perforatum</i>	FLS	[58]
	<i>H. confertum</i>	LE- FL	
	<i>H. thymifolium</i>	LE- FL	
	<i>H. linarioides</i>	LE- FL	
	<i>H. pruinatum</i>	ST- LE- FL	
	<i>H. olivieri</i>	LE- FL	

	<i>H. scabrum</i>	FL	
	<i>H. lydium</i>	LE- FL	
	<i>H. richeri</i>	AE	[61]
	<i>H. perforatum</i>	AE	
	<i>H. barbatum</i>	AE	
	<i>H. acutum</i>	AE	
	<i>H. maculatum</i>	AE	
Terpenoid derivatives			
2-benzoyl-3,3-dimethyl-4 <i>R</i> ,6 <i>S</i> -bis-(3-methylbut-2-enyl)-cyclohexanone	<i>H. galioides</i>	AE	[59]
2-benzoyl-3,3-dimethyl-4 <i>S</i> ,6 <i>R</i> -bis-(3-methylbut-2-enyl)-cyclohexanone	<i>H. galioides</i>	AE	
betulinic acid	<i>H. lanceolatum</i>	STB	[69]
	<i>H. lanceolatum</i>	STB - TW	[70]
ursolic acid	<i>H. lanceolatum</i>	STB - TW	
Benzophenones			
2,2',5,6'-tetrahydroxybenzophenone	<i>H. lanceolatum</i>	STB	[69]
isogarcinol	<i>H. lanceolatum</i>	FL	[67]
Hyperibone	<i>H. coadunatum</i>	AE	[68]
Xanthone			
5-hydroxy-3-methoxyxanthone	<i>H. lanceolatum</i>	STB	[69]
6,7-dihydroxy-1,3-dimethoxyxanthone	<i>H. lanceolatum</i>	STB	[60]
1,3,5,6-tetrahydroxy-4-prenylxanthone	<i>H. lanceolatum</i>	STB	
3-hydroxy-5-methoxyxanthone	<i>H. lanceolatum</i>	STB	[69]
	<i>H. lanceolatum</i>	STB	[60]
1,6-dihydroxyxanthone	<i>H. lanceolatum</i>	STB - TW	[70]
euxanthone	<i>H. lanceolatum</i>	FL	[67]
norathyriol	<i>H. lanceolatum</i>	STB - TW	[70]

1,7-dihydroxyxanthone	<i>H. laricifolium</i>	AE	[66]
1,3,8-trihydroxy-2-methoxyxanthone	<i>H. laricifolium</i>	AE	
1,3-dihydroxy-2-methoxyxanthone	<i>H. laricifolium</i>	AE	
1,3-dihydroxy-6-methoxy-xanthone	<i>H. laricifolium</i>	AE	[71]
2,8-dihydroxy-1-methoxyxanthone	<i>H. laricifolium</i>	AE	[66]
3,8-dihydroxy-1,2-dimethoxyxanthone	<i>H. laricifolium</i>	AE	
1-hydroxy-7-methoxyxanthone	<i>H. laricifolium</i>	AE	[71]
2-hydroxy-xanthone	<i>H. laricifolium</i>	AE	
6-deoxyisojacareubin	<i>H. laricifolium</i>	AE	
1,5,6-trihydroxy-7-methoxy-xanthone	<i>H. laricifolium</i>	AE	
hyperixanthone A	<i>H. trichocaulon</i>	AE	[51]
5-O-methyl-2-deprenylrheediaxanthone B	<i>H. roeperanum</i>	RO	[72]
5-O-methylisojacareubin	<i>H. roeperanum</i>	RO	
5-O-demethylpaxanthonin	<i>H. roeperanum</i>	RO	
roeperanone	<i>H. roeperanum</i>	RO	
Dimethylmangiferin	<i>H. perforatum</i>	FLS	[56]
3,6-Dihydroxy-1,5,7-trimethoxy-xanthone	<i>H. perforatum</i>	RO	
Cadensin C	<i>H. perforatum</i>	RO	
Cadensin C isomer	<i>H. perforatum</i>	RO	
γ -Mangostin	<i>H. perforatum</i>	RO	
5-O-Methyl-2-deprenylrheediaxanthone B	<i>H. perforatum</i>	RO	
Cadensin G	<i>H. perforatum</i>	RO	
Garcinone C	<i>H. perforatum</i>	RO	

AE: Aerial parts; ST: Stem; FL: flowers; LE: leaves; FR: Fruits; TW: twigs; RO: roots; STB: stem bark; FLS: Flower shoot.

5. Antimicrobial activity of *Hypericum* species

Microorganisms are present in the environment. Therefore, they can easily access food [73] with the problem of developing resistance, other problems such as high cost and side effects motivated researchers to look for alternative sources of antimicrobial agents, especially plants and plant products [74]. The

genus *Hypericum*. These days, drug-resistant bacteria are rapidly emerging all over the world and pose a threat to the efficiency of antibiotics [75]. Along has been found to produce compounds that have antimicrobial properties [15]. Many studies have been conducted in many countries showing the antibacterial effect of this genus (table 3).

Methanolic extracts of six *Hypericum* species (*H. perforatum*, *H. neurocalycinum*, *H. spectabile*, *H. thymbrifolium*, *H. malatyanum* and *H. pseudolaeve*) were screened for antibacterial effect against many microorganisms. All extracts displayed antibacterial effect against *Staphylococcus aureus* ATCC 6538, *Staphylococcus epidermidis* ATCC 12228 and Methicillin-resistant *Staphylococcus aureus* ATCC33591, with MIC values ranged from 4.8 µg/ml to 156 µg/ml.

Boga et al., (2016) [76] evaluated the antibacterial effect of many extracts of the whole plant of *H. capitatum* var. *capitatum* against *Staphylococcus aureus* ATCC25923, *Escherichia coli* ATCC 25922, *Streptococcus pyogenes* ATCC19615, and *Pseudomonas aeruginosa* ATCC27853. The acetone and methanolic extracts were active on all tested microorganisms. The methanolic extract showed highest effect against *Escherichia coli* with an MIC value of 10 µg/ml [76].

Tchakam et al., (2012) [77] investigated the antibacterial effect of the methanolic extract of *H. lanceolatum* leaves. The extract showed antimicrobial activity against *Pseudomonas aeruginosa* ATCC 27853, *Klebsiella pneumonia* ATCC 13883, *Salmonella typhi* ATCC 6539, *Enterococcus faecalis* ATCC 10541, with MIC values ranged from 32 µg/ml to 64 µg/ml [77]. According to the published literature, the antibacterial activity varies according to the plant part. This is because the concentration of secondary metabolites accumulated in the plant cells varies according to the plant parts [78]. Antimicrobial activity of different parts (leaf, stem and flower) of *H. montbretii* and *H. bupleuroides* was determined using disc diffusion methods against several microbial species (*Bacillus subtilis* ATCC 6633, *Staphylococcus epidermidis* ATCC 12228, *Escherichia coli* ATCC 25922, and *Staphylococcus aureus* ATCC 25923). The results showed that the extracts displayed good-moderate antimicrobial effect, with MIC values ranged between (0.20-100)µg/ml.

Table 3. Antibacterial effect of some *Hypericum* Species against various bacterial strains

Species	Extract/ plant part	Micro-organisms	MIC (mg/ml)	Ref.
<i>H. perforatum</i>	Total methanol / AE	<i>S. aureus</i> (ATCC 6538)	0.0048	[75]
		Methicillin-resistant <i>S. aureus</i> (ATCC33591)	0.039	
		<i>S. epidermidis</i> (ATCC 12228)	0.078	
<i>H. spectabile</i>	Total methanol / AE	<i>S. aureus</i> (ATCC 6538)	0.0048	
		Methicillin-resistant <i>S. aureus</i> (ATCC33591)	0.0048	
		<i>S. epidermidis</i> (ATCC 12228)	0.039	
<i>H. pseudolaeve</i>	Total methanol / AE	<i>S. aureus</i> (ATCC 6538)	0.019	
		Methicillin-resistant <i>S. aureus</i> (ATCC33591)	0.156	
		<i>S. epidermidis</i> (ATCC 12228)	0.078	
<i>H. thymbrifolium</i>	Total methanol / AE	<i>S. aureus</i> (ATCC 6538)	0.039	
		Methicillin-resistant <i>S. aureus</i> (ATCC33591)	0.0048	
		<i>S. epidermidis</i> (ATCC 12228)	0.0048	
<i>H. neurocalycinum</i>	Total methanol / AE	<i>S. aureus</i> (ATCC 6538)	0.0048	
		Methicillin-resistant <i>S. aureus</i> (ATCC33591)	0.0048	
		<i>S. epidermidis</i> (ATCC 12228)	0.0048	
<i>H. malatyanum</i>	Total methanol / AE	<i>S. aureus</i> (ATCC 6538)	0.0048	
		Methicillin-resistant <i>S. aureus</i> (ATCC33591)	0.0048	
		<i>S. epidermidis</i> (ATCC 12228)	0.0048	
<i>H. jovis</i>	EO/	<i>B. cereus</i> (human isolate)	0.0075	

	AE	<i>S. aureus</i> (ATCC 11632)	0.015	[79]
		<i>E. coli</i> (ATCC 25922)	0.0015	
		<i>P. aeruginosa</i> (ATCC 27853)	0.0015	
<i>H. empetrifolium</i>	EO/ AE	<i>B. cereus</i> (human isolate)	0.015	
		<i>S. aureus</i> (ATCC 11632)	0.030	
		<i>E. coli</i> (ATCC 25922)	0.010	
		<i>P. aeruginosa</i> (ATCC 27853)	0.005	
<i>H. amblycalyx</i>	EO/ AE	<i>B. cereus</i> (human isolate)	0.0025	
		<i>S. aureus</i> (ATCC 11632)	0.015	
		<i>E. coli</i> (ATCC 25922)	0.0025	
		<i>P. aeruginosa</i> (ATCC 27853)	0.0025	
<i>H. androsaemum</i>	Methanol (100%)/ AE	<i>S. aureus</i> (NCIMB 8625)	0.1	[80]
		<i>B. subtilis</i> (NCIMB 1026)	0.1	
		<i>E. coli</i> (B 81)	0.1	
		<i>P. aeruginosa</i> (NCIMB 1039)	0.1	
		(ATCC 13048) <i>E. aerogenes</i>	0.1	
<i>H. ericoides</i>	Methanol (100%)/ AE	<i>S. aureus</i> (NCIMB 8625)	0.1	
		<i>B. subtilis</i> (NCIMB 1026)	0.1	
		<i>E. coli</i> (B 81)	0.1	
		<i>P. aeruginosa</i> (NCIMB 1039)	0.1	
		(ATCC 13048) <i>E. aerogenes</i>	0.1	
<i>H. xmoserianum</i>	Methanol (100%)/ AE	<i>S. aureus</i> (NCIMB 8625)	0.1	
		<i>B. subtilis</i> (NCIMB 1026)	0.1	
		<i>E. coli</i> (B 81)	0.1	
		<i>P. aeruginosa</i> (NCIMB 1039)	0.1	
		(ATCC 13048) <i>E. aerogenes</i>	0.1	
<i>H. olympicum</i>	Methanol (100%)/ AE	<i>S. aureus</i> (NCIMB 8625)	0.025	
		<i>B. subtilis</i> (NCIMB 1026)	0.025	
		<i>E. coli</i> (B 81)	0.025	
		<i>P. aeruginosa</i> (NCIMB 1039)	0.05	
		(ATCC 13048) <i>E. aerogenes</i>	0.05	
<i>H. scabrum</i>	EO/ AE	<i>P. aeruginosa</i> (CIP 82118)	> 0.512	[81]
		<i>S. aureus</i> (ATCC 29213)	> 0.512	
<i>H. lanceolatum</i>	Methanol /LE	<i>K. pneumonia</i> (ATCC 13883)	0.064	[77]
		<i>P. aeruginosa</i> (ATCC 27853)	0.064	
		<i>S. typhi</i> (ATCC 6539)	0.032	
		<i>E. faecalis</i> (ATCC 10541)	0.064	
<i>H. humifusum</i>	Ethanol/ AE	<i>S. aureus</i> (ATCC 49444)	0.078	[16]
		<i>L. monocytogenes</i> (ATCC 19114)	0.078	
		<i>B. cereus</i> (ATCC 11778)	0.62	
		<i>P. aeruginosa</i> (ATCC 27853)	0.62	
		<i>S. typhimurium</i> (ATCC 14028)	1.25	
		<i>E. coli</i> (ATCC 25922)	0.62	
<i>H. bupleuroides</i>	Methanol/ LE	<i>B. subtilis</i> (ATCC 6633)	0.00156	[82]
		<i>S. aureus</i> (ATCC 25923)	0.00125	
		<i>S. epidermidis</i> (ATCC 12228)	0.0125	
		<i>E. coli</i> (ATCC 25922)	0.1	
	Methanol/ FL	<i>B. subtilis</i> (ATCC 6633)	0.0002	
		<i>S. aureus</i> (ATCC 25923)	0.025	
		<i>S. epidermidis</i> (ATCC 12228)	0.0125	
		<i>E. coli</i> (ATCC 25922)	0.1	
<i>H. montbretii</i>	Methanol/ LE	<i>B. subtilis</i> (ATCC 6633)	0.0002	
		<i>S. epidermidis</i> (ATCC 12228)	0.00625	
	Methanol ST	<i>B. subtilis</i> (ATCC 6633)	0.0002	
		<i>E. coli</i> (ATCC 25922)	0.1	
<i>Hypericum capitatum</i> var.	Petroleum ether/ WP	<i>S. pyogenes</i> (ATCC 19615)	2	[76]
		<i>S. aureus</i> (ATCC 25923)	-	

<i>capitatum</i>		<i>P. aeruginosa</i> (ATCC 27853)	2.2
		<i>E. coli</i> (ATCC 25922)	-
	Acetone/ WP	<i>S. pyogenes</i> (ATCC 19615)	0.28
		<i>S. aureus</i> (ATCC 25923)	0.25
		<i>P. aeruginosa</i> (ATCC 27853)	0.3
		<i>E. coli</i> (ATCC 25922)	0.015
	Methanol/ WP	<i>S. pyogenes</i> (ATCC19615)	0.25
		<i>S. aureus</i> (ATCC 25923)	0.3
		<i>P. aeruginosa</i> (ATCC 27853)	0.26
		<i>E. coli</i> (ATCC 25922)	0.01

AE: Aerial parts; ST: Stem; FL: flower; LE: leaf; WP: whole plant; *S. aureus*: *Staphylococcus aureus*; *S. epidermidis*: *Staphylococcus epidermidis*; *E. coli*: *Escherichia coli*; *P. aeruginosa*: *Pseudomonas aeruginosa*; *S. pyogenes*: *Streptococcus pyogenes*; *B. subtilis*: *Bacillus subtilis*; *B. cereus*: *Bacillus cereus*; *E. aerogenes*: *Enterobacter aerogenes*; *K. pneumonia*: *Klebsiella pneumonia*; *S. typhimurium*: *Salmonella typhimurium*; *S. typhi*: *Salmonella typhi*; *E. faecalis*: *Enterococcus faecalis*; *L. monocytogenes*: *Listeria monocytogenes*.

6. Antioxidant activity of *Hypericum* species

Oxidative stress is the main cause of many diseases such as atherosclerosis, arthritis, cancer, as well as neurodegenerative diseases [17]. Because some synthetic antioxidants such as BHA and BHT are now suspected of being harmful to human health [83], extensive research is being done to isolate phytochemicals that can act as antioxidants [17]. Many herbs used in complementary medicine have antioxidant potential [84]. *Hypericum* species have been reported to contain many phenolic compounds and are good sources of antioxidants, making them possible to use in ethnomedicine [85].

The antioxidant capacity of *Hypericum* species has been well documented (Table 4). The antioxidant effect of the whole plant extracts of *H. capitatum* var. *capitatum* was tested. The results indicated that the methanolic and water extracts displayed moderate lipid peroxidation inhibitory effect in β -carotene bleaching test and strong inhibition in ABTS test. The methanolic extract also displayed stronger effect than

α -Toc and BHT standards in DPPH test [76].

The antioxidant capacities of the methanolic extracts of three *Hypericum* species (*H. aviculariifolium*, *H. salsugineum*, *H. perforatum*) were evaluated. The methanolic extract of *H. salsugineum* showed the highest antioxidant effect (DPPH inhibition=88.29%) among the extracts [14].

The antioxidant effect of *H. scabrum* and *H. origanifolium* aerial parts extracts was tested. The species showed stronger activities in DPPH test than ascorbic acid and butylated hydroxytoluene standards [19].

When testing the antioxidant activity of any plant extract, one should take into account that this activity varies according to the plant part. Methanolic extracts of different plant parts of *H. hookerianum* (leaf, flower and aerial parts) were tested for antioxidant effect using different methods. The extract from leaf had the strongest antioxidant activity in comparison with the extracts from flowering tops and aerial parts [86].

Table 4. Antioxidant activity of some *Hypericum* species

Species	extract/ plant part	Technique	Concentration (mg/ml)	Results	Ref.
<i>H. perforatum</i>	Methanol (FL)	DPPH	-	scavenging %=32	[2]
<i>H. perforatum</i>	Methanol	DPPH	0.5	scavenging %=81.21 \pm 0.58	[14]
<i>H. salsugineum</i>	Methanol	DPPH	0.5	scavenging %=88.29 \pm 0.96	

<i>H. aviculariifolium</i> subsp. <i>depilatum</i> var. <i>depilatum</i>	Methanol	DPPH	0.5	scavenging % =86.88 ±0.87	
<i>H. spectabile</i>	Ethanol (AE)	DPPH	2	scavenging %=86.74	[84]
<i>H. spectabile</i>	Methanol (AE)	DPPH	-	mg/ml 0.028 0.567 ± EC50=	[87]
		FRAP	5	reducing power= 2.66 ± 0.031 mM Fe 2+	
<i>H. pseudolaeve</i>	Methanol (AE)	DPPH	-	0.036 mg/ml EC50=0.916 ±	
		FRAP	5	reducing power= 2.21 ± 0.015 mM Fe 2+	
<i>H. thymbrifolium</i>	Methanol (AE)	DPPH	-	EC50=0.622 ± 0.051 mg/ml	
		FRAP	5	0.036 reducing power= 2.58 ± mM Fe 2+	
<i>H. neurocalycinum</i>	Methanol (AE)	Anti-LPO	-	EC50= 2.49 ± 0.09 mg/ml	[87]
		DPPH	-	EC50= 0.251 ± 0.01 mg/ml	
		Superoxide anion	-	EC50= 0.613 ± 0.05 mg/ml	
		FRAP	5	FRAP value= 2.39 ± 0.039 mM Fe 2+	
<i>H. malatyanum</i>	Methanol (AE)	Anti-LPO	-	EC50= 4.82 ± 0.03 mg/ml	[86]
		DPPH	-	EC50= 1.54 ± 0.04 mg/ml	
		Superoxide anion	-	EC50= 0.802 ± 0.03 mg/ml	
		FRAP	5	FRAP value= 2.23 ± 0.013 mM Fe 2+	
<i>H. hookerianum</i>	Methanol (LE)	ABTS	-	IC50= 5.60±0.30 µg/ml	[86]
		DPPH	-	IC50= 3.30±0.05 µg/ml	
		Hydrogen peroxide	-	IC50= 58.06±1.95 µg/ml	
		Lipid per oxidation	-	IC50= 58.00±1.42 µg/ml	
	Methanol (FLT)	ABTS	-	IC50= 7.65±0.26 µg/ml	
		DPPH	-	IC50= 5.10±0.12 µg/ml	
		Hydrogen peroxide	-	IC50=117.64±5.14 µg/ml	
		Lipid per oxidation	-	IC50= 295.00±11.2 µg/ml	
	Methanol (AE)	ABTS	-	IC50= 10.90±0.32	
		DPPH	-	IC50= 5.25±0.11 µg/ml	
		Hydrogen peroxide	-	IC50= 74.14±3.82 µg/ml	
		Lipid per oxidation	-	IC50= 850.00±28.4 µg/ml	
<i>H. androsaemum</i>	Methanol (AE)	DPPH	-	IC50= 92.70±2.85 µg/ml	[17]
		anion (super oxide)	0.5	Anion Scavenging%=33.20±1.22	
		anti-lipid peroxidation	0.5	Anti-lipid peroxidation%= 11.13±1.02	
<i>H. androsaemum</i>	Methanol (AE)	DPPH	-	IC50= 0.093 mg/ml	[80]
<i>H. ericoides</i>	Ethyl acetate (AE)	DPPH	-	IC50= 0.295 mg/ml	
<i>H. x moserianum</i>	Ethyl acetate	DPPH	-	IC50= 0.13 mg/ml	

	(AE)				
<i>H. olympicum</i>	Methanol (AE)	DPPH	-	IC50= 0.098 mg/ml	
<i>H. origanifolium</i>	Ethyl acetate (LE)	DPPH	-	IC50= 3.37 ± 0.26 µg/ml	[88]
<i>H. origanifolium</i>	Ethanol-water (AE)	DPPH	-	IC50= 3.79 ± 0.27 µg/ml	[19]
<i>H. scabrum</i>	Ethanol-water (AE)	DPPH	-	IC50= 3.65±0.40 µg/ml	
<i>H. aucheri</i>	Methanol (AE)	DPPH	0.01	scavenging %=58.2 ± 0.1	[6]
		ABTS	0.01	scavenging %=65.6 ± 0.2	
<i>H. barbatum</i>	Methanol (AE)	DPPH	0.01	scavenging %= 31.9 ± 0.1	
		ABTS	0.01	scavenging %= 34.5 ± 0.2	
<i>H. cerastoides</i>	Methanol (AE)	DPPH	0.01	scavenging %=84.2 ± 0.3	
		ABTS	0.01	scavenging %=90.2 ± 0.1	
<i>H. elegans</i>	Methanol (AE)	DPPH	0.01	scavenging %=25.9 ± 0.1	
		ABTS	0.01	scavenging %=31.9 ± 0.2	
<i>H. linarioides</i>	Methanol (AE)	DPPH	0.01	scavenging %=54.2 ± 0.1	
		ABTS	0.01	scavenging %=57.8 ± 0.4	
<i>H. maculatum</i>	Methanol (AE)	DPPH	0.01	scavenging %=56.2 ± 0.2	
		ABTS	0.01	scavenging %=61.9 ± 0.2	
<i>H. richeri</i>	Methanol (AE)	DPPH	0.01	scavenging %= 49.9 ± 0.1	
		ABTS	0.01	scavenging %=55.7 ± 0.2	
<i>H. rumeliacum</i>	Methanol (AE)	DPPH	0.01	scavenging %= 58.9 ± 0.2	
		ABTS	0.01	scavenging %=65.2 ± 0.1	
<i>H. tetrapterum</i>	Methanol (AE)	DPPH	0.01	scavenging %=61.8 ± 0.2	
		ABTS	0.01	scavenging activity%=68.3 ± 0.2	
<i>H. umbellatum</i>	Methanol (AE)	DPPH	0.01	scavenging activity%= 49.1 ± 0.2	
		ABTS	0.01	scavenging activity%=55.9 ± 0.2	
<i>H. humifusum</i>	Ethanol (AE)	DPPH	-	IC50=18.51 ± 4.94 µg/ml	[16]
<i>H. bupleuroides</i>	Methanol (LE)	DPPH	-	IC50=0.06 mg/ml	[82]
<i>H. montbretii</i>	Methanol (LE)			IC50=0.06 mg/ml	
<i>H. capitatum</i> var. <i>capitatum</i>	petroleum ether (WP)	Lipid peroxidation	-	IC50= >200 µg/ml	[76]
		DPPH	-	IC50= >200 µg/ml	
		ABTS	-	IC50= >200 µg/ml	
	acetone (WP)	Lipid peroxidation	-	IC50= >200 µg/ml	
		DPPH	-	IC50= >200 µg/ml	
		ABTS	-	IC50= 88.84 ± 1.40 µg/ml	
	methanol (WP)	Lipid peroxidation	-	IC50= 41.69 ± 1.29 µg/ml	
		DPPH	-	IC50= 16.82 ± 0.58 µg/ml	
		ABTS	-	IC50= 9.24 ± 0.28 µg/ml	
	water (WP)	Lipid peroxidation	-	IC50= 92.85 ± 1.05 µg/ml	
		DPPH	-	IC50= >200 µg/ml	
		ABTS	-	IC50=9.76 ± 0.14 µg/ml	

AE: Aerial parts; ST: Stem; FL: flower; LE: leaf; WP: whole plant; FLT: Flowering tops; Dw: dry weight

7. Conclusions

There are only few reports about the chemical composition of the genus *Hypericum*. In the current study, we compared the chemical composition of EOs

isolated from different *Hypericum* species. This comparison can contribute to the addition of new criteria for chemotaxonomy to distinguish between *Hypericum* species. We have also reviewed several

studies that included the isolation of chemical compounds from *Hypericum* species. We documented and classified about 100 different natural products isolated from *Hypericum* species. The isolated compounds are in the class of phenolic compounds, flavonoids and its glycosides, phloroglucinols, triterpenoid, benzophenones, naphthodianthrone and xanthenes. Antibacterial and antioxidant activities of the genus have also been reviewed. There may be a need for other studies to accurately determine the components responsible for these biological activities, and to determine how to isolate and prepare them.

Conflict of Interest

There is no conflict of interest.

Author contribution

The author was responsible for conception and design of the study, acquisition of data, analysis and interpretation of data, in addition to drafting and revising the manuscript and approving it for submission

References

1. Chassagne, F., Samarakoon, T., Porras, G., Lyles, J. T., Dettweiler, M., Marquez, L., Salam, A., Shabih, S., Farrokhi, D., and Quave, C., (2021) A Systematic Review of Plants with Antibacterial Activities: A Taxonomic and Phylogenetic Perspective, *Frontiers in Pharmacology*, 11, 1–29. doi: 10.3389/fphar.2020.586548.
2. Okmen, G., and Balpınar, N. (2017) The Biological Activities of *Hypericum perforatum* L., *African Journal of Traditional, Complementary and Alternative Medicines*, 14(1):213–218. doi: 10.21010/ajtcam.v14i1.23.
3. Krishnaiah, D., Sarbatly, R., and Nithyanandam, R. (2011) A review of the antioxidant potential of medicinal plant species, *Food and Bioproducts Processing*, 89:217–233. doi:10.1016/j.fbp.2010.04.008.
4. Mimica-Dukić, N., and Božin, B., (2007) Natural Product Communications Essential Oils from Lamiaceae Species as Promising Antioxidant and Antimicrobial Agents, *Natural Product Communications*, 2(4), 445-452.
5. Bhatt, I. D., Rawat, S. and Rawal, R. (2013) Antioxidants in Medicinal Plants. Chapter 13, 295-299. doi: 10.1007/978-3-642-29974-2
- Zheleva-Dimitrova, D., Nedialkov, P., and Kitanov, G. (2010) Radical scavenging and antioxidant activities of methanolic extracts from *Hypericum* species growing in Bulgaria, *Pharmacognosy Magazine*, 6(22):74–78. doi: 10.4103/0973-1296.62889.
2. Vaou, N., Stavropoulou, E., Voidarou, C., Tsigalou, C., and Bezirtzoglou, E. (2021) Towards Advances in Medicinal Plant Antimicrobial Activity: A Review Study on Challenges and Future Perspectives, *Microorganisms*, 9(10), 2041. doi:10.3390/microorganisms9102041.
3. Kebede, T., Gadisa, E., and Tufa, A. (2021) Antimicrobial activities evaluation and phytochemical screening of some selected medicinal plants: A possible alternative in the treatment of multidrug-resistant microbes. *Plos One*, 16(3):1–16. doi:10.1371/journal.pone.0249253.
4. Endes, Z., Uslu, N., Gümüşçü, A., and Özcan, M. (2015) Antioxidant activity of some traditionally medicinal plants from Turkey, *Journal of Agroalimentary Processes and Technologies*, 21(3):213–215.-
5. Ion, V., Ielciu, I., Cârje A., Muntean, D., Crisan, G., and Păltinean, R., (2022) *Hypericum* spp. -An Overview of the Extraction Methods and Analysis of Compounds, *Separations*, 9, 17:1–21. doi.org/10.3390/separations9010017.
6. Silva, A., Taofiq, O., Ferreira, I., and Barros, L., (2021) *Hypericum* genus cosmeceutical application– A decade comprehensive review on its multifunctional biological properties, *Industrial Crops and Products*, 159, doi.org/10.1016/j.indcrop.2020.113053.
7. Gedik, O., (2022) Fixed and essential oil composition of *Hypericum venustum*, *Hypericum scabrum* and *Hypericum spectabile* species, *Progress in Nutrition*, 24(2):1–8. doi:10.23751/pn.v24i2.12172.
8. Rojas, J., Buitrago, A., Rojas, L., and Morales, A., (2013) Chemical Composition of *Hypericum laricifolium* Juss. Essential Oil Collected from Mérida-Venezuela. *Medicinal*

- & *Aromatic Plants*, 2:132. doi:10.4172/2167-0412.1000132.
9. Maltas, E., Uysal, A., Yildiztugay, E., Aladag, M., Yildiz, S., and Kucukoduk, M., (2013) Investigation of Antioxidant and Antibacterial Activities of Some *Hypericum* Species, *Fresenius Environmental Bulletin*, 22, 3a, 862–869.
 10. Bejaoui, A., Salem, I., Rokbeni, N., M'rabet, Y., Boussaida M., and Boulila A., (2017) Bioactive compounds from *Hypericum humifusum* and *Hypericum perforatum*: inhibition potential of polyphenols with acetylcholinesterase and key enzymes linked to type-2 diabetes, *Pharmaceutical Biology*, 55(1):906–911. doi.org/10.1080/13880209.2016.1270973.
 11. Toiu, A., Vlase, L., Drăgoi, C., Vodnar, D., and Oniga, I., (2016) Phytochemical Analysis, Antioxidant and Antibacterial Activities of *Hypericum humifusum* L. (Hypericaceae), *FARMACIA*, 64(5):663–667.
 12. Saddiqe, Z., Maimoona, A., Abbas, G., Naem, I., and Shahzad, M., (2016) Pharmacological screening of *Hypericum androsaemum* extracts for antioxidant, anti-lipid peroxidation, antiglycation and cytotoxicity activity, *Pakistan Journal of Pharmaceutical Sciences*, 29(2):397–405.
 13. Rouis, Z., Elaissi, A., Abid, N., Lassoued, M., Cioni, P., Flamini, G., and Aouni, M., (2011) Volatile Components of *Hypericum humifusum*, *Hypericum Perfoliatum* and *Hypericum ericoides* by HS-SPME- GC and HS-SPME-GCMS Using NANO Scale Injection Techniques, *Digest Journal of Nanomaterials and Biostructures*, 6(4):1919–1928.
 14. Seyrekoglua, F., Temizb, H., Esera, F., and Yildirim, C., (2022) Comparison of the antioxidant activities and major constituents of three *Hypericum* species (*H. perforatum*, *H. scabrum* and *H. origanifolium*) from Turkey, *South African Journal of Botany*, 146:723–727. doi:10.1016/j.sajb.2021.12.012.
 15. Demirci, F., and Baser, K., (2006) Volatiles of *Hypericum bupleuroides* Griseb., *Journal of Essential Oil Research*, 18, 650–651.
 16. Caldeira, G., Gouveia, L., Serrano, R. and Silva, O. (2022) *Hypericum* Genus as a Natural Source for Biologically Active Compounds, *Plants*, 11, 2509, 1–16. doi.org/10.3390/plants11192509.
 17. Jaimand, K., Rezaee, M., Mirza M., Naderi M., Mozaffrian V., Azadi R., Golipoor M., and Karimi S., (2013) Essential Oil Composition of Eight *Hypericum* species (Hypericaceae) from Iran: Part II, *Journal of Medicinal Plants and By-products*, 1:61–68.
 18. Balikci, N., Sarimahmut, M., Ari, F., Aztopal, N., Özel, M., Ulukaya, E., Celikler, S., (2020) Toxicity assessment of *Hypericum olympicum* subsp. *olympicum* L. on human lymphocytes and breast cancer cell lines, *Journal of Applied Biomedicine*. 18(1), 1–8. doi:10.32725/jab.2020.002.
 19. Anna, B., Maria, G., Veaceslav, K. Ion, D., and Anatolie, N., (2013) Essential Oil Chemical Composition Biodiversity in The *Hypericum* L. Species from The Spontaneous Flora of the Republic of Moldova, Muzeul Olteniei Craiova. *Oltenia. Studii si comunicari. Stiintele Naturii*, 29(2):47–52.
 20. Çakir, A., Duru, M., Harmandar, M., Ciriminna, R., Passannanti, S., and Piozzi, F. (1998) Comparison of the volatile oils of *Hypericum scabrum* L. and *Hypericum perforatum* L. from Turkey, *Flavour and Fragrance Journal*, 12(4), 285–287. doi.org/10.1002/(SICI)1099-1026(199707).
 21. Özkan E., Demirci, B, Gürer, Ç., Kültür, Ş, Mat, A., and Başer, K., (2013) Essential Oil Composition of Five Endemic *Hypericum* Species from Turkey. *Medicinal & Aromatic Plants*, 2:121, 1-4. doi:10.4172/2167-0412.1000121.
 22. Yüce-Babacan, and E., Bagci, E., (2017) Essential oil composition of *Hypericum uniglandulosum* Hausskn. ex Bornm. and *Hypericum lydium* Boiss. from Turkey, *International Journal of Nature and Life Sciences*, 1(1): 12–16.
 23. Koç E, Arabaci T. (2021) Antioxidant Capacity and Essential Oil Composition of *Hypericum thymopsis* Boiss. (Hypericaceae) from Turkey. *Hacettepe J Biol Chem*. 49(4):385–390.
 24. Đorđević, A., (2015) Chemical Composition of

- Hypericum perforatum* L. Essential Oil, *Advanced technologies*, 4(1):64–68.
25. Jaimand, K., Rezaee, M., Naderi, M., Mozaffarian, V., Azadi, R., Karimi, S., and Gholipour, M. (2012) Chemical Composition of the Essential Oils of Six *Hypericum* Species (Hypericaceae) from Iran, *Journal of Medicinal Plants and By-products*, 1:7–11.
 26. Sajjadi, S., Mehregan, I. and Taheri, M. (2015) Essential oil composition of *Hypericum triquetrifolium* Turra growing wild in Iran, *Research in Pharmaceutical Sciences*, 10(1):90–94.
 27. Jerkovića, I., Marasovića, M., Marijanovićb, Z., Pilepić, K., Malešć, Ž., and Miloša, M., (2013) Chemical Composition of *Hypericum richeri* subsp. *grisebachii* Essential Oil from Croatia, *Natural Product Communications*, 8(2):231–233. doi:10.1177/1934578X1300800226.
 28. Morshedloo, M., Yazdani, D., and Aghayi, F. (2014) Chemical Composition of the Essential Oil from *Hypericum patulum* Thunb. Cultivated in Iran, *Journal of Essential Oil Bearing Plants*, 17(1), 131-135. doi.org/10.1080/0972060X.2013.862076.
 29. Radulović, N., and Blagojević P. (2012) Chemical Composition of *Hypericum rumeliacum* Boiss. Essential Oil. A New Chemotype of This Pharmacologically Valuable Species?, *Chemistry and Biodiversity*, 9(10), 2324-2341. doi.org/10.1002/cbdv.201200139.
 30. Djordjevic A, Mitic V, Stankov-Jovanovic V, and Stojanovic G. (2020) Chemical Composition, Antimicrobial Activity and Chemotaxonomy of Essential Oil of Aerial Parts of *Hypericum rumeliacum* Boiss. Species, *Journal of Essential Oil Bearing Plants*, 23(2):246-254.
 31. Couladis M, Chinou I, Tzakou O, Petrakis P V, Cooperation G. (2003) Composition and Antimicrobial Activity of the Essential Oil of *Hypericum rumeliacum* subsp. *apollinis* (Boiss. & Heldr.). *Phyther Res.*17:152–154.
 32. Khorshidi J, Morshedloo M, Moradi S. Essential oil composition of three Iranian *Hypericum* species collected from different habitat conditions, Biocatalysis and Agricultural Biotechnology, 2020;28: 101755.
 33. Bagci, E., and Babacan, E. (2013) The Essential Oils of the Aerial Parts of Two *Hypericum* (*H. pseudolaeve* Robson and *H. thymbrifolium* Boiss. & Noë) (Guttiferae) Species from Turkey, *Journal of essential oil-bearing plants*, 13(4):390-397. DOI:10.1080/0972060X.2010.10643840.
 34. Grafakou M, Barda C, Karikas GA, Skaltsa H. (2022) *Hypericum* Essential Oils-Composition and Bioactivities: An Update (2012–2022). *molecules*. 27(5246):1–21.
 35. Patiño-bayona WR, Plazas E, Bustos-cortes J, Prieto- rodríguez JA, Patiño-Ladino O. Essential Oils of Three *Hypericum* Species from Colombia: Chemical Composition, Insecticidal and Repellent Activity Against *Sitophilus zeamais* Motsch. (Coleoptera: Curculionidae). *Rec Nat Prod*. 2020;1–11.
 36. Senkal B, Uskutoglu T. Investigation of the change in essential oil components with the composition of fatty acids from *Hypericum heterophyllum* Vent., an endemic species in Turkey, *Authorea*. 2020.
 37. Babacan EY. (2019) Volatile oils composition of *Hypericum amblysepalum* Hochst., *Hypericum spectabile* Jaub. & Spach. and *Hypericum helianthemoides* (Spach) Boiss. growing in Turkey. *Pharm Chem J*. 6(6):114–20.
 38. Demirci B, Can Baser H, Crockett S, Khan I. (2005) Analysis of the Volatile Constituents of Asian *Hypericum* L. (Clusiaceae, Hyperidoideae) Species, *Journal of Essential Oil Research*, 17(6):659-663.
 39. Bagci, E., and Babacan, E. (2011) Composition of the essential oil of *Hypericum salsolifolium* Hand.- Mazz. and *Hypericum retusum* Aucher from Turkey, *Acta Botanica Gallica*, 158:2, 169-173, DOI: [10.1080/12538078.2011.10516264](https://doi.org/10.1080/12538078.2011.10516264).
 40. Gudžić B, Djokovic D, Vajs V, Palić R, Stojanovic G. (2002) Composition and antimicrobial activity of the essential oil of *Hypericum maculatum* Crantz, Flavour and Fragrance Journal, 17(5):392 -394.
 41. Santos P, Figueiredo A, Barroso J, Pedro L, Scheffer J. Composition of the essential oil of *Hypericum foliosum* Aiton from five Azorean

- islands, Flavour and Fragrance Journal, 1999;14(5):283-286.
42. Abreu IN, Reis MG, Marsaioli AJ, Mazzafera P. (2004) Essential oil composition of *Hypericum brasiliense* choise. Flavour Fragr J. 19:80–82.
 43. Hajdari, A., Mustafa, B., Nebija, D., Kashtanjeva, A., Widelski, J., Glowniak, K., and Novak, J., (2014) Essential oil composition and variability of *Hypericum perforatum* L. from wild population in Kosovo, Current Issues in Pharmacy and Medical Sciences, 27(1), 51–54. doi: 10.2478/cipms-2014- 0013.
 44. Sanna C, Scognamiglio M, Fiorentino A, Corona A, Graziani V, Caredda A, et al. (2018) Prenylated phloroglucinols from *Hypericum scruglii*, an endemic species of Sardinia (Italy), as new dual HIV-1 inhibitors effective on HIV-1 replication. PLoS ONE 13(3): e0195168. doi.org/10.1371/journal.pone.0195168
 45. Crockett S, Wenzig E, Kunert O, and Bauer R. (2008) Anti-inflammatory phloroglucinol derivatives from *Hypericum empetrifolium*, Phytochemistry Letters, 1(1):37-43. doi.org/10.1016/j.phytol.2007.12.003.
 46. Daskalaki A, Grafakou M, Barda C, Kyriotakis Z, Heilmann J, Skaltsa H. (2021) Secondary metabolites from *Hypericum trichocaulon* Boiss. & Heldr., growing wild in the island of crete, 1-5. doi.org/10.1016/j.bse.2021.104294.
 47. Bridia H, Pustay A, Bordignonc S, Picoli S, Poser G, and Ferraz A. (2022) Antimicrobial activity of dimeric acylphloroglucinols isolated from southern Brazilian *Hypericum* species against to resistant bacterial, 36(24):6448-6452. DOI: 10.1080/14786419.2022.2038596.
 48. Rocha L, Marston A, Potterat O, Kaplan M, Stoeckli- Evans H, and Hostettmann K. (1995) Antibacterial phloroglucinols and flavonoids from *Hypericum brasiliense*, Phytochemistry, 40(5):1447-1452. [https://doi.org/10.1016/0031-9422\(95\)00507-4](https://doi.org/10.1016/0031-9422(95)00507-4).
 49. Han Q, Shu G, Cheng H, Wang S, Zhou T, et al. (2022) New Acylphloroglucinols and geranyl- α - pyrones from *Hypericum hengshanense*, Fitoterapia, 162. doi.org/10.1016/j.fitote.2022.105253.
 50. Cossuta D, Vatai T, Báthori M, Hohmann J, Keve T, Simándi B. (2011) Extraction of Hyperforin and Hypericin from ST. John's Wort (*Hypericum Perforatum* L.) With Different Solvents, Journal of Food Process Engineering, 35(2):222-235. DOI: 10.1111/j.1745-4530.2010.00583.x.
 51. Tusevski O, Krstikj M, Stanoeva JP, Stefova M, Simic SG. (2018) Phenolic profile and biological activity of *Hypericum perforatum* L.: Can roots be considered as a new source of natural compounds? South African J Bot. 117:301–310. <https://doi.org/10.1016/j.sajb.2018.05.030>.
 52. Béjaoui A, Salem I, Rokbeni N, M'rabet Y, Boussaid M, and Boulila A. (2017) Bioactive compounds from *Hypericum humifusum* and *Hypericum perforatum*: inhibition potential of polyphenols with acetylcholinesterase and key enzymes linked to type-2 diabetes, Pharm Biol. 55(1):906–911. doi: 10.1080/13880209.2016.1270973.
 53. Camas N, Radusiene J, Ivanauskas L, Jakstas V, Kayikci S, and Cirak C. (2014) Chemical composition of *Hypericum* species from the Taeniocarpium and Drosanthe sections. Plant Syst Evol. 300:953–960. DOI 10.1007/s00606-013-0934- 3.
 54. Crockett SL, Kunert O, Pferschy-Wenzig E-M, Jacob M, Schuehly W and Bauer R (2016) Phloroglucinol and Terpenoid Derivatives from *Hypericum cistifolium* and *H. galioides* (Hypericaceae). Front. Plant Sci. 7:961. doi: 10.3389/fpls.2016.00961.
 55. Kowa TK, Tchokouaha LRY, Nyemb JN, Wabo HK, Tchinda AT, Appiah-opong R, et al. (2020) Antipromastigote and cytotoxic activities of some chemical constituents of *Hypericum lanceolatum* Lam. (Guttifereae). J Appl Pharm Sci. 10(1):26–32. DOI: 10.7324/JAPS.2020.101004.
 56. Zdunica G, Godjevack D, Savikina K and Petrovic S. (2017) Comparative Analysis of Phenolic Compounds in Seven *Hypericum* Species and Their Antioxidant Properties. Nat Prod Commun. 12(11):1805–1811.

57. Alzoubi K, Abdel-Hafiz L, Khabour O, El-Elimat T, Alzubi M, Alali F. (2020) Evaluation of the Effect of *Hypericum triquetrifolium* Turra on Memory Impairment Induced by Chronic Psychosocial Stress in Rats: Role of BDNF, 14:5299-5314.
58. Kırmızıbekmeza H, Bassarellob C, Piacenteb S, Celepa E, Ataya I, Mercanoğluc G and Yeşilada E. (2009) Phenolic Compounds from *Hypericum calycinum* and Their Antioxidant Activity. Nat Prod Commun. 4(4):531–534.
59. Bernardi A, López-Alarcón C, Aspee A, Rech S, Von Poser G.L, Bride R, and Lissp E. (2007). Antioxidant activity of flavonoids isolated from *Hypericum ternum*. Journal of the Chilean Chemical Society, 52(4), 1326-1329. dx.doi.org/10.4067/S0717-97072007000400013.
60. Cakir A, Mavi A, Yıldırım A, Durub M, Harmandarb M, and Kazaz C. (2003) Isolation and characterization of antioxidant phenolic compounds from the aerial parts of *Hypericum hyssopifolium* L. by activity-guided fractionation. Journal of Ethnopharmacology, 87:73–83. doi:10.1016/S0378-8741(03)00112-0.
61. Miguel M, Moiséla M, Brel O, Bourdy G, Cruza M, Dionicio I, Carraz M, Jullian V. (2018) Xanthones from *Hypericum laricifolium* Juss., and Their Antiproliferative Activity Against HEP3B Cells. Rev Soc Quím Perú. 84(4):428–435.
62. Nguemo RBT, Njouendou AJ, Mpetga JDS, Kengne IC, Nago RDT, Fobofou SAT, et al. (2020) Antimicrobial activities of chemical constituents from the flowers of *Hypericum lanceolatum* Lam. (Hypericaceae). Int J Appl Microbiol Biotechnol Res. 8:66–72. doi.org/10.33500/ijambr.2020.08.007.
63. Díaz JG. (2022) Chemical Composition of *Hypericum Coadunatum* Chr. from the Canary Islands. J Mol Struct. 1248:1–13. doi.org/10.1016/j.molstruc.2021.131447.
64. Zofou D, Kowa T.K, Wabo H.K. et al. *Hypericum lanceolatum* (Hypericaceae) as a potential source of new anti-malarial agents: a bioassay-guided fractionation of the stem bark. Malar J 10, 167 (2011). <https://doi.org/10.1186/1475-2875-10-167>.
65. Happi GM, Ahmed SA, Kemayou G, Salau S, Dzouemo LC, et al. (2023) Bioassay-Guided Isolation of Antiplasmodial Compounds from *Hypericum lanceolatum* Lam. (Hypericaceae) and Their Cytotoxicity and Molecular Docking. Biomed Res Int. 1–9. <https://doi.org/10.1155/2023/4693765>.
66. Ramírez-gonzález I, Amaro-luis JM, and Bahsas A. (2013) Xanthones from aerial parts of *Hypericum laricifolium* Juss. Nat Prod Commun. 8(12):1731–2. doi.org/10.1177/1934578X1300801218.
67. Rath G, Potterat O, Mavi S, Hostettmann K. (1996) Xanthones from *Hypericum roeperanum*, Phytochemistry, 43(2):513-520. doi.org/10.1016/0031-9422(96)00284-1.
68. Gonelimali, F., Lin, J., Miao, W., Xuan, J., Charles, F., Chen, M., and Hatab, S. (2018) Antimicrobial Properties and Mechanism of Action of Some Plant Extracts Against Food Pathogens and Spoilage Microorganisms. *Frontiers in Microbiology*. 9(1639):1-9. doi:10.3389/fmicb.2018.01639.
69. Anusha, K., Ganashree, N., Kruthika, G., Poojashree, R., Sushmitha, K., and Prashith, T. R. (2015) Antimicrobial and antioxidant activity of leaf and flower of *Hypericum mysorensense*, *Journal of Chemical and Pharmaceutical Research*, 7(5):981–987.
70. Özkan, E., Çelîk, B., and Mat, A., (2019) Antimicrobial activities of five endemic *Hypericum* species from Anatolia compared with *Hypericum perforatum*, *Journal of Research in Pharmacy*, 23(1):114–119. doi.org/10.12991/jrp.2018.115.
71. Boga, M., Ertas, A., Eroglu-Ozkan, E., Kizil, M., Ceken, B., and Topcu, G. (2016) Phytochemical analysis, antioxidant, antimicrobial, anticholinesterase and DNA protective effects of *Hypericum capitatum* var. *capitatum* extracts, *South African Journal of Botany*. 104:249–257. doi: 10.1016/j.sajb.2016.02.204.
72. Tchakam, P., Lunga, P., Kowa, T., Lonfouo,

- A., Wabo, H., Taponjdjou, L., Tane, P., and Kuiate, J., (2012) Antimicrobial and antioxidant activities of the extracts and compounds from the leaves of *Psorospermum aurantiacum* Engl. and *Hypericum lanceolatum* Lam., *BMC Complementary and Alternative Medicine*, 12, 136:1–8. doi:10.1186/1472-6882-12-136.
73. Selvamohan, T., Ramadas V., and Kishore, S. (2012) Antimicrobial activity of selected medicinal plants against some selected human pathogenic bacteria, *Advances in Applied Science Research*, 3(5):3374–3381.
74. Grafakou, M., Diamanti, A., Antaloudaki, E., Kyriotakis, Z., Cirić, A., Soković, M., and Skaltsa, H. (2020) Chemical Composition and Antimicrobial Activity of the Essential Oils of Three Closely Related *Hypericum* Species Growing Wild on the Island of Crete, Greece, *Applied sciences*, 10, 2823:1–12. doi:10.3390/app10082823.
75. Saddiqe, Z., Naeem, I., Hellio, C., Patel, A., and Abbas, G. (2020) Phytochemical profile, antioxidant and antibacterial activity of four *Hypericum* species from the UK, *South African Journal of Botany*, 133(3):45-53. doi.org/10.1016/j.sajb.2020.05.018.
76. Fahed, L., Beyrouthy, M., Ouaini, N., Eparvier, V., Stien, D., Vitalini, S., and Iriti, M. (2021) Antimicrobial Activity and Synergy Investigation of *Hypericum scabrum* Essential Oil with Antifungal Drugs. *Molecules*, 26, 6545, 1-8. doi.org/10.3390/molecules26216545.
77. Ceylan, S., Harşit, B., Saral, Ö., Özcan, M., and Demir, I. (2020) Comparison of antioxidant and antimicrobial activities of plant extracts Centaury (*Hypericum montbretii* and *Hypericum bupleuroides*) and Blackthorn (*Paliurus spina-christi* Mill) in Artvin, Giresun region of Turkey, *Journal of Forestry Faculty*, 21(1):13–20. doi:10.17474/artvinofd.579494.
78. El Ouariachi, E., Lahhit, N., Bouyanzer, A., Hammouti, B., Paolini, J., Majidi, L., et al. (2014) Chemical composition and antioxidant activity of essential oils and solvent extracts of *Foeniculum vulgare* Mill. from Morocco. *Journal of Chemical Pharmaceutical Research*, 6(4):743–748.
79. Mohammed, F. S., Şabik, A., Dogan, M., Selamoglu, Z., and Sevindik, M. (2020) Antioxidant Potential of *Hypericum spectabile* JAUB. ET SPACH, *Bulletin of Biotechnology*, 1(2):43–45.
80. Ozkan, E., Ozden, T., Ozsoy, N., and Mat, A. (2018) Evaluation of chemical composition, antioxidant and anti-acetylcholinesterase activities of *Hypericum neurocalycinum* and *Hypericum malatyanum*, *South African Journal of Botany*, 114, 104-110. doi.org/10.1016/j.sajb.2017.10.022.
81. Chandrashekhar, H., Venkatesh, P., Ponnusankar, S., and Vijayan, P. (2009) Antioxidant activity of *Hypericum hookerianum* Wight and Arn, *Natural Product Research*, 23(13), 1240–1251. doi: 10.1080/14786410902885070.
82. Ozkan, E., Ozsoy, N., Ozden, T., Ozhan, G., and Mat, A. (2018) Evaluation of Chemical Composition and In-vitro Biological Activities of Three Endemic *Hypericum* Species from Anatolia (*H. thymbriifolium*, *H. spectabile* and *H. pseudolaeve*), *Iranian Journal of Pharmaceutical Research*, 17(3), 1036–1046.
83. Öztürk, N., Tunçel M., and Potoğlu-Erkara, İ. (2009) Phenolic compounds and antioxidant activities of some *Hypericum* species: A comparative study with *H. perforatum*, *Pharmaceutical Biology*, 47(2), 120–127. doi:10.1080/13880200802437073.