

Effect of organic and biostimulants on Yield and quality of evening primrose oil (*Oenothera biennis* L.)

El-Ghawwas, E. O. * and Soliman, M. M. **

*Medicinal and Aromatic Plants Research Department, Horticulture Research Institute, Agriculture Research Center, Giza, Egypt. marwasw@yahoo.com.

**Botanical Gardens Department, Horticulture Research Institute, Agriculture Research Center, Giza, Egypt. m.soliman85@yahoo.com

*Corresponding author: El-Ghawwas, E. O. marwasw@yahoo.com.

Received: 12 November 2023; in revised form: 20 November 2023 / Accepted: 5 December 2023 / Published: 30 December 2023

DOI: [10.21608/jpfs.2023.258502.1002](https://doi.org/10.21608/jpfs.2023.258502.1002)

ABSTRACT

New biotechnologies that would enable a decrease in the usage of chemical inputs without adversely affecting crop output or farmers' income are being sought after by modern agriculture. Vermicompost and seaweed extract are used as nutrient supplements, organic fertilizers and biostimulants in agriculture and horticulture to increase plant growth and yield. This study was conducted at private farm in Beni-Suef, during the two successive years of 2020/2021 and 2021/2022 to investigate the effects of vermicompost (VC), seaweed extract (SW), and chemical fertilizers on the growth, physiological and yield characteristics of evening primrose. The obtained results show that the combination treatment of vermicompost at 2 tons/fed and seaweed extract at 400 ppm gave the highest values of growth characteristics, seed yield and linoleic acid content of oil.

Key words: Evening primrose (*Oenothera biennis* L.), vermicompost, seaweed extract.

INTRODUCTION

During the Green Revolution era, high-yielding varieties were introduced, irrigated areas were expanded, high analysis NPK fertilizers were used, and cropping intensity increased. During this process, the proportion of organic manures to chemical fertilizers as a source of plant nutrients decreased significantly. Additionally, it has been observed that insect pests are becoming more resistant to chemical insecticides. Numerous studies have documented the presence of multinutrient deficits and an overall decrease in the soil's productive capacity as a result of careless fertilizer application. Such concerns and problems posed by modern-day agriculture gave birth to new concepts in farming, such as organic farming. (Madhusudhan, 2017).

Vermicompost improves growth and development of plants when added even in little amounts, also it is effectively utilized in both floriculture and horticulture. In addition to making plants qualified, healthy, and productive, vermicompost also organizes plant development by providing hormones and humic acid. Additionally, it improves soil quality and fertility by promoting microbial activity and preventing the spread of diseases and pests that are carried by the soil. Doklega and Imryed (2020)

Recent years, biostimulants, *i.e.* algal and plant extracts or secondary plant metabolites, have got a lot of attention in the last decay since it acts as synthetic products that capable to stimulate plant growth, stress resistance and increase yield, hence being integrated in agricultural production system (Brown and Saa, 2015). Many reports mentioned that alga

extracts are ecology-oriented products that increase growth and biological yield of different vegetable species (Sharma *et al.*, 2014) by increasing plant resistance against stresses and diseases. Likewise, multicellular alga “Seaweed” is one of the most marine resources that have been used as a biofertilizer since sixteens of this century. It contains many bioactive substances and plant growth regulators such as auxins, cytokinins, and betaines, as well as considers a cost-effective source for minerals and micronutrients (Craigie, 2011).

The use of medicinal herbs for curing diseases has been documented in the history of all civilizations. Herbs are the main source of medicine for a large portion of the global population. Nowadays, with all of its scientific and technological developments, the world is starting to rediscover the health advantages of herbs and spices that previous generations were so familiar with. These advantages are closely related to a greater understanding of how eating affects one's health and well-being. Numerous medications have been created in pharmacology using herbs and their active ingredients.

Evening primrose (*Oenothera biennis* L.) is a plant belonging to the Onagraceae family. It has been indicated that *Oenothera biennis* L. is beneficial in the treatment of many diseases. (Murphy *et al.*, 2004). In recent years, there has been increased interest in plants that produce gamma-linolenic acid (GLA). Evening primrose (*Oenothera biennis* L.) is an important medicinal plant native to North America and South America and is cultivated as an oil seed crop because its seeds are rich in γ -linolenic acid (GLA), which is used in both nutritional and pharmaceutical applications. The seeds contain about 20-30 % oil with approximately 7-10 % GLA, which is transformed in the body into the important prostaglandin that is essential for the proper functioning of cells. GLA intake has been associated with the improvement in some chronic diseases such as rheumatic arthritis, atopic eczema, cardiovascular disease, high blood cholesterol and high blood pressure (Timoszuk *et al.*, 2018).

The main aim of this research was to examine the effects of seaweed extract and vermicompost on yield and quality of evening primrose plants.

MATERIALS AND METHODS

This study was conducted at private farm in Beni-Suef, 29°04'N 31°05'E during the two seasons of 2020/2021 and 2021/2022.

Oenothera biennis L. seeds were obtained from Agriculture research center. The seeds were sown on 20th October 2020/2021 and 2021/2022 in rows of nursery beds inside a greenhouse. After two months of seed sowing, when the seedlings were 15-20 cm in height, seedlings were transplanted to the plots (2.5 × 3 m) that were prepared in the experimental field, with each plot containing three rows. The rows within each plot were 60 cm apart and 2.50 meters in length. Each plot contained 12 seedlings, which were cultivated on three rows, at a spacing of 60 cm between rows, and 60 cm between plants within each row. The physical and chemical properties of the soil of the experimental field are shown in Table (1).

Table 1. Physical and chemical analysis of the soil used for growing Evening Primroseplants during the 2020/2021 and 2021/2022 seasons.

Chemical & physical characteristics	First season	Second season
Clay%	41.36	40.21
Silt%	23.25	23.32
Fine sand%	33.21	34.14
Coarse sand%	2.18	2.33
Soil type	Clay sand	Clay sand
pH	7.11	7.24
N (ppm)	22.4	21.8
P ₂ O ₅ (ppm)	104	106
K ₂ O (ppm)	167	162
Zn (ppm)	5.22	5.19
Fe (ppm)	2.51	2.88
B (ppm)	2.24	2.27
Mn (ppm)	0.67	0.69
Cu (ppm)	0.48	0.41

In both seasons, the plants were supplied with nine different fertilization treatments. The tested treatments included NPK fertilization, as well as adding vermicompost and Seaweed extract, as follows:

1. Control (the recommended NPK fertilization).
2. vermicompost at 2 ton/fed. (VC1).
3. vermicompost at 4 ton/fed. (VC2).
4. Seaweed extract at 200 mg/l (SW1).
5. Seaweed extract at 400 mg/l (SW2).
6. vermicompost at 2 ton/fed. (VC1). + Seaweed extract at 200mg/l (SW1).
7. vermicompost at 4 ton/fed. (VC2). + Seaweed extract at 200 mg/l(SW1).
8. vermicompost at 2 ton/fed. (VC1). + Seaweed extract at 400 mg/l(SW2).
9. vermicompost at 4 ton/fed. (VC2). + Seaweed extract at 400 mg/l(Sw2).

The recommended NPK fertilization (as described by **David et al., 2009**) consisted of ammonium nitrate (33%), calcium superphosphate (15.5%) and potassium sulphate (48%), at the rates of 150, 60 and 60 kg/fed., respectively. The ammonium nitrate was divided into two equal doses (one applied 2 weeks after planting the seedlings, and the other applied 4 weeks later), while both calcium superphosphate and potassium sulphate were mixed into the soil one day before planting.

Vermicompost properties:

Vermicompost was added during soil preparation. Chemical analysis of vermicompost are shown in Table 2

Table 2. Chemical analysis of vermicompost:

OM%	OC%	N%	C/N	P%	K%	pH	EC dS m ⁻¹
35.11	21.08	1.72	1:16	2.18	1.43	7.62	2.17

OM: Organic matter**OC: Organic carbon**

Seaweed extract treatments were sprayed at rate 200 and 400 ppm (SW1 and SW2) until the dripping point. The first seaweed application was added after month from planting, and the treatments were repeated every three weeks throughout the season. The chemical and biochemical analyses of seaweed extract are shown in table (3).

Table 3. Chemical and biochemical analyses of seaweed extract:

Organic matter				Growth regulators			Macro and micro elements						
Amino acid%	Carbohydrates %	Alginic acid%	Manitol %	IAA%	Cytokinins % (Adenine)	N%	P%	K %	Ca %	S %	Mg %	Fe ppm	Zn ppm
13%	32%	8%	2%	0.06%	0.04%	4.12	2.76	4.8	0.2	3.71	0.63	123	62

Layout of the experiment:

According to **Snedecor and Cochran (1989)**, the experiment's design was a randomized complete blocks layout, with 9 fertilization treatments and three replicates (blocks).

1. Recorded data:**a. Vegetative growth characteristics (at the stage of 50% flowering)**

In both seasons, the following growth characteristics were recorded when approximately half of the flowers on each plant were open:

1. Plant height.
2. Number of branches/plant.
3. Plant fresh and dry weights (g). Plant dry weight was recorded after drying the plants in an oven at 70° C until a constant weight was reached (after approximately 48 hours).

b. Seeds production:

The Capsules were collected by hand at the ripening stage, and the following characteristics were recorded:

1. Number of capsules/ plant
2. Capsules dry weight/ plant (g) was recorded after sun-drying of the fruits.
3. Seed weight/plant (g).
4. Seeds weight/ fed. (kg) was calculated by multiplying the seed weight/plant by the number of plants/fed. (11111 plants/fed.).

c. Fixed Oil production:

1. The fixed oil was extracted from the seeds using the Soxhlet method (as described in the **A.O.A.C., 1970**), and the following characteristics were calculated:
2. Oil yield (g/plant) was calculated by multiplying the seeds dry weight by the oil percentage.
3. Oil yield (kg/fed.) was calculated by multiplying the oil yield/plant by the number of plants /feddan.
4. The fatty acid content in the oil was determined using gas chromatography (GC) according to (**Adams, 2007**).

d. Determination of some chemical constituents:

1. Determination of total chlorophyll (mg/g/f.w.) according to **Rodriguez and Miller (2000)**.
2. **NPK determinations:** Mineral content included, nitrogen percentage using kjeldhal method described by **Hach et al., (1985)**, phosphorus percentage was estimated according to **A.O.A.C. (1970)** and potassium percentage was determined by flame photometer using the method described by **Brown and Lilleland, (1946)**.

RESULTS AND DISCUSSION**a. Growth characteristics:**

Results recorded during both seasons (Table, 4) demonstrate that chemical NPK fertilization had a more pronounced effect on increasing growth characteristics of *Oenothera biennis* plants (plant height, number of branches per plant, fresh and dry weight/plant) in comparison to the addition of Vermicompost or seaweed extract individually.

However, combining vermicompost and seaweed extract gave better results than those from NPK fertilization. In both seasons, using the combination of VC1 and SW2 gave the tallest plants 111 cm and 123.21 cm, the highest branches number per plant 30.50 and 32.94 branches/plant, fresh weight of plant 1190.63 and 1798.57g and dry weight of plant 401.72 and 547.30 g in the first and the second seasons, respectively.

Furthermore, the increase in the level of organic matter increases the biological activity in the soil which caused the increase in concentration of available nutrients for absorption by plant roots, hence reflected on positive effects on chlorophyll formation, vegetative growth and yield quantity. These results concur with those attained by (**Arancon and Edwards, 2005; Tejada et al., 2006 and. Frasetya et al.,2019**)

In connection, similar results indicated that foliar applications of seaweed extract increased growth of canola (**Ferreira and Lourens, 2002**) and sweet pepper (**Arthur et al., 2003**). Many reports mentioned that biostimulants improve growth and development of sprayed plants by increasing carbon and nitrogen metabolism (**Gonzalez et al., 2014**), photosynthesis as well as enhancing nutrients uptake from the soil (**Halpern et al., 2015**).

Table 4. Effect of NPK, vermicompost and seaweed extract on growth characteristics of evening primrose plants throughout the 2020/2021 and 2021/2022 seasons.

Treatments	Plant height (cm)		Branches number/plant		Fresh weight/plant (g)		Dry weight/plant (g)	
	First season	Second season	First season	Second season	First season	Second season	First season	Second season
NPK	101.01	113.22	21.96	24.40	796.65	1179.35	215.05	308.59
VC1	88.80	100.64	12.20	14.24	633.93	882.90	179.64	247.12
VC2	93.24	103.97	16.67	20.74	693.43	936.39	194.83	271.26
SW1	84.36	89.17	7.32	10.58	253.12	477.23	67.90	126.35
SW2	87.69	95.83	10.98	16.26	358.58	622.63	96.11	171.65
VC1+SW1	103.23	118.03	26.84	30.90	1023.02	1563.92	339.31	528.18
VC2+SW1	107.30	122.47	29.28	31.72	1159.38	1756.02	384.46	569.08
VC1+SW2	111.00	123.21	30.50	32.94	1190.63	1798.57	401.72	603.04
VC2+SW2	112.08	119.51	28.06	29.28	1106.27	1720.99	361.34	547.30
L.S.D. at 0.05	3.32	4.11	2.32	2.76	19.23	21.09	7.72	10.56

NPK= ammonium nitrate, calcium superphosphate and potassium sulphate at 150, 60 and 60 kg/fed., respectively - VC1, VC2= Vermicompost at 2 and 4 ton /fed., respectively - SW1, SW2= Seaweed extract at 200 and 400 mg/L, respectively

b. Seeds production:

Results observed in the two seasons (Table 5) showed that fertilization of evening primrose plants with NPK had a significantly better effect on increasing capsules number per plant with values 616.37 and 629.07g, capsules dry weight per plant 124.76 and 139.46 g, seed weight per plant 111.88 and 126.91 g and seed yield per feddan 1243.10 and 1410.10 kg in both seasons respectively, compared to vermicompost or seaweed extract individually.

While, capsules number per plant, capsules dry weight per plant and seed weight per plant were significantly increased (in both seasons) by supplying the plants with combinations of VC1 + SW1, VC2 + SW1, VC1+ SW2 or VC2+SW2 compared to NPK fertilization. Among the different tested double treatment combinations, the most effective one for increasing capsules number per plant 869.86 and 906.45, capsules weight per plant 276.79 and 317.60 g, seeds weight per plant 193.06 and 219.47g and seed yield per feddan 2145.09 and 2438.53 kg were supplying the plants with VC1 + SW2 in both seasons.

In our investigation Seaweed extract application promoting evening primrose vegetative growth, yield and its components, these enrichments in the previous characters due to seaweed extract content with high levels of organic matter, microelements, vitamins and amino acids and also has a high concentration of growth regulators like gibberellins, cytokinins, and auxins. (Khan et al., 2009). It has previously been demonstrated that exogenous application of seaweed extract increases plant growth, yield, and quality, as well Shehata et al., (2011) on Celeriac plant. Vermicompost provides plants with easily absorbed nutrients. It provides auxins, which are plant growth hormones that can be produced during the fermentation process and may be the reason for the growth increases and seed yield. These results came from a similar point of view to those reported by

Edwards et al., (2007) and Massoud et al., (2022). The beneficial effects of vermicompost and seaweed extract on *Oenothera biennis* seed production may be attributed to the release of nutrients from vermicompost into the soil. In addition, seaweed has a promotive effect on plants; it has been reported to be an enriched source for cytokinins, vitamins, enzymes, amino acids and minerals. It also releases CO₂, which promotes better plant growth and increases net photosynthesis in plants. **Karimi et al., (2011)** reported that application of vermicompost significantly increased corn yield compared with control treatment. Furthermore, **Amin et al. (2010)** showed that vermicompost is a more effective organic fertilizer than other ones for increasing the grain yield of castor beans.

Table 5. Effect of NPK, vermicompost and seaweed extract on number of capsules/ plant (g), capsules dry weight/ plant (g), seed weight/plant (g) and seed yield/fed. (kg) on evening primrose plants throughout the 2020/2021 and 2021/2022 seasons.

Treatments	Number of capsules/ plant (g)		Capsules dry weight/ plant (g)		Seed weight/plant (g)		Seed yield/fed. (kg)	
	First season	Second season	First season	Second season	First season	Second season	First season	Second season
NPK	616.37	629.07	124.76	139.46	111.88	126.91	1243.10	1410.10
VC1	392.00	401.33	66.88	92.85	59.30	68.83	658.88	764.77
VC2	437.54	493.92	98.51	104.77	70.60	83.06	784.44	922.88
SW1	210.56	222.13	29.75	34.83	21.26	26.13	236.22	290.33
SW2	291.20	299.04	54.32	58.65	32.34	37.17	359.33	413.00
VC1+SW1	688.80	730.24	205.93	232.34	138.97	162.10	1544.10	1801.09
VC2+SW1	804.53	845.23	248.24	285.25	178.59	196.10	1984.31	2178.87
VC1+SW2	869.86	906.45	276.79	317.60	193.06	219.47	2145.09	2438.53
VC2+SW2	772.05	833.28	224.35	249.10	160.96	190.52	1788.43	2116.87
L.S.D. at 0.05	15.83	11.31	7.22	9.53	4.89	8.61	102.59	112.16

NPK= ammonium nitrate, calcium superphosphate and potassium sulphate at 150, 60 and 60 kg/fed., respectively - VC1, VC2= Vermicompost at 2 and 4 ton /fed., respectively - SW1, SW2= Seaweed extract at 200 and 400 mg/L, respectively

c. Fixed Oil production:

Data reported in Table (6) revealed that fertilization of evening primrose plants with NPK gave better effect on increasing fixed oil percentage with values 16.46% and 18.18 %, oil yield per plant 18.42 and 23.07 g and oil yield per feddan 204.66 and 256.33 kg in both seasons, in comparison to vermicompost or seaweed extract independently.

The combination of VC1 and SW2 gave higher fixed oil percentage 22% and 23.71%, oil productivity per plant of 42.47 and 52.03 g/plant and oil yield per fedaan 471.88 and 578.11 kg compared to NPK treatment in the first and second seasons, respectively. The results are in agreement with the findings of **Badr et al. (2013)** on *Helianthus annuus*, who discovered that plants that got organic fertilization produced the highest seed oil yield per plant. Vermicompost performs well as a bio-control agent and organic fertilizer because it provides organic nutrients and promotes plant growth. (**Simsek, 2011**). Oil yield and content can be increased by 10 ton ha⁻¹ vermicompost (**Mohammadi et al., 2012**). Sesame oil yield also increases because of vermicomposting (**Sajadi et al., 2011**). **Ardebili et al., (2012)** discovered that applying seaweed extract at appropriate doses as a source of amino acids improved oil content.

Table 6. Effect of NPK, vermicompost and seaweed extract on fixed oil percentage oil yield/plant (g) and oil yield/fed. (kg) on evening primrose plants throughout the 2020/2021 and 2021/2022 seasons.

Treatments	Fixed oil percentage		Oil yield/ plant (g)		Oil yield/fed. (kg)	
	First season	Second season	First season	Second season	First season	Second season
NPK	16.46	18.18	18.42	23.07	204.66	256.33
VC1	14.90	16.53	8.84	11.38	98.22	126.44
VC2	15.89	15.80	11.21	13.13	124.55	145.89
SW1	12.19	12.88	2.59	3.37	28.78	37.44
SW2	13.10	13.59	4.24	5.05	47.11	56.11
VC1+SW1	19.92	22.01	27.68	35.68	307.55	396.44
VC2+SW1	21.11	22.91	37.69	44.92	418.77	499.11
VC1+SW2	22.00	23.71	42.47	52.03	471.88	578.11
VC2+SW2	20.21	21.05	32.54	40.11	361.55	445.66
L.S.D. at 0.05	0.65	0.53	1.67	1.83	8.21	9.34

NPK= ammonium nitrate, calcium superphosphate and potassium sulphate at 150, 60 and 60 kg/fed., respectively - VC1, VC2= Vermicompost at 2 and 4 ton /fed., respectively - SW1, SW2= Seaweed extract at 200 and 400 mg/L, respectively

Fatty acid analysis with GC:

The kind and amount of fatty acids in oilseed reflect the quality of oil. Evening primrose is a good source of oleic or monolinoleic and linolenic acid or polyunsaturated fatty acids.

Results of the chromatographic analysis of fixed oil samples extracted from (*Oenothera biennis* L.) plants in the second season (Table 7 and Figures 1 to 9) show that, linoleic acid was the most important fixed oil component (with contents of 63.30 – 66.20%), followed by γ -linolenic acid (with contents of 7.22 -14.61%), and oleic acid% (with contents of 10.16 - 12.32%). Also, the saturated and unsaturated fatty acids in *Oenothera biennis* L. oil were higher with NPK treatment, compared to VC or SW individually. Combining VC1 and SW2 gave the highest unsaturated fatty acid content (92.40%), but gave the lowest saturated fatty acid content (4.65%).

On the other hand, vermicompost at rate 4 ton/fed., gave higher linoleic acid (65.13%) than all concentrations of seaweed extract. Regarding the effect of double treatment combinations, it can be seen that the combination of VC1+SW gave the highest unsaturated fatty acid content (91.41%). In most cases, the linoleic acid and γ -linolenic acid contents in the oil of *Oenothera biennis* were higher with the NPK treatment (with values 64.71% and 10.40% respectively), compared to VC or SW independently. The best results were recorded in plants treated with the VC1+SW2, which gave higher linoleic acid and γ -linolenic acid contents (66.20% and 14.61%, respectively), compared to other treatments. According to **Mohammadi et al., (2011)**, linoleic and oleic acid content of rapeseeds was considerably raised by applying vermicompost as opposed to artificial fertilizers. **Monir et al., (2007)** found that the increase in the content of linoleic acid in comparison to the control is less pronounced, as in the variant with 40 t/decare of vermicompost it reaches 76.70%.

Table (7). Effect of NPK, vermicompost and seaweed extract on fatty acids of evening primrose oil throughout the first season (2020/2021).

Treatments	Component							Total fatty acids (%)
	Saturated fatty acids (%)			Unsaturated fatty acids (%)				
	Palmetic acid	Stearic acid	Total	Oleic acid	Linoleic acid	γ-Linolenic acid	Total	
NPK	8.10	1.67	9.77	10.16	64.71	10.40	85.27	95.04
VC1	8.47	0.95	9.42	11.16	63.67	7.53	82.35	91.78
VC2	5.98	2.07	8.04	11.60	65.13	7.22	83.95	91.99
SW1	7.61	1.47	9.08	10.53	63.30	8.54	82.37	91.45
SW2	7.08	0.74	7.81	10.64	64.62	8.46	83.71	91.53
VC1+SW1	5.57	1.03	6.60	11.42	64.86	12.95	89.23	95.83
VC2+SW1	5.15	0.66	5.81	12.32	65.06	13.36	90.75	96.56
VC1+SW2	3.45	1.20	4.65	11.59	66.20	14.61	92.40	97.05
VC2+SW2	3.92	1.51	5.44	11.98	64.56	13.58	90.12	95.56

NPK= ammonium nitrate, calcium superphosphate and potassium sulphate at 150, 60 and 60 kg/fed., respectively - VC1, VC2= Vermicompost at 2 and 4 ton /fed., respectively - SW1, SW2= Seaweed extract at 200 and 400 mg/L, respectively




Fig.1. Chromatogram of the fatty acid of plants treated with NPK.




Fig.2. Chromatogram of the fatty acid of plants treated with VC1.




Fig.3. Chromatogram of the fatty acid of plants treated with VC2.




Fig.4. Chromatogram of the fatty acid of plants treated with SW1.




Fig.5. Chromatogram of the fatty acid of plants treated with SW2.




Fig.6. Chromatogram of the fatty acid of plants treated with VC1+SW1.




Fig.7. Chromatogram of the fatty acid of plants treated with

Fig.8. Chromatogram of the fatty acid of plants treated with

Fig.9. Chromatogram of the fatty acid of plants treated with

1. Palmitic acid (C16:0) 2. Stearic acid (C18:0) 3. Oleic acid (C18:1)
4. Linoleic acid (C18:2) 5. γ -Linolenic acid (C18:3)

d. Chemical constituents:

1. Total chlorophyll (mg/g F.W.) in leaves

According to the data presented in Table (8) and, NPK fertilizer increased total chlorophyll content more effectively of *Oenothera biennis* L. plants with values 1.75 and 1.96 mg/g F.W in both seasons, compared to the addition of VC or SW independently.

Also, the addition of double combination of VC and SW gave better results than the NPK treatment. The double combination of VC1+SW2 gave the highest total chlorophyll content, with values of 1.15 and 1.62 mg/g F.W. in the first season and the second seasons, respectively. Similar results were obtained by **El-Shayeb (2009)** and **El-Hanafy et al., (2016)** on *Oenothera biennis* L. and **Mathivanan et al. (2012)** on *Arachis hypogaea*, who found that organic manure increased chlorophyll content. Also, seaweed extract helps treated plants to accumulate more photosynthetic pigments **Taha and Abdelaziz (2015)**.

2. Percentages of nitrogen, phosphorus and potassium:

The data recorded in both seasons (Table, 8) reveal that, the nitrogen, phosphorus and potassium content varied considerably among plants receiving the different treatments. In both seasons, NPK gave the highest N, P and K content in leaves, with values of 3.37%, 3.41%, 0.47%, 0.46%, 2.84% and 2.74% in the first and the second seasons, respectively. On the other hand, the lowest N, P and K contents (with values of 2.17%, 2.15%, 0.34, 0.37, 1.42% and 1.60%, in the first and the second seasons) was obtained by SW1.

It can also be seen that, the double treatment combination of VC1+SW2 gave a higher N, P and K contents which recorded 3.74%, 4.07%, 0.54%, 0.55%, 3.40% and 3.48% during in the first and the second seasons, respectively, compared to NPK or any other treatment.

Vermicompost increases the rate of mineralization and the amount of elements that are available for plant roots to absorb from the soil. It also raises the concentration of CO₂ in the soil solution, which lowers the soil's alkalinity and promotes growth by enabling roots to absorb more nutrients, which increases the amount of N, P, and K that are subsequently absorbed by

the plant tissues. Furthermore, an increase in chlorophyll content increases the efficiency of the photosynthesis process. These results confirm those obtained by **Frasetya et al., 2019**. Exogenous application of seaweed extract has already been shown to enhance plant growth, stimulation of chlorophylls biosynthesis, stimulation the uptake of N, P, K, Mg, Ca, Zn, Fe and Cu by plants (**Mahdy et al., 2022**).

Table (8). Effect of NPK, vermicompost and seaweed extract on total chlorophyll (mg/g. F.W) in leaves, N, P and K% in dry matter of evening primrose throughout the 2020/2021 and 2021/2022 seasons.

Treatments	Total chlorophyll (mg/g.f.w.)		N% of dry matter		P% of dry matter		K% of dry matter	
	First season	Second season	First season	Second season	First season	Second season	First season	Second season
NPK	1.75	1.96	3.37	3.41	0.47	0.46	2.84	2.74
VC1	1.58	1.61	2.50	2.49	0.38	0.39	1.81	1.90
VC2	1.54	1.57	2.70	2.87	0.41	0.40	2.09	2.17
SW1	1.35	1.37	2.17	2.15	0.34	0.37	1.42	1.60
SW2	1.39	1.42	2.33	2.41	0.31	0.36	1.57	1.69
VC1+SW1	1.91	2.33	3.48	3.70	0.47	0.50	3.36	3.00
VC2+SW1	2.03	2.48	3.63	3.83	0.49	0.53	3.35	3.05
VC1+SW2	2.15	2.62	3.74	4.07	0.54	0.55	3.40	3.48
VC2+SW2	2.01	2.45	3.45	3.73	0.52	0.54	3.19	3.21
L.S.D. at 0.05	0.24	0.36	0.26	0.38	0.03	0.05	0.43	0.56

NPK= ammonium nitrate, calcium superphosphate and potassium sulphate at 150, 60 and 60 kg/fed., respectively - VC1, VC2= Vermicompost at 2 and 4 ton /fed., respectively - SW1, SW2= Seaweed extract at 200 and 400 mg/L, respectively

Conclusions:

It could be recommended that, for commercial production, Evening Primrose (*Oenothera biennis* L.) plants should be fertilized with a combination of vermicompost at the rate of 2 ton/fed. and seaweed extract 400 ppm. This combination had a generally favourable effect on the seed yield, oil productivity, as well as the Linoleic acid and γ -Linolenic acid contents.

REFERENCES

- Adams, R.P. (2007).** In "Identification of Essential Oil Components by Gas Chromatography/mass Spectroscopy". 4th Edition. Allured, Carol Stream, Illinois and USA.
- Amin, G. A., Moghddam, R. P. and Mehallati, N. M. (2010).** Effect of organic manure on yield and yield components of *Ricinus sativus*. *First National Congress of Sustainable Agriculture and Development and Safe Crop Production. Isfahan.* (in persian).
- A.O.A.C. (1970).** Official Methods of Analysis. Association of Official Agricultural Chemists, Washington, DC., USA. pp. 6-12.
- Arancon, N. and Edwards, C. A. (2005).** Effects of vermicomposts on plant growth. International Symposium Workshop on Vermitechnology. Philippines.
- Ardebili, Z. O., Moghadam, A. R. L., Ardebili, N. O. and Pashaie, A. R. (2012).** The Induced Physiological Changes by Foliar Application of Amino Acids in Aloe vera L. Plants. *Plant Omics J.*, 5(3): 279 - 284.

- Arthur, G. D., Stirk, W. A. and van Staden, J. (2003).** Effect of seaweed concentrate on the growth and yield of three varieties of *Capsicum annum*. *South Afri. J. Bot.* 69 (2): 207-211.
- Badr, E. A., Wali, A. M. and Amin, G. A. (2013).** Effect of Microbine inoculation, nitrogen fertilizer and organic manure on productivity of sunflower plants. *J. Appl. Sci. Res.*, 9(8): 5318-5322.
- Brown, P. and Saa, S. (2015).** Biostimulants in agriculture. *Front. Plant Sci.* 6:671, 21-32.
- Brown, J.D. and Lilleland, O. (1946).** Rapid determination of potassium and sodium in plant material and soil extracts by flame photometry. *Proc. Amer. Soc. Hort. Sci.*, 48: 341-346.
- Carigie, J. (2011).** Seaweed extract stimuli in plant science and agriculture. *J. Appl. Phycol.* 23: 371-393.
- David, G., Nita, S. and Borcean, A. (2009).** *Oenothera biennis* L. in south western Banat. *Res. J. Agric. Sci.*, 41(2): 512-517.
- Doklega, S. M. A. and Imryed, Y. F. E. (2020).** Effect of Vermicompost and Nitrogen Levels Fertilization on Yield and Quality of Head Lettuce. *J. of Plant Production, Mansoura Univ.*, 11 (12):1495 - 1499, 2020
- Edwards, C.A., Arancon, N. Q., Emerson, E. and Pulliam, R. (2007).** Suppressing plant parasitic nematodes and arthropod pests with vermicompost teas. *Biocycle*, 48(12): 38-39.
- El-Hanafy, S. H., Mansour, El-Ghawwas, E. O. and Abd El-Khalek, S. N. (2016).** Effect of Cattle Manure, Active Dry Yeast and Humic Substances on Growth, Seed Yield and Oil Productivity of Evening Primrose (*Oenothera biennis*) Plants. *International Journal of PharmTech Research*,9(7): 56-66.
- El-Shayeb, N. S. A. (2009).** Physiological Studies on *Oenothera biennis* Plants. Ph.D. Thesis, Fac. Agric., Benha Univ., Egypt. 241 p.
- Ferreira, M. I., and Lourens, A. F. (2002).** The efficacy of liquid seaweed extract on the yield of canola plants. *South Afr. J. Plant Soil.* 19 (3):159-161.
- Frasetya, B., Harisman; K., Maulid, S. and Ginandjar, S. (2019).** The effect of vermicompost application on the growth of lettuce plant (*Lactuca sativa* L.) *J. Phys.: Conf. Ser.* 1402 033050.
- González, A., Contreras, R. A., Zúiga, G. and Moenne, A. (2014).** Oligo-carrageenan kappa-induced reducing redox status and activation of TRR/TRX system increase the level of indole-3-acetic acid, gibberellin A3 and trans-zeatin in *Eucalyptus globulus* trees. *Molecules.* 19: 12690-12698.10.3390/molecules190812690.
- Hach, S., Brayton, V. and Alan, B. K. (1985).** A powerful Kjeldahl nitrogen method using peroxyanion sulfuric acid. *J. Agric. Food Chem.*, 33: 1117-1120.
- Halpern, M., Bar-Tal, A., Ofek, M., Minz, D., Muller, T. and Yermiyahu, U. (2015).** The use of biostimulants for enhancing nutrient uptake. *Adv. Agron.* 130: 141-174. 10.1016/bs.agron.2014.10.001.
- Jithesh, M. N., Khan, W., Rayirath, U. P., Subramanian, S., Rayorath, P., Hodges, D. M., Critchley, A. T., Craigie, J. S., Norrie, J. and Prithiviraj B. (2009)** Seaweed extracts as biostimulants of plant growth and development. *Journal of Plant Growth Regulation*, 28(4):386-99. doi:10.1007/s00344-009-9103-x.

- Karimi, H., Mazaheri, D., Peyghambari, S.A. and Mirabzadeh Ardakani, M. (2011).** Effect of application of organic and mineral fertilizers on yield and yield components of maize single cross 704. *Iranian Journal of crop Sciences* 2, 611–626.
- Khan, W., Rayirath, U. P., Subramanian, S., Jithesh, M. N., Rayorath, P., Prithivira, B. (2009).** Seaweed extracts as bio stimulants of plant growth and development”, *J. Plant Growth Regul.*, 28: 386-399.
- Madhusudhan, L. (2017)** Organic Farming-Ecofriendly Agriculture. *J Ecosys Ecograph* 6: 209. 1-4 doi:[10.4172/2157-7625.1000209](https://doi.org/10.4172/2157-7625.1000209)
- Mahdy, H. A. A., Zaki, M. F., Abd El-Rheem, Kh. M. and Ibrahim, H. A. (2022).** Physiological Studies on the Effect of Seaweed Extract and Potassium Humate on the Tolerance of Green Bean Plants to Saline Water Irrigation. *Middle East J. Agric. Res.*, 11(4): 1307-1316.
- Massoud; H. Y. A., Abd El-Baset, M. M. and Abd El-Hady, A. B. B. (2022).** Effect of Ground Treatments (Vermicompost) and Foliar Application with some Plant Growth Stimulants on Growth, Chemical Composition, and Oil Yield of Italian Basil (*Ocimum basilicum* L. var. Genovese). *J. of Plant Production, Mansoura Univ.*, 13 (9):733-741.
- Mathivanan, S.; Chidambaram, A. A.; Sundaramoorthy, P. and Bakiyaraj, R. (2012).** Effect of vermicompost on growth and yield of groundnut (*Arachis hypogaea* L.). *Internat. J. Environ. Biol.*, 2(1): 7-11.
- Mohammadi, K.H., Pasari, B. Rokhzadi, A., Ghalavand, A., Aghaalikhani, M. and Eskandari, M. (2011).** Response of grain yield and canola quality to different resources of farmyard manure, vermicompost and bio-fertilizers in Kurdistan region. *Elec. J. Crop Prod.* 4(2), 81–102.
- Mohammadi, K. H., Pasary, B., Rakhzady, A., Ghalavand, A., Alikhani, M. A. and Eskandari, M. (2012).** Response of rapeseed yield and quality of manure, compost and fertilizer in Kurdistan. *Electronic Journal of Crop Production*, 4(2), 101–81.
- Monir, M. A., Malik, M. A. and Saleem, M. F. (2007).** Impact of integration of crop manuring and nitrogen application on growth, yield and quality of spring planted sunflower (*Helianthus annuus* L.), *Pak. J. Bot.*, 39, 441–449.
- Murphy, C. L., McKenny, C.B., Auld, D.L. (2004).** Hopper NW. Field production of texas native evening primrose (*Oenothera* spp.) as a source of gamma-linolenic acid. *Acta Horticulturae*, 629:283-8.
- Rodriguez, I. R. and Miller, G. L. (2000).** Using a chlorophyll Meter to Determine the chlorophyll concentration, nitrogen concentration and visual quality of St. Augustine grass. *HortScience*, 35(4): 751-754.
- Sajadi N. R., Yadavi, A.R., Balouchi, H.R. and Farajee, H. (2011).** Effect of chemical (Urea), organic (vermicompost) and biological (nitroxin) fertilizers on quantity and quality yield of Sesame (*Sesamum indicum* L.). *J. Agric. Sci. Sustain. Prod.* 21(2), 87–101.
- Sharma, H.S., Fleming, C., Selby, C., Rao, J. H. and Martin, T. (2014).** Plant biostimulants: a review on the processing of macroalgae and use of extracts for crop management to reduce abiotic and biotic stresses. *J. Appl. Phycol.*, 26:465-490.
- Shehata, S.M., Abdel-Azem, H. S., Abou El-Yazied, H. S. and El-Gizawy, A. M. (2011).** Effect of foliar spraying with amino acids and seaweed extract on growth chemical constituents, yield and its quality of celeriac plant”, *European J. of Sci. Res.*, 58(2): 257-265.

- Simsek, E. Y. (2011).** The Use of vermicompost Products to Control Plant Diseases and Pests. In: Karaca A (ed) *Biology of Earthworms, Soil Biology* Springer-Verlag Berlin Heidelberg.
- Snedecor, G. W. and Cochran, W. G. (1989).** *Statistical Methods*, 8th Ed., Iowa State University Press, 153 p.
- Taha, S. S. and Abdelaziz M.E. (2015).** Effect of different concentrations of seaweed extract on growth, yield and quality of two carrot (*Daucus carota* L.) cultivars. *Curr. Sci. Int.*, 7(4): 750-759.
- Tejada, M., Garcia, C., Gonzalez, J. L., Hernandez, M. T. (2006).** Use of organic amendment as a strategy for saline soil remediation: influence on the physical, chemical and biological properties of soil. *Soil Biol. Biochem.*, 38(6), 1413-1421.
- Timoszuk, M., Bielawska, K. and Skrzydlewska, E. (2018).** Evening Primrose (*Oenothera biennis*) Biological Activity Dependent on Chemical Composition. *Antioxidants J.*, 7, 108; 1-11. doi:10.3390/antiox7080108