

Production of broth cubes from microwave-dried meet and chicken and comparison with their commercial counterparts Submitted by

إنتاج مكعبات المرق من اللحم البقري والدجاج المجفف بالميكروويف
ومقارنتها بنظيرتها التجارية

الباحثة / مروة أحمد بدوي أحمد

باحثة ماجستير تخصص (التغذية وعلوم الأطعمة) قسم الاقتصاد المنزلي

كلية التربية النوعية ، جامعه اسيوط

د.ا / هبه عطية يسى

د.ا / ناريمان سعيد اسحق

أستاذة التغذية وعلوم الأطعمة ووكيل الكلية أستاذ الطب الشرعي والسموم الإكلينيكية

كلية الطب - جامعه اسيوط

لشئون الدراسات العليا والبحوث

بكلية التربية النوعية

جامعة اسيوط

المجلد السابع - العدد ٢١ - أبريل ٢٠٢٤ - الجزء ١

الترقيم الدولي

P-ISSN: ٢٥٣٥-٢٢٢٩

O - ISSN: ٣٠٠٩-٦٠١٤

موقع المجلة عبر بنك المعرفة المصري/ <https://hgg.journals.ekb.eg/>

العنوان: كلية التربية النوعية - جامعة اسيوط - جمهورية مصر العربية



Add: Faculty of Specific Education-Nile street- Assiut

العنوان : كلية التربية النوعية - شارع النيل - اسيوط

Print ISSN: 2535-2229

Office / Fax 088/2143535

فاكس / مباشر :

On Line ISSN: 3009-6014

Tel 088/2143536

تليفون :

<https://hgg.journals.ekb.eg>

Mob 01027753777

موبايل :

Production of broth cubes from microwave-dried meet and chicken and comparison with their commercial counterparts

Abstract:

This study aimed to produce beef and chicken stock cubes and compare them with their commercial counterparts. It is prepared from chicken and beef dried in the home microwave, adding some healthy spices to it, and comparing the home product with commercial samples of cubed beef and chicken. The chemical composition was also verified, and the sensory evaluation was studied, as well as the effect of shelf life on the moisture percentage. The results obtained indicate that all homemade samples were higher in nutritional value compared to commercial samples. The percentage of protein in chicken and beef samples was (51.04%, 50.46%), respectively, and in commercial samples (7.33%, 7.75%), respectively. The commercial samples were higher in (moisture, ash, raw fat, and total carbohydrates) than the home-made samples. The results also indicated that there were statistically significant differences in sensory evaluation between the home and commercial samples in favor of the home sample. The results indicated a loss in the percentage of moisture in all samples during the specified period of 30 days, except for the commercial chicken broth cubes, which did not cause any loss in moisture during the aforementioned period. It showed that the yield of homemade products (1.2) and the number of serving was 1 pieces. The results showed a color change for the homemade samples during the storage period of (0.90) day. A color change was observed in the homemade samples from (dark brown and brown to light brown and yellow), respectively.

Key words: Microwave, beef, chicken, broth.

Introduction

Foods are organic substances which are consumed for nutritional purposes. Foods undergo spoilage due to microbial, chemical, or physical actions. Nutritional values, color, texture, and edibility of foods are susceptible to spoilage (Rahman., ٢٠٠٧). Food preservation is defined as the processes or techniques undertaken in order to maintain internal and external factors which may cause food spoilage. The principal objective of food preservation is to increase its shelf life retaining original nutritional values, color, texture, and flavor (Rahman., ٢٠١٤).

Red meat is very important for nutrition thanks its protein, lipid, mineral, vitamin and bioactive contents. (Aykın and Erbas., ٢٠١٤). Chicken meat is a substantial food for consumers all over the world. It is rich in essential amino acids and involves large amount of proteins (Cui et al., ٢٠٠٩). Chicken breast meat (a type of white meat) was selected because it has a lower fat content and is normally perceived as a healthier meat option by consumers.(Charlton et al., ٢٠٠٨). The chicken breast can be dehydrated and then is made into a famous natural seasoning (chicken powders) at home and abroad. Chicken powders can be healthy, delicious, and rich in nutritional ingredient (Ran et al., ٢٠١٩).

Drying and dehydration Is the removal of water like volatile materials from the solid materials, for the aim of slowing the growth of microorganisms or chemical reactions (Radoiu., ٢٠٢٠). Drying extends the shelf-life of meat and meat products because of the reduction in water activity (Modi et al., ٢٠٠٧). In addition, the drying process reduces the microbial load and results in lighter volumes and weights, which decrease storage and transportation costs (Doymaz et al., ٢٠١٦).

Drying meat and produce involves the simple process of exposing (Erbaş and Aykın., ٢٠١٦). Meat products are recognized as a highly perishable food with a limited shelf-life during processing, transportation, and storage. Meat should be subjected to preservation methods to increase its shelf-life (Modi et al., ٢٠٠٧). The main advantages of microwave drying are: i) short drying time, ii) more uniform heating) good energy efficiency (almost all electromagnetic energy is converted into heat) improved product quality, and v) flexibility in the production of a wide variety of products (Haghi and Amanifard., ٢٠٠٨ and Monteiro et al., ٢٠١٥). Microwave dried meat products have higher

microbial quality as compared to dried by conventional methods as the destruction of bacteria occurs by microwaves (Duan et al., ٢٠١١).

Broth is important for palatable dishes of various cuisines requiring broth. (Lee and Kim, ٢٠١٥). the manufacturing process of the cubes consists in mixing ingredients (salt, monosodium glutamate, palm oil, caramel, spices such as soybean, locust bean gum, onion, tomato, natural proteins and starch), coloring and flavoring agents (Raw Materials Research and Development Council., ٢٠٠٣ and Rodrigues et al., ٢٠١٦). The nutritional value of a food is related to the qualitative and quantitative composition of the nutrients present in it. Among these, both lipid fraction and sodium content have an enormous relevance because they play important roles on the maintenance of human health (Karsulínova et al., ٢٠٠٧).

Materials and Methods

Materials:

Beef meat was obtained from a local butcher in Assiut, Egypt. The fat of beef meat is extracted by the butcher. Chicken breast meat was purchased from a local market in Assiut, Egypt. Four chicken and beef stock products (CM) were obtained from the local grocery store (market, assiut , Egypt), and an online. All packaged samples were ١٠ g of stock cubes. All samples were kept in a dry and cool place.

Methods:

Preparation of homemade broth

A piece of red beef weighing ٢٠٠ g was washed well and then cut with a sharp knife into pieces of dimensions (length * width * thickness) (١,٥mm*١,٥mm*١mm) (Marlene., ٢٠١٩). A piece of chicken breast meat weighing ٢١٠ g was washed well and then cut with a sharp knife into pieces of dimensions (length*width*thickness) (٢٠mm*٢٠mm *٧mm) for the raw samples to protect its quality (Hii et al., ٢٠١٤). After obtaining the meat and chicken powder, they are mixed with all the ingredients mentioned in (Table ١) and kneaded well and then formed into a cube shape as shown in the markets. The samples were placed in the dish and then dried using the microwave - assisted hot air drying method (Drying experiments were

performed in a microwave oven Bosch GmbH, Munich, (HMT^{٧٢}G^{٤٢٠} model, Robert Bosch Hausgerate Germany), which has a maximum power output of ٨٠٠ W working at ٢٤٥٠ MHz Siemens HB^{٨٦}K^{٥٧٥}, Germany) was set to a fixed air speed (١,٠ m/s). **Table (١): The ingredients of homemade chicken and beef broth cubes.**

The ingredients	Quantity (gm)
garlic powder	١,٦
Ginger	٠,٥
Curry	٠,٧٥
Turmeric	١,٢٥
Paprika	١
Clove	٠,٢٥
Cinnamon	٠,١٥
Cardamom	٠,١٥
Zaatar	٠,١٥
laureate paper	١,٥
onion powder	١,٦
Coriander	٠,٢٥
parsley powder	٠,٥
Iodized Salt	٠,٦٢٥
Caramel	٠,٥
Cumin	٠,٣٧٥
black pepper	٠,٥
lemon salt	١
Starch	١
Precise	٢
Yeast	٠,١٢٥
Corn oil	٣٠ ml
Low fat milk	٣٠ ml
Egg whites	٥ ml
Water	١٠ ml

In this study, a meat and chicken cube was manufactured by mixing the ingredients of (Table ١). Natural spices and herbs were used instead of artificial colorings and monosodium glutamate to

obtain the same color and flavor and to avoid health risks resulting from the chemicals present in the commercial cube. Meat powder (beef and chicken) dried in the microwave was used instead of bone meal and meat flavor to increase the nutritional value of the cube. Use corn oil instead of palm oil. And low-fat liquid milk instead of powdered milk. And liquid egg whites instead of powdered egg whites. Then the dry ingredients were mixed together. Liquids were added (milk, water, egg whites, corn oil). He started mixing the ingredients together and getting a cohesive mixture. The mixture is cut into square shapes with known dimensions ($\text{cm}^2 * \text{cm}^2$). The produced cubes were stored in a plastic container at room temperature until used for sensory evaluation and laboratory analysis, and the artificial preservatives were replaced by adding lemon salt as a natural preservative. All ingredients of the stock cubes (beef, chicken) are detailed in (Table ١).

Chemical analysis of chicken and beef broth cubes:

Moisture was determined according to the methods of AOAC (٢٠٠٠) by taking (٣g) of sample in a pre-weighed porcelain crucible, which was dried to constant weight at 100°C for ١٤ hours. Loss in weight was taken as the moisture content of the sample. The protein was determined by Micro-Kjeldahl according to the methods of (AOAC, ٢٠١٠), ١g of sample was placed in a digestion tube; ٠,٢g CuSO_4 , ١g K_2SO_4 , and ٢٠ml concentrated H_2SO_4 was added to the tube. The sample was let digested on digestion block until white fumes can be seen and continued heated for about ٦٠ - ٩٠ minutes until cleared with no charred material remaining. The tube was placed in the distillation apparatus and ٥٠ml NaOH ٣٢% was added. The ammonia in the sample was steam distilled for ٥ minutes into a receiving flash containing ٤% boric acid the sample was titrated with H_2SO_4 ٠,١N solution. The protein was calculated by the equation: %Nitrogen x ٦,٢٥. The crude fiber was determined with an enzymatic-gravimetric (procedure according to AOAC Method ٩٩١,٤٣. (AOAC, ٢٠١٠)

Crude fiber was determined following the approved method ٩٦٢,٠٩. Crude fiber is lost on ignition of dried residue remaining after digestion of the sample with ١,٢٥% H_2SO_4 and ١,٢٥% NaOH solutions under specific conditions. ٢g of each sample was extracted with ether or petroleum ether and transferred to beakers of ceramic fiber mixture. Two beakers of ceramic fiber mixture for each sample were prepared as follows: ١,٥ g dry weight of the sample was added to each ١٠٠ ml beaker, then ٦٠-٧٥ ml ٠,٢٥٥N H_2SO_4 was added to each beaker and allowed to soak. Beakers were placed on the digestion apparatus with a pre-adjusted hot

plate and boiled for exactly 30 minutes. were filtered through a Buchner funnel (pre-coated with ceramic fiber if extremely fine materials are being analyzed). Beaker was rinsed with 50-70 ml boiling H₂O and washed through Buchner the funnel. The Residue was removed before 200 ml 1.25% NaOH was added and boiled for exactly 30 minutes. Contents were filtered and then washed with 20 ml boiling 1.25% H₂SO₄, 50 ml H₂O and 20 ml alcohol. The residue was transferred to the ashing dish and dried for 2 hours at 130 ± 20 C Then, it was cooled in a desiccator and weighed. The residue was ignited for 30 minutes at 600 ± 100C and cooled in a desiccator before being reweighed. % Crude fiber in ground sample = C = (Loss in weight on ignition loss in weight of ceramic fiber blank) x 100- weight sample AOAC (2010). Fat was determined by Soxhlet extractor according to AOAC (2010), 3g of sample with hexane using Soxhlet apparatus for 6 hours. The residual hexane was removed from the extracted sample by evaporation. The extracted fat was then dried and weighed and ash was determined according to the method of AOAC (2010), the crucibles containing 0g of sample was charred on a heater before being kept in the muffle furnace at 550C for 4 hours until only white matters can be seen. Then, the crucible with ash content was then cooled in a desiccator and weighed accurately to a constant weight.

All these analyses were determined in the Central Laboratory for Chemical Analysis, Faculty of Agriculture, Assiut University. The proximate analysis was carried out in triplicate to obtain a mean value for each nutrient. Carbohydrate content was calculated by difference. Total carbohydrates % = 100 - (moisture% + ash % +fat + % protein + % fiber content). Total calories were calculated using the equation: E = (4.12 × % protein) + (8.37 × % fat) + (4.2 × % carbohydrate) (Crisan and Sands., 1978). Where: E =Energy as calories per 100 grams sample.

Organ leptic evaluation of chicken and beef stock cubes:

Chicken and beef stock cubes were evaluated in terms of (color, taste, flavour, texture and appearance). All these attributes were evaluated by 50 persons (specialists in Nutrition and food science (30 persons) and ordinary consumers (20 persons)) on a hedonic scale ranging from 1 to 9 (1 is very bad and 9 for excellent) according to. (1 =dislike extremely, 2= dislike very much, 3= dislike moderately, 4 = dislike slightly, 5 = neither like a nor dislike, 6 = like slightly, 7 = like moderately, 8 = like very much and 9 = like extremely) according to (Watts et al ., 1989). Scores

were collated and analyzed statistically.

Yield factor:

The calculated food productivity was calculated from the cooked ratio and the results were expressed in the number of servings of 100 grams of beef and chicken broth cubes and powder (Wendy et al., 2015).

Shelf life of beef and chicken broth cubes:

Shelf life in terms of moisture acquisition was conducted by storing the beef and chicken broth cubes at room temperature (30°C). The moisture content was determined beef and chicken broth cubes: (from initial known moisture content) every 10 day.

Statistical analysis:

Data were presented as the mean of duplicate \pm standard deviation (mean \pm SD). A test was used to establish the significance of differences among mean values at ($p < 0.05$). Analysis of variance (ANOVA) was carried out using Proc Mixed of SAS package version 9.2 (SAS, 2008) and means were compared by the Duncan test at a 5% level of significance (Steel and Trrie., 1981).

Results and Discussion:

From data presented in Table (2), showed the gross chemical composition of the commercial and home-made beef and chicken broth cubes. There were significant differences at ($P < 0.05$). (HM) beef broth cube was higher in moisture, crude fiber (28.6%, 0.912% respectively) than (CM) beef bouillon cube (1.37%, 0.92%). It may be due to the ability of fibers to attract and absorb water (Chawla and Patil., 2010). These results are consistent with (Antonio et al., 2017). Also, HM beef broth cube was higher in crude protein content (6.6%) than beef broth cube (CM) (4.04%). The low protein content found in the samples, especially, This may be a result of the amount of protein source used in the broth formulation in the analyzed samples, the beef appeared only from the seventh ingredient (Antonio et al., 2017 & Fatima., 2013).

However, the data showed that carbohydrate content (4.24%) and crude fat content (12.7%) were lower in (HM) beef bouillon cube than in CM content (18.7%, 13.36%, respectively), They had a significant difference ($P < 0.05$). The energy of the (CM) beef broth cube was higher in CM (212.66 kJ/cal) than the HM beef broth cube (142.99 kJ/cal) with a significant difference ($P < 0.05$). These results are in agreement with that reported by (Antonio., 2017). The results in Table (2) showed that there were significant differences ($p \leq 0.05$) between chicken broth cube (CM and HM) in protein content of chicken broth cubes (HM) and the protein content of commercial samples (CM). Protein content of (HM) was significantly higher than (CM), where (HM) recorded (01.04) protein, while (CM) recorded (7.33). This indicated that the chicken broth cubes are rich in protein. (Fatima, 2013) studied the protein content of mage cube (vegetable, chicken and beef) he found that the protein content of the chicken cube was the highest protein content (8.6%), vegetable cube had the lowest protein content (4.8%) on dry weight basis. From the same table it could be noticed that there were significant differences ($p \leq 0.05$) between chicken broth cube (CM and HM) in HM chicken broth cube was higher in moisture. The moisture loss or gain in foods is an important factor that alters the food qualities and organoleptic characteristics; therefore, water retention is an essential component for maintaining food quality (Roudaut and Debeaufort., 2011), Crude fiber (25.19%, 2.32% respectively) than (CM) beef broth cube (0.19%, 0.12% respectively) with a significant difference ($P < 0.05$). While (Anderson et al., 2018) recorded different results. The results showed a difference in the percentage of fiber (25.9%), which is a higher percentage than shown here. Ash is a reflection of the total inorganic matter present in a good sample and it was ranged in chicken broth cubes (HM) and (CM) from (0.862, 28% respectively) dry matter with a significant difference ($P < 0.05$) These results are in agreement with that reported by (Fatima, 2013).

The results showed there were significant differences ($p \leq 0.05$) between chicken broth cube (CM and HM) in Crude fat For samples (HM), (CM) were (11.6%, 5.7% respectively) These results are in agreement with that reported by (Caponio et al., 2002). Crude fat in CM cubes (vegetable, chicken and beef) products and mushroom broth were found to be fairly high ranging between 2.9 to 4.7% respectively. Carbohydrates content of (HM) were significantly lower than (CM), where (HM) recorded (0.76), while

(CM) recorded (25.38) (Fatima, 2013) also recorded an increase in the percentage of carbohydrates in the (CM) samples. The highest percentage of energy was recorded in commercial samples compared to home-made (121.30, 154.30 respectively).

Table (2) The chemical composition of the commercial and home-made beef and chicken broth cubes:

Sample	Moistur -e %	Ash %	Protei n %	Crude Fat %	Crude Fiber %	Total carboh yd- rates %	Energy (kcal/ 100g
Chicken broth cubes (HM)	25.19 ±1.0	5.08 ± 1.0	51.04 ±1.0	11.6 ± 1.0	1.326 ±1.0	5.76 ± 1.0	121.30 ±1.0
Chicken broth cubes (CM)	0.19 ± 0.01	61.28 ±1.0	7.33 ±1.0	5.7 ± 1.0	0.119 ± 0.001	25.38 ± 1.0	154.31 ±1.0
p.valuo	**	*	**	*	*	*	*
Meat broth cubes (CM)	1.37 ± 1.0	58.72 ±1.0	7.75 ± 1.0	18.7 ± 1.0	0.092 ± 0.001	13.37 ± 1.0	212.66 ±1.0
Meat broth cubes (HM)	28.60 ± 1.0	3.09 ± 1.0	50.46 ±1.0	12.7 ± 1.0	0.913 ± 0.001	4.24 ± 1.0	142.09 ±1.0
p.valuo	**	*	**	*	*	*	*

Values are mean ± standard deviation. Means in the same column with different superscripts are significantly different ($p < 0.05$).

The Organ leptic results from Table (3) revealed that there were significant differences ($p \leq 0.05$) chicken and beef broth cube (CM, HM) in color/appearance, taste, mouth feel, flavor, consistency and overall acceptability. The highest mean score

obtained for chicken and beef broth cubes (HM) (9.08, 8.82 respectively), while chicken and beef broth cube (CM) was (6.24, 5.42 respectively). There was significantly decrease in color/appearance, taste, mouth-feel, flavor, consistency and overall acceptability in (CM). The highest scores of chicken and beef broth cubes (HM) may be able to due probably to the high fat and protein, compared to other (CM). The relatively lower mean scores of the chicken and beef broth cube (CM) are probably due to the small protein content and high salt content (Na Cl). All the broth cubes (HM) were still acceptable to the panelists as indicated by their mean score for overall acceptability. These results are consistent with (Fatima., 2013). Increase in meat flavor is caused by a greater activity of the Millard reaction and associated reacting in valuing muscle protein, carbohydrates and lipids and their degradation products (Ekanem and Solomon., 1997). Spices are known to enhance and improve the flavor and color of foods (Ekanem and Achirnewhu., 1998). They vary in their effects in that the aroma and flavor principles in spices are based on the essential contained in them. It would be assumed that the oils present in magi cube and some of the spices may have produced some modification of the flavor, taste and color thus making the meat samples generally accepted by the panelists (Fatima, 2013).

Table (3) Organ leptic evaluation for commercial (CM) and homemade (HM) beef and chicken broth cubes:

Samples	sensory properties					
	Appearance (10)	Texture (10)	Color (10)	Taste (10)	Flavor (10)	Overall acceptability (10)
Chicken cubes (CM)	5.84 ± 1.0	5.94 ± 1.0	6.4 ± 1.0	6.06 ± 1.0	6.06 ± 1.0	6.24 ± 1.0
Chicken cubes (HM)	8.72 ± 1.0	8.46 ± 1.0	8.56 ± 1.0	8.68 ± 1.0	8.58 ± 0.577	9.08 ± 1.0
p.valuo	*	*	*	*	*	*
Beef cubes (CM)	5.76 ± 1.0	6.28 ± 1.0	6.08 ± 1.0	6.22 ± 1.0	5.8 ± 1.0	5.42 ± 1.258

Beef cubes (HM)	٨,٩ ± ١,٠	٩,٠٢ ± ١,٠	٨,٧٤ ± ١,٠	٨,٦٦ ± ١,٠	٨,٨٤ ± ١,٠	٨,٨٢ ± ١,٠
p.valuo	*	*	*	*	*	*

Values are mean ± standard deviation. Means in the same column with different superscripts are significantly different ($p < 0,05$).

Table(٤) showed that the yield of homemade products between Chicken broth cubes and beef broth cubes are Nan significantly. The results revealed that the yield factor in the home-made beef and chicken broth cubes was at isotropic (١,٢), the number of serving was ٦ pieces and the weight of the piece was ٨ grams. Table(٤) Yield of homemade products beef and chicken broth cubes.

Description	Chicken broth cubes(^)g	beef broth cubes(^)g
Raw material (mg)	٤٠,٠٧ ± ٠,٠١	٤٠,٠٧ ± ٠,٠١
p.valuo	N.S	N.S
Factory products (mg)	٤٨ ± ١,٠	٤٨ ± ١,٠
p.valuo	N.S	N.S
Yield factor	١,٢ ± ٠,١	١,٢ ± ٠,١
p.valuo	N.S	N.S
Number of servings	٦ ± ١,٠	٦ ± ١,٠
p.valuo	N.S	N.S

Values are mean ± standard deviation. Means in the same column with different superscripts are Nan significantly.


The results in Table (٥) showed that there were significant differences ($p \leq 0,05$) between chicken and beef broth cube (CM, HM), the moisture content in the beef broth cubes (CM) high significantly decreased ($P < 0,05$) at room temperature (30°C) (open). After 30 days, it became not edible. These results are in agreement with that reported by (Steele, 2004) while beef broth cubes (HM) The rate of moisture loss was constant at ($4,0\%$) during (10 days and 30 days), respectively. was noticed high significantly decreased ($P < 0,05$) on Chicken broth cubes (HM) moisture content was ($4,0\text{g}, 4\text{g}$) during (10 days and 30 days), respectively at room temperature (30°C) (open). This is consistent with (Yang-Ju Son., 2023) moisture is lost with the length of storage period, while Chicken broth cubes (CM) There was no loss of moisture content in it. These results are in agreement with those obtained by (Sandra et al., (2009) and Elizabeth and Judy., (2009). Adriana et al., (2012).

Table (٥): Effect of shelf life on moisture content of commercial (CM) and homemade (HM) beef and Chicken broth cubes at the room temperature (30°C) (open).

Samples	Day(0) (g)	Day(10) (g)	Day(30) (g)
Chicken broth cubes(HM)	$8 \pm 0,1$	$4,0 \pm 0,1$	$4 \pm 0,1$
Chicken broth cubes(CM)	$8 \pm 0,1$	$8 \pm 0,1$	$8 \pm 0,1$
p.valuo	*	*	**
Meat broth cubes(CM)	$8 \pm 0,1$	$4,0 \pm 0,1$	Not edible
Meat broth cubes(HM)	$8 \pm 0,1$	$4,0 \pm 0,1$	$4,0 \pm 0,1$

p.valuo	*	*	**
---------	---	---	----

Values are mean \pm standard deviation. Means in the same column with different superscripts are significantly different ($p < 0.05$).



Fig(1)

Conclusion:

Analysis of homemade beef and chicken stock cubes showed high moisture and protein content and low levels of dietary fiber, carbohydrates, fat and fixed mineral residues. There was also a difference in energy compared to those commercial samples. The shelf life of household samples was determined based on their moisture content, and the results showed a continuous decrease over the specified period. Through sensory evaluation, it was noted that consumers accept the home product more than the commercial

product. The results also reinforce the importance of the consumer checking the nutritional composition mentioned on product labels. The results also showed the effect of shelf life on the shelf life in moisture loss. The loss was at a higher rate in samples of homemade chicken stock cubes compared to the foreign product. When monitoring the color change of the home-made samples during the storage period, a color change was observed in the home-made samples. The home microwave oven was used to dry samples of beef and chicken because it preserves the nutritional value of the foods and gives a better color compared to regular ovens.

REFERENCES

AOAC. (٢٠١٠) : Official Methods of Analysis of Association of Official Analytical Chemists. ١٨th Edition, Washington, DC.

Anderson M. V., Sandra. M.L, Marlene. D,(٢٠١٨) : Physicochemical characterization and comparison of labels of beef bouillon cubes

[Food Science and Technology \(Campinas\)](#) ٣٨(١)

DOI:[10.1590/fst.11.017](https://doi.org/10.1590/fst.11.017)

Antonio al. (٢٠١٧) :Physicochemical characterization and comparison of labels of beef bouillon cubes DDOI:
<https://doi.org/10.1590/fst.11.017>.

Adjangba KM^{١*}, Agbodji Y^١, Agoro S^١ and Amouzou EKS. (٢٠٢٠) : Cooking Salt Content in Food Cubes Commonly used in Northern Togo. Open Journal of Nutrition and Food Sciences ٢٠٢٠ | Volume ٢ | Article ١٠١٢٠٥٨ © ٢٠٢٠ - Medtext Publications. All Rights Reserved. ISSN ٢٦٩٠-٢٤٧٨ Cooking Salt Content in Food Cubes Commonly used in Northern Togo.

Aykin E., Erbas M.(٢٠١٦) : Quality properties and adsorption behavior of freeze-dried beef meat from the Biceps femoris and Semimembranosus muscles. Meat Sci. ٢٠١٦;١٢١:٢٧٢-٢٧٧. doi: ١٠.١٠١٦/j.meatsci.٢٠١٦.٠٦.٠٣٠. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)].

Buket Er Demirhan, Burak Demirhan, Ceren Sönmez, Hilal Torul, Uğur Tamer & Gülderen Yentür. (٢٠١٥) : Monosodium glutamate in chicken and beef stock cubes using highperformance liquid chromatography, Food Additives & Contaminants: Part B, ١:١, ٦٣-٦٦, DOI: ١٠.١٠٨٠/١٩٣٩٣٢١٠,٢٠١٤,٩٩١٣٥٥ To link to this article: <https://doi.org/١٠.١٠٨٠/١٩٣٩٣٢١٠,٢٠١٤,٩٩١٣٥٥>.

Blum D(٢٠١٢). Food that lasts forever, in TIME Magazine. ٢٠١٢.

Boada L.D., Henríquez-Hernández L.A., and Luzardo O.P(٢٠١٦): The impact of red and processed meat consumption on cancer and other health outcomes: epidemiological evidences. Food Chem. Toxicol. ٩٢:٢٣٦-٢٤٤. doi: ١٠.١٠١٦/j.fct.٢٠١٦.٠٤.٠٠٨ [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)].

Caponio, F., T. Gomes and D. Delcuratolo, (٢٠٠٢) : Quantitative and qualitative characterization of the lipid fraction of bouillon cubes. European Food Res. and Tec., ٢١٥: ٢٠٠-٢٠٣

Charlton, K.E.; Probst, Y.C.; Tapsell, L.C.; Blackhall, P.J.(٢٠٠٨) :Food, Health and Nutrition: Where Does Chicken Fit?; Australian Chicken Meat Federation Inc.: North Sydney, Australia, ٢٠٠٨.

Chawla R. and Patil G. (٢٠١٠): Soluble Dietary Fiber. <https://doi.org/١٠.١١١١/j.١٥٤١-٤٣٣٧>.

Crisan E. and Sands A. (١٩٧٨): Nutrition value In: Biology and cultivation of Edibles Mushrooms, edited by Chang ST and Hayes WA, New York, Academic Press. Pp: ١٣٧-١٦٨.

Croitoru MD, Fülöp I, Kincses AM, Dudutz G, Craciun O, Dogaru MT.(٢٠١٠) : Glutamate determination in foodstuffs with a very simple HPLC-UV method. Acta Alimentaria. ٣٩:٢٣٩-٢٤٧.

Cui, C.; Zhou, X.; Zhao, M.; Yang, B.(٢٠٠٩) : Effect of Thermal Treatment on the Enzymatic Hydrolysis of Chicken Proteins. Innov. Food Sci. Emerg. Technol. ٢٠٠٩, ١٠, ٣٧-٤١. DOI: ١٠.١٠١٦/j.ifset.٢٠٠٨.٠٩.٠٠٣.

Doymaz I., Karasu S., Baslar M.(٢٠١٦) : Effects of infrared heating on drying kinetics, antioxidant activity, phenolic content, and color of jujube fruit. J. Food Meas. Charact. ٢٠١٦;١٠:٢٨٣-٢٩١. doi: ١٠.١٠٠٧/s١١٦٩٤-٠١٦-٩٣٠٥-٤. [[CrossRef](#)] [[Google Scholar](#)].

Duan Zhen-Hua., Jiang Li-Na Wang, Ju-Lan Yu Xiao-Yang. and Wang T. (٢٠١١) :Drying and quality characteristics of Tilapia fish fillets dried with hot air-microwave heating. Food and Bioproducts Processing. ٨٩: ٤٧٢-٤٧٦.

Ekanem, E. O. and Solomon, I. P. (١٩٩٧) :Chemical in the preservation and storage of selected animal food products. In Application of chemistry (I.O, Essien ed.), MEF publication Ltd. Uyo. Nigeria.

Ekanem, E. O. and Achinewhu, S. C. (١٩٩٨) :Effects of spices application on microbiological quality and acceptance of West African clam meat processed by intermediate-moisture technology. Proc. Silver Jubilee Anniversary conference. O.Oduguwa, A.O Fanimu and O.A Osinowo (ed) Nig. Soc. Anim Prod.

Fatmagul MD, Osman I (٢٠١٧) Investigation of the effect of a hybrid drying method on the color quality of nectarine slices and energy consumption. Stud U Babes-Bol Chemia LXII(١):٢٣٧-٢٤٨.

Fatima, M. (٢٠١٣):Evaluation of mushrooms broth cube and its compared with Maggi broth cube products in Saudi Arabia. Journal of American Science, ٢(٥), ٢٥٠-٢٥٥.

Haghi, A. K., & Amanifard, N. (٢٠٠٨) :Analysis of heat and mass transfer during microwave drying of food products. Brazilian Journal of Chemical Engineering, ٢٥(٣), ٤٩١-٥٠١. <http://dx.doi.org/١٠.١٥٩٠/S٠١٠٤-٦٦٣٢٢٠٠٨٠٠٠٣٠٠٠٠٧>

Karšulínová, L., Folprechtová, B., Doležal, M., Dostálová, J., & Velíšek, J. (٢٠٠٧) : Analysis of the lipid fractions of coffee creamers, cream aerosols, and bouillon cubes for their health risk associated constituents. Czech Journal of Food Sciences, ٢٥(٥), ٢٥٧-٢٦٤. [http:// dx.doi.org/١٠,١٧٢٢١/٦٧٩-CJFS](http://dx.doi.org/10.17221/679-CJFS) .

Klurfeld D.M.(٢٠١٥) : Research gaps in evaluating the relationship of meat and health. Meat Sci. ١٠٩:٨٦-٩٥. doi: ١٠,١٠١٦/j.meatsci.٢٠١٥,٠٥,٠٢٢ [\[PubMed\]](#) [\[CrossRef\]](#) [\[Google Scholar\]](#).

Lee, C., Lee, S., & Kim, K. (٢٠١٥) :Use of consumer acceptability as a tool to determine the level of sodium reduction: A case study on beef soup substituted with potassium chloride and soy-sauce odor. Journal of Food Science, ٨٠(١١), S٢٥٧٠-S٢٥٧٦.

Modi V.K., Sachindra N.M., Nagegowda P., Mahendrakar N.S., Narasimha Rao D.(٢٠٠٧) :Quality changes during the storage of dehydrated chicken kebab mix. Int. J.

FoodSciTechnol. ٢٠٠٧;٤٢:٨٢٧-٨٣٥doi: ١٠,١١١١/j.١٣٦٥-٢٦٢١,٢٠٠٧,٠١٢٩١.x [\[CrossRef\]](#) [\[Google Scholar\]](#)

Monteiro, R. L., Carciofi, B. A. M., Marsaioli, A. Jr, & Laurindo, J. B. (٢٠١٥) : How to make a microwave vacuum dryer with turntable. Journal of Food Engineering, ١٦٦, ٢٧٦-٢٨٤. [http://dx.doi.org/١٠,١٠١٦/j.jfoodeng.٢٠١٥,٠٦,٠٢٩](http://dx.doi.org/10.1016/j.jfoodeng.2015.06.029).

Populin T, Moret S, Truant S, Conte LS.(٢٠٠٧) : A survey on the presence of free glutamic acid in foodstuffs, with and without added monosodium glutamate. Food Chem. ١٠٤:١٧١٢-١٧١٧ ٦٢/ Skurray GR, Pucar N. ١٩٨٨. L-glutamic acid content of fresh and processed food. Food Chem. ٢٧:١٧٧-١٨٠.

Radoiu Marilena. (٢٠٢٠) : Microwave drying process scale-up ١٥٥:١٠٨٠٨٨ [Chemical Engineering and Processing - Process Intensification](#) https://www.researchgate.net/publication/3433019376_Microwave_drying_process_scale-up. DOI:[10,١٠١٦/j.cep.٢٠٢٠,١٠,٨٠٨٨](https://doi.org/10.1016/j.cep.2020.108088)

Ran, X. L., Zhang, M., Wang, Y., & Liu, Y. (٢٠١٩) : A comparative study of three drying methods on drying time and physicochemical properties of chicken powder. *Drying Technology*, ٣٧(٣), ٣٧٣- ٣٨٦

<https://doi.org/10.1080/07373937.2018.1450873>, [Google Scholar]

Rahman MS (eds). (٢٠٠٧) : Handbook of food preservation. ٢nd ed. Food science and technology. Boca Raton: CRC Press; ٢٠٠٧.
Rahman.R. (٢٠١٤): Food preservation.

http://en.banglapedia.org/index.php?title=Food_Preservation

Raw Materials Research and Development Council – RMRDC. (٢٠٠٣, December) : Multi-disciplinary committee report of the techno-economic survey of food, beverage and tobacco sector - ٤ Update. Nigeria: Federal Ministry of Science and Technology Abuja.

Rodrigues, É. D. F., Pantoja, L. D. A., Soares, M. B., Nelson, D. L., & Santos, A. S. D. (٢٠١٦) : Development of bouillon cubes from souari nut pulp: formulation and physicochemical and sensorial evaluations. *Brazilian Journal of Food Technology*, ١٩(٠), ١-٨.
<http://dx.doi.org/10.1590/1981-6723.0410>

Rossi, D. M.; Flores, S. H.; Heck, J. X.; Ayub, M. A. Z. (٢٠٠٩) : Production of High-Protein Hydrolysate from Poultry Industry Residue and Their Molecular Profiles. *Food Biotechnol.* ٢٠٠٩, ٢٣, ٢٢٩-٢٤٢. DOI: 10.1080/08905430903102828.

Roudaut, G.; Debeaufort, F. (٢٠١١) : Moisture loss, gain and migration in foods. In *Food and Beverage Stability and Shelf Life*; Kilcast, D., Subramaniam, P., Eds.; Woodhead Publishing: Sawston, UK, ٢٠١١; pp. ٦٣-١٠٥

SAS. (٢٠٠٨) : Static Analysis, ١٥th International Symposium, Valencia, Spain, Proceedings. *Lecture Notes in Computer Science* ٥٠٧٩, ISBN ٩٧٨-٣-٥٤٠-٦٩١٦٣-١

Steel R. and Torrie J. (١٩٨٠): Principles and Procedures of Statistics. ٢nd Ed. New York: McGraw-Hill.

Steele R. Understanding and measuring the shelf-life of food, 1st ed. Woodhead Publishing Limited; ٢٠٠٤.

Watts B., Ylimaki G., Jeffery L., and Elias, L. (١٩٨٩): IDRC-٢٧٧e. Basic sensory methods for food evaluation. Ottawa, Ont., UOC: ٦٦٤,٠٠١,٥:٣٣٩,٤ ISBN: ٠-٨٨٩٣٦-٥٦٣-٦.

Wendy et al ,٢٠١٥ BCCooArticulation Committee. (٢٠١٥) : Basic kitchen and food service Management. Victoria B.C retrieved from <http://opentextbc.ca/basick>.

Yang-Ju Son(٢٠٢٣) : The Physical, Sensory, and Microbial Qualities of Broth Gels Enclosing Food Cubes and Their Changes during Cold Storage [Processes](#) ١١(٨):٢٣٢٧
DOI: [10.3390/pr11082327](https://doi.org/10.3390/pr11082327) License [CC BY](#) ٤,٠.

إنتاج مكعبات المرق من اللحم البقري والدجاج المجفف بالميكروويف ومقارنتها بنظيرتها التجارية

مستخلص البحث:

هدفت هذه الدراسة إلى إنتاج مكعبات مرقة اللحم البقري والدجاج ومقارنتها مع نظيراتها التجارية. يتم تحضيره من الدجاج ولحم البقر المجفف في الميكروويف المنزلي، مع إضافة بعض البهارات الصحية إليه، ومقارنة المنتج المنزلي مع العينات التجارية من مكعبات اللحم البقري والدجاج. كما تم التحقق من التركيب الكيميائي ودراسة التقييم الحسي وكذلك

تأثير مدة الصلاحية على نسبة الرطوبة. وتشير النتائج التي تم الحصول عليها إلى أن جميع العينات محلية الصنع كانت أعلى في القيمة الغذائية مقارنة بالعينات التجارية. بلغت نسبة البروتين في عينات الدجاج ولحم البقر (٥١,٠٤%، ٥٠,٤٦%) على التوالي، وفي العينات التجارية (٧,٣٣%، ٧,٧٥%) على التوالي. وكانت العينات التجارية أعلى في (الرطوبة والرماد والدهون الخام والكربوهيدرات الكلية) من العينات محلية الصنع. كما أشارت النتائج إلى وجود فروق ذات دلالة إحصائية في التقييم الحسي بين العينة المنزلية والتجارية لصالح العينة المنزلية. أشارت النتائج إلى فقدان في نسبة الرطوبة في جميع العينات خلال الفترة المحددة بـ ٣٠ يوما باستثناء مكعبات مرق الدجاج التجارية التي لم تسبب أي فقدان في الرطوبة خلال الفترة المذكورة. وأظهرت أن محصول المنتجات محلية الصنع (١,٢) وكان عدد الحصص ٦ قطع. وأظهرت النتائج التغير اللون للعينات محلية الصنع خلال فترة التخزين من (٠,٩٠) يوم لوحظ تغير اللون في العينات محلية الصنع (اللحم البقري والدجاج) من اللون (البنّي الغامق والبنّي وصولاً للون البنّي الفاتح والاصفر) على التوالي.

الكلمات المفتاحية: الميكروويف، لحم البقر، الدجاج، المرق.