



Studying The possibility of Improving Post-weaning Body Weight Traits of New Zealand White Rabbits Using Different Selection Indices

Shehab El-Din, M. I.¹, M. G. Gharib², Ahmed A. Ramzy¹ and Esteftah M. El-Komy³

¹ Department of Animal Production, Faculty of Agriculture, Al-Azhar University, Cairo, Egypt.

² Animal Production Research Institute (APRI), Agriculture Research Center (ARC), Egypt.

³ Animal Production Department, Agriculture and Biology Research Institute, National Research Centre, El Buhouth St., 12622, Dokki, Cairo, Egypt.

Abstract

THE article aims to improve post-weaning body weight traits of New Zealand white rabbits (NZW). A total of 1,904 progenies dataset of rabbits belonging to a research rabbit farm of the Department of Animal Production, Faculty of Agriculture, Al-Azhar University, for three generations from 2019 to 2022 was used to estimate genetic parameters of body weights and to establish different selection indices with desired gains. Body weights (BW) in grams were measured at 4 (BW4), 6 (BW6), 8 (BW8), 10 (BW10) and 12 (BW12) weeks of age (referred to in the selection indices with X1, X2, X3, X4 and X5, respectively). To analyze the data, a multi-trait animal model was applied. Heritability estimates ranged from 0.12 for BW4 to 0.23 for BW12. The expected genetic gain per generation (ΔG) ranged from 5 to 50.4 g for BW4, 8.1 to 70.4 g for BW6, 13.8 to 98 g for BW8, 22.1 to 78.1 g for BW10 and from -9.47 to 130 g for BW12. To maximize the genetic response to selection, four indices were suggested based on their relative efficiency (RIH): I13: $-8.31 X1 + 5.14 X2 + 1.22 X5$ (RIH = 0.64), I6: $-1.47 X2 + 1.30 X4$ (RIH = 0.59), I17: $-1.894 X2 + 3.976 X3 - 1.343 X5$ (RIH = 0.58) and I10: $0.62 X2 + 0.078 X5$ (RIH = 0.57). Regarding characteristics related to body weight in NZW rabbits it is possible to utilize selected programs aimed at genetic enhancement. However, it is important to gather additional data sources and expand to enhance indices accuracy.

Keywords: Body weight, genetic gain, Rabbits, Selection indices.

Introduction

Under Egyptian conditions, the expected economic indicators for red meat until 2025 indicate an expected increase in consumption, the number of imports and the food gap, accompanied by a decrease in average per capita and an expected decrease in the self-sufficiency rate from 62% in 2020 to 57.5% in 2025 [1]. To face this shortage, attention to animal genetic resources as the basic biological capital for the development of livestock is necessary to achieve food security in the long term, which requires optimal utilization of livestock in all its forms and alternatives and the rapid adoption of a more effective process of genetic improvement to meet the large increase in demand for animal products [2].

In this regard, rabbit breeds represent an important solution in addressing the lack of meat production in developing countries [3], especially the

NZW rabbit breed, which has shown wide adaptation despite the varying conditions of the Egyptian environment or different conditions of care [4,5].

One of rabbit meat production's most crucial economic traits is body weight. Understanding these traits can be a key step in creating suitable genetic selection strategies for rabbits and increasing productivity [6]. To obtain this goal, the selection index is the most efficient tool for the exploitation of available information sources for the effective genetic improvement of several traits simultaneously [7].

The main goals of this research were to: (1) evaluate the genetic and phenotypic factors that affect the body weight performance of NZW rabbits and (2) develop a set of selection indices that can be used to improve body weight traits in NZW rabbits.

*Corresponding author: Esteftah M. El-Komy, E-mail: emelkomy@gmail.com.

(Received 22/04/2024, accepted 19/05/2024)

DOI: 10.21608/EJVS.2024.284394.2025

©2024 National Information and Documentation Center (NIDOC)

Material and Methods

Source of data

In the present investigation, a total of 1,904 progenies dataset of NZW rabbits, fathered by 75 sires and mothered by 209 dams, were maintained at a rabbit research farm for rabbits belonging to the Animal Production Department, Faculty of Agriculture, Al-Azhar University, Cairo, Egypt, from November 2019 until October 2022.

Management and traits considered

Breeding dams and sires were kept apart in hierarchical rabbit batteries with a suitable conventional dimension, arranged in two rows along the farm with a suitable service entrance. The animals were maintained under uniform housing, feeding, medical and other management practices. Three dams were randomly allocated to each sire according to a breeding plan, with the restriction that all forms of inbreeding be avoided. Weaned young rabbits were transferred to conventional progeny wire cages with feeding hoppers and drinking nipples at the age of 4 weeks. They are also ear-tagged, sexed and relocated to groups of 3-4 rabbits per cage. After that, weighing is done every two weeks.

Traits studied

The body weights in grams at 4 (BW4), 6 (BW6), 8 (BW8), 10 (BW10) and 12 (BW12) weeks of age were recorded for each rabbit (referred to in the selection indices with X_1 , X_2 , X_3 , X_4 and X_5 , respectively).

Data analysis

Genetic parameters

The data was analyzed using a multi-trait animal model of body weights using the MTDFREML programs of Boldman *et al.* [8]. Variances computed using the REML approach of the VARCOMP procedure of SAS [9]. were used as starting values for estimating variance components. The general model was used to conduct the analyses:

$$y = Xb + Za + Wpe + e$$

where: y is the vector of phenotypic observations; b is the vector of fixed effects including sex (2 levels); year (3 levels); season (4 levels), parity (5 levels) a is the vector of random additive genetic effects of the progeny; pe is the vector of random permanent environmental effects of the dam; e is the vector of residual effects and X , Z and W incidence matrices that, respectively, link phenotypic observations to fixed and random additive genetic effects.

Selection Indices(I)

Estimates of genetic and phenotypic (co) variances and the relative economic weight (REV) of the studied body weight traits in different

combinations were used to construct twenty-two selection indices according to Hazel *et al.* [10].

REV was calculated as $1/\sigma_p$, where σ_p is the phenotypic standard deviation of each trait [11]. Set the REV of the lowest age as a reference unit for other traits, since under Egyptian conditions, selling at the lowest age as breeding animals has the same profitability as selling at W12 as marketing meat, with a decrease in time and effort in favor of selling at the lowest age.

The correlation between the calculated index and the total aggregate genotype (RIH) was estimated as $RIH = \sigma_i / \sigma_h$. where σ_i is the index standard deviation and σ_h is the genotype standard deviation. To compare indices and decide which traits combine best into an index, relative efficiency (RE) was calculated for each index based on R_{IH} compared to the best index with a high R_{IH} .

The expected genetic gain, gm (ΔG) for each trait after one generation of selection on the index was computed according to Hazel *et al.* [10]. All arithmetic operations performed on matrices were based on the Minitab 17.0 software package (State College, Pennsylvania).

The indices consisted of the full index (I_{22}), which included all sources of information for the five traits, and the reduced indices, which included all possible combinations, from two to four traits (from I_1 to I_{21}).

Results and Discussion

Means

Table 1 summarizes the overall means, standard deviation (SD) and coefficient of variation (CV) for BW traits. The means for BW of NZW rabbits ranged from 804.1 g at 4 weeks of age to 1564.8 g at 12 weeks of age; these means were within the range found by [12-16]. The values of CV for BW at different ages decreased with the advancement of age (Table 1). Comparable results were obtained by [14-17]. This pattern might indicate that the kits become less sensitive to non-genetic maternal effects as they age (in terms of lactation, mothering ability, litter size and weight) which decreases with age [18].

Genetic parameter estimates

Table 2. summarizes the estimates of heritability (h^2a), phenotypic (rp) and genetic (rg) correlations for the body weight traits.

Heritability (h^2a)

Estimates of h^2a for BW in Table 2 ranged from 0.12 at BW4 of age to 0.23 at BW12 of age. Currently, available estimates of moderate h^2a for BW were comparable to those found by [15-17,19]. While the estimates of h^2a for BW traits were lower than those obtained by Farouk *et al.* [15], they were 0.23, 0.24, 0.31 and 0.34 for BW4, BW6, BW8 and

BW10, respectively. The moderate estimates of h^2a for body weight traits of NZW rabbits (Table 2) showed that selection might improve BW in a short amount of time and these features could be employed as selection criteria. According to the current study, h^2a estimates for the studied traits increased with age. Like those obtained by Farouk et al. [15], a study based on 625 NZW rabbits in Egypt found that h^2a estimates for BW at 4, 6, 8 and 10 weeks of age were 0.23, 0.24, 0.31 and 0.34, respectively. Furthermore, Anous [20], indicated that selection for BW at younger ages may be useful for improving early rabbit body weight. Youssef et al. [12] observed increasing h^2a with advancing age in BR and NZW rabbits. Furthermore, advances in age may be accompanied by a decline in non-additive genetic effects, such as common litter effects.

Phenotypic (r_p) and genetic (r_g) correlations

All estimates of phenotypic correlation (r_p) for body weight traits were high and significantly positive and ranged from 0.748 to 0.969 (Table 2). The current findings suggest that a large BW at any body weight would result in a large BW at the final marketing weight (BW12). In the same trend, current estimates concur with those found by [7, 15,16,21], which varied from 0.02 to 0.90. Given that r_p includes the value of both genetic and environmental correlation, this could be the cause of the disparity between the two, since r_p magnitude was higher than r_g . This also suggests that the environmental conditions to which the herd is exposed positively impact showing optimal productivity, particularly for traits with a negative r_g (BW12 in combination with other traits).

From a genetic standpoint, strong positive correlations (r_g) and high (ranging from 0.692 to 0.963) between sequential body weight stages from BW4 to BW12 weeks of age were detected in the current study (Table 2). These results might be explained by the selection for higher BW at weaning (BW4) would be an associated with increase in marketing body weight. The magnitude of r_g between pairs of BW4, BW6, BW8, BW10 and BW12 traits generally reported a wide variation ranging from medium positive to high for the NZW breed [16, 21-24].

Construction of selection indices

Table 3 lists the selection index weighting factors (b), index accuracy (RIH) and efficiency (RE%) of various indices in comparison to the reference index (I_{13}) and the estimated genetic gain (G) for each trait.

In terms of RIH, the triple index (I_{13} : RIH = 0.64) with BW4, BW6 and BW12 included produced the best results, followed by the binary index (I_6 : RIH = 0.59) with both BW6 and BW10. They were followed in the descending order of preference by a binary (I_{17} : RIH = 0.58) and a triple (I_7 : RIH = 57) index, the

similarity of both being that they had both BW6 and BW12. From another point of view, indices not including BW4 and BW6 (I_{12} , I_{18} , I_{11} and I_{12}) showed reduced accuracy of 0.19, 0.27, 0.30 and 0.37, respectively (Table 3). While including BW4 and/or BW6 caused a rise in the accuracy of the index (accuracy ranged from 0.57–0.64), this seems to be due to the higher r_g between them ($rg = 0.818$, Table 2). In this regard, it would therefore be preferable to include BW4 and BW6 in an index that incorporates body weight traits in NZW rabbits.

Consistent with the current results, Khalil et al. [23], El-Fiky et al. [25] and El-Dehadi and Ibrahim [18] reported that selection indices at the early age stage after weaning are promising compared to the late stages. In contrast, Moustafa et al. [26] found that selection based on marketing body weight is preferable to that based on BW4 and daily gain for improving body weight performance.

The expected genetic change per generation (ΔG) ranged from 5 to 50.4 g for BW4, from 8.1 to 70.4 g for BW6, from 13.8 to 98 g for BW8, from 22.1 to 78.1 g for BW10 and from -9.47 to 130 kg for BW12. In this respect, Anous [20] found comparable results using an index that included BW4 and BW6 of age, with ΔG of 57.5 and 1.3 g, respectively.

From an applied point of view, among the top four indices, I_{13} combining both BW4 and BW6 can be proposed for NWZ rabbit commercial farms to improve body weight traits at an early age. Whereas, to improve the final marketing weight, it is recommended to use I_7 .

Conclusion

Considering the findings of the current study, it is possible to use the four selection indices that have been identified as the best ones, which are: I_{13} , I_6 , I_{17} and I_7 . These indices can be used as an effective tool for breeders to achieve rapid genetic improvement of body weight traits to reach the highest possible economic return for such traits. There is a need for more data sources and to broaden the scope of research in both genetic and non-genetic components to increase the accuracy of the indices.

Acknowledgment

Appreciation is extended to the Department of Animal Production, Faculty of Agriculture, Al-Azhar University for their provision of facilities.

Ethical Considerations

The present study has been conducted by the guidelines of the Ethics Committee of the International Animal Care and use Committee (ARC-IACUC) Agricultural Research Center Approval number ARC ABRI 67- 2024.

Conflicts of interest

No Fund.

The authors declare that there is no conflict of interest.

Author's contributions

All Authors collaborate equally.

Funding statement

TABLE 1. Actual means, standard deviations (SD) and coefficients of variation (CV%) for body weight traits in NZW rabbits.

Body weight(g)	Number	Mean	SD	CV (%)
BW4	1904	408.10	92.26	22.60
BW6	1737	670.77	107.43	16.01
BW8	1627	962.80	128.51	13.34
BW10	1571	1260.44	133.73	10.61
BW12	1564	1554.80	145.73	9.37

BW4, BW6, BW8, BW10 and BW12= body weights at 4, 6, 8, 10 and 12 weeks of age.

TABLE 2. Estimates of heritability \pm standard error (SE) (bold on diagonal), genetic \pm SE (below diagonal) and phenotypic correlations (above diagonal) among post-weaning growth traits in NZW rabbits.

	BW4	BW6	BW8	BW10	BW12
BW4	0.12 \pm 0.02	0.868	0.798	0.804	0.748
BW6	0.818 \pm 0.07	0.14 \pm 0.05	0.879	0.863	0.808
BW8	0.731 \pm 0.06	0.799 \pm 0.05	0.18 \pm 0.06	0.935	0.893
BW10	0.727 \pm 0.05	0.768 \pm 0.06	0.884 \pm 0.04	0.21 \pm 0.03	0.969
BW12	0.692 \pm 0.06	0.712 \pm 0.07	0.828 \pm 0.06	0.963 \pm 0.03	0.23 \pm 0.06

BW4, BW6, BW8, BW10 and BW12= body weights at 4, 6, 8, 10 and 12 weeks

TABLE 3. Weighing factors (*b* values), the efficiency of the index (RIH), the relative efficiency (RE) of different indices and the expected genetic gain per generation (Δ G) for body weight traits in NZW rabbits.

Index no.	<i>b</i> values	RIH	Re%	Expected genetic gain, gm (Δ G)				
				BW4	BW6	BW8	BW10	BW12
<u>Binary indices</u>								
I ₁	0.126 X ₁ + 0.139 X ₂	0.38	59.2	5	8.1	-	-	-
I ₂	1.047 X ₁ - 0.514 X ₃	0.19	30.3	18	-	13.8	-	-
I ₃	0.013 X ₁ + 0.282 X ₄	0.49	76.1	30	-	-	53.8	-
I ₄	0.488 X ₁ + 0.006 X ₅	0.42	65.2	43.5	-	-	-	84.2
I ₅	0.112 X ₂ + 0.121 X ₃	0.36	57.1	-	16.3	40	-	-
I ₆	-1.47 X ₂ + 1.300 X ₄	0.59	91.6	-	37.6	-	70.2	-
I ₇	0.620 X ₂ + 0.078 X ₅	0.57	88.5	-	34	-	-	130
I ₈	2.26 X ₃ - 1.570 X ₄	0.27	41.9	-	-	98	48	-
I ₉	0.11 X ₃ - 0.22 X ₅	0.46	70.8	-	-	19.7	63.7	-
I ₁₀	1.69 X ₄ - 1.08 X ₅	0.38	59.8	-	-	-	46.7	70.8
<u>Triple indices</u>								
I ₁₁	-0.019 X ₁ + 1.49 X ₂ - 0.880 X ₃	0.30	46.6	8	19.4	39	-	-
I ₁₂	1.35 X ₁ - 0.892 X ₂ + 0.135 X ₄	0.37	57.5	16	24.2	-	63.3	-
I ₁₃ *	-8.31 X ₁ + 5.14 X ₂ + 1.22 X ₅	0.64	100.0	83	70.4	-	-	-9.47
I ₁₄	-0.639 X ₁ + 0.598 X ₃ + 0.234 X ₅	0.45	69.9	50.4	-	69.4	-	116.5
I ₁₅	0.761 X ₁ - 0.0544 X ₄ + 0.463 X ₅	0.46	71.4	13.5	-	-	22.1	32.9
I ₁₆	1.75 X ₂ - 0.952 X ₃ - 0.0224 X ₄	0.37	58.7	-	16.8	31.4	56.9	-
I ₁₇	-1.894 X ₂ + 3.976 X ₃ - 1.343 X ₅	0.58	90.2	-	53.1	17	-	114.2
I ₁₈	-0.530 X ₃ + 3.321 X ₄ - 1.900 X ₅	0.43	66.8	-	-	21.1	78.1	125
<u>Quad indices</u>								
I ₁₉	-0.262 X ₁ + 1.30 X ₂ - 0.277 X ₃ - 0.0901 X ₄	0.36	56.4	15.1	21.9	32.9	52.4	-
I ₂₀	1.198 X ₁ - 0.164 X ₂ - 0.0695 X ₃ - 0.017 X ₅	0.40	62.6	14	16.8	21.5	-	40.3
I ₂₁	1.76 X ₂ + 3.16 X ₃ - 6.82 X ₄ + 2.95 X ₅	0.38	59.0	-	23.3	25.3	35.5	45.7
<u>Full index</u>								
I ₂₂	1.584 X ₁ + 0.0735 X ₂ - 0.749 X ₃ - 0.232 X ₄ + 0.435 X ₅	0.18	29.3	19.5	24.6	36.1	54.8	83.5

X₁=BW4; X₂=BW6; X₃=BW8; X₄=BW10 and X₅ = BW12; Reference index for calculating relative efficiency (RE)

References

1. El-Shaheed, A., Emad, M. and Swefy, A.A. The economic dimension of the food gap and food security of red meat in Egypt. *Egyptian Journal of Agricultural Research*, **100**(1), 96-106 (2022). DOI: 10.21608/EJAR.2022.105571.1170
2. Gitz, V., Meybeck, A., Lipper, L., Young, C. D. and Braatz, S. Climate change and food security: risks and responses. Food and Agriculture Organization of the United Nations (FAO) Report, **110**(1), 2-4 (2016). DOI 9789251089989
3. Mousa-Balabel, T.M. Effect of heat stress on New Zealand White rabbits behavior and performance. *Minufiya Veterinary Journal*, **3**(1), 125-134 (2004). DOI:10.21608/bvmj.2018.38761
4. Khalil, M.H. and Baselga, M. The Giza White Rabbits (Egypt). *Options Méditerranéennes: Série B. Etudes et Recherches*, **38** (1), 27-36 (2002).
5. Shehab El-Din, M.I., Ahmed, R.A. and El-Komy, E. M. Assessing the Genetic Response of Crossing on Post-Weaning Growth Traits in Rabbits. *Egyptian Journal of Veterinary Sciences*, **55**(5), 1327-1335 (2024).
6. Suárez-López, J.R., Becerril-Pérez, C.M., Pro-Martínez, A., Cuca-García, J.M., Torres-Hernández, G. and Sosa-Montes, E. Productive performance of six genotypes of fattening rabbits fed with seven commercial feeds. 9th World Rabbit Congress, Verona, Italy 10-13 Jun (2008).
7. Choudhury, S.A. and Goswami, R.N. Selection indices for improving body weight in New Zealand White breed of rabbit maintained at the hilly Indian State of Meghalaya. In Proc.: 10th World Rabbit Congress, 3-6 September, Sharm El- Sheikh, Egypt, 25-29 (2012). DOI: 10.21608/EPSJ.2014.32587.
8. Boldman, K.G., Kriese, L.A., Van Vleck, L.D., Van Tassell, C.P. and Kachman, S.D. A manual for the use of MTDFREML. Agricultural Research Service, USDA, Clay Center, NE, USA (1995).
9. SAS. Statistical analysis system. SAS Release 9.1.3. For windows, *SAS Institute Inc. Cary, NC, USA* (2003).
10. Hazel, L.N., Dickerson, G.E. and Freeman, A.E. The selection index-Then, now, and for the future. *Journal of Dairy Science*, **77**(10), 3236-3251 (1994). [https://doi.org/10.3168/jds.S0022-0302\(94\)77265-9](https://doi.org/10.3168/jds.S0022-0302(94)77265-9)
11. Falconer, D.S. and Mackay, T.F.C. Introduction to quantitative genetics. *Essex. UK: Longman Group* (1996).
12. Youssef, M.K., Farid, A., Gad-Alla, S.A. and Abo-Warda, M.A. Genetic evaluation for post weaning body weight traits in three genetic groups of rabbits under Egyptian conditions. *5th International Poultry Conference, Taba Egypt*, 10-13 March (2009).
13. Nowier, A.A. Using of different genetic techniques to improve some economic traits in animal. *Ph.D. Thesis, Fac. of Agric., Tanta Univ., Egypt* (2012).
14. Gouda, G.F. Consequences of selection for post-weaning growth performance traits on fat partition traits in rabbits. *International Journal of Veterinary Science*, **11**(4), 461-466 (2022). DOI:10.47278/journal.ijvs/2022.136
15. Farouk, S.M., Khattab, A.S., Noweir, A. and Ghavi Hossein-Zadeh, N. Genetic analysis of some productive and reproductive traits in New Zealand White rabbits. *World Rabbit Science*, **30**(2), 141-146 (2022). DOI: <https://doi.org/10.4995/wrs.2022.15939>
16. El-Deghadi, A.S. Yonan G.E., El-Naser, M.I. and Seif El-Naser, M.G. Gharib Genetic Analysis of Productive Performance in New Zealand White Rabbits. *International Journal of Life Science and Agriculture Research*, **2**(6), 95-101 (2023). DOI:<https://doi.org/10.55677/ijlsar/V02I06Y2023-01>
17. El-Deghadi, A.S. and Ibrahim, M.K. Selection indices for improving body weight in Gabali rabbits. *Egyptian Poultry Science Journal*, **38**(4), 1115-1126 (2018). <https://doi.org/10.21608/epsj.2018.22904>.
18. Khalil, M., Owen, J. and Afifi, E. Selection indices for rabbit improvement. *The Journal of Agricultural Science*, **107**(3), 537-548 (1986). <https://doi.org/10.1017/S0021859600069690>
19. Soliman, H.A.M., Gad, S.M.A. and Ismail, I.M. Genetic parameters for post-weaning growth traits of Gabali rabbit in Egypt. *Egyptian Poultry Science Journal*, **34** (2), 655- 664 (2014).
20. Anous, R.M. Selection indexes for improved weaning and marketing body weights of the Burundi local rabbits. *Archives Animal Breeding*, **44**(5) 561-567 (2001). <https://aab.copernicus.org/articles/44/561/2001/>
21. Rabie, T., Nower, A., Abou Zeid, A. and Khattab, A.S. Assessment of genetic capability for post-weaning growth traits of reciprocal cross between Gabali and V- Line rabbits by animal model. *World's Veterinary Journal*, **10**(3), 405-413 (2020). <https://www.scopus.com/inward/record.uri?eid=2-s2>
22. Iraqi, M.M. Estimation and evaluation of genetic parameters for body weight traits of New Zealand White rabbits in Egypt using different multivariate

- animal models. *Livestock Research for Rural Development*. **15**(1), 6-12 (2003).
http://www.cipav.org.co/lrrd/lrrd15/6/iraq156.htm
23. El-Deighadi, A. and Ibrahim, I.K. Genetic aspects of post-weaning for growth traits in New Zealand white rabbits. *Egyptian Journal of Rabbit Science*, **27**(2), 507-521 (2017).
24. Gouda, G.F. Correlated response in carcass traits when using selection indexes involving marketing body weight and daily gain in New Zealand white rabbits. *Egyptian Journal of Animal Production*, **45**(1), 45-54 (2008).
25. El-Fiky, F.A., Aboul-Hassan, M.A., Attalah, G. E.Y. and Bata, S.S. Selection indices for improving body weight in Baladi rabbit. *Egyptian Poultry Science*, **21**(2), 305-318 (2001).
26. Moustafa, H., El-Raffa, A., Shebl, M.K., El-Delebs hany, A. and ElSayed, N.A. Genetic evaluation of some economic traits in a maternal line of rabbits. *Egyptian Poultry Science Journal*, **34**(1), 85-98 (2014).
https://doi.org/10.21608/epsj.2014.5308.

دراسة إمكانية تحسين صفات وزن الجسم بعد الفطام في الأرانب النيوزلندي الأبيض باستخدام أدلة انتخابية مختلفة

محمد إبراهيم شهاب الدين¹، ومحمود غريب غريب²، أحمد رمزي عبد الله³ وإستفتاح محمد الكومي³

¹قسم الإنتاج الحيواني - كلية الزراعة - جامعة الأزهر - القاهرة - مصر.

²معهد بحوث الإنتاج الحيواني - مركز البحوث الزراعية - مصر.

³قسم الانتاج الحيواني- معهد البحوث الزراعية والبيولوجية - المركز القومي للبحوث - الجيزة - مصر.

الملخص

يهدف البحث إلى تحسين صفات وزن الجسم بعد الفطام للأرانب من سلالة النيوزلندي الأبيض. تم استخدام إجمالي بيانات وزنية 1904 مجموعة من السجلات الانتاجية لمزرعة الارانب البحثية التابعة لقسم الإنتاج الحيواني بكلية الزراعة جامعة الأزهر بالقاهرة، في الفترة من 2019 إلى 2022 لتقدير المقاييس الوراثية لأوزان الجسم وإنشاء مجموعة مختلفة من الأدلة الانتخابية لتقدير العوائد الوراثية المتوقعة لكل دليل. تم قياس أوزان الجسم بالجرام عند 4، 6، 8، 10 و 12 أسبوع من العمر وتحليل البيانات، تم تطبيق النموذج الحيواني متعدد الصفات بطريقة معظمة الاحتمال غير المقيدة وغير المعتمدة على حساب المشتقات التفاضلية وقد اشتمل النموذج الإحصائي على تأثير سنة و موسم الولادة وترتيب البطن وجنس الارنب كعوامل ثابتة كما شمل تأثير الحيوان والتأثير البيئي الدائم كعوامل عشوائية. وتم تكوين أدلة الانتخاب لوزن الجسم عند 4، 6، 8، 10 و 12 أسبوع. وتراوح وزن الجسم بين 408.1 جم لوزن الجسم عند 4 أسابيع إلى 1554.8 جم لوزن الجسم عند 12 أسبوع. تراوحت قيم المكافئ الوراثي من 0.12 لوزن الجسم عند 4 أسابيع إلى 0.23 لوزن الجسم عند الأسبوع 12. تراوح العائد الوراثي المتوقع لكل جيل (ΔG) من 5 إلى 50.4 جم لوزن الجسم عند 4 اسابيع، ومن 8.1 إلى 70.4 جم لوزن الجسم عند 6 اسابيع، ومن 13.8 إلى 98 جم لوزن الجسم عند 8 اسابيع، ومن 22.1 إلى 78.1 جم لوزن الجسم عند 10 اسابيع، ومن - 9.47 إلى 130 كجم لوزن الجسم عند 12 أسبوع، لتعظيم الاستجابة الوراثية للانتخاب، تم اختيار أربعة ادلة انتخابية بناءً على كفاءتها النسبية (RIH) وهي I_{13} ، I_6 ، I_{17} ، و I_{10} وتراوحت كفاءتهم النسبية بين 64% وحتى 0.57%. ولذا و فيما يتعلق بصفات وزن الجسم بعد الفطام للأرانب من سلالة النيوزلندي الأبيض، يمكن استخدام الانتخاب كأداة للتحسين الوراثي مع الأخذ في الاعتبار الحاجة إلى المزيد من مصادر البيانات وتوسيع نطاقها لزيادة دقة الأدلة الانتخابية.

كلمات مفتاحية: وزن الجسم، العائد الوراثي، الأرانب، الأدلة الانتخابية.