

POSSIBLE EFFECTS OF SUPPLEMENTATION WITH NIGELLA SATIVA CAKE ON REPRODUCTIVE PERFORMANCE, OVARIAN RESPONSE, AND EMBRYO RECOVERY OF FEMALE BALADI GOATS

S. A. BADAWY* ; G. M. DARWISH** and G.M. RAKHA***

* Dep. Anim. Rep. And Artif. Insemin., National Research Center, Dokki, Cairo, Egypt.

** Dept. AI and ET. Animal Reproduction Research, Institute, ARC, Giza, Egypt.

*** Dept. of Medicin, Fac. Vet. Med. Cairo Univ., Egypt.

Received: 15.5.2001.

Accepted: 29.7.2001.

SUMMARY

This study is one of many recent attempts that search for economic feed resources with the aim to improve reproductive efficiency of the local breeds of farm animals in Egypt. Thirty female baladi goats were used in two experiments. In experiment 1, twelve immature females weighed 8-11 kg and aged 4-5 months were allotted into two groups: control (n = 4) fed on basal control ration and NSc group (n = 8) fed on the same basal ration supplemented with 100gm nigella sativa cake. All females were fed till maturity and run continuously with fertile bucks. Jugular blood samples were collected twice weekly for progesterone assay to detect sexual maturity, ovarian cyclicity, conception and pregnancy. The parameters

of the reproductive characteristics in the pattern of continuous/total, lambing/continuous, conception, fertility, and prolificacy were calculated. Initial and final body weights were individually recorded and body weight gain was calculated. Five months postfeeding, three females from each group were slaughtered and blood samples were collected for determination of plasma protein and lipid indicators. Samples of rumen contents were taken for determination of pH, ammonia, volatile fatty acids (total and individual) and protozoal activity. Right and left ovaries were desiccated immediately after slaughtering and ovarian weights and structures were recorded. In experiment 2, eighteen mature female baladi goats of 1.5-2 years were allocated into two groups: control (n=8) and NSc (n=10) receiving

the same rations as in the experiment 1. The oestrus was synchronized by withdrawing progestagen-impregnated pessaries 14 days after their intravaginal insertion. Two days prior to pessaries removal, each doe received 800-1000 i.u. equine Chorionic Gonadotrophin. Twenty four hours, does were observed for heat, then naturally mated with fertile buck for 2-3 times during the estrus period. Assessment of the superovulatory response was carried out on day 6 following estrus. Both ovaries were examined morphologically and the number of corpora lutea and follicles >5 were recorded. The uterine horns were flushed and the recovered embryos / ova were counted, evaluated, and classified according to their stage and morphology. Plasma total protein, albumin, total lipids, and triglycerides were not significantly changed. However, plasma urea and cholesterol were significantly decreased on supplementation with *Nigella sativa* cake. Analysis of rumen juice revealed that the total volatile fatty acids, the molar proportion of propionate were increased, while the molar proportion of acetate, and accordingly the A/P ratio were decreased in (NSc) group. However, the values of pH, ammonia, the molar proportion of butyrate and the protozoal activity were not affected by the addition of *Nigella sativa* cake. There was a marked increase in the reproductive indexes of does supplemented with *Nigella sativa* cake, where the percentages of conception, fertility and prolificacy were 37.5, 37.5, and 100 % versus Zero in control does. The final body weight was increased significantly in

does of (NSc) group 15.04 ± 0.82 Kg than controls. Consequently the weight gain is significantly higher in (NSc) groups than control. The number of corpora lutea was significantly increase in (NSc) group than the control one. Similarly, the total ovarian responses (TOR) was also significantly increased in (NSc) group than the controls. However, the ovulation rates among the groups were not significantly different. The total percentage of transferable embryos was significantly higher in (NSc) group compared to control one. Meanwhile the percentage of non-transferable embryos (Fair) was significantly ($P < 0.001$) lower in treated group (38.98 %) compared to control one (80.00 %). In addition the recovery rate for transferable embryos was significantly higher in treated group (28.57 %) than in control one (11.76 %). The total embryo recovered was not significantly affected by dietary treatment

Key wards: *Nigella sativa* , goats , Reproductive performance , blood and rumen profiles

INTRODUCTION

Energy availability influences reproduction fitness. Changes in an animal's energetic status can result in modulation of the hypothalamic-pituitary - gonadal axis (Cunningham et al., 1999). Suppression of pulsatile LH secretion has been documented after fasting or feed restriction in myriad species, including sheep and goats (Thomas et al.,

1990). In addition to its effect on reproductive hormonal profiles, feed restriction can delay the timing of puberty onset (Foster and Olster, 1985 and Hall et al., 1990) and directly affect the other reproductive processes (Dickerman et al., 1993 and Gill and Rissmar, 1997).

Many factors affect the ovarian response to superovulatory treatments, recovery rate and embryo quality. The level of dietary energy pre and postmating play an important role in ovarian response to gonadotrophins. Mani et al. (1992 & 1993) concluded that low level of feeding in goats pre and postmating resulted in the delay and suppression of estrus, reduced ovulation rate, decreased incidence of multiple ovulation and increased loss of embryos.

The mechanisms by which ovulation and ovulation rate are modulated by nutrition is a matter of speculation. Increased nutritional levels may result in increased hepatic mixed function oxidase activity, thereby increasing the rate of degradation of steroids and lessening the inhibition of the hypothalamic-pituitary axis and resulting in increased gonadotrophin secretion (Kirkwood and Aherne, 1985 and Thomas and Williams, 1996)

Poor reproductive efficiency of local breeds of farm animals in Egypt is mainly due to nutritional factors. Large shortage in animal's feedstuffs obligate workers to search for new cheaper uncommon feed sources to fulfill the nutrients require-

ments and energy demands.

Nigella sativa seed is one of the most important medicinal plants. After oil extraction, considerable amounts of Nigella sativa cake become available. This cake is a suitable meal to be included in the diets of farm animals as it contains some remained oil and more than 30% protein (Abdel-Aal and Attia, 1993; Khalifah, 1995; and Zeweil, 1996).

Previous studies investigated the effects of feeding diets containing Nigella sativa cake on productive performance of sheep (Awadalla, 1997 and Gabr et al., 1998), buffaloes (Youssef et al., 1998), and poultry (El-Ghamry et al., 1997). They recorded improved feed conversion, quantity and quality of produced meat, and increased economic efficiency.

These beneficial positive effects make it important to fully investigate the possible effects of Nigella sativa cake on reproductive potentials of farm animals. The present study was carried out with the aim to clarify the effect of Nigella sativa cake as a supplement on: (1) Plasma and Rumens metabolic profiles, and subsequent reproductive performance of growing female baladi goats (2) superovulatory responses of baladi does treated with equine Chorionic Gonadotrophin (eCG) with special focus on ovarian responses and embryo quality.

MATERIALS AND METHODS

A total number of thirty female baladi goats were used in two experiments, carried out at April in the experimental farm of National Research Center, in Abi-Rawash.

Experiment 1

Twelve immature females aged 4-5 months and weighed 8-12 kg were randomly allotted into two groups, control (N=4) fed on 750 g maintenance ration /head daily recommended by NRC (1989), and treated one (N=8) given the same basal ration supplemented with 100 g / animal of *Nigella sativa* cake. The chemical composition, ingredients, and daily intake of experimental rations are illustrated in table 1. All females were fed in an open shed till maturity and run continuously with the bucks. Jugular blood samples were collected twice weekly for progesterone assay to detect sexual maturity and ensure ovarian cyclicity, conception and pregnancy (Continuous P4 profile > 1 ng/ml over at least 25 days post mating as recorded by Pope et al., 1989). The parameters of the reproductive performance and frequency of cyclicity in the pattern of continuous/total, lambed/continuous, conception , fertility, and prolificacy were calculated. Initial and final body weights of all animals were individually recorded and body weight change (gain) was calculated. Five months postfeeding, three females from each group were slaughtered and blood samples were collected for determination of plasma protein and lipid indica-

tors. Samples of rumen contents were taken for determination of pH value, ammonia production, volatile fatty acids (total and individual) and protozoal activity. Right and left ovaries were desiccated immediately after slaughtering and ovarian weights and structures were recorded.

Experiment 2

Eighteen mature female baladi goats of 1.5-2 years were allocated into two groups and received the same rations as recorded in the experiment 1. The oestrus symptoms and duration of oestrus were recorded for all does. Synchronization of oestrus was achieved by withdrawing progestagen-impregnated pessaries (Containing 60 mg Medroxy progesterone acetate, veramix, Upjohn, USA) 14 days after their intravaginal insertion. Two days prior to pessaries removal, each doe received 800-1000 i.u. (eCG) equine chorionic gonadotrophin (Folligon, Intervet, Holland). Twenty four hours, does were observed for heat, then naturally mated with fertile buck for 2-3 times during the estrus period. Assessment of the superovulatory response was carried out at laparotomy on day 6 following estrus (McKelvey et al., 1989). The animals were generally anaesthetized with xylazine (Rompn, Bayer), 0.2 mg/kg intramuscularly and followed by injection of sodium thiopental 15-20 mg/kg bw (Dorn and Kraemer, 1987). Both ovaries were examined morphologically and the number of corpora lutea (Functional and Regressed) and follicles >5 were recorded. The uterine horns were flushed with 40

ml Dulbecco's phosphate buffer saline (PBS) enriched with 2% Bovine Serum Albumin (BSA) using silicon two way folly catheter. The recovered embryos / ova were counted, evaluated, and classified according to their stage and morphology under stereomicroscope.

Data Collection techniques

Plasma levels of total protein, albumin, urea, triglycerides, and cholesterol were determined using Stanbio kits (USA). Determination of plasma total lipids was performed using kits of CDI (Cal-Tech Diagnostics, INC, USA). Plasma Progesterone was measured in all samples (RIA) by the Coat-A-Count technique using kits of DPC (Diagnostic Products Corporation, Los Anglos, USA). Several parameters of rumen metabolic profile were determined as recorded by Badawy (1992). Reproductive efficiency indexes including conception rate (no. of does lambed/no. of does exposed) x 100, fertility percentage (no. of lambs born/no. of does exposed) x 100, and prolificacy percentage (no. of lambs born/ no. of does lambed) x 100 were calculated according to Doroliya et al. (1990)

Statistical analysis

One way analysis of variance (ANOVA) was used for comparing means of rumen metabolic profile, body weight, ovarian weight, and ovarian responses to superovulation in both control and treated groups according to steel and Torrie (1980) using the linear models procedure avail-

ble in SAS software (SAS, 1986). Meanwhile the embryo recovery data was analysed using Microstate computer program, copyright (C) 1984 Escosoft, Inc., USA.

RESULTS

Experiment 1

The effect of the *Nigella sativa* cake supplementation on some plasma biochemical parameters are represented in Table 2. Plasma total protein and albumin are not significantly changed. However, plasma urea is significantly decreased in (NSc) group than controls. At the same time plasma total lipids and triglycerides are not affected, while plasma cholesterol is significantly decreased on supplementation with *Nigella sativa* cake.

Rumen metabolic profile of baladi goats supplemented with *Nigella sativa* cake is represented in Table 3. Analysis of rumen juice revealed that the total volatile fatty acids (TVFAs) is significantly increased in (NSc) group than control, while the values of pH and ammonia are not changed. Supplementation with *Nigella sativa* cake increase the molar proportion of propionate, decrease the molar proportion of acetate, and accordingly decrease the A/P ratio, but not affect the molar proportion of butyrate. The protozoal activity in the pattern of number, motility and viability are not affected by the addition of *Nigella sativa* cake.

The effect of *Nigella sativa* cake supplementation on the reproductive performance and cyclicity of female baladi goats during sexual maturity are presented in Table (4). Data of the reproductive characteristics accompanied with the progesterone profile indicated that the number and percentage of does that showed regular changes in progesterone concentration coincide with the ovarian activity, where three does of (NSc) group showed estrus behaviour and accepted the male. However, continuous basal progesterone level (below 1ng/ml) of control does indicated ovarian inactivity. These results were confirmed by the gross weight and appearance of the ovaries obtained from slaughtered females, where there is a significant increase in the ovarian weights and activities in (NSc) group compared to control. Moreover, percentage of does showed normal and continuous cycling is 75 % in (NSc) group and 50 % in control one.

There is a marked increase in the reproductive indexes of does supplemented with *Nigella sativa* cake, where the percentages of conception, fertility and prolificacy were 37.5, 37.5, and 100% versus Zero in control does (Table 4). Concerning the body weight gain the initial body weight is not significantly different between groups. However the final body weight is increased significantly in does of (NSc) group 15.04 ± 0.82 Kg than controls (12.5 ± 0.50 Kg). Consequently the weight gain is significantly higher in (NSc) groups than

control (4.58 ± 0.8 and 1.50 ± 0.3 Kg respectively).

Experiment 2

The data represented in the Table 5. revealed that the number of unovulated follicles was higher but not significantly different by feeding regimens ($P < 0.05$) between (NSc) and control groups (9.56 ± 2.43 and 3.86 ± 1.61 resp.). The number of corpora lutea was significantly increase ($P < 0.05$) in (NSc) group (14.00 ± 2.43) than the control one (4.86 ± 1.88). Similarly, the total ovarian responses (TOR) was significantly ($P < 0.01$) increased in (NSc) group than the controls (23.56 ± 2.91 and 8.86 ± 2.89 resp.), however, the ovulation rates among the groups were not significantly different.

The total percentage of transferable embryos (Table 6) was significantly ($P < 0.05$) higher in (NSc) group compared to control one (61.02 and 20.00 % resp.). Meanwhile the percentage of non-transferable embryos (Fair) was significantly ($P < 0.001$) lower in treated group (38.98 %) compared to control one (80.00 %). In addition the recovery rate for transferable embryos was significantly ($P < 0.001$) higher in treated group (28.57 %) than in control one (11.76 %). The total embryo recovered was not significantly affected by dietary treatment. Although the (NSc) group had lowest recovery rate (46.83 %), it had highest

percentage of transferable embryos (61.02 %) and lowest percentage of fair embryos (38.98 %). On the contrary, the control group had highest recov-

ery rate (58 %) but it had highest percentage of fair embryos (80 %) and lowest percentage of transferable embryos (20%).

Table 1: Ingredients (%), chemical composition, and daily intake of basal (control) and Nigella sativa cake supplemented (NSc) rations.

Items	Control ration	(NSc) ration
Ingredient (%)		
Conc. Mix	42.86	40.54
Wheat straw	57.14	54.06
Nigella Sativa oil seed meal	----	5.40
Total	100	100
Chemical compositions		
(DN (%))	92.30	91.95
Crude protein (%)	8.47	9.34
TDN (%)	52.86	54.06
EE (%)	1.470	2.299
Daily intake/head/day		
DM intake (g)	1560	1660
Crude Protein (g)	132.13	155.04
TDN (Kg)	82.46	87.74
EE	22.93	38.16

Table 2: Plasma biochemical parameters of Baladi does in control (C) and Nigella sativa (NSc) groups.

Variables	Treatments	
	C	NSc
Total protein	8.85 ± 0.30	8.78 ± 0.15
Albumin	6.54 ± 0.19	6.21 ± 0.12
Urea	22.1 ± 3.1**	13.4 ± 1.3
Total lipids	386.5 ± 11.07	378.1 ± 6.1
Cholesterol	145.3 ± 4.5**	127.6 ± 4.2
Triglycerides	99.6 ± 5.5	99.8 ± 6.6

** Significant at P < 0.05.

Table 3: Ruminal metabolic profile of Baladi goats in control (C) and Mige La saliva supplemented (MSc) groups.

Group	PH	Ammonia (mg/dL)	TVFAS (mg/dL)	Individual VFAs (%)						Protozoal Activity		
				Acetic	Propionic	A/P ratio	Butyric	Isovaleric	Number	Mortality	Viability	
C	6.45 ± 0.40	14.35 ± 1.48	9.13 ± 5.16	48.73** ± 7.52	34.04 ± 3.16	1.43** ± 0.12	14.78 ± 2.11	2.45 ± 0.15	+++	+++	75%	
MSc	6.23 ± 0.41	12.55 ± 1.76	11.57 ± 7.53	34.17 ± 5.89	42.27** ± 4.47	0.80 ± 0.15	20.99 ± 2.75	2.57 ± 0.11	+++	+++	80%	

** P < 0.01

* P < 0.05

Table 4: Subsequent reproductive performance and Frequencies of Cyclicity of Baladi goats in control (C) and Nigella sativa (NSc) groups.

Variables	Treatments	
	C	NSc
No. of does exposed	4	8
Continuous/Total ¹	2/4 (50%)	6/8 (75%)
No. of does lambded	Zero	3
Lambded/continuous ²	Zero	3/6 (50%)
Conception (%) ³	Zero	37.5
Fertility (%) ⁴	Zero	37.5
Prolificacy (%) ⁵	Zero	100
Ovarian weight (g)		
Right ovary	0.30 ± 0.06	0.60 ± 0.05
Left ovary	0.23 ± 0.03	0.51 ± 0.04
Ovarian structures	Small smooth ovaries, however small follicles were seen	Multifollicles (3-4 large and medium sized follicles)
Body wieght		
Initial B wt. (Kg)	11.0 ± 0.76	10.46 ± 0.59
Final B wt (kg)	12.5 ± 0.50	15.04 ± 0.82**
B wt change	1.50 ± 0.3	4.58 ± 0.8**

1: No. of does with continuous cycling/total No. of does.

2: No. of does lambded/No. of does with continuous cycling

3: No. of does lambded/No. of does exposed.

4: No. of lambded born/No. of does exposed.

5: No. of lambded born/No. of does lambded.

** Signivicant at P<0.05.

Table 5: Ovarian responses in superovulated Balady does in control and Nigella Sativa cake supplemented (NSc) rations.

Animal group	No. of does	Unovulated follicle	Corpora lutea (CL)	Total Ovarian Responses (TOR)	CL/TOR
NSc	9	9.56 ± 2.43 ^a	14.00 ± 2.43 ^a	23.56 ± 2.91 ^a	0.58 ± 0.09 ^a
Control	7	3.86 ± 1.61 ^a	4.86 ± 1.88 ^b	8.86 ± 2.89 ^b	0.55 ± 0.16 ^a
Ttoal	16	7.06 ± 1.66	10.00 ± 1.94	17.13 ± 2.75	0.57 ± 0.08

a,b: Values having the same letters in the same column are not significantly different

Table 6: Embryo quality and quantity in superovulated Balady does in Control and Nigella Sativa cake supplemented (NSc) rations.

Animal group	Transferable embryo stages No. (%)			Recovery rate for transferable embryo (%)	Non-transferable embryo No. (%) (Fair)	Total embryo recovered	Recovery rate* (%)
	Excellent	Good	Total				
NSc	24 (40.68) ^a	12 (20.34) ^a	36 (16.02) ^a	28.57 ^a	23 (38.98) ^a	59	46.83 ^a
Control	3 (15.00) ^b	1 (5.00) ^b	4 (20.00) ^b	11.76 ^b	16 (80.00) ^b	20	58.82 ^a
Ttoal	27 (34.18)	13 (16.46)	40 (50.63)	25.00	39 (49.37)	79	49.38
P value	P<0.02	P<0.05	P<0.001	P<0.02	P<0.001		P<0.12

a,b: Values having the same letters in the same column are not significantly different

* Recovery rate = Number of recovered embryos/number of CL.

DISCUSSION

Experiment 1

Feeding of *Nigella sativa* cake to Baladi goats had no effect on plasma total proteins and albumin. This result agrees with those obtained by El-Ekhnawy et al. (1999). Other investigators (Abdel Aal and Attia, 1993, Khodary et al., 1996, Nassar, 1997, and Youssef et al., 1998) used *Nigella sativa* seeds and found a significant increase in the plasma total proteins, which might be due to the increase in the globulins level. Otherwise, Nassar (1997) reported no significant changes in plasma albumin level on feeding of different forms of *Nigella sativa* in the ration of baladi cockerels. Plasma urea showed no significant difference between groups. Dissimilar result was obtained by El-Ekhnawy et al. (1999) who reported an increase in plasma urea with the supplementation of 150-250 g *Nigella sativa* cake. The present data showed a significant decrease in plasma cholesterol level, which was previously recorded by El-Dakhkhny et al. (1997), which might be attributed to the high content of unsaturated fatty acids, mainly linolinic (Barowicz et al., 1997)

Rumen juice analysis revealed that TVFAs had a significant increase in (NSc) group, which could be attributed to the high organic matter content (90.5 %) of *Nigella sativa* cake (Youssef et al., 1998). The low ammonia level due to *Nigella sativa* cake supplementation indicated slow release

of protein degradation in spite of the relative high level of crude protein content (29.3%) of *Nigella sativa* cake (Gabr et al., 1998).

The present study revealed that the most consistent action of *Nigella sativa* cake on rumen fermentation is its ability to increase the molar proportion of propionate at the expense of acetate, and thus theoretically increased the efficiency of converting feed energy available to the animal. The concept that the propionic acid fermentation was energetically more efficient than either the acetic acid or butyric acid fermentations based on many factors. Smith (1971) reported that propionate was utilized by the ruminants tissue more efficiently than acetate. Another possible advantage of propionate was that it was more flexible as an energy source than acetate. Propionate enjoyed the luxury of having the potential to be used for glucineogenesis in addition to direct oxidation by the citric acid cycle (Badawy, 1992). Moreover, the efficiency of microbial protein synthesis was markedly higher with the propionate as opposed to the acetate fermentation pattern as recorded by Ishaque et al. (1971).

The present data indicated that the supplementation with *Nigella sativa* cake enhance the sexual maturity and improve fertility parameters of growing baladi goats. Improved reproductive performance and fertility with supplementation of *Nigella sativa* cake was previously recorded in buffaloes by Youssef et al. (1998) and in Barki

ewes by El-Ekhnawy et al. (1999). Also, final body weight gain was significantly higher in does fed supplemented *Nigella sativa* cake ration. This result coincide with those obtained by Sharobeem (1996) in the albino, however, other investigators (Zeweil, 1996, El-Gohary, 1997, Awadalla, 1997) observed no effect of *Nigella sativa* cake on body weight and growth rate. Increased body weight and growth of (NSc) group might be attributed to high nutrient content and antimicrobial effect of *Nigella sativa* cake which act as a growth promoter agent (Rathee et al., 1982).

The current study in addition to the previous investigation, lead us to suspect that the positive reproductive impact of NS cake in growing female baladi goats may be attributed to :

- 1- Increased ruminal production of propionate due to feeding of *Nigella sativa* cake caused elevation of plasma insulin and glucose turnover which are important for the proper function of the reproductive processes (Ebling et al., 1990 and Pope and Hallford, 1991).
- 2- The higher content of unsaturated fatty acids especially linoleic acid originated from *Nigella sativa* (Al-Gaby, 1992), was recorded to improve reproductive response and fertility (Filley et al. 2000).
- 3- Increasing the energy content of the ration due to high content of crude protein and ether extract of NS cake, which raised the growth rate and body weight gain that enhance puberty and sexual maturity (Robinson, 1990 and Adam and Robinson, 1994).

Experiment 2

In the current study, the percentage of corpora lutea (CL) was significantly higher in (NSc) group which may explained by the higher ovulation rate. In this respect, *Nigella sativa* known to contain high amount of propionate which cause elevation of LH concentration and pulsation that lead to increase the ovulation rate (DiCostanzo et al. 1999). The total ovarian responses (TOR) was also significantly higher in (NSc) group than the control one, which may be attributed to high fat content of *Nigella sativa* cake. These results coincided with those obtained by Ryan et al. (1992) and Lammoglia et al. (1996) in dairy and beef cows, who found that the changes of follicular development associated with feeding of high fat ration reflected its ability to enhance follicular development. It could alter specific aspects of ovarian steroidogenetic potential and increase the population of medium sized follicular theoretically available to respond to gonadotrophin treatment in superovulation regimens (Wehrman et al., 1991). However, there is a significant elevation in the formed number of unovulated follicles. This result was confirmed by Houghton et al. (1995) who recorded that the treatment with high oil diet enhanced the fertility and conception rate. Moreover, Ryan et al. (1992) reported that the unsaturated fatty acids constituted 84.82 % of total acids of *Nigella sativa* increased medium sized follicles in cattle.

The recovery rate of transferable embryos was higher in (NSc) supplemented does than the control which may be due to high fat content of NS cake and consequently elevation of the plasma progesterone level as well (Hawkins et al,1995). It was found that the level of progesterone at the day of recovery can affect the number of recovered embryos (Ismail et al., 1992 and Mansour, 1993) and embryo quality (Jensen et al., 1982, Lindsell et al., 1986). The percentage of fair (non-transferable) embryos was significantly lower in (NSc) group than control. This result was confirmed by Gunn (1983) who concluded that embryos of ewes kept in poor condition and low nutrition during breeding were to be detrimental to embryo survival.

REFERENCES

- Abdel-Aal, M.S.E. and Attia, S.R. (1993): Characterization of black cumin (*Nigella sativa* L. seeds). *Alex. Sci. Exch.*, 14:467.
- Adam, C.L. and Robinson, J. J. (1994): The role of nutrition and photoperiod in the timing of puberty. *Proc. Nut. Soc.* 53:89-102.
- Al-Gaby, A. M. (1992): Biochemical studies on Egyptian *Nigella Sativa* L. oils. *Egypt. J. Appl. Sci.*, 5:739-748.
- Awadalla, L. M. (1997): The use of black cumin seed (*nigella sativa*) cake in rations of growing sheep. *Egypt. J. Nutr. and feeds*, 1:243-249.
- Badawy, S.A. (1992): Some biological effects of rumen fermentation modifier (monensin) on buffalo heifers with special reference to sexual puberty. Ph. D. Vet. Thesis, Cairo Univ., Egypt.
- Barowicz, T.; Brzoska, F.; Peitras, m.; Gasior, R. (1997): Hypocholesterolemic effect of full-fat flaxseeds in the diets of growing pigs. *Medycyna-Weterynaryjna*, 53:164-167.
- Cunningham, M.J.; Clifton, D.K.; and Steiner, R.A. (1999): Leptin's action on the reproductive axis. *Biol. of Reprod.*, 60: 216-222.
- Dickerman, R.W. and Li-HY, Wade, G.N.(1993): Decreased availability of metabolic fuels suppresses estrous behavior in Syrian hamsters. *Am. J. Physiol.*, 264: 568-572.
- DiCostanzo, A.; Williams, J. E. and Keisler, D. H. (1999): Effects of short- or long- term infusions of acetate or propionate on Luteinizing hormone, insulin and metabolite concentrations in beef heifers. *J. Anim. Sci.* 77:3050-3056.
- Dorn, C.G. and Kraemer, D.C. (1987): Bovine embryo grading. Texas A&M University, College of Vet. Med., Dep. Of Physiol. And Pharm.
- Ebling F. J. P.; Wood R. I., Karsch, F. J. Vannerson, L.A.; Suttie, J. M.; Bucholtz D. C.; Schall R. E. and Foster, D. L. (1990): Metabolic interfaces between growth and reproduction. III. Central mechanisms controlling pulsatile luteinizing hormone secretion in the nutritionally growth-limited female lamb. *Endocrinology* 126:2719-2727.
- El-Dakhkhny, M.; Madi, M.N.; and Abdel Halim, M. (1997): Effect of *Nigella sativa* L. oil on some liver function tests, protective against induced hepatotoxicity and lipid profile in rats. Cited by Nassar (1997).
- El-Ekhnawy, K.E.; Olteifa, A.M.; Omaima, H. Ezzo and Hegazy, M. A. (1999): Post weaning reproductive activi-

- ty of Barki ewes lambing in spring fed *Nigella sativa* oil seed meal. *Assiut Vet. Med. J.* 40 (80): 1-18.
- El-Ghamry, L. D.; Abdel-Samee; L. D.; and Ibrahim, M. (1997): Effect of feeding black cumin (*Nigella sativa*) meal and rice polishing at different levels in the laying performance and some blood plasma constituents. *Egypt. J. Nutr. and Feeds.* 1:311-320.
- El-Gohary, E.S.H.I. (1997): Studies on sheep feeding under Egyptian conditions (performance of lambs fed on diets containing *Nigella sativa* meal). M. Agric. Sc. Thesis, Mansoura Univ., Egypt.
- Filley, S.J.; Turner, H.A.; and Stormashak, F. (2000): Plasma fatty acids, prostaglandin $F_2\alpha$ metabolite, and reproductive response in postpartum heifers fed rumen bypass fat. *J. Anim. Sci.* 78: 139-144.
- Foster, D.L. and Olster, D.H. (1985): Effect of restricted nutrition on puberty in the lamb: patterns of tonic luteinizing hormone (LH) secretion and competency of the LH surge system. *Endocrinology*, 116:375-381.
- Gabr, A. A. ; El-Syouty, S. A.; Zaki, A. A.; Abou Ammo, F. F.; and El-Ghary, E.S.I. (1998): Productive performance of lambs fed diets containing *Nigella sativa* meal. *Egypt. J. Nutr. and feeds*, 2:97-107.
- Gill, C.J. and Rissman, E.F. (1997): Female sexual behavior is inhibited by short and long-term food restriction. *Physiol. Behav.* 61: 387-394.
- Gunn, R.G. (1983): The influence of nutrition on the reproductive performance of ewes. In Waresign, W. (Ed) *Sheep reproduction*, P: 99, Butterwiths, UK.
- Hall, J. B.; Schillo, K. K.; Fitzgerald, B. P.; Hileman, S. M.; Estienne, M. J.; Bradley, N. W.; and Boling, J. A. (1990): Effects of bovine somatotropin and dietary energy on LH secretion, follicular growth and onset of puberty in beef heifers. *J. Anim. Sci.* 68 (suppl. 1) 437 (abstr).
- Hawkins, D. E. Niswender, K. D., Oss, G. M.; Moeller, C. L.; Odde, K. G., Sawyer, H. R. and Niswender, G.D. (1995): An increase in serum lipids increases luteal lipid content and alters the disappearance rate of progesterone in cows. *J. Anim. Sci.* 73:541.
- Houghton, P.J.; Zarka, R.; Deles Heras, B.; and Hoult, J.R.S. (1995): Fixed oil of *Nigella sativa* and derived thymoquinone inhibit Eicosanid generation in leukocytes and membrane lipid peroxidation. *Planta Medica*, 61:33.
- Ishaque, M.; Thomas, P.C.; and Rook, J.A.F. (1971): Consequences to the host of changes in rumen microbial activity. *Nature (London) New Biol.*, 231:253.
- Ismail, S. T.; Abou-Ahmed, M. M.; and El-Beley, M. S. (1992): The relationship between plasma progesterone concentration and embryo production in dairy cows superovulated with FSH or PMSG. *Vet. Med. J.*, 3: 63-72.
- Jensen, A.M.; Greve, T.; Modej, A.; and Edquist, L.E. (1982): Endocrine profiles and Embryo quality in PMSG-PGF $_2$ alpha treated cows. *Theriogenology*, 18, 33-44.
- Khalifah, M.M. (1995): *Nigella* seed oil meal as a protein supplement in broiler diets. M.Sc. Thesis, Fac. Of Agric., Alex. Univ., Egypt.
- Khodary, R.M.; El-Ezzawy, M.H.; and Hamdy, I.R. (1996): Effect of *Nigella sativa* on egg production, hatchability percentage and some biochemical values in laying hens with reference to fertility in cockerels. 7th Sci. Cong., Fac. Vet. Med. Assiut Univ., 17-19 Nov., 1996, Assiut Egypt.
- Kirkwood, R.n. and Aherne, F.X. (1985): Energy intake, body composition, and reproductive performance of the

- gilt. *J. Anim. Sci.*, 60: 1518-1529.
- Lammoglia, M.A.; Willard, S.T.; Oldham, J.R.; and Randel, R.D. (1996): Effect of dietary fat and season on steroid hormonal profile before parturition and on hormonal, cholesterol, triglycerides, follicular patterns and postpartum reproductive in Brahman cows. *J. Anim. Sci.*, 2253.
- Lindsell, C. E. Murphy, B. D. and Mapletoft, R. J. (1986): Superovulatory and endocrine responses in heifers treated with FSH-P at different stages of oestrus cycle. *Theriogenology*, 26:209-219.
- Mani, A. U.; McKelvey, W. A. C.; and Watson E. D. (1992): The effect of low of feeding on response to synchronization of estrus, ovulation rate and embryo loss in goats. *Theriogenology*, 36: 1013-1022.
- Mani, A. U.; Watson E. D.; and McKelvey, W. A. C. (1993): The effect of sub-nutrition on the components of the gravid uterus in doe. *Theriogenology*, 40:287-294
- Mansour, M. M. (1993): Embryo productin and related hormonal changes in Rahmani ewes . Ph. D. Thesis, Vet. Med., Cairo. Univ.
- McKelvey, W.A.C.; Wallace, J.M.; Robinson, J.J.; and Aitken, R.P.(1989): Studies on increasing breeding frequency in the ewe. I. The fertilization of ova during the early post-partum period. *Anim. Reprod. Sci.*, 18: 1-12.
- Nassar, H.M. (1997): Pharmacological and toxicological studies on *Nigella sativa*. Ph. D. Vet. Thesis, Alex. Univ., Egypt.
- NRC (1989): Nutrient requirment of dairy cattle (6th Ed). Washington, D.C. National Academy Press.
- Pope, T. C. and Hallford, D. M. (1991): Effect of insulin on the concentration of glucose and hormones in serum and milk yeild of postpartum ewes. *Can. J. Anim. Sci.* 71:1257-1260
- Pope, W.F.; McClure, K.E.; Hogue, D.E.; and Day, M.I. (1989): Effect of season and lactation on postpartum fertility of Targhee ewes. *J. Anim. Sci.*, 67:1167
- Rathee, P.S. ; Mishra, S.H.; and Haushal, R. (1982): Antimicrobial activity of essential oil fixed and unsaponifiable matter of *Nigella sativa* . *Ind. J. Pharm. Sci.*, 44:8
- Robinson, J.J. (1990): Nutrition in the reproduction of farm animals. *Nutr. Res. Rev.*, 3: 253-276.
- Ryan, D. P.; Spoon, R. A.; and Williams, G. L. (1992): Ovarian follicular characteristics, embryo recovery, and embryo viability in heifers fed high-fat diets and treated with FSH. *J. Anim. Sci.* 70: 3505-3513.
- SAS Institute (1986): SAS/STAT guide for personal computers. Version 6 (Ed. SAS Inst. Inc.) Cary, NC.
- Sharobeem, S.F. (1996): Chemical characterization of *Nigella* cake protein and its biological effects on albino rats. *Zagazig J. Agric. Res.*, 3:355-368
- Smith, G.E. (1971): Energy metabolism. Cited by Badawy (1992).
- Steel, R.G.D. and Torrie, J.H. (1980): Principals and procedures of statistics. New York: McGraw-Hill
- Thomas, M. G. and Williams, G. L. (1996): Metabolic hormone secretion and FSH induced superovulatory responses of beef heifers fed dietary fat supplements containing predominantly saturated or polyunsaturated fatty acids. *Theriogenology*, 45:451.
- Thomas, G.B.; Mercer, J.E.; Karalis, T.; Rao, A.; Cummins, J.T.; and Clarke, I.J. (1990): Effect of restricted feeding on the concentrations of growth hormone (GH), Gonadotropins, and prolactin (PRL) in plasma, and on the amounts of messenger ribonucleic acid for GH, gonadotropin subunits, and PrL in the pituitary glands of adult ovariectomized ewes. *Endocrinology*, 126:1361-1367.

Youssef, M.M.; Abdiene, A. M.; Khattab, R. M.; and Darwish, S. A. (1998): Effect of feeding *Nigella Sativa* cake on productive and reproductive performance of buffaloes. *Egypt. J. Nutr. and Feeds*, 2: 73-85.

Wehrman, M. E.; Welsh, T. H., J R.; and Williams, G. L.

(1991): Diet-induced hyper lipidemia in cattle modifies the intrafollicular cholesterol environment, modulates ovarian follicular dynamics, and hastens the onset of post-partum luteal activity. *Biol. Reprod.* 45:514-522.