



Partial Replacment of Yellow Maize Grains by Sun-Dried Orange Pulp in Summer Rations of Lactating Buffalo



Mohammed H. Bakr¹, Sabbah M. Allam¹, Eman H. Elsabaawy², Ebtehal

H. Al-Adawy^{1*} and Fatma Abou-Hashim¹

¹Animal Production Department, Faculty of Agriculture, Cairo University, 12613 Giza, Egypt.

²Animal Production Department, Agricultural and Biological Research Institute, National Research Centre, 12622, Cairo, Egypt.

Abstract

TO study the partial replacement of yellow maize grains (YMG) with sun-dried orange pulp (SDOP) during the summer months, 27 Egyptian lactating buffaloes (body weight: 600 ± 50 kg; parity: 1st to 4th) were allotted randomly to 3 experimental groups. Investigations were done on specific blood parameters, feed consumption, nutrient digestibility, milk composition and yield, feed and economic efficiency. Sun-dried orange pulp replaced YMG at 0 Control (R1), 30%SDOP (R2), and 60% SDOP (R3). Comparing to Control, the dry matter intake (DMI) in 30%SDOP and 60%SDOP was decreased by 0.63 and 0.83 kg, respectively. Raising the replacement level of SDOP led to an enhancement in the digestibility of crude fiber (CF; $P < 0.05$), while the rest of nutrients remained unchanged except for crude protein (CP) and digestible CP, which dramatically reduced. Statistically, no significant variations were observed across all blood parameters analyzed ($P > 0.05$). Concerning fat-corrected milk (FCM) and daily milk yield (DMY), no significant changes ($P > 0.05$) were observed, however. Compared to Control, the cost of concentrate feed mixture (CFM; LE/kg) was lower by 6.6% and 13.17%, meanwhile, feed efficiency was improved by 16.67 and 4.17% in 30% SDOP and 60% SDOP, respectively, which was strongly reflected in reducing feeding cost to produce 1 kg of milk and enhancing feed and economic efficiency. The results of the study imply that feed producers and ruminant holders may be able to reduce feeding expenses without affecting neither animal welfare nor performance in a way that is still profitable. Moreover, Egyptian buffalo is ideal for Egypt's summer climate.

Keywords: Buffalo Feeding, Sun Dried Orange Pulp, Heat Stress, feed digestibility and Economic Efficiency.

Introduction

Total digestible nutrients (TDN) an estimated 4.2 million tons are needed annually for animal production systems in Egypt; however, the amount of feed that is needed and what is available differs significantly [1]. This gap is seen to pose a serious threat to Egypt's animal husbandry activities, especially in the area of concentrated feedstuffs.

Grain maize accounts for a substantial portion of the CFM for livestock (around 0.50 ± 0.10) [2]. Egypt imported over than 11 million tons of corn grains in 2021–2022 compared to 64,000 tons in 1969, a 172-double increase that required a

substantial amount of foreign currency annually (roughly \$3 billion in 2022) beside the ongoing increase in concentrate feed ingredient costs [3]. However, the 2021/22 season saw a decrease in grain imports due to Russia and Ukraine ongoing conflict. Hence, shipping disturbance occurred in the Black Sea and inactive ports in Ukraine, which consequently prompted a sharp increase in grain prices [4]. These days, feeding dairy ruminants a grain-based diet doesn't seem like a cost-effective approach and this is because of the low price of milk and high feed cost [5]. So that, it is inevitable to find a new feed strategies to reduce the proportion of grains used in livestock feeding to reduce feed costs without sacrificing the animals' health or

*Corresponding author: Ebtehal Habib Al-Adawy, E-mail: ebtehalaladawy26.me@gmail.com, Tel.: 01007656831

(Received 23 June 2024, accepted 19 August 2024)

DOI: 10.21608/EJVS.2024.298859.2189

©2025 National Information and Documentation Center (NIDOC)

performance.

Heat stress (HS) is a major contributor to summertime losses for farms raising dairy and meat because it impairs animal immunity, growth rates, milk production, and reproductive efficiency [6]. From this perspective, domestic ruminants are less susceptible to HS than buffaloes [7].

Wind speed, atmospheric pressure, ambient temperature (AM), and relative humidity (RH) are environment-related elements that influence livestock and cause HS [8]. The detrimental effects of HS on cattle could be scaled by the Temperature Humidity Index (THI), where a decrease in milk production and fertility is linked to a high THI of 75 or higher [9-10]. Omran *et al.* [11] suggest that feed supplements may reduce the climate sensitivity of buffaloes.

Moreover, the occurrence of global warming highlights the necessity of locating alternative feed sources for cattle raised in heat-stress environments that either lowering heat increment produced from metabolism processes or supplementation of specific additives with alleviating heat stress properties, such as vitamin E and C, or other antioxidants [12].

Moreover, 48.6 million tons of oranges were expected to be produced worldwide in 2020–2021 with 50–60% of the processed fruit may be discarded. Additionally, Egypt is recognized as one of the top manufacturing and exporting nations [13-14].

Excellent nonconventional feeds for ruminants could be obtained as orange by-products such as DOP, citrus silage, citrus molasses and liquor of citrus peel [2-15-16], being an energy supplement to cover ruminant requirements [17]. Moreover, bioactive materials with antioxidant qualities like sugar, fiber, organic acids, phenolic compounds, and vitamin E can be found in citrus by-products (like pulp) [18-20].

After fresh oranges are pressed to obtain drink for human consumption, what's left over is called orange pulp (entire fruits without juice) [21-22]. Due to the significant amount of rapidly fermenting carbohydrates in DOP and various energy compounds for the ruminal microbiota, such as soluble carbohydrates and digestible neutral detergent fiber (NDF), DOP appears to be a high quality feedstuff and has a high nutritive value [2].

Prior research suggested that substituting DOP for grains in a dairy goat's diet did not lead to any negative effect on milk production, feed consumption, or digestibility [23-25]. Moreover, adding 40 and 80% levels of DOP to grains increased the quantity of vitamin E and phenols in lactating

goat milk and enhanced the growth performance of the goat kids [26]. Breastfeeding cows given the DOP 75% substitution of YMG did not experience any negative health effects [16]. Lactating buffaloes fed a diet made up of a combination of DOP and dried tomato pomace (1:1) showed increased milk output [27]. In the same way, a study discovered that feeding lactation buffaloes a diet containing 20% of beet pulp and 10% DOP increased milk output and nutrient digestibility [28].

Research on the use of DOP and the advantageous effects of its bioactive components to alleviate the effect heat stress on lactating buffalo performance is still lacking. Thus, the aim of the current study was to investigate how partial replacement of YMG by DOP in the summer diet of Egyptian lactating buffalo affected the nutrients' digestibility, some blood parameters, milk production, composition, and economic efficiency.

Material and Methods

All fieldwork has been done at the Experimental and Agricultural Research Station (Buffalo Research Unit). Chemical analyses of feed ingredients, feces and blood measures were completed by Cairo University Research Park (CURP; Hormonal Laboratory) and the Department of Animal Production at the Faculty of Agriculture - Cairo University (Egypt). This study was accomplished during the summer months of July through September 2018. During this time, at the afternoon hours (between 13:00 and 14:00 h), it was noted that ambient temperatures ranged between 35 and 38°C and humidity levels between 54 and 55%.

Animals, experimental diets, and ethical clearance

For the experiment, 27 Egyptian buffaloes (600±50 kg; 1st to 4th parity) were used, and arbitrarily divided into 3 groups. The buffaloes were fed the experimental and control rations, in proper amounts, twice daily, at 7:00 and 19:00 h, to ensure that their nutritional demands were met. The lactating buffaloes were fed daily 8 kg of rice straw, 25 kg of Drawa, the green corn plants, and 8 kg of CFM, with a roughage concentrate ratio (R:C) of 65.60:34.40% for each. The average daily milk production per head at the beginning of the research was 6.43 kg ± 0.1 kg. The experimental animals were housed in open-air pens. Clean and fresh water was available freely throughout the day.

The CFMs consisted of yellow maize grains, soybean meal (44%), wheat bran, SDOP, sunflower meal (36%), and some additional feed additives. Three experimental rations were used, as Table 1 illustrates: R1, which consisted of 500 kg of YMG (Control); R2, which consisted of 350 kg of YMG plus 150 kg of SDOP (30% SDOP); and R3, which

consisted of 200 kg of YMG plus 300 kg of SDOP (60% SDOP).

The current study was ethically approved and issued with number: CU/II/F/20/21 for the experimental design by Cairo University - the Institutional Animal Care and Use Committee (IACUC)- Giza, Egypt.

Digestibility trial and chemical composition of feeds and feces

Digestion studies were carried out in the last week of the experiment to ascertain the nutritional value and digestibility of the experimental ration. Each experimental animal's rectum was sampled three times in the morning and evening before feeding to obtain feces. In accordance with Van Kullen and Young [29], an internal marker, acid-insoluble ash (AIA), was utilized to assess the digestion coefficient of nutrients. The following nutrients, according to AOAC, were examined in fecal samples and experimental diets: dry matter (DM), crude protein (CP), organic matter (OM), ether extract (EE), and crude fiber (CF) [30]. The variation utilized in the nitrogen-free extract (NFE) formula. The following equations from Edwards and Allan [31] were used to determine total energy (GE; MJ/kg DM): [Carbohydrates (g) $\times$ 17.20] + [CP (g) $\times$ 23.60] + [EE (g) $\times$ 39.50]. TDN (nutritive values) was determined using the standard equation of McDonald et al. [32]. As per Van Kullen and Young [29], the estimate of nutrients digestibility was calculated as follows:

$$\text{Total digestible nutrients (\%)} = [(\text{Digestible EE (\%)} \times 2.25) + \text{Digestible CF (\%)} + \text{Digestible NFE (\%)} + \text{Digestible CP (\%)}].$$

$$\text{Digestibility of DM (\%)} = 100 - \left(100 \times \frac{\text{Marker\% in feed}}{\text{Marker\% in feces}}\right).$$

$$\text{Digestibility of nutrients (\%)} = 100 - \left(100 \times \frac{\% \text{ nutrient in feces} \times \text{Marker\% of feed}}{\% \text{ nutrient in feed} \times \text{Marker\% of feces}}\right).$$

Blood biochemistry

On day 90 of the trial 10-ml blood sample was taken from each animal's jugular vein. Samples were collected and placed in heparinized tubes. Then, the blood plasma blood was separated by centrifuging (4000 rpm - 15 min). Afterwards, blood samples were stored at -20 °C until analyzed.

Alanine transaminase (ALT) and aspartate transaminase (AST) activity (RFU/ml) was assayed using the Reitman and Frankel [33] method. The methods supplied by Cannon [34] and Doumas et al. [35] were followed for measuring total protein and albumin concentrations, respectively.

Total lipid, triglyceride, and cholesterol levels

were measured following Eisemann et al. [36] methodology. A spectrophotometer (Jenway 6300; U.K.) was utilized to analyze all plasma parameters calorimetrically. The Stat lab szsl60 – Spectrum device was used to assess total antioxidant capacity, in accordance with Koracevic et al. [37] description. The levels of cortisol, immunoglobulin G (IgG), thyroxine (T4), and triiodothyronine (T3) were determined by ELISA (BIOTEK ELX808; USA).

Thermo-humidity index

Once a week, the humidity and ambient temperature (AT) were monitored during the hottest daytime (13:00 to 14:00 h). A dry bulb thermometer was used to measure the ambient temperature, and the wet and dry bulb thermometers were used to calculate RH. As a severity gauge of the HS in animals, a study by Mader et al. [38], concluded the temperature humidity index (THI) was determined using the following formula:

$$\text{THI} = 46.4 + \left[\left(\frac{\text{RH}}{100}\right) + (\text{AT}; ^\circ\text{C} - 14.4)\right] + (0.8 \times \text{AT}; ^\circ\text{C}).$$

Milk production, composition, and sampling

Manually, the buffalos were milked 2 times a day considering the first 3 weeks of the trial as an adaptation period, and milk yield was monitored every day until the experiment's finished. On the last 2 days of the experiments, representative and random samples of milk were taken and conserved under -20 °C until analyzed. A milko scan (type 10900 FOSS electric; 130 series; Denmark) was used to analyze the total solid -TS- content, fat, solid-not-fat (SNF), protein and lactose. Meanwhile, daily milk production (actual) was corrected into 4% FCM following Gaines [39] formula: [(fat yield, kg) $\times$ 15] + 0.4 $\times$ (milk production, kg).

Statistical analysis

All results were subjected to one-way ANOVA by SPSS software (V.15), using the statistical model:

$$Y_{ij} = \mu + T_i + E_{ij},$$

where: Y_{ij} = the dependent variable; μ = Over all mean; T_i = Treatment effect; E_{ij} = residual error. The mean $\pm$ SEM was used to express the results. After performing Tukey's multiple comparisons, the variations between groups were considered significant at $P < 0.05$.

Results and Discussion

In the current investigation, animals in R2 and R3 groups were fed CFMs that partially replaced YMG with 30% and 60% of SDOP, respectively, while animals in control (R1) got CFM without SDOP. Together with their prices, Table 1 displays the experimental CFMs' formula

When it came to feed costs, the price of CFM

decreased as replacement levels of SDOP increased. This was because, during the trial period in July to September 2018, the average cost of 1 kg of YMG was 4.15 LE, but 2.20 LE per kg of SDOP. As a result, compared to Control, the CFM cost for 1 kg was lower in 30%SDOP and 60%SDOP by 0.293 and 0.585 LE, respectively (that is, it was lower by 6.6 and 13.17%, respectively, compared to control). This is the first indication that the work can be applied in buffalo feed manufacturers and farms of ruminant production to reduce the CFM cost and enhance profits.

Our findings were in accordance with those of Kemmere *et al.* [40], who fed dairy cows about 50% of the standard herd grain-based ration and 50% dried grapefruit meal. About 17.2% less was spent on a ton of rations because of this partial replacement. Additionally, Allam *et al.* [16] found that raising the replacement levels of YMG by DOP substitution level in dairy cow diets up to roughly 17% reduced the cost of CFM.

Composition of chemicals in the feed ingredients and CFMs used in the experiment

Chemical composition of feedstuffs and CFMs is reported in Table 2. Results showed that the CP and EE contents of SDOP and YMG were nearly identical. If not, SDOP's OM, DM, and NFE values were lower than YMG's. Nonetheless, SDOP's ash and CF values were greater than YMG's.

The contents of OM and CP in SDOP were determined to be 91–92 for OM and 7.2% for CP, which were almost identical to those reported by several studies [16 - 17]. Value of Ether extract in SDOP (4.77%) was quite similar to the values reported by Allam *et al.* [41]: 4.5% and the NRC [42]: 4.9%. Table 2 indicates that the CFMs were roughly isonitrogenous and isocaloric, with CP levels being roughly equal (17.33 to 17.60%).

Digestibility of nutrients and other nutritive values

Table 3 shows the results of digestion coefficients, the nutritive value expressed in digestible crude protein, and TDN for rations of the experiment containing different levels of SDOP substitution (30% and 60% of YMG) compared to the control diet. No statistically significant difference in the experimental digestibility of DM, OM, EE, and NFE when substituting SDOP for YMG is found ($P > 0.05$). These results were consistent with those of other studies [43], [44], [45], who examined the effects of substituting dried citrus pulp (DCP) for barley grains at varying levels in the diets of goats, sheep, and fattening male calves, and found no discernible variation in the DM digestibility. Additionally, DCP was used in place of YMG in small ruminant rations by Allam *et al.* [41] and

Gilaverte *et al.* [46], who concluded no significant difference in the overall digestibility of nutrients. Additionally, Shdaifat *et al.* [47] found no significant variations in the DM and OM digestion coefficients after substituting 20% of the barley grains in the dairy ewes' diet with DCP.

Comparison with control revealed a substantial decrease in CP digestibility concurrent with the SDOP substitution level ($P < 0.05$). However, Allam *et al.* [16] and Ben-Ghedalia *et al.* [45] observed that CP digestibility increased as the amount of DOP in ruminant diets increased. This might be explained by DOP that could improve N flow into the intestine by creating favorable ruminal conditions.

When the SDOP substitution level was increased in 30%SDOP and 60%SDOP relative to Control, a significant improvement in digestibility of CF was observed ($P < 0.05$). Grigelmo-Miguel and Mart  n-Belloso [48] suggested that the reduced lignin content in DOP was linked to the improvement in CF digestibility. Furthermore, even at high dietary inclusion, it might also be connected to the DOP's capacity to promote ruminal fibrolytic bacteria-friendly environments [45], thereby, enhancing CF digestion. Allam *et al.* [16] concurred with these results.

In contrast to control and 30%SDOP, 60%SDOP recorded the lowest value for NFE digestibility (76.91 vs 69.3%). However, differences were not significant ($P > 0.05$). Maybe it due to the high variability.

Furthermore, TDN in the control group was numerically higher than 30%SDOP and 60%SDOP, with no significant differences found ($P > 0.05$). Additionally, because of decreasing CP digestibility, digestible CP was considerably reduced by raising the SDOP substitution amount ($P < 0.05$).

Effects on blood parameters

The blood biochemical indicators for the experimental animals are illustrated in Figure 1 and 2, and all of the parameters were confirmed to be within the normal ranges for adult buffalo.

Regarding blood parameters examined no discernible change was showed between groups. Comparable levels of total protein, albumin, hemoglobin, and liver enzyme activity (ALT and AST) were seen overall the experimental groups. Figure 1 indicate that total lipid, triglyceride, and cholesterol values did not differ significantly from the control. In the same context, Figure 2 showed no significant variation was seen for cortisol, T3, and T4 hormone levels. Additionally, the TAC values for control, 30%SDOP, and 60%SDOP were very similar 2.11, 2.05, and 2.09 mmol/L, respectively

without reporting any significance ($P > 0.05$). The concentration of IgG was somewhat higher in R2 and R3 (9.97 and 9.90 mg/mL, respectively) when compared to control (9.54 mg/mL).

These results were consistent with those of Belibasakis and Tsirgogianni [49], who found that the blood total protein and albumin of dairy cows were not affected in a significant manner by the addition of dried beet pulp and ground maize to their diet when DCP replaced 20% of CFM. Further research on the effects of DOP replacement up to 50% of YMG in small ruminant rations on ALT, AST, and total protein levels was done by Allam et al. [41], who also showed that the blood parameters evaluated were within the normal range and that there was no significant variation between groups. Furthermore, Allam et al. [16] found that all blood measures were also within normal ranges after examining the impact of substituting DOP for up to 75% of YMG in the diets of dairy cows. On the other hand, Oni et al. [50] investigated the effects on several blood markers of adding DCP to the goat ration at varying percentages (up to 75%) in place of the dried brewers grains. They found that, in comparison to the control group, total protein and hemoglobin significantly increased with increasing substitution levels.

Milk yield and composition

This trial had a 12-weeks duration, with the first three weeks serving as an adaptation period during which SDOP was introduced, and the remaining 9 weeks being the experimental phase. Because SDCP's seeds and peels contain limonin, which gives them a bitter taste, buffalo needed to adapt during this time [51]. In addition, it's important to gradually introduce citrus by-products into animal diets so that animals become used to their unique flavor and aroma [52]. Additionally, it might alter the ruminal microbiota, which is closely linked to dietary item variation and may therefore have an impact on the welfare and productivity of animals. Figure 3 displays the actual DMY and FCM results for the 9-weeks experiment and lists the chemical composition of milk.

In terms of DMY, no significant differences were reported among the groups; however, the highest value (6.06 kg/h/d) was for R2. According to many previous studies [53-55], our results were in agreement with theirs. Additionally, as illustrated in Figure 3, milk production as a 4% FCM increased in 30% SDOP but remained the same in 60% SDOP and control.

As SDOP replacement levels in the diets of lactating buffalo increased, the content of protein, lactose, and total solid in the milk increased as well,

although the significance level was not reached ($P > 0.05$; Figure 3). However, the R2 and R3 groups showed a non-significant drop in milk fat content; nevertheless, R2 had the lowest value. Similarly, between the treatments and the control group, there was no change in milk ash ($P > 0.05$). These results were in line with those of Lima et al. [53], Assis et al. [54], and Wing [55], who discovered no significant change in the fat content of cows fed rations containing varying amounts of DCP.

These findings were discussed in light of earlier research that found adding DCP to the diets of lactating cows notably influenced milk composition, particularly the amount of fat in it. Santos et al. [18] found that feeding diets with 3% soybean oil, 9, and 18% DCP significantly decreased total solid and milk fat contents. Furthermore, Shdaifat et al. [47] looked at replacing 20% of barley grains with DCP in small ruminant diets and discovered that the milk fat had the tendency to be higher comparing to the control group (7.42 vs. 6.21%; $P < 0.08$).

All experimental groups showed no significant differences in the milk protein. These findings were consistent with those of Belibasakis and Tsirgogianni [49], who demonstrated that 20% of CFM, dried beet pulp, and ground maize were replaced with DCP in dairy cows without affecting the contents of milk protein. Additionally, when Assis et al. [54] replaced maize grains in the rations of lactating cows at percentages of 33%, 67%, and 100% with DCP, there was no significant change in the milk protein.

Similarly, there was no significant variation in the lactose content across the experimental groups. These results corroborated those made by Wing [55], who determined that adding DCP to the diets of lactating cattle had no notable effects. Furthermore, DCP substitution of 20% of CFM in lactating cows was studied by Belibasakis and Tsirgogianni [49] and found no discernible effect on lactose content. Moreover, the influence of substituting up to 18% of DCP in the diet of dairy cows was examined by Santos et al. [18], and discovered no markable change in the lactose content of the milk. When Lima et al. [53] investigated DCP supplementation effect as an antioxidant in canulated lactation cows, they found the same results.

In our study, the TS content of milk was found to be unaffected by the treatments ($P > 0.05$; Figure 3). Similar to these results, Belibasakis and Tsirgogianni [49] discovered that milk TS was not significantly impacted substituting 20% of CFM by DCP in lactating cows. Additionally, Assis et al. [54] revealed no notable changes in milk TS when they fed dairy cow diets that included DCP (33–100%) in place of maize grains.

Efficiency of feed and economics

Table 4 shows the effects on feed and economic efficiencies of partially substituting SDOP for YMG in the diets of lactating buffalo. Regarding the feed intake, raising the SDOP substitution level resulted in a decrease in the overall dry matter intake. In comparison to control, this decline was accounted for by 0.63 and 0.83 kg/h/d, in the 30%SDOP and 60%SDOP, respectively. This was represented by a reduction in daily feed cost as well as the CFM price.

As the SDOP level rose, feed efficiency improved and control, 30%SDOP, and 60%SDOP representing 0.24, 0.29, and 0.25, respectively. Raising the DOP replacement levels in rations significantly not only reduced the cost of CFM but also total feed cost for producing 1 kg of milk. Thus, as the DOP replacement levels increased, so did the income (LE) and relative economic efficiency (REE).

In other words, elevating the YMG replacement level by SDOP, lowered the cost of daily feed (LE/head/day) to be 46.22, 42.58, and 39.9 for the control, 30%SDOP, and 60%SDOP, respectively. In addition, compared to the control group, the cost of feed (LE) per kg of milk decreased by 1.80 and 1.09 LE, or 20.4% and 12.36%, in 30%SDOP, and 60%SDOP, respectively. Additionally, prior gains resulted in an increase in REE for producing 1 kg of milk 30%SDOP, and 60%SDOP registering 2.11 and 1.48 times, respectively, over control.

These results were consistent with those obtained by Miron *et al.* [56], who found that feeding high-producing dairy cows feeds partially substituted with DCP (10–20%), instead of maize grains, increased the efficiency of feed. Additionally, Allam *et al.* [16] reported that the REE (%) of replacing YMG by DOP in the rations of lactating cows were 116.43, 120.65, and 126.08% at 25, 50, and 75% of replacing levels, respectively.

Environmental physical aspects

Figure 4 displays the THI, relative humidity, and ambient temperature that were noted throughout the summertime experimental period. The degree of heat stress was measured using the scale: Normal THI is 75 or less; alert is 75–78; danger is 79–83; emergency is 84 or more [57]. The buffaloes in the current study had a mean THI value of 89.9, which indicated that they were in the emergency category and indicated a stressed condition based on the prior index.

Once THI ranged between 72–89, buffalo's milk production, lipid content, and protein levels decreased [9–57]. The cortisol level in buffalo increased during heat stress, as stated by Kumar *et al.* [7] and Comin *et al.* [58]. Consequently, the cortisol

result in the current study followed the same trend. Additionally, T3 concentration in buffalo was dramatically declining as the THI index rose [59].

According to our research, giving lactating buffaloes diets containing SDOP may prevent the summertime decline in milk yield and composition. Additionally, feeding SDOP did not lead hormones to become stressed out in response to heat stress; this could be clarified by the fact that DOP has a higher concentration of several bioactive anti-heat stressed chemicals than YMG, such as vitamin C.

Conclusion

According to the conditions of our study, it is possible to conclude that adding SDOP instead of YMG to the rations of lactating buffaloes improved the digestibility of CF while decreasing the digestibility of CP and its nutritional value as a digestible CP. Despite the fact that this experiment was conducted in the summer, none of the blood parameters in 30%SDOP or 60%SDOP were impacted by the substitution. Furthermore, there was no significant variation in the amount or composition of milk produced, meanwhile, feed efficiency and profitability were enhanced by SDOP inclusion into the diet. Moreover, Egyptian buffalo are well-suited to summertime living. Finally, the study found that farm profitability increased when SDOP was substituted for up to 60% of the YMG. This feeding strategy of reducing feeding costs without compromising animal performance or health is commercially viable among producers of ruminant feed and animal production farms

Acknowledgments

This paper is a component of the Scientific Project No. 57-16, which Cairo University provided funding from 2017 to 2020. We sincerely thank the Scientific Research Project Coordination Unit at Cairo University in Egypt for providing financial support for this work.

Funding statement

This article is apart from the Scientific Project No 57-16 that was financially supported from Cairo University (2017-2020)

Conflicts of interest

Authors declare that there were no conflicts of interest.




Fig. 1. Some blood indicators measured for the experimental buffaloes given SDOP at levels of 0, 30 and 60% as a partial substitute of YMG. Error bars refer to standard error of means. Symbols a, b, c (if given) indicate difference at $P < 0.05$. Control: SDOP-free CFM; 30%SDOP: A 30% partial replacement of YMG in the CFM; 60%SDOP: A 60% partial replacement of YMG in the CFM. IgG: immunoglobulin G; ALT: Alanine transaminase; AST: Aspartate transaminase.




Fig. 2. Some hormonal indicators (triiodothyronine, thyroxine and cortisol) and total antioxidant capacity measured for the experimental buffaloes given SDOP at levels of 0, 30 and 60% as a partial substitute of YMG. Error bars refer to standard error of means. Symbols a, b, c (if given) indicate difference at $P < 0.05$. Control: SDOP-free CFM; 30%SDOP: A 30% partial replacement of YMG in the CFM; 60%SDOP: A 60% partial replacement of YMG in the CFM.




Fig. 3. Milk yield and composition for the experiment buffaloes given SDOP at levels of 0, 30 and 60% as a partial substitute of YMG. Error bars refer to standard error of means. Symbols a, b, c (if given) indicate difference at $P < 0.05$. Control: SDOP-free CFM; 30%SDOP: A 30% partial replacement of YMG in the CFM; 60%SDOP: A 60% partial replacement of YMG in the CFM. FCM: fat corrected milk.




Fig. 4. Environmental aspects recorded during the experimental period. Error bars refer to standard error of means

TABLE 1. Formulation (g/kg) and price (LE/kg) of the experimental concentrate feed mixtures

Items	Feed mixtures concentrate		
	Control (R1)	30%SDOP (R2)	60%SDOP (R3)
Feed ingredients, g/kg			
Sun-Dried orange pulp	0	150	300
Yellow maize grains	500	350	200
Bran of wheat	200	200	200
Sunflower meal (36%)	100	100	100
Soybean meal (44%)	150	150	150
Sodium chloride	10	10	10
Limestone	21	21	21
Sodium bicarbonate	10	10	10
Dicalcium phosphate	4	4	4
Vitamins and minerals premix ¹	3	3	3
Yeast	1	1	1
Mycotoxins binder	1	1	1
Price (LE/kg)²	4.440	4.148	3.855

¹ Vitamins and minerals premix was formulated to fit lactating cattle requirements and produced by Dakahlia Company, May 2018 Sadat City, Egypt). ² The average price (LE/kg) market of feed ingredients from July to September (2018) were: YMG (4150) and SDOP (2200). Control: SDOP-free CFM; 30%SDOP: A 30% partial replacement of YMG in the CFM; 60%SDOP: A 60% partial replacement of YMG in the CFM.

TABLE 2. Chemicals composition of the concentrate feeds mixtures and experimental rations (g/kg, except for GE) ¹

Items	Chemical composition (g/kg)							
	DM	OM	Ash	CF	CP	EE	NFE	GE
Feed ingredients								
Sun-Dried orange pulp	846.1	916.8	74.8	123.2	74.3	47.7	680.3	17.70
Yellow maize grains	906.3	986.9	13.1	25.5	80.1	42.7	838.6	18.70
Wheat bran	903.5	946.6	53.4	103.0	131.1	29.7	682.8	18.02
Sunflower meal	827.2	928.6	71.4	241.6	350.7	9.2	327.1	18.59
Soybean meal	917.5	936.2	63.8	41.7	439.2	15.1	440.2	19.39
Experimental CFMs								
Control	909.7	964.3	35.7	67.1	176.0	32.1	689.1	18.66
30% SDOP	901.9	955.4	44.6	82.4	174.7	32.1	666.2	18.49
60% SDOP	894.2	946.5	53.5	97.6	173.3	32.2	643.4	18.33

¹ Dry matter (DM), organic matter (OM), crude fiber (CF), crude protein (CP), ether extract (EE), nitrogen-free extract (NFE), gross energy (GE; MJ/kg DM; calculated). The CFM without SDOP was the control; 30% SDOP was the partial replacement of YMG in the CFM; 60% SDOP was the partial replacement of YMG in the CFM.

TABLE 3. The nutritive value as TDN and digestible CP, as well as the digestion coefficients of nutrients for the experimental rations (Control, 30%SDOP, and 60%SDOP) ¹

Items	Treatments			SEM
	Control (R1)	30%SDOP (R2)	60%SDOP (R3)	
Nutrients digestibility (%)				
DM	60.51a	61.18a	61.38a	0.507
OM	66.70a	68.87a	66.96a	0.358
CF	52.32a	56.47b	57.88c	0.509
CP	68.65a	64.87b	63.15c	0.347
EE	71.30a	70.75a	70.46a	0.310
NFE	76.91a	76.91a	69.30a	6.308
Nutritive values (%)				
Digestible CP	12.08a	11.33b	10.94c	0.06
TDN	73.71a	72.33a	66.28a	4.26

¹ Dry matter (DM), organic matter (OM), crude fiber (CF), crude protein (CP), ether extract (EE), nitrogen-free extract (NFE), and total digestible nutrients (TDN). Different symbols a, b, and c in the same row indicate differ at $P < 0.05$. The CFM without SDOP was the control; 30% SDOP was the partial replacement of YMG in the CFM; 60% SDOP was the partial replacement of YMG in the CFM.

TABLE 4. Feeding and milk yield efficiency and profitability for the experimental animals fed SDOP-containing diets at 0, 30 and 60% replacement of YMG.

Items	Treatments		
	Control (R1)	30%SDOP (R2)	60%SDOP (R3)
Milk production (kg/head/day)	5.24	6.06	5.16
Daily DMI (kg/head)	21.43	20.80	20.60
FE	0.24	0.29	0.25
Cost of feed (LE/ head/d)	46.22	42.58	39.9
Cost of feed per kg milk (LE)	8.82	7.02	7.73
Total income of milk production (LE/day)	57.65	66.64	56.78
Feeding net revenue (LE)	11.43	24.06	16.88
Economic feed efficiency %	24.7%	56.5%	42.3%
REE	1.00	2.11	1.48

Dry matter intake (DMI); Feed efficiency (FE) = milk produced (kg) / dry matter consumed (kg). Cost of feed (LE/h/d) = price of 1 kg CFM*8 + 8 kg rice straw * 0.4 LE + 25 kg Drawa*0.3 LE. Production feed cost (LE/ kg milk) = Daily feed cost (LE/h/d) / Average DMY (kg/h). Total income of milk production (LE/buffalo): Average DMY (kg/buffalo) * market price for 1 kg of buffalo milk (11 LE in September 2018). Feeding net revenue (LE) = Total income of milk production (LE) - Daily feed cost (LE). Relative economic efficiency (REE) = feeding net revenue (LE) per buffalo in each group relative to the control group.

References

- Shoukry, M.M. The possibility of using some unconventional feed resources as a source of energy and protein in feeding livestock. *Egyptian Journal of Nutrition and Feeds*, **22**(2) special issue. (2019).
- Bakr., M.H. Citrus pulp as an innovative feed ingredient in ruminant nutrition. A review. *Egyptian Journal of Animal Production*, **57**(Suppl. Issue), 73-80 (2020).
- USDA. Grain and Feed Annual U.S by Ahmed Wally, Agricultural Specialist, and Jenny Morgan, Agricultural Attaché, Cairo, Egypt. Report N. EG2023-0003(2023)
- Ali , A.A, Azaroual, F., Bourhriba, O. and Dadush, U. The economic implications of the war in Ukraine for Africa and Morocco. Policy center for the new south. PB-11/22, February(2022).
- Ertl, P., Zebeli, Q., Zollitsch, W. and Knaus, W. Feeding of by-products completely replaced cereals and pulses in dairy cows and enhanced edible feed conversion ratio. *Journal of Dairy Science*, **98**(2),1225-1233 (2015).
- Dash, S., Chakravarty, A.K., Singh, A., Upadhyay, A., Singh, M. and Yousuf, S. Effect of heat stress on reproductive performances of dairy cattle and buffaloes. A review. *Veterinary World*, **9**(3), 35-244(2016).
- Kumar, B., Sahoo, A.K., Ray, P.K., Chandran ,P.C., Taraphder, S., Das, A.K., Batabyal, S., Dayal, S. Evaluation of environmental heat stress on physical and hormonal parameters in murrah buffalo. *Journal of Animal Health and Production*, **7**(1), 21-24 (2019)
- Hahn, G.L., Mader, T.L. and Eigenberg, R.A. Perspectives on development of thermal indices for animal studies and management In Proceeding Symposium. Interactions between Climate and Animal Production. *EAAP Technical Series*, **7**(1),31-44 (2003)
- Pawar, H.N., Kumar, G.R. and Narang, R. Effect of heat stress on milk production and composition in Murrah buffaloes. *Journal of Buffalo Science*, **2**(2), 98-103 (2013).
- Dash, S., Chakravarty, A.K., Sah, V., Jamuna, V., Behera, R., Kashyap, N. and Deshmukh, B. Influence of temperature and humidity on pregnancy rate of Murrah buffaloes under subtropical climate. *Asian-Australasian Journal of Animal Sciences*, **28**(7), 943-950 (2015)
- Omran, F.I, Mahgoub, A.A.S. and Fooda, T.A. Effect of biological additive on physiological and production performance of buffaloes during cold weaves stress in mid of the Egyptian valley. *Egypt Journal of Agricultural Research*, **95**(4),1-12 (2017).
- Godde, C.M., Mason-D'Croz, D., Mayberry, D.E., Thornton, P.K. and Herrero, M. Impacts of climate change on the livestock food supply chain. *Global Food Security*, **28**(1),100488-100505 (2021).
- Jiménez-Castro, M.P., Buller, L.S., Sganzerla, W.G. and Forster-Carneiro, T. Bioenergy production from orange industrial waste a case study. *Biofuels, Bioproducts and Biorefining*, **14**(6),1239-1253(2020).
- USDA. Citrus World Markets and Trade U.S. Production and exports forecast down despite global gains. SW Washington DC, USA. Available (2021) at: <https://apps.fas.usda.gov/psdonline/circulars/citrus.pdf> (accessed on 13 August 2021).
- Ensminger, M.E., Oldfield, J.E. and Heinemann, W.W. Feeds and Nutrition, 2nd ed., The Ensminger Publishing Company Clovis. CA USA. (1990).
- Allam, S.M., El-Bedawy, T.M., Bakr, M.H. and Mahmoud, A.E.M. Effect of feeding dried orange pulp to lactating dairy cows on nutrients digestibility, blood constituents, plasma antioxidant biomarker, and pathogenic fecal bacteria. *Pakistan Journal of Zoology*, **52**(1),79-86 (2020).
- Bampidis, V.A. and Robinson, P.H. Citrus by-products as ruminant feeds. A review. *Animal Feed Science and Technology*, **128**(3-4),175-217 (2006).
- Santos, G.T, Lima, L.S, Schogor, A.L.B, Romero, J.V., De Marchi, F.E., Grande, P.A., Santos, N.W., Santos, F.S. and Kazam, R. Citrus pulp as a dietary source of antioxidants for lactating holstein cows fed highly polyunsaturated fatty acid diets. *Asian-Australasian Journal of Animal Sciences*, **27**(8),1104-1113 (2014).
- Luciano, G., Roscini, V., Mattioli, S., Ruggeri, S., Gravador, R.S., Natalello, A., Lanza, M., De Angelis, A. and Priolo, A. Vitamin E is the major contributor to the antioxidant capacity in lambs fed whole dried citrus pulp. *Animal*, **11**(4),411-417 (2017).
- Mahato, N., Sharma, K., Sinha, M. and Cho, M.H. Citrus waste derived nutra-pharmaceuticals for health benefits: Current trends and future perspectives. *Journal of Functional Foods*, **40**, 307-316 (2018).
- Heuzé, V., Tran, G., Hassoun, P. and Lebas, F. Citrus Pulp dried Feedipedia a programme by INRA CIRAD AFZ and FAO. 2017. available at: <https://www.feedipedia.org/node/680> (accessed on 27 August 2023)
- Cypriano, D.Z., Lopes da Silva, L. and Tasic, L. High value-added products from the orange juice industry. *Waste Management*, **79**(1), 71-78 (2018).
- López, M.C., Estellés, F., Moya, V.J. and Fernández, C. Use of dry citrus pulp or soybean hulls as a replacement for corn grain in energy and nitrogen partitioning, methane emissions and milk performance in lactating Murciano-Granadina goats. *Journal of Dairy Science*, **97**(12), 7821-7832 (2014).
- Ibáñez, C., Criscioni, P., Arriaga, H., Merino, P., Espino's, F.J. and Fernández, C. Murciano-Granadina goat performance and methane emission after replacing barley grain with fibrous by-products. *PLoS ONE*, **11**(1), 1-21(2016)
- Zedan, A.H. and Khattab, A.R. Effect of replacement corn grains by dried orange pulp in dairy goat rations on their productive performance. *Egyptian Journal of Nutrition and Feeds*, **20**(3),399-340 (2017).
- Guzmán, J.L, Perez-Ecija, A., Zarazaga, L.A., Martín-García, A.I., Horcada, A., Delgado-Pertíñez, M. Using dried orange pulp in the diet of dairy goats. effects on milk yield and composition and blood parameters of

- dams and growth performance and carcass quality of kids. *Animal*, **14**(10), 2212-2220 (2020).
27. Abdel Gawad, A.M., Hanafy, M.H., Mahmoud, A.E.M. and Al-Slibi, Y.H. Effect of tomato pomace citrus and beet pulp on productive performance and milk quality of egyptian buffaloes. *Pakistan Journal of Biological Sciences*, **23**(9), 1210-1219 (2020).
 28. El-Garhy, G.M., El-Mola, A.M.A., Azzaz, H.H. and Mousa, G.A. Influence of using pectinase enzymes in the ration on nutrient digestibility blood chemistry milk composition and economics of lactating buffaloes. *Journal of Animal Health and Production*, **8**(4), 158-167 (2020).
 29. Van Keulen, J.V. and Young, B.A. Evaluation of acid insoluble ash as a natural marker in ruminant digestibility studies. *Journal of Animal Science*, **44**(2), 282-287 (1977).
 30. AOAC. "Official Methods of Analysis" 14th ed., Published by the Association of Official Analytical Chemists Washington. DC, USA, pp. 684 (2000)
 31. Edwards, P. and Allan, G.L. Feeds and feeding for inland aquaculture in mekong region countries. ACIAR Technical Reports. No 56 P.136 (2004).
 32. McDonald, P., Edwards, R.A., Greenhalgh, F.D. and Morgan, C.A. Animal nutrition 5th Ed. Prentices Hall, London, UK. (1995).
 33. Reitman, A. and Frankel, S.A. Colorimetric method for the determination of serum glutamic oxalacetic and glutamic pyruvic transaminases. *American Journal of Clinical Pathology*, **28**, 56-63 (1957).
 34. Doumas, B.T., Weston, W. and Biggs, H.G. Albumin standards and the measurement of serum albumin with bromocresol green. *Clinica Chimica Acta*, **31**(1), 87-96 (1971).
 35. Cannon, D. C. In clinical chemistry - principles and techniques 2nd Ed Harper and Row Hagerstown MD. **27**(3), 411-421 (1974).
 36. Eisemann, J.H., Hammond, A.C., Bauman, D.E., Renolds, P.J., Maculchoen, S.N., Tyrrel, H.F. and Haaland, G.L. Effect of bovine growth hormone administration on metabolism of growing Hereford heifers: protein and lipid metabolism and plasma concentrations of metabolites and hormones. *The Journal of Nutrition*, **116**(12), 2504-2515 (1986).
 37. Koracevic, D., Koracevic, G., Djordjevic, V., Andrejevic, S. and Cosic, V., Method for the measurement of antioxidant activity in human fluids. *Journal of Clinical Pathology*, **54**(5), 356 -361 (2001).
 38. Mader, T.L., Davis, M.S. and Brown-Brandl, T. Environmental factors influencing heat stress in feedlot cattle. *Journal of Animal Science*, **84**(3), 712-719 (2006).
 39. Gaines, W.L. The energy basis of measuring energy milk in dairy cows. Bulletin University of Illinois (Urbana-Champaign campus). Agricultural Experiment Station. No.308 (1928) .
 40. Kemmerer, A.R., Harland, F.G. and Davi, R.N. Dried citrus by-products as feeds in the Rations of dairy cows and calves. *General Bulletin*, **218**(1), 1-7 (1948).
 41. Allam, S.M., Abou-ward, G.A., Tawila, M.A., Ali, M.A., Al-Hosseiny, H.M., Sayed, S.K. and El-Naggar, S.I. Nutritional and economical impact of using dried orange pulp as energy source in growing lambs rations . *Egyptian Journal of Nutrition and Feeds*, **14**(3), 337-347 (2011).
 42. NRC. National Research Council. Nutrient requirements of dairy cattle 7th rev Ed. National Academy Press. Washington, DC. USA. (2001).
 43. Ahooei, G.R., Foroughi, A.R., Tahmasbi, A.M., Shahdadi, A.R. and Vakili, R. Effects of different levels of dried citrus pulp and urea on performance of fattening male calves. *Journal of Animal and Veterinary Advances*, **10**(14), 1811- 1816 (2011).
 44. Gholizadeh, H. and Naserian, A.A. The effects of replacing dried citrus pulp with barley grain on the performance of Iranian Saanen kids. *Journal of Animal and Veterinary Advances*, **9**(15), 2053-2056 (2010).
 45. Ben-Ghedalia, D.E., Yosef, E., Miron, J. and Est, Y. The effects of starch- and pectin-rich diets on quantitative aspects of digestion in sheep. *Animal Feed Science and Technology*, **24**(4), 289-298 (1989).
 46. Gilaverte, S., Susin, L., Pires, A.V., Ferreira, E.M., Mendes, C.Q., Gentil, R.S., Biehl, M.V. and Rodrigues, G.H. Diet digestibility, ruminal parameters and performance of Santa Ines sheep fed dried citrus pulp and wet brewer grain. *Revista Brasileira de Zootecnia*, **40**(3), 639-647 (2011).
 47. Shdaifat, M.M., Al-Barakah, F.S., Kanana, A.Q. and Obeidat, B.S. The effect of feeding agricultural by-products on performance of lactating Awassi ewes. *Small Ruminant Research*, **113**(1), 11-14 (2013).
 48. Grigelmo-Miguel, N. and Martin-Belloso, O. Characterization of dietary fiber from orange juice extraction. *Food Research International*, **31**(5), 355-361 (1999).
 49. Belibasakis, N.G. and Tsigogianni, D. Effects of dried citrus pulp on milk yield, milk composition and blood components of dairy cows. *Animal Feed Science and Technology*, **60**(2), 87-92 (1996).
 50. Oni, A.O., Onwuka, C.F.I., Oduguwa, O.O., Onifade, O.S. and Arigbede, O.M. Utilization of citrus pulp based diets and Enterolobium cyclocarpum (JACQ. GRISEB) foliage by West African dwarf goats. *Livestock Science*, **117**(2-3), 184-191 (2008).
 51. Crawshaw, R. Co-product Feeds. Animal Feeds from the Food and Drinks Industries. Nottingham University Press. (2004).
 52. Bath, D.L., Dunbar, J.R., King, J.M., Berry, S.L., Leonard, R.O. and Olbrich, S.E. By-products and unusual feedstuffs in livestock rations. Western Regional Extension Publication. No. 39. USDA-ARS, Washington, DC, USA (1980).
 53. Lima, L.S., Santos, G.T., Schogor, A.B., Damasceno, J.C., Marchi, F.E., Santos, N.W., Santos, F.S. and Petit, H.V. Effect of abomasal or ruminal supplementation of citrus pulp and soybean oil on nutrient digestibility and ruminal fermentation of dairy cows. *Animal Feed Science and Technology*, **189**, 123-129 (2014).

54. Assis, A.J., Campos, J.M.S., Filho, S.C.V., Queiroz, A.C., Lana, R.P., Euclides, R.F., Neto, J.M., Magalhães, A.L.R. and Mendonça, S.S. Citrus pulp in diets for milking cows 1 intake of nutrients, milk production and composition. *Revista Brasileira de Zootecnia*, **33**(1), 242-250 (2004).
55. Wing, J.M. Effect of physical form and amount of citrus pulp on utilization of complete feeds for dairy cattle. *Journal of Dairy Science*, **58**(1) 63-66 (1975).
56. Miron, J., Yosef, E., Ben-Ghedalia, D., Chase, L.E., Bauman, D.E. and Solomon, R. Digestibility by dairy cows of monosaccharide constituents in total mixed rations containing citrus pulp. *Journal of Dairy Science*, **85**(1), 89-94 (2002).
57. LCI. Patterns of transit losses: Livestock Conservation, Inc. Omaha, Nebraska, USA. (1970).
58. Comin, A., Prandi, A., Peric, T., Corazzin, M., Dovier, S. and Bovolenta, S. Hair cortisol levels in dairy cows from winter housing to summer highland grazing. *Livestock Science*, **138**(1-3), 69-73 (2011).
59. Chaudhary, S.S., Singh, V.K., Upadhyay, R.C., Puri, G., Odedara, A.B. and Patel, P.A. Evaluation of physiological and biochemical responses in different seasons in Surti buffaloes. *Veterinary World*, **8**(6), 727-731 (2015).

أثر الاستبدال الجزئي لحبوب الذرة الصفراء بتفل البرتقال الجاف علي أداء الجاموس الحلاب خلال فصل الصيف

محمد حسن بكر¹، صباح علام¹، ايمان سيعاوى²، ابتهاج العدوي¹ وفاطمة ابو هاشم¹

¹ كلية الزراعة، جامعة القاهرة، مصر.

² المركز القومى للبحوث، مصر.

الملخص

في هذه الدراسة تم تقسيم سبعة وعشرون من الجاموس المصري الحلاب (متوسط وزن الجسم 600 ± 50 كجم، في الموسم الأول إلى الرابع من الحليب) عشوائيًا إلى ثلاث مجموعات مماثلة لدراسة الاستبدال الجزئي لحبوب الذرة الصفراء بتفل البرتقال المجفف خلال موسم الصيف. تم دراسة استهلاك العلف، هضم العناصر الغذائية، إنتاج اللبن وتركيبه، بعض مؤشرات الدم والكفاءة الاقتصادية. وتم استبدال حبوب الذرة الصفراء بتفل البرتقال عند مستويات 0 و 30 و 60% في R1 (المقارنة) و R2 و R3 على التوالي. لوحظ انخفاض في استهلاك المادة الجافة بمقدار 0.63 و 0.83 كجم في R2 و R3، على التوالي، مقارنة بمجموعة المقارنة. لم تتأثر معاملات هضم المادة الجافة، المادة العضوية، الدهن الخام المستخلص الخالي من الزيت. وبالمثل، لم تتأثر قيمة المركبات الكلية المهضومة (TDN) بالمعاملة ($P > 0.05$). بخلاف ذلك، انخفض معامل هضم البروتين وكذلك البروتين المهضوم بشكل ملحوظ عن طريق زيادة مستوى استبدال تفل البرتقال المجفف. وعلى العكس من ذلك، ازداد معامل هضم الألياف الخام في R2 و R3 مقارنة بـ R1 ($P < 0.05$). لم يكن هناك اختلاف كبير في مؤشرات الدم (مستويات الهيموجلوبين، البروتين الكلي، الألبومين، ALT و AST، الدهون الكلية، الكوليسترول، الدهون الثلاثية، T3، T4، الكورتيزول، إجمالي القدرة المضادة للأكسدة (TAC) و IgG؛ $P > 0.05$). أيضًا، لم يتم العثور على فروق ذات دلالة إحصائية بين المجموعات التجريبية، لا في متوسط إنتاج اللبن اليومي ولا في اللبن معدل محتوى الدهون (FCM). وكذلك لم تصل الفروق في محتوى اللبن من بروتين والدهن واللاكتوز والمواد الصلبة الكلية والرماد إلى مستوى معنوي بين المجموعات التجريبية ($P > 0.05$). علاوة على ذلك، انخفضت تكلفة مخلوط العلف المركز (جنيه/طن) بنسبة 6.6% و 13.2% في R2 و R3 على التوالي، مقارنة بـ R1 مما يؤثر إيجاباً على تكاليف التغذية لإنتاج 1 كجم حليب.

الكلمات الدالة: تغذية الجاموس، تفل البرتقال المجفف، الإجهاد الحراري، الكفاءة الغذائية، الكفاءة الاقتصادية.