



Egyptian Mathematical Society
Journal of the Egyptian Mathematical Society

www.etms-eg.org
 www.elsevier.com/locate/joems



ORIGINAL ARTICLE

Solitary wave solutions Zakharov–Kuznetsov–Benjamin–Bona–Mahony (ZK–BBM) equation



Ayyaz Ali, Muhammad Asad Iqbal, Syed Tauseef Mohyud-Din *

Department of Mathematics, Faculty of Sciences, HITEC University, Taxila, Pakistan

Received 17 July 2014; revised 17 October 2014; accepted 29 October 2014
 Available online 20 February 2015

KEYWORDS

Exp-function method;
 Solitary solution;
 Zakharov–Kuznetsov–Benjamin–Bona–Mahony equation

Abstract In this paper, exp-function method is used to construct generalized solitary solutions of the Zakharov–Kuznetsov–Benjamin–Bona–Mahony equation. It is shown that the exp-function method, with the help of symbolic computation, provides a straightforward and powerful mathematical tool to solve such nonlinear equations. The performance of the method is reliable, efficient and it gives useful exact solutions.

MATHEMATICS SUBJECT CLASSIFICATION: 35Cxx; 35Dxx; 35Qxx

© 2015 Production and hosting by Elsevier B.V. on behalf of Egyptian Mathematical Society.

1. Introduction

The nonlinear partial differential equations play a pivotal role in the mathematical modeling of diversified physical phenomena. Finding exact solutions [1–27] of nonlinear evolution equations (NLEEs) has become one of the most exciting and extremely active areas of research investigation. The investigation of exact travelling wave solutions to nonlinear evolution equations plays a vital role in the study of nonlinear physical phenomena. The wave phenomena are observed in fluid dynamics, plasma, elastic media, optical fibers, etc. Many effective methods have been presented such as variational iteration method [1], homotopy perturbation method [2], Adomian's decomposition method [3]

and others [4]. The aim of the present paper was to extend the exp-function method to find new solitary solutions and periodic solutions for Zakharov–Kuznetsov–Benjamin–Bona–Mahony equation. Recently, Shakeel and Mohyud-Din [5] used the (G'/G) -expansion method to obtain solutions of Zakharov–Kuznetsov–Benjamin–Bona–Mahony equation.

2. Exp-function method

Consider the general nonlinear partial differential equation of the type

$$P(u, u_t, u_x, u_{tt}, u_{xx}, u_{xxx}, \dots) = 0 \quad (1)$$

Using a transformation

$$\eta = kx + \omega t, \quad (2)$$

where k and ω are constants, we can rewrite Eq. (1) in the following nonlinear ODE,

$$Q(u, u', u'', u''', u'''' , \dots) = 0 \quad (3)$$

* Corresponding author.

E-mail address: syedtauseefs@hotmail.com (S.T. Mohyud-Din).

Peer review under responsibility of Egyptian Mathematical Society.



Production and hosting by Elsevier

where the prime denotes derivative with respect to η .

According to the exp-function method, which was developed by He and Wu, we assume that the wave solutions can be expressed in the following form

$$u(\eta) = \frac{\sum_{n=-c}^d a_n \exp(m\eta)}{\sum_{m=-p}^q b_m \exp(m\eta)} \quad (4)$$

where p, q, c and d are positive integers which are known to be further determined, a_n and b_m are unknown constants. We can rewrite Eq. (4) in the following equivalent form

$$u(\eta) = \frac{a_c \exp(c\eta) + \dots + a_{-d} \exp(-d\eta)}{b_p \exp(p\eta) + \dots + b_{-q} \exp(-q\eta)} \quad (5)$$

To determine the value of c and p , we balance the linear term of highest order of Eq. (4) with the highest order nonlinear term. Similarly, to determine the value of d and q , we balance the linear term of lowest order of Eq. (3) with lowest order nonlinear term.

3. Solution procedure

3.1. Zakharov–Kuznetsov–Benjamin–Bona–Mahony equation

Consider the following Zakharov–Kuznetsov–Benjamin–Bona–Mahony (ZK–BBM) equation

$$u_t + u_x - 2auu_x - bu_{xx} = 0. \quad (6)$$

Introducing a transformation as $\eta = kx + \omega t$ we can convert Eq. (6) into ordinary differential equations

$$\omega u' + ku' - 2akuu' - bk^2 u u''' = 0, \quad (7)$$

where the prime denotes the derivative with respect to η . The trial solution of the Eq. (7) can be expressed as follows,

$$u(\eta) = \frac{a_c \exp(c\eta) + \dots + a_{-d} \exp(-d\eta)}{b_p \exp(p\eta) + \dots + b_{-q} \exp(-q\eta)}.$$

To determine the value of c and p we balance the linear term of highest order of Eq. (7) with the highest order nonlinear term and to determine the value of d and q we balance the linear term of lowest order of Eq. (7) with the lowest order nonlinear term. We obtain $p = c$ and $d = q$.

3.1.1. Case 3.1.1

we can freely choose the values of c and p , we balance the linear term of highest order of Eq. (7) with the highest order nonlinear term, but we will illustrate that the final solution does not strongly depend upon the choice of values of c and d . For simplicity, we set $p = c = 1$ and $d = q = 1$ Eq. (5) reduces to

$$u(\eta) = \frac{a_1 \exp(\eta) + a_0 + a_{-1} \exp(-\eta)}{b_1 \exp(\eta) + a_0 + b_{-1} \exp(-\eta)} \quad (8)$$

Substituting Eq. (8) into Eq. (7), we have

$$\begin{aligned} C_1 = & \omega a_1 b_0^2 + k a_1 b_0^2 - \alpha k a_1^2 b_{-1} + 2\omega a_0 b_1 b_0 + 2k a_0 b_1 b_0 \\ & - \alpha k a_0^2 b_1 + k a_{-1} b_1^2 + \omega a_{-1} b_1^2 - 2\alpha k a_{-1} a_1 b_1 - 4\beta \omega k^2 a_{-1} b_1^2 \\ & + \beta \omega k^2 a_0 b_1 b_0 - 2\alpha k a_0 a_1 b_0 - \beta \omega k^2 a_1 b_0^2 + 4\beta \omega k^2 a_1 b_1 b_{-1} \\ & + 2\omega a_1 b_1 b_{-1} + 2k a_1 b_1 b_{-1} \end{aligned}$$

$$\begin{aligned} C_{-1} = & \omega a_{-1} b_0^2 + k a_{-1} b_0^2 - \alpha k a_{-1}^2 b_{-1} + 2\omega a_0 b_{-1} b_0 + 2k a_0 b_{-1} b_0 \\ & - \alpha k a_0^2 b_{-1} + k a_1 b_{-1}^2 + \omega a_1 b_{-1}^2 - 2\alpha k a_{-1} a_1 b_{-1} \\ & - 4\beta \omega k^2 a_1 b_{-1}^2 + \beta \omega k^2 a_0 b_{-1} b_0 - 2\alpha k a_0 a_{-1} b_0 \\ & - \beta \omega k^2 a_{-1} b_0^2 + 4\beta \omega k^2 a_{-1} b_1 b_{-1} + 2\omega a_{-1} b_1 b_{-1} \\ & + 2k a_{-1} b_1 b_{-1} \end{aligned}$$

$$\begin{aligned} C_2 = & -\alpha k a_1^2 b_0 + 2\omega a_1 b_1 b_0 + 2k a_1 b_1 b_0 + k a_0 b_1^2 + \omega a_0 b_1^2 \\ & - \beta \omega k^2 a_0 b_1^2 + \beta \omega k^2 a_1 b_1 b_0 - 2\alpha k a_0 a_1 b_1 \end{aligned}$$

$$\begin{aligned} C_{-2} = & -\alpha k a_{-1}^2 b_0 + 2\omega a_{-1} b_{-1} b_0 + 2k a_{-1} b_{-1} b_0 + k a_0 b_{-1}^2 + \omega a_0 b_{-1}^2 \\ & - \beta \omega k^2 a_0 b_{-1}^2 + \beta \omega k^2 a_{-1} b_{-1} b_0 - 2\alpha k a_0 a_{-1} b_{-1} \end{aligned}$$

$$C_3 = -\alpha k a_1^2 b_1 + k a_1 b_1^2 + \omega a_1 b_1^2$$

$$C_{-3} = -\alpha k a_{-1}^2 b_{-1} + k a_{-1} b_{-1}^2 + \omega a_{-1} b_{-1}^2$$

$$\begin{aligned} C_0 = & -\alpha k a_0^2 b_0 + 2\omega a_{-1} b_1 b_0 + 2\omega a_0 b_1 b_{-1} + 2\omega a_1 b_{-1} b_0 \\ & + 2k a_{-1} b_1 b_0 + 2k a_0 b_1 b_{-1} + 2k a_1 b_{-1} b_0 + k a_0 b_0^2 \\ & + \omega a_0 b_0^2 - 3\beta \omega k^2 a_1 b_{-1} b_0 - 3\beta \omega k^2 a_{-1} b_1 b_0 \\ & + 6\beta \omega k^2 a_0 b_1 b_{-1} - 2\alpha k a_0 a_{-1} b_1 - 2\alpha k a_1 a_{-1} b_0 \\ & - 2\alpha k a_0 a_1 b_{-1} \end{aligned} \quad (9)$$

are constants obtained by Maple 15. Equating the coefficients of $\exp(m\eta)$ to be zero, we obtain

$$\{C_3 = 0, C_2 = 0, C_1 = 0, C_{-3} = 0, C_{-2} = 0, C_{-1} = 0, C_0 = 0\} \quad (10)$$

Solution of (10) we have following solution sets satisfy the given equation,

3.1.1.1. 1st Solution set.

$$\left\{ \begin{aligned} \omega = \frac{k}{-1+k^2}, \quad a_{-1} = 0, \quad a_0 = a_0, \quad a_1 = 0, \quad b_{-1} = \frac{1}{36} \frac{a_0^2(1-2k^2+k^4)}{k^4 b_1}, \\ b_0 = \frac{1}{3} \frac{a_0(-1+k^2)}{k^2}, \quad b_1 = b_1 \end{aligned} \right\} \quad (11)$$




Figure 3.1a Singular Kink wave solution of Eq. (12).

We therefore, obtained the following generalized solitary solution $u(x, t)$ of Eq. (6) (see Fig. 3.1a)

$$u(x, t) = \frac{36a_0k^4b_1}{\left(36b_1^2e^{-\frac{k(-x+xk^2+t)}{-1+k^2}}k^4 + \left\{a_0(-1+k^2)e^{-\frac{k(-x+xk^2+t)}{-1+k^2}} + 12k^2b_1\right\}a_0(-1+k^2)\right)} \quad (12)$$

3.1.1.2. 2nd Solution set.

$$\left\{ \begin{array}{l} \omega = \frac{k}{-1+k^2}, \quad a_{-1} = \frac{1}{16} \frac{a_0^2(1+k^2)}{k^2b_1}, \quad a_0 = a_0, \quad a_1 = \frac{k^2b_1}{1+k^2}, \\ b_{-1} = \frac{1}{16} \frac{a_0^2(1+2k^2+k^4)}{k^4b_1}, \quad b_0 = -\frac{1}{2} \frac{a_0(1+k^2)}{k^2}, \quad b_1 = b_1 \end{array} \right\} \quad (13)$$

$$u(x, t) = \frac{(36a_1e^{-\frac{k(-x+xk^2+t)}{(k+1)(k-1)}}k^4b_0)}{\left(a_1^2(k-1)^2(k+1)^2e^{-\frac{2k(-x+xk^2+t)}{(k-1)(k+1)}} + 36k^2b_0\left\{a_1\left(-\frac{1}{3} + \frac{1}{3}k^2\right)e^{-\frac{k(-x+xk^2+t)}{(k-1)(k+1)}} + k^2b_0\right\}\right)} \quad (19)$$

We therefore, obtained the following generalized solitary solution $u(x, t)$ of Eq. (6) (see Fig. 3.1b)

$$u(x, t) = \frac{36a_0k^4b_1}{\left(36b_1^2e^{-\frac{k(-x+xk^2+t)}{-1+k^2}}k^4 + a_0\left\{a_0(-1+k^2)e^{-\frac{k(-x+xk^2+t)}{-1+k^2}} + 12k^2b_1\right\}(-1+k^2)\right)} \quad (14)$$

3.1.2. Case 3.1.2

If $p = c = 2$ and $q = d = 2$ then Eq. (5) reduces to

$$u(\eta) = \frac{a_2 \exp(2\eta) + a_1 \exp(\eta) + a_0 + a_{-1} \exp(-\eta)}{b_2 \exp(2\eta) + b_1 \exp(\eta) + b_0 + b_{-1} \exp(-\eta)} \quad (15)$$

Proceeding as before, we obtain

3.1.2.1. 1st Solution set.

$$\left\{ \begin{array}{l} \omega = \frac{k}{-1+k^2}, \quad a_{-2} = 0, \quad a_{-1} = 0, \quad a_0 = 0, \quad a_1 = a_1, \quad a_2 = 0, \quad b_{-2} = 0, \\ b_{-1} = 0, \quad b_0 = b_0, \quad b_1 = \frac{1}{3} \frac{a_1(-1+k^2)}{k^2}, \quad b_2 = \frac{1}{36} \frac{a_1^2(1-2k^2+k^4)}{k^4b_0} \end{array} \right\} \quad (16)$$

Hence we get the generalized solitary wave solution of Eq. (6) as follows (see Fig. 3.1c)

$$u(x, t) = \frac{(36a_1e^{-\frac{k(-x+xk^2+t)}{(k+1)(k-1)}}k^4b_0)}{\left(a_1^2(k-1)^2(k+1)^2e^{-\frac{2k(-x+xk^2+t)}{(k-1)(k+1)}} + 36k^2b_0\left\{a_1\left(-\frac{1}{3} + \frac{1}{3}k^2\right)e^{-\frac{k(-x+xk^2+t)}{(k-1)(k+1)}} + k^2b_0\right\}\right)} \quad (17)$$

3.1.2.2. 2nd Solution set. Therefore, proceeding as before, we obtain

$$\left\{ \begin{array}{l} \omega = -\frac{k}{1+k^2}, \quad a_{-2} = 0, \quad a_{-1} = 0, \quad a_0 = a_0, \quad a_1 = -\frac{2bk^2}{1+k^2}, \\ a_2 = \frac{1}{4} \frac{k^4b_1^2}{a_0(1+k^2)^2}, \quad b_{-2} = 0, \quad b_{-1} = 0, \quad b_0 = \frac{a_0(1+k^2)}{k^2}, \\ b_1 = b_1, \quad b_2 = \frac{1}{4} \frac{b_1^2}{a_0(1+k^2)} \end{array} \right\} \quad (18)$$

Hence we get the generalized solitary wave solution of Eq. (6) as follows (see Fig. 3.1d)

3.1.2.3. 3rd Solution set.

$$\left\{ \begin{array}{l} \omega = \frac{k}{-1+k^2}, \quad a_{-2} = 0, \quad a_{-1} = 0, \quad a_0 = a_0, \quad a_1 = a_1, \\ a_2 = 0, \quad b_{-2} = 0, \quad b_2 = b_2, \\ b_0 = \frac{\left(a_1^2 - 2a_1^2k^2 + a_1^2k^4 - 12k^2a_0b_2 + 12a_0b_2k^4\right)}{36b_2k^4}, \quad b_{-1} = \frac{1}{36} \frac{a_0a_1(1-2k^2+k^4)}{b_2k^4}, \\ b_2 = b_2, \quad b_1 = \frac{1}{3} \frac{-a_1^2 + a_1^2k^2 + 3k^2a_0b_2}{a_1k^2} \end{array} \right\} \quad (20)$$

We therefore, obtained the following generalized solitary solution $u(x, t)$ of Eq. (6) (see Fig. 3.1e)

$$u(x, t) = \frac{(36a_1e^{-\frac{k(-x+xk^2+t)}{(k+1)(k-1)}}k^4b_0)}{\left(a_1^2(k-1)^2(k+1)^2e^{-\frac{2k(-x+xk^2+t)}{(k-1)(k+1)}} + 36k^2b_0\left\{a_1\left(-\frac{1}{3} + \frac{1}{3}k^2\right)e^{-\frac{k(-x+xk^2+t)}{(k-1)(k+1)}} + k^2b_0\right\}\right)} \quad (21)$$

3.1.2.4. 4th Solution set.

$$\left\{ \begin{array}{l} \omega = -\frac{k}{1+k^2}, \quad a_{-2} = 0, \quad a_{-1} = 0, \quad a_0 = \frac{b_0k^2}{1+k^2}, \quad a_1 = a_1, \quad a_2 = \frac{1}{16} \frac{a_1^2(1+k^2)}{b_0k^2}, \\ b_{-2} = 0, \quad b_{-1} = 0, \quad b_0 = b_0, \quad b_1 = -\frac{1}{2} \frac{a_1(1+k^2)}{k^2}, \quad b_2 = \frac{1}{16} \frac{a_1^2}{b_0k^4} \end{array} \right\} \quad (22)$$

We therefore, obtained the following generalized solitary solution $u(x, t)$ of Eq. (6) (see Fig. 3.1f)

$$u(x, t) = \frac{(36a_1 e^{-\frac{k(-x+xk^2+t)}{(k+1)(k-1)}} k^4 b_0)}{\left(a_1^2 (k-1)^2 (k+1)^2 e^{-\frac{2k(-x+xk^2+t)}{(k-1)(k+1)}} + 36k^2 b_0 \left\{ a_1 \left(-\frac{1}{3} + \frac{1}{3} k^2 \right) e^{-\frac{k(-x+xk^2+t)}{(k-1)(k+1)}} + k^2 b_0 \right\} \right)} \tag{23}$$




Figure 3.1b Singular Kink wave solution of Eq. (14).




Figure 3.1e Singular Kink wave solution of Eq. (21).




Figure 3.1c Singular Kink wave solution of Eq. (17).




Figure 3.1f Singular Kink wave solution of Eq. (23).




Figure 3.1d Singular Kink wave solution of Eq. (19).

In both cases, for different choices of c, p, d and q we do not get the same solitary solutions which clearly illustrate that final solution does not strongly depends upon these parameters.

4. Conclusion

The exp-function method has been used to obtain generalized solitary solutions of the Zakharov–Kuznetsov–Benjamin–Bona–Mahony equation. This method can also be extended to other nonlinear evolution equations (NLEEs). The exp-function method is a promising and powerful method for problems arising in mathematical physics. It is also observed that the exp-function method is useful for finding solutions of a wide class of nonlinear problems.

Acknowledgment

The authors are highly grateful to the unknown referee for valuable comments which really helped to improve the quality of work.

References

- [1] S.T. Mohyud-Din, A. Yldrm, S.A. Sezer, M. Usman, Modified variational iteration method for free-convective boundary-layer equation using padé approximation, *Math. Prob. Eng.* 2010 (2010) 1–11 (Article ID 318298).
- [2] B. Raftari, A. Yldrm, The application of homotopy perturbation method for MHD flows of UCM fluids above porous stretching sheets, *Comput. Math. Appl.* 59 (10) (2010) 3328–3337.
- [3] T. Özip, A. Yldrm, Comparison between Adomian's method and He's homotopy perturbation method, *Comput. Math. Appl.* 56 (5) (2008) 1216–1224.
- [4] A. Yldrm, S.T. Mohyud-Din, D.H. Zhang, Analytical solutions to the pulsed Klein–Gordon equation using modified variational iteration method (MVIM) and Boubaker polynomials expansion scheme (BPES), *Comput. Math. Appl.* 59 (8) (2010) 2473–2477.
- [5] M. Shakeel, S.T. Mohyud-Dinusing, (G'/G) -expansion method and its application to the Zakharov–Kuznetsov–Benjamin–Bona–Mahony equation, *J. Assoc. Arab Uni. Basic Appl. Sci.* (2014).
- [6] S.T. Mohyud-Din, M.A. Noor, K.I. Noor, Some relatively new techniques for nonlinear problems, *Math. Prob. Eng.* 2009 (2009), <http://dx.doi.org/10.1155/2009/234849> (Article ID 234849).
- [7] S.T. Mohyud-Din, Solution of nonlinear differential equations by exp-function method, *World Appl. Sci. J.* 7 (2009) 116–147.
- [8] J.H. He, X.H. Wu, Exp-function method for nonlinear wave equations, *Chaos Solit. Fract.* 30 (3) (2006) 700–708.
- [9] T.B. Benjamin, J.L. Bona, J. Mahony, Model equations for waves in nonlinear dispersive systems, *J. Philos. Trans. R. Soc. Lond.* 227 (1978) 47–78.
- [10] S. Zhang, Exp-function method for solving Maccari's system, *Phys. Lett. A* 371 (2007) 65–71.
- [11] X.H. Wu, J.H. He, Solitary solutions, periodic solutions and compacton-like solutions using the exp-function method, *Comput. Math. Appl.* 54 (2007) 966–986.
- [12] S. Zhu, Discrete $(2 + 1)$ -dimensional Toda lattice equation via exp-function method, *Phys. Lett. A* 372 (2008) 654–657.
- [13] H. Naher, F.A. Abdullah, M.A. Akbar, The Exp-function method for new exact solutions of the nonlinear partial differential equations, *Int. J. Phys. Sci.* 6 (29) (2011) 6706–6716.
- [14] J.H. He, M.A. Abdou, New periodic solutions for nonlinear evolution equations using exp-function method, *Chaos Solit. Fract.* 34 (2007) 1421–1429.
- [15] A.M. Wazwaz, The extended tanh-method for new compact and non-compact solutions for the KP–BBM and the ZK–BBM equations, *Chaos Solit. Fract.* 38 (2008) 1505–1516.
- [16] H. Naher, F.A. Abdullah, M.A. Akbar, New travelling wave solutions of the higher dimensional nonlinear partial differential equation by the exp-function method, *J. Appl. Math.* 14 (2012).
- [17] M.A. Noor, S.T. Mohyud-Din, A. Waheed, Exp-function method for solving Kuramoto–Sivashinsky and Boussinesq equations, *J. Appl. Math. Comput.* 29 (2008) 1–13, <http://dx.doi.org/10.1007/s12190-008-0083-y>.
- [18] S.D. Zhu, Exp-function method for the hybrid–lattice system, *Int. J. Nonlin. Sci. Numer. Simul.* 8 (2007) 461–464.
- [19] X.H. Wu, J.H. He, Solitary solutions, periodic solutions and compacton like solutions using the exp-function method, *Comput. Math. Appl.* 54 (2007) 966–986.
- [20] M.A. Abdou, A.A. Soliman, S.T. Basyony, New application of exp-function method for improved Boussinesq equation, *Phys. Lett. A* 369 (2007) 469–475.
- [21] A. Bekir, A. Boz, Exact solutions for nonlinear evolution equation using Exp-function method, *Phys. Lett. A* 372 (2008) 1619–1625.
- [22] X.H. Wu, J.H. He, Exp-function method and its application to nonlinear equations, *Chaos Solit. Fract.* 38 (2008) 903–910.
- [23] H. Naher, F.A. Abdullah, The improved (G'/G) -expansion method to the $(2 + 1)$ -dimensional breaking soliton equation, *J. Comput. Anal. Appl.* 16 (2) (2014) 220–235.
- [24] H. Naher, F.A. Abdullah, New generalized and improved (G'/G) -expansion method for nonlinear evolution equations in mathematical physics, *J. Egypt. Math. Soc.* 22 (3) (2014) 390–395.
- [25] H. Naher, F.A. Abdullah, New approach of (G/G') -expansion method and new approach of generalized (G/G') -expansion method for nonlinear evolution equation, *AIP Adv.* 3 (2013) 032116.
- [26] H. Naher, F.A. Abdullah, The basic (G'/G) -expansion method for the fourth order Boussinesq equation, *Comp. Sci. Commun.* 3 (10) (2012) 1144–1152.
- [27] H. Naher, F.A. Abdullah, The modified Benjamin–Bona–Mahony equation via the extended generalized Riccati equation mapping method, *Appl. Math. Sci.* 6 (111) (2012) 5495–5512.