



Original Article

# New stability and boundedness results to Volterra integro-differential equations with delay



Cemil Tunç\*

Department of Mathematics, Faculty of Sciences, Yüzüncü Yıl University, 65080 Van, Turkey

Received 1 June 2015; accepted 13 August 2015

Available online 9 September 2015

## Keywords

Non-linear;  
Volterra integro-differential equation;  
Stability;  
Boundedness;  
Lyapunov functional

**Abstract** In this paper, we consider a certain non-linear Volterra integro-differential equations with delay. We study stability and boundedness of solutions. The technique of proof involves defining suitable Lyapunov functionals. Our results improve and extend the results obtained in literature.

**2010 Mathematics Subject Classifications:** 34D05; 34K20; 45J05

Copyright 2015, Egyptian Mathematical Society. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 1. Introduction

In the last years, the qualitative properties of Volterra integro-differential equations without delay have been discussed by many researches. In particular, the reader can refer to the papers of Becker [1], Burton [2,3], Burton and Mahfoud [4,5], Diamandescu [6], Hara et al. [7], Miller [8], Staffans [9], Tunç [10], Vanualailai and Nakagiri [11] and the books of Burton [12], Corduneanu [13], Gripenberg et al. [14] and the references cited therein for some works done on qualitative properties of various Volterra integro-differential equations without delay. An

important tool to discuss the qualitative properties of solutions of ordinary and functional differential equations and integro-differential equations is the Lyapunov's direct method. Theoretically this method is very appealing, and there are numerous applications where it is natural to use it. The key requirement of the method is to find a positive definite function or functional which is non-increasing along solutions.

However, it is a quite difficult task to find a suitable Lyapunov function or functional for a non-linear ordinary or functional differential equation and a non-linear functional Volterra integro-differential equation. The situation becomes more difficult when we replace an ordinary or a functional differential equation with a functional integro-differential equation. By this time, the construction of Lyapunov functions and functionals for non-linear differential and integro-differential systems remains as an open problem in the literature. Besides, in the literature, there are a few papers on the qualitative behaviors of Volterra integro-differential equations with delay. See, for ex-

\* Tel.: +5375517167.

E-mail address: [cemtunc@yahoo.com](mailto:cemtunc@yahoo.com), [tunccemil@gmail.com](mailto:tunccemil@gmail.com)

Peer review under responsibility of Egyptian Mathematical Society.



Production and hosting by Elsevier

ample, the recent papers of Adıvar and Raffoul [15], Graef and Tunc [16], Raffoul [17] and Raffoul and Unal [18].

In 2003, Vanualailai and Nakagiri [11] considered the nonlinear Volterra integro-differential equation without delay,

$$\frac{d}{dt}[x(t)] = A(t)f(x(t)) + \int_0^t B(t, s)g(x(s))ds, \tag{1}$$

where  $t \geq 0, x \in \mathfrak{R}, A(t) : [0, \infty) \rightarrow (-\infty, 0), f, g : \mathfrak{R} \rightarrow \mathfrak{R}$  are continuous functions, and  $B(t, s)$  is a continuous function for  $0 \leq s \leq t < \infty$ . Vanualailai and Nakagiri [11] studied the stability of solutions of Eq. (1) by defining a suitable Lyapunov functional.

In this paper, we consider the nonlinear the Volterra integro-differential equation with delay

$$x'(t) = -a(t)f(x(t)) + \int_{t-\tau}^t B(t, s)g(x(s))ds + p(t), \tag{2}$$

where  $t \geq 0, \tau$  is a positive constant, fixed delay,  $x \in \mathfrak{R}, a(t) : [0, \infty) \rightarrow (0, \infty), p : [0, \infty) \rightarrow \mathfrak{R}, f, g : \mathfrak{R} \rightarrow \mathfrak{R}$  are continuous functions with  $f(0) = g(0) = 0$ , and  $B(t, s)$  is a continuous function for  $0 \leq s \leq t < \infty$ .

We investigate the stability of zero solution and boundedness of all solutions of Eq. (2) by defining new suitable Lyapunov functionals, when  $p(t) \equiv 0$  and  $p(t) \neq 0$ , respectively.

It follows that Vanualailai and Nakagiri [11] considered a Volterra integro-differential equation without delay. However, in this paper, we consider a Volterra integro-differential equation with delay. Besides, Vanualailai and Nakagiri [11] discussed the stability of the zero solution of Eq. (1). However, beside the stability of zero solution, we also discuss the boundedness of solutions of Eq. (2), when  $p(t) \equiv 0$  and  $p(t) \neq 0$ , respectively. Further, Eq. (2) includes and extends the equations discussed by Vanualailai and Nakagiri [11], when  $\tau = 0$ .

We give some basic information related Eq. (2).

We use the following notation throughout this paper.

For any  $t_0 \geq 0$  and initial function  $\varphi \in [t_0 - \tau, t_0]$ , let  $x(t) = x(t, t_0, \varphi)$  denote the solution of Eq. (2) on  $[t_0 - \tau, \infty)$  such that  $x(t) = \varphi(t)$  on  $\varphi \in [t_0 - \tau, t_0]$ .

Let  $C[t_0, t_1]$  and  $C[t_0, \infty)$  denote the set of all continuous real-valued functions on  $[t_0, t_1]$  and  $[t_0, \infty)$ , respectively.

For  $\varphi \in C[0, t_0], |\varphi|_{t_0} := \sup\{|\varphi(t)| : 0 \leq t \leq t_0\}$ .

**Definition.** The zero solution of Eq. (2) is stable if for each  $\varepsilon > 0$  and each  $t_0 \geq 0$ , there exists a  $\delta = \delta(\varepsilon, t_0) > 0$  such that  $\varphi \in C[0, t_0]$  with  $|\varphi(t)|_{t_0} < \delta$  implies that  $|x(t, t_0, \varphi)| < \varepsilon$  for all  $t \geq t_0$ .

Let  $p(t) = 0$  in Eq. (1).

The following theorem is need for the stability result of this theorem.

**Theorem 1** (Driver [19]). *If there exists a functional  $V(t, \varphi(\cdot))$ , defined whenever  $t \geq t_0 \geq 0$  and  $\varphi \in C([0, t], \mathfrak{R})$ , such that*

- (i)  $V(t, 0) \equiv 0, V$  is continuous in  $t$  and locally Lipschitz in  $\varphi$ ,
- (ii)  $V(t, \varphi(\cdot)) \geq W(|\varphi(t)|), W : [0, \infty) \rightarrow [0, \infty)$  is a continuous function with  $W(0) = 0, W(r) > 0$  if  $r > 0$ , and  $W$  strictly increasing (positive definiteness), and
- (iii)  $V'(t, \varphi(\cdot)) \leq 0$ ,

then the zero solution of Eq. (2) is stable, and

$$V(t, \varphi(\cdot)) = V(t, \varphi(s) : 0 \leq s \leq t)$$

is called a Lyapunov functional for Eq. (2).

## 2. The main results

We state some assumptions on the functions that are appearing in Eq. (2).

### A. Assumptions

- (A1) There exist positive constants  $\alpha, m, J, M$  and  $N$  such that  $f(0) = 0, g(0) = 0, g^2(x) \leq m^2 f^2(x)$  if  $|x| \leq M, \alpha > 4$  such that  $4x^2 \leq (\alpha - 4)f^2(x)$  if  $|x| \leq N$ .
- (A2)  $a(t) > 0$  for  $t \geq 0, B(t, s)$  is continuous for  $0 \leq s \leq t < \infty,$   
 $J \geq 1, \frac{1}{4a(t)} \int_{t-\tau}^t |B(t, s)|ds < \frac{1}{J}$  for every  $t \geq s - \tau \geq 0,$   
 $\int_{t-\tau}^\infty |B(u + \tau, s)|du$  is defined and continuous for  $0 \leq s - \tau \leq t < \infty,$   
 $a(t) - k \int_{t-\tau}^\infty |B(u + \tau, t)|du \geq 0$  for every  $t \geq s - \tau \geq 0.$

For the case  $p(t) = 0$  in Eq. (2), we have the following result.

**Theorem 2.** *Assume conditions (A1) and (A2) hold. If  $k = \frac{m^2(1+\alpha)}{J}$ , then the zero solution of Eq. (2) is stable.*

**Proof.** We introduce a functional  $V_0 = V_0(t) = V_0(t, x(t))$  defined by

$$V_0 = \frac{1}{2}x^2 + \sqrt{\alpha} \int_0^x \sqrt{f(u)}udu + \frac{1}{2}\alpha \int_0^x f(u)du + k \int_0^t \int_{t-\tau}^\infty |B(u + \tau, s)|du f^2(x(s))ds, \tag{3}$$

where  $k$  is a positive constant to be determined later in the proof.  $\square$

It is clear that the functional  $V_0$  is positive definite.

Differentiating the functional  $V_0$  with respect to  $t$ , we obtain from (3) that

$$V_0' = xx' + \sqrt{\alpha}\sqrt{f(x)}xx' + \frac{1}{2}\alpha f(x)x' + k \int_{t-\tau}^\infty |B(u + \tau, t)|du f^2(x) - k \int_0^t |B(t, s)|f^2(x(s))ds. \tag{4}$$

Then, it is clear that

$$\begin{aligned} xx' &= -a(t)xf(x) + x \int_{t-\tau}^t B(t, s)g(x(s))ds \\ &= -a(t)xf(x) - \left[ \sqrt{a(t)}x - \frac{1}{2\sqrt{a(t)}} \int_{t-\tau}^t B(t, s)g(x(s))ds \right]^2 \\ &\quad + a(t)x^2 + \frac{1}{4a(t)} \left[ \int_{t-\tau}^t B(t, s)g(x(s))ds \right]^2 \\ &\leq -a(t)xf(x) + a(t)x^2 + \frac{1}{4a(t)} \left[ \int_{t-\tau}^t B(t, s)g(x(s))ds \right]^2 \\ &\leq -a(t)xf(x) + a(t)x^2 \\ &\quad + \frac{1}{4a(t)} \int_{t-\tau}^t |B(t, s)|ds \int_{t-\tau}^t |B(t, s)|g^2(x(s))ds \\ &\leq -a(t)xf(x) + a(t)x^2 \end{aligned}$$

$$\begin{aligned} & + \frac{m^2}{4a(t)} \int_{t-\tau}^t |B(t, s)| ds \int_{t-\tau}^t |B(t, s)| f^2(x(s)) ds \\ & \leq -a(t)x f(x) + \frac{1}{4} \alpha a(t) f^2(x) - a(t) f^2(x) \\ & + \frac{m^2}{J} \int_{t-\tau}^t |B(t, s)| f^2(x(s)) ds \end{aligned}$$

by the assumption of [Theorem 2](#) and the Schwarz inequality, that is,

$$(A2) \text{ and } [\int_{t-\tau}^t B(t, s)g(x(s))ds]^2 \leq \int_{t-\tau}^t |B(t, s)| ds \int_{t-\tau}^t |B(t, s)| g^2(x(s)) ds.$$

Besides, as before, by the assumptions of [Theorem 2](#) and the Schwarz inequality, we have

$$\begin{aligned} & \sqrt{\alpha} \sqrt{f(x)} x' \\ & = - \left[ \frac{\sqrt{\alpha}}{2\sqrt{a(t)}} x' - \sqrt{a(t)} \sqrt{f(x)} x \right]^2 + \frac{\alpha}{4a(t)} (x')^2 + a(t) f(x) x \\ & \leq \frac{\alpha}{4a(t)} (x')^2 + a(t) f(x) x \\ & = a(t) f(x) x + \frac{\alpha}{4} a(t) f^2(x) - \frac{1}{2} \alpha f(x) \int_{t-\tau}^t B(t, s) g(x(s)) ds \\ & + \frac{\alpha}{4a(t)} \left[ \int_{t-\tau}^t B(t, s) g(x(s)) ds \right]^2 \\ & \leq a(t) f(x) x + \frac{\alpha}{4} a(t) f^2(x) - \frac{1}{2} \alpha f(x) \int_{t-\tau}^t B(t, s) g(x(s)) ds \\ & + \frac{\alpha m^2}{J} \int_{t-\tau}^t |B(t, s)| f^2(x(s)) ds \end{aligned}$$

and

$$\frac{1}{2} \alpha f(x) x' = -\frac{1}{2} \alpha a(t) f^2(x) + \frac{1}{2} \alpha f(x) \int_{t-\tau}^t B(t, s) g(x(s)) ds.$$

On gathering the above discussion into (4), it follows that

$$\begin{aligned} V'_0 & \leq - \left[ k - \frac{m^2(1+\alpha)}{J} \right] \int_0^t |B(t, s)| f^2(x(s)) ds \\ & - \frac{m^2}{J} \int_0^{t-\tau} |B(t, s)| f^2(x(s)) ds \\ & - \frac{\alpha m^2}{J} \int_0^{t-\tau} |B(t, s)| f^2(x(s)) ds \\ & - \left[ a(t) - k \int_{t-\tau}^\infty |B(u + \tau, t)| du \right] f^2(x) \\ & \leq - \left[ k - \frac{m^2(1+\alpha)}{J} \right] \int_0^t |B(t, s)| f^2(x(s)) ds \\ & - \left[ a(t) - k \int_{t-\tau}^\infty |B(u + \tau, t)| du \right] f^2(x). \end{aligned}$$

Let  $k = \frac{m^2(1+\alpha)}{J}$ . Hence, we have

$$V'_0 \leq - \left[ a(t) - k \int_{t-\tau}^\infty |B(u + \tau, t)| du \right] f^2(x).$$

If

$$a(t) - k \int_{t-\tau}^\infty |B(u + \tau, t)| du \geq 0,$$

then it follows that

$$V'_0 \leq 0.$$

Therefore, we can conclude that the zero solution of [Eq. \(2\)](#) is stable.

Let  $p(t) \neq 0$  and

$$\rho(t) = \alpha_0 a(t) - \frac{1}{2} \int_{t-\tau}^t |B(t, s)| ds - \frac{1}{2} m^2 \int_{t-\tau}^\infty |B(u + \tau, t)| du.$$

### B. Assumptions

(H1) There exist positive constants  $\alpha_0$ ,  $m$  and  $M$  such that  $g(0) = 0$ ,  $g^2(x) \leq m^2 x^2$  if  $|x| \leq M$ ,  $f(0) = 0$ ,  $\frac{f(x)}{x} \geq \alpha_0 > 0$ , when  $x \neq 0$ ,

(H2)  $a(t) > 0$  for  $t \geq 0$ ,  $B(t, s)$  is continuous for  $0 \leq s \leq t < \infty$ ,  $\int_{t-\tau}^\infty |B(u + \tau, s)| du$  is defined and continuous for  $0 \leq s - \tau \leq t < \infty$ , and  $\rho(t) \geq 0$  for every  $t \geq s - \tau \geq 0$ .

We prove the following theorem.

**Theorem 3.** In addition to assumptions (H1) and (H2), if  $p \in L^1(0, \infty)$  and  $\lambda = \frac{1}{2} m^2$ , then all solutions of [Eq. \(2\)](#) are bounded.

**Proof.** We define a functional  $V_1 = V_1(t) = V_1(t, x(t))$  by

$$V_1 = \frac{1}{2} x^2 + \lambda \int_0^t \int_{t-\tau}^\infty |B(u + \tau, s)| du x^2(s) ds, \tag{5}$$

where  $\lambda$  is a positive constant to be determined later in the proof.  $\square$

It is clear that the functional  $V_1$  is positive definite.

Calculating the time derivative of the functional  $V_1$  we get from (5) that

$$\begin{aligned} V'_1 & = -a(t) f(x) x + x \int_{t-\tau}^t B(t, s) g(x(s)) ds + x p(t) \\ & + \lambda \int_{t-\tau}^\infty |B(u + \tau, t)| du x^2 - \lambda \int_0^t |B(t, s)| x^2(s) ds. \end{aligned} \tag{6}$$

In view of the assumptions of [Theorem 3](#) and the estimate  $|\alpha\beta| \leq 2^{-1}(\alpha^2 + \beta^2)$ , it is obvious that

$$\begin{aligned} V'_1 & \leq -a(t) f(x) x + \frac{1}{2} \int_{t-\tau}^t |B(t, s)| (x^2(t) + g^2(x(s))) ds \\ & + |p(t)| + |p(t)| x^2 + \lambda \int_{t-\tau}^\infty |B(u + \tau, t)| du x^2 \\ & - \lambda \int_0^t |B(t, s)| x^2(s) ds \\ & = - \left[ a(t) \frac{f(x)}{x} - \frac{1}{2} \int_{t-\tau}^t |B(t, s)| ds - \lambda \int_{t-\tau}^\infty |B(u + \tau, t)| du \right] x^2 \\ & + |p(t)| + |p(t)| x^2 + \frac{1}{2} \int_{t-\tau}^t |B(t, s)| g^2(x(s)) ds \end{aligned}$$

$$\begin{aligned}
 & -\lambda \int_0^t |B(t, s)|x^2(s)ds \\
 \leq & -\left[\alpha_0 a(t) - \frac{1}{2} \int_{t-\tau}^t |B(t, s)|ds - \lambda \int_{t-\tau}^\infty |B(u + \tau, t)|du\right]x^2 \\
 & + |p(t)|(1 + x^2) + \frac{1}{2} \int_{t-\tau}^t |B(t, s)|g^2(x(s))ds \\
 & -\lambda \int_0^t |B(t, s)|x^2(s)ds \\
 \leq & -\left[\alpha_0 a(t) - \frac{1}{2} \int_{t-\tau}^t |B(t, s)|ds - \lambda \int_{t-\tau}^\infty |B(u + \tau, t)|du\right]x^2 \\
 & -\left(\lambda - \frac{1}{2}m^2\right) \int_0^t |B(t, s)|x^2(s)ds + |p(t)|(1 + x^2) \\
 & -\frac{1}{2}m^2 \int_0^{t-\tau} |B(t, s)|x^2(s)ds \\
 \leq & -\left[\alpha_0 a(t) - \frac{1}{2} \int_{t-\tau}^t |B(t, s)|ds - \lambda \int_{t-\tau}^\infty |B(u + \tau, t)|du\right]x^2 \\
 & -\left(\lambda - \frac{1}{2}m^2\right) \int_0^t |B(t, s)|x^2(s)ds + |p(t)|(1 + x^2).
 \end{aligned}$$

Let  $\lambda = \frac{1}{2}m^2$ . Then, we have

$$\begin{aligned}
 V_1' & \leq -\rho(t)x^2 + |p(t)| + |p(t)|x^2 \leq |p(t)| + |p(t)|x^2 \\
 & \leq (1 + 2V_1)|p(t)|.
 \end{aligned}$$

Integrating the last estimate from zero  $t_0$  to  $t$ , we have

$$V_1(t) \leq V_1(t_0) + \int_0^t |p(s)|ds + 2 \int_0^t V_1(s)|p(s)|ds.$$

Hence, an application of Gronwall’s inequality bounds  $V_1$ . Hence we can conclude that all solutions of Eq. (2) are bounded.

**Remark.** By Theorem 2, we improve and extend a stability result obtained for a Volterra integro-differential equation without delay to its delay form (see Vanualailai and Nakagiri [11, Theorem 3.2]). Besides, Theorems 2 and 3 complement to the papers in the references, and they have a contribution to the papers of Becker [1], Burton [3], Tunc [10], Adivar and Raffoul [15] and Raffoul [17]. By this way, we also mean that the Volterra integro-differential equation considered and the established assumptions here are different from that in the mentioned papers above. Theorem 3 gives an additional result, the boundedness of solutions, to that of Vanualailai and Nakagiri [11, Theorem 3.2]. This paper may be useful for researchers working on the qualitative behaviors of solutions of functional Volterra integro-differential equations. These cases show the novelty and originality of the present paper.

### 3. Conclusion

A non-linear Volterra integro-differential equation of first order with delay is considered. The stability and boundedness of

solutions are discussed by using the Lyapunov’s functional approach. The obtained results improve and extend some results in the literature.

### References

- [1] L.C. Becker, Uniformly continuous  $L^{-1}$ - solutions of Volterra equations and global asymptotic stability, *Cubo* 11 (3) (2009) 1–24.
- [2] T.A. Burton, Stability theory for Volterra equations, *J. Differ. Eq.* 32 (1) (1979) 101–118.
- [3] T.A. Burton, Construction of Liapunov functionals for Volterra equations, *J. Math. Anal. Appl.* 85 (1) (1982) 90–105.
- [4] T.A. Burton, W.E. Mahfoud, Stability criteria for Volterra equations, *Trans. Amer. Math. Soc* 279 (1) (1983) 143–174.
- [5] T.A. Burton, W.E. Mahfoud, Stability by decompositions for Volterra equations, *Tohoku Math. J.* (2) 37 (4) (1985) 489–511.
- [6] A. Diamandescu, On the strong stability of a nonlinear Volterra integro-differential system, *Acta Math. Univ. Comenian. (N.S.)* 75 (2) (2006) 153–162.
- [7] T. Hara, T. Yoneyama, T. Itoh, Asymptotic stability criteria for nonlinear Volterra integro-differential equations, *Funkcial. Ekvac.* 33 (1) (1990) 39–57.
- [8] R.K. Miller, Asymptotic stability properties of linear Volterra integro-differential equations, *J. Differ. Eq.* 10 (1971) 485–506.
- [9] O.J. Staffans, A direct Lyapunov approach to Volterra integro-differential equations, *SIAM J. Math. Anal.* 19 (4) (1988) 879–901.
- [10] C. Tunc, A note on the qualitative behaviors of non-linear Volterra integro-differential equation, *J. Egyptian Math. Soc.* (2015) (in press).
- [11] J. Vanualailai, S. Nakagiri, Stability of a system of Volterra integro-differential equations, *J. Math. Anal. Appl.* 281 (2) (2003) 602–619.
- [12] T.A. Burton, Volterra integral and differential equations, *Mathematics in Science and Engineering*, 202 (2005).
- [13] C. Corduneanu, Integral equations and stability of feedback systems, *Mathematics in Science and Engineering*, 104, Academic Press, New York–London, 1973.
- [14] G. Gripenberg, S.Q. Londen, O. Staffans, Volterra integral and functional equations, *Encyclopedia of Mathematics and its Applications*, Cambridge University Press, Cambridge, 1990.
- [15] M. Adivar, Y.N. Raffoul, Inequalities and exponential stability and instability in finite delay Volterra integro-differential equations, *Rend. Circ. Mat. Palermo* 61 (3) (2012) 321–330.
- [16] John R. Graef, C. Tunc, Continuability and boundedness of multi-delay functional integro-differential equations of the second order, *Rev. R. Acad. Cienc. Exactas Fis. Nat. Ser. A Math. RACSAM* 109 (1) (2015) 169–173.
- [17] Y. Raffoul, Exponential stability and instability in finite delay nonlinear Volterra integro-differential equations, *Dyn. Contin. Discrete Impuls. Syst. Ser. A Math. Anal.* 20 (1) (2013) 95–106.
- [18] Y. Raffoul, M. Unal, Stability in nonlinear delay Volterra integro-differential systems, *J. Nonlinear Sci. Appl* 7 (2014) 422–428.
- [19] R.D. Driver, Existence and stability of solutions of a delay-differential system, *Arch. Ration. Mech. Anal.* 10 (1962) 401–426.