



Egyptian Society of Anesthesiologists
Egyptian Journal of Anaesthesia

www.elsevier.com/locate/egja
www.sciencedirect.com



Research Article

A comparative study between magnesium sulphate and dexmedetomidine for deliberate hypotension during middle ear surgery

Ossama H. Aboushanab ^a, Ahmed M. El-Shaarawy ^a, Ahmed M. Omar ^{b,*},
Hisham H. Abdelwahab ^a

^a Department of Anesthesia, Faculty of Medicine, Cairo University, Cairo, Egypt

^b King Faisal Specialist Hospital & Research Center, Riyadh, Saudi Arabia

Received 27 May 2011; revised 29 July 2011; accepted 31 July 2011

Available online 23 August 2011

KEYWORDS

Deliberate hypotension;
Magnesium sulphate;
Dexmedetomidine

Abstract *Background:* This study was designed to compare magnesium sulphate with dexmedetomidine, regarding their efficiency in inducing deliberate hypotension and providing a better surgical field exposure during middle ear surgery. It also compared the influence of their use on postoperative pain and recovery time.

Methods: Eighty-eight adult patients undergoing middle ear surgery were included. Patients were randomly divided into two equal groups. Patients were assigned to receive either magnesium sulphate (M group) or dexmedetomidine (D group). Anaesthesia was induced by propofol 2 mg/kg iv and fentanyl 1 µg/kg. Patients in the M group received an iv bolus of magnesium sulphate 50 mg/kg in a total of 100 ml saline over 10 min followed by infusion of 15 mg/kg/h until the end of surgery. Similarly, patients in the D group received dexmedetomidine 1 µg/kg over 10 min followed by 0.4–0.8 µg/kg/h until the end of operation. The target MAP during operation was

* Corresponding author. Address: Anesthesia Department, MBC 22, King Faisal Specialist Hospital & Research Center. PO Box 3354, Riyadh, Saudi Arabia. Tel.: +966 569527320.

E-mail addresses: ossamaaboushanab@yahoo.com (O.H. Aboushanab), drshaarawy@yahoo.com (A.M. El-Shaarawy), ahmedmommer@yahoo.com (A.M. Omar), hishamhosny@yahoo.com (H.H. Abdelwahab).

1110-1849 © 2011 Egyptian Society of Anesthesiologists. Production and hosting by Elsevier B.V. Open access under [CC BY-NC-ND license](https://creativecommons.org/licenses/by-nc-nd/4.0/).

Peer review under responsibility of Egyptian Society of Anesthesiologists.
doi:10.1016/j.egja.2011.07.008



Production and hosting by Elsevier

between 60 and 70 mmHg. The surgeon who was blinded of the selected hypotensive agent was asked to assess the quality of the surgical field. In the postanesthesia care unit (PACU), postoperative pain was assessed and recovery time was recorded.

Results: Both study drugs succeeded to reach the target MAP. The quality of the surgical field was not different between the two groups. Postoperative pain was not different between the two groups and only eight patients in the M group and seven patients in the D group required analgesics. Recovery time was significantly longer for the patients in group D ($p < 0.05$).

Conclusion: We concluded that both magnesium sulphate and dexmedetomidine successfully induced deliberate hypotension in patients undergoing middle ear surgery but magnesium sulphate was associated with shorter recovery time and earlier discharge from the PACU.

© 2011 Egyptian Society of Anesthesiologists. Production and hosting by Elsevier B.V.

Open access under [CC BY-NC-ND license](#).

1. Introduction

Deliberate hypotension is a technique which has been used to decrease intraoperative bleeding and improve the quality of the surgical field for better visualization during middle ear surgery and other head and neck operations [1–4].

Several pharmacologic agents have been used to produce deliberate hypotension. Direct vasodilators as nitroprusside and nitroglycerine, α -2 adrenergic agonists as clonidine and dexmedetomidine, beta adrenergic antagonists as propranolol and esmolol, α and beta adrenergic antagonists as labetalol, inhalational anaesthetics as isoflurane and sevoflurane, μ -receptors agonist as remifentanyl, and *N*-methyl *D*-aspartate antagonist as magnesium were all used to induce hypotension during middle ear surgery [5–11].

Magnesium is the fourth most common salt in the human body after phosphorus, calcium and potassium [12]. Magnesium sulphate was previously used to induce deliberate hypotension [13,14]. It produces its hypotensive effect by limiting the outflow of calcium from the sarcoplasmic reticulum and produces a vasodilating effect by increasing the synthesis of prostacyclin and inhibiting angiotensin converting enzyme activity [15]. Hypotension induced by magnesium during surgery is also explained by its powerful analgesic effect [16]. The analgesic action of magnesium is explained by its antagonistic effect at *N*-methyl *D*-aspartate receptors [17].

Dexmedetomidine is an α -2 adrenoreceptor agonist. It decreases heart rate and arterial blood pressure by reducing norepinephrine and epinephrine plasma levels [18]. Activation of postsynaptic receptors in the central nervous system by α -2-agonists inhibits the sympathetic activity and decreases heart rate and blood pressure and causes sedation.

No studies have been published to compare magnesium sulphate and dexmedetomidine regarding their hypotensive effects in patients undergoing middle ear surgery. The main objective of this prospective randomized study was to compare magnesium sulphate with dexmedetomidine as regards their efficacy in inducing deliberate hypotension and providing better quality of the surgical field during middle ear surgery. The effect of both drugs on postoperative pain and recovery time was also compared.

2. Methods

In this study, 88 adults ASA physical status I and II patients undergoing middle ear surgery were included. A written informed consent was obtained from all patients after approval

by the Local Ethics Committee. The enrollment period lasted from January 2010 to March 2011 in Kasr El Eini University Hospital in Cairo (Egypt). Patients with hepatic, renal, cardiovascular, neuromuscular or haematological disorders were excluded. Those with history of using sedatives, narcotics or anticoagulants were also excluded. Patients were randomly assigned to receive either magnesium sulphate (M) or dexmedetomidine (D). Randomization was accomplished by using computerized randomization tables. Anaesthesia was induced by propofol 2 mg/kg iv and fentanyl 1 μ g/kg. Patients in the M group received an iv bolus of magnesium sulphate 50 mg/kg in a total of 100 ml saline over 10 min followed by infusion of 15 mg/kg/h until the end of operation. Similarly, patients in the D group received dexmedetomidine 1 μ g/kg over 10 min followed by 0.4–0.8 μ g/kg/h until the end of surgical procedure. The study drugs were infused using an infusion pump (Module DPS + IS3-Fresenius vial, Le Grand Chemin-38590, Brezins, France). When the eyelid reflex was lost, rocuronium 0.6 mg/kg was injected. A peripheral nerve stimulator (Plexygon nerve stimulator Vygon I Talia S.r.L Via P. Dona, II, Padova, Italy, Ref. 7501031 S/N0507PB202) was used at the wrist to monitor neuromuscular block and the trachea was intubated when T1 was 0%. Additional doses of rocuronium were not administered to allow intraoperative monitoring of facial nerve. When T1 was 25%, muscle relaxation was reversed with atropine 0.02 mg/kg and neostigmine 0.04 mg/kg iv. Anaesthesia was maintained with 1–2 MAC sevoflurane and lungs were ventilated with oxygen and medical air at a ratio of 1:1. Ringer's solution was infused continuously at a rate of 5 ml/kg/h. Mean arterial pressure (MAP) and heart rate (HR) were recorded before induction of anaesthesia (baseline), 1 min after induction of anaesthesia, 1 min before intubation, 1 min after intubation then every 15 min intraoperatively, 1 min before extubation and 5 min after extubation. (Drager Medical systems, Inc., Telford, PA 18969, USA). Blood samples for serum magnesium were obtained before and after surgery. The target MAP during operation was between 60 and 70 mmHg. If hypertension or tachycardia more than 20% of the preoperative value occurred, fentanyl 1 μ g/kg was given iv. Sevoflurane concentration was manipulated according to targeted MAP and its end-tidal concentration was recorded every 15 min while the patients were intubated. If hypotension or bradycardia more than 20% of the preoperative value occurred, 5 mg ephedrine for hypotension or 0.5 mg atropine for bradycardia was given iv. Before skin closure all patients were given ondansetron 4 mg iv as an antiemetic prophylaxis. The surgeon who was not aware of the selected hypotensive agent was asked to assess the quality

of the surgical field according to the quality scale proposed by Fromme and colleagues [19].

- 0 = no bleeding.
- 1 = slight bleeding – blood evacuation not necessary.
- 2 = slight bleeding – sometimes blood has to be evacuated.
- 3 = low bleeding – blood has to be often evacuated. Operative field is visible for some seconds after evacuation.
- 4 = average bleeding – blood has to be often evacuated. Operative field is visible only right after evacuation.
- 5 = high bleeding – constant blood evacuation is needed. Sometimes bleeding exceeds evacuation. Surgery is hardly possible.

At the end of surgery, study drugs and sevoflurane were discontinued. Patients, after the end of surgery, were transferred to the postanesthesia care unit (PACU). In the PACU, MAP and HR were recorded on admission and then 15 and 30 min later. Pain was evaluated using visual analogue scale (VAS) starting from 0 for no pain to 100 for worst pain. If VAS was > 40, ketorolac 30 mg iv was administered. Patients complaining of postoperative nausea and vomiting were given metoclopramide 10 mg iv. Patients were discharged when fulfilling a score of 9 using a modified Aldrete scoring system [20]. Nurses of the PACU were blinded to patients' group allocation.

The primary outcome was the quality of surgical field assessed by the surgeon. A sample size of 44 in each group was based on a pilot study we have conducted before. We assumed that a 30% difference in the score of surgical field quality would be clinically significant. A sample size was calculated to be 40 at an alpha error of 0.05 and a beta error of 0.2. Because the primary outcome on which we calculated the sample size was ordinal, we increased the sample size by 10%.

Data were statistically described in terms of mean (\pm SD), frequencies (number of cases) and percentages when appropriate. Data were tested first for normal distribution by Kolmogorov–Smirnov test. Comparison of quantitative variables between the study groups was done using Student *t* test for independent samples if normally distributed. Mann–Whitney U test was used for non-normally distributed quantitative and ordinal data. For comparing categorical data, Chi square (χ^2) test was performed. Exact test was used instead when the expected frequency is less than 5. A probability value (*p* value) less than 0.05 was considered statistically significant. All statistical calculations were done using computer programs Microsoft Excel 2007 (Microsoft Corporation, NY, USA) and SPSS (Statistical Package for the Social Science; SPSS Inc., Chicago, IL, USA) version 15 for Microsoft Windows.

Table 1 Patients characteristics. Data are mean (SD) or number.

	Group M (<i>n</i> = 44)	Group D (<i>n</i> = 44)
Age (years)	32.4 (7.3)	34.1 (6.4)
Gender (male/female)	26/18	24/20
Weight (kg)	72.5 (9.7)	72.2 (9.1)
Height (cm)	164.2 (9.5)	163.0 (8.4)
ASA (I/II)	31/13	29/15

Group M: Magnesium sulphate, Group D: Dexmedetomidine.

Table 2 Pre- and postoperative serum magnesium levels. Data are mean (SD).

	Group M (<i>n</i> = 44)	Group D (<i>n</i> = 44)
Preoperative magnesium (mmol/l)	0.94 (0.14)	0.95 (0.16)
Postoperative magnesium (mmol/l)	1.3 (0.15)	0.91 (0.15)*

Group M: Magnesium sulphate, Group D: Dexmedetomidine.
* *p* = 0.008.

3. Results

One hundred and forty-one patients were found eligible for the study. Forty-three patients met our exclusion criteria and 10 patients refused participation. Eighty-eight patients were randomized to two groups: group M (*n* = 44) and group D (*n* = 44). No patient was excluded from the study.

There were no significant differences between the two groups with regard to age, gender, weight, height and ASA classification as shown in Table 1. Postoperative magnesium levels were significantly higher in group M than in group D as shown in Table 2.

There were no significant differences between the two groups neither in MAP as shown in Fig. 1 nor in HR as shown in Fig. 2.

The two study groups were similar in terms of duration of surgery and anaesthesia, type of surgery, bleeding quality score and the need to atropine, ephedrine and fentanyl (Table 3)

Again the end-tidal concentrations of sevoflurane were similar in the two study groups (Table 4)

Table 5 shows the difference between the two groups in the PACU. Discharge from the PACU required a significantly longer time for the patients in the D group (*p* < 0.001).

4. Discussion

This study shows the ability of both magnesium sulphate and dexmedetomidine to induce deliberate hypotension in patients undergoing middle ear surgery. Mean arterial pressure was successfully reduced to the target values in both groups. Both drugs were equally successful to produce satisfactory surgical field. However the discharge time from PACU in dexmedetomidine group was almost double that in magnesium sulphate group.

Magnesium sulphate is known to augment the action of non-depolarizing neuromuscular blocking drugs and reduce their consumption [21]. However, facial nerve monitoring is required during middle ear surgery and neuromuscular blocking agents were reversed shortly after the beginning of the operation.

Recovery in patients receiving dexmedetomidine was prolonged most probably due its sedative-sparing effects via central actions in the locus coeruleus [22,23]. Other studies observed that recovery time was prolonged in patients who received dexmedetomidine for induced hypotension when compared with control groups [6,24]. Another study observed that there was a prolonged recovery time in patients received dexmedetomidine when compared with those who received remifentanyl during gynaecologic laparoscopic surgery [25].




Figure. 1 Mean (SD) of mean arterial blood pressure (MAP, mmHg) between the study groups over the study period. M, Magnesium sulphate group; D, Dexmedetomidine group; PACU, postanaesthesia care unit.




Figure. 2 Mean (SD) of heart rate (HR, beat/min) between the study groups over the study period. M, Magnesium sulphate group; D, Dexmedetomidine group; PACU, postanaesthesia care unit.

Table 3 Intraoperative data. Data are mean (SD), median (range) or number.

	Group M (n = 44)	Group D (n = 44)
Duration of surgery (min)	116 (18)	118 (17)
<i>Type of surgery</i>		
Tympanoplasty	27	25
Mastoidectomy	17	19
Duration of anaesthesia	132 (18)	135 (19)
Surgical field quality score	2.2 (1–4)	2.4 (1–4)
Atropine (no. of patients)	7	9
Ephedrine (no. of patients)	6	8
Fentanyl (no. of patients)	2	1

Group M: Magnesium sulphate, Group D: Dexmedetomidine.

Several studies have reported the ability of both drugs in providing deliberate hypotension. Ryu and colleagues compared magnesium sulphate and remifentanyl and proved that both drugs can induce adequate deliberate hypotension for

middle ear surgery but magnesium sulphate was associated with better postoperative analgesia and less postoperative nausea, vomiting and shivering [26]. Elsharnouby and colleagues performed a placebo controlled study and concluded that magnesium sulphate reduced MAP, HR and blood loss and was associated with less anaesthetic requirements and emergence time for patients underwent functional endoscopic sinus surgery [13]. Yosri and colleagues designed a comparative study between magnesium sulphate and nitroprusside and concluded that magnesium sulphate could produce deliberate hypotension and provided good surgical conditions for resection of choroidal melanoma with no need for adding potent hypotensive agents [14]. Durmus and colleagues have proved in a placebo controlled study that dexmedetomidine decreased bleeding and anaesthetic requirements in patients undergoing tympanoplasty and septorhinoplasty [24]. Ayoglu and colleagues in a placebo controlled study concluded that dexmedetomidine reduced bleeding and improved visibility and decreased analgesic requirements in septoplasty operations [27]. Although magnesium sulphate and dexmedetomidine

Table 4 End-tidal concentrations of sevoflurane. Data are mean (SD).

	Group M (n = 44)	Group D (n = 44)
15 min	2.6 (0.4)	2.5 (0.3)
30 min	2.3 (0.4)	2.3 (0.3)
45 min	2.4 (0.3)	2.4 (0.3)
60 min	2.3 (0.4)	2.2 (0.4)
75 min	2.2 (0.3)	2.2 (0.3)
90 min	2.3 (0.3)	2.3 (0.4)
105 min	2.4 (0.4)	2.2 (0.3)
120 min	2.3 (0.3)	2.3 (0.4)
135 min	1.4 (0.3)	1.3 (0.4)
150 min	0.7 (0.3)	0.8 (0.3)

Group M: Magnesium sulphate, Group D: Dexmedetomidine.

Table 5 PACU data. Data are mean (SD) or number.

	Group M (n = 44)	Group D (n = 44)
Time for discharge (min)	38.3 (5.2)	60.7 (6.8)*
VAS 0	34.8 (7.9)	34.4 (7.6)
VAS 15	34.6 (7.3)	35.8 (6.4)
VAS 30	27 (6.1)	28 (6.4)
Analgesic received	8	7
Antiemetic received	12	11

Group M: Magnesium sulphate, Group D: Dexmedetomidine.
VAS: visual analogue score of pain.

* $p < 0.001$.

were previously investigated as hypotensive agents, there were no studies designed to compare between them as regards their efficiency in controlling MAP and HR, analgesic effect and emergence from general anaesthesia.

The target MAP between 60 and 70 mmHg was decided to achieve hypotensive anaesthesia and bloodless field without subjecting the patients to peripheral tissue ischaemia. This target MAP was determined after revising previous studies in which metabolic and hormonal responses were observed in patients subjected to induced hypotension. Yoshikawa and colleagues induced hypotensive anaesthesia using a similar target MAP of 60–70 mmHg in patients undergoing mandibular osteotomy. They measured blood pyruvate, lactate and glucose and observed insignificant increase in their levels. They concluded that hypotension can be induced safely, if MAP is maintained between 60 and 70 mmHg [28]. Newton and colleagues targeted a MAP of 55 mmHg in patients undergoing middle ear surgery. They also observed insignificant increase in blood lactate and pyruvate levels [29].

Our study was limited by the unavailability of two monitors. We used a subjective scoring system to evaluate the quality of the surgical field instead of a Laser Doppler flowmetry which was used by Degoute and colleagues to measure the middle ear blood flow [9]. We could not assess the depth of anaesthesia as we lacked a bispectral index (BIS) monitor. However the lack of any significant differences in end-tidal sevoflurane concentrations between the two groups suggests that hypotensive properties of the two study drugs are similar in the doses that we used.

Only 16% of patients in the magnesium group and 14% of patients in the dexmedetomidine group required rescue analge-

sics. These results can be explained by the analgesic properties of both study drugs [17,18]

We concluded that both magnesium sulphate and dexmedetomidine successfully induced deliberate hypotension and good surgical field visibility but magnesium sulphate was associated with shorter recovery time and earlier discharge from the PACU.

References

- [1] Boezaart AP, Merwee J, Coetzee A. Comparison of sodium nitroprusside and esmolol induced hypotension for functional endoscopic sinus surgery. *Can J Anaesth* 1995;42:373–6.
- [2] Coursin DB, Coursin DB, Maccioli GA. Dexmedetomidine. *Curr Opin Crit Care* 2001;7:221–6.
- [3] Marchal JM, Gomez-Luque A, Martos-Crespo F. Clonidine decreases intraoperative bleeding in middle ear microsurgery. *Acta Anaesthesiol Scand* 2001;45:627–33.
- [4] Degoute CS, Ray MJ, Manchon M. Remifentanyl and controlled hypotension; comparison with nitroprusside or esmolol during tympanoplasty. *Can J Anaesth* 2001;48:20–7.
- [5] Degoute CS, Ray MJ, Gueugniaud PY, Dubreuil C. Remifentanyl induces consistent and sustained controlled hypotension in children during middle ear surgery. *Can J Anaesth* 2003;50(3):270–6.
- [6] Richa F, Yazigi A, Sleilaty G, Yazbek P. Comparison between dexmedetomidine and remifentanyl for controlled hypotension during tympanoplasty. *Eur J Anaesthesiol* 2008;25(5):369–74.
- [7] Dietrich GV, Heesen M, Boldt J, Hempelmann G. Platelet function and adrenoceptors during and after induced hypotension using nitroprusside. *Anesthesiology* 1996;85(6):1334–40.
- [8] Tobias JD. Controlled hypotension in children: a critical review of available agents. *Paediatr Drugs* 2002;4:439–53.
- [9] Degoute CS. Controlled hypotension: a guide to drug choice. *Drugs* 2007;67:1053–76.
- [10] Testa LD, Tobias JD. Pharmacologic drugs for controlled hypotension. *J Clin Anesth* 1995;7:326–37.
- [11] Delhumeau A, Granry JC, Monrignal J-P. Therapeutic use of magnesium in anaesthesia and intensive care. *Ann Fr Anesth Reanim* 1995;14:406–16.
- [12] Saris NE, Mervaala E, Karppanen H. Magnesium. An update on physiological, clinical and analytical aspects. *Clin Chim Acta* 2000;294:1–26.
- [13] Elsharnouby NM, Elsharnouby MM. Magnesium sulphate as a technique of hypotensive anaesthesia. *Br J Anaesth* 2006;96:727–31.
- [14] Yosri M, Othman IS. Controlled hypotension in adults undergoing choroidal melanoma resection: comparison between the efficacy of nitroprusside and magnesium sulphate. *Eur J Anaesthesiol* 2008;25:891–6.
- [15] Sanders GM, Sim KM. Is it feasible to use magnesium sulphate as a hypotensive agent in oral and maxillofacial surgery? *Ann Acad Med Singapore* 1998;27:780–5.
- [16] Buvanendran A, McCarthy RJ, Kroin JS. Intrathecal magnesium prolongs fentanyl analgesia: a prospective, randomized, controlled trial. *Anesth Analg* 2002;95:661–6.
- [17] Dube L, Granry JC. The therapeutic use of magnesium in anaesthesiology, intensive care and emergency medicine: a review. *Can J Anaesth* 2003;50:732–46.
- [18] Bloor BC, Ward DS, Belleville JP. Effects of intravenous dexmedetomidine in humans II. Hemodynamic changes. *Anesthesiology* 1992;77:1134–42.
- [19] Fromme GA, Mackenzie RA, Gould Jr AB, Lund BA, Offord KP. Controlled hypotension for orthognathic surgery. *Anesth Analg* 1986;65(6):683–6.

- [20] Thomas WF, Macario A. The postanesthesia care unit. In: Miller RD, editor. *Anesthesia*. Philadelphia: Churchill Livingstone; 2005. p. 2708–9.
- [21] Herroeder S, Schonherr ME, De Hert SG, Hollmann MW. Magnesium-essentials for anesthesiologists. *Anesthesiology* 2011;114:971–93.
- [22] Maze M, Segal I, Bloor B. Clonidine and other alpha₂ adrenergic agonists: strategies for the rational use of these novel anesthetic agents. *J Clin Anesth* 1988;1:146–57.
- [23] Guo TZ, Jiang JY, Butterman AE, Maze M. Dexmedetomidine injection into the locus coeruleus produces antinociception. *Anesthesiology* 1996;84:873–81.
- [24] Durmus M, But AK, Dogan Z. Effect of dexmedetomidine on bleeding during tympanoplasty or septorhinoplasty. *Eur J Anaesthesiol* 2007;24:447–53.
- [25] Bulow NM, Barbosa NV, Rocha JB. Opioid consumption in total intravenous anesthesia is reduced with dexmedetomidine: a comparative study with remifentanyl in gynecologic videolaparoscopic surgery. *J Clin Anesth* 2007;19:280–5.
- [26] Ryu JH, Sohn IS, Do SH. Controlled hypotension for middle ear surgery: a comparison between remifentanyl and magnesium sulphate. *Br J Anaesth* 2009;103(4):490–5.
- [27] Ayoglu H, Yapakci O, Ugur MB, Uzun L, Altunkaya H, Ozer Y, et al.. Effectiveness of dexmedetomidine in reducing bleeding during septoplasty and tympanoplasty operations. *J Clin Anesth* 2008;20:437–41.
- [28] Yashikawa F, Kohase H, Umino M, Fukayama H. Blood loss and endocrine responses in hypotensive anaesthesia with sodium nitroprusside and nitroglycerin for mandibular osteotomy. *Int J Oral Maxillofac Surg* 2009;38:1159–64.
- [29] Newton MC, Chadd GD, O'donghue B, Sapsed-Byrne SM, Hall GM. Metabolic and hormonal responses to induced hypotension for middle ear surgery. *Br J Anaesth* 1996;76(3):352–7.