

EFFECT OF CERTAIN CHEMICAL COMPONENTS AND SOURCE OF RICE PLANT ON ITS RESISTANCE TO RICE STEM BORER, *CHILO AGAMEMNON* BLES. AND RICE LEAF MINER, *HYDRELLIA PROSTERNALIS*, DEEM .

ALI M. SOLIMAN¹, W.M. EL-ATTAR², REFAAT G.ABOU ELELA³
AND AHMED E.ABDEL-WAHAB⁴

1 Plant Protection Research Institute, Agricultural Research Centre, Dokki, Egypt .

2 Faculty of Agriculture., Zagazig University.

3 Faculty of Science., Cairo Univerity.

4 Field Crops Research Institute, Agricultural Research Centre, Giza.

(Manuscript received 21 October, 1996)

Abstract

Different rice entries, of known sources, were planted. Rice Stem Borer (RSB) and Rice Leaf Miner (RLM) infestation was evaluated. Total protein, crude silica and fifteen amino acid contents were determined. The correlation between chemical contents and insect infestation showed certain results.

Protein increased RSB infestation in most rice entries with positive correlation ($r=0.45$), while decreased RLM infestation in most rice entries with negative correlation ($r=-0.77$). Silica decreased infestation of both RSB and RLM and RLM with negative correlations ($r=-0.13$ and -0.23 , respectively). Free amino acids showed different effects. Some of them decreased RSB infestation as alanine, glycine, histidine + arginine, aspartic+serine and valine, and some decreased RLM infestation as ornithine + lysine, histidine + arginine and methionine. Other amino acids increased insect infestation.

Japonica entries showed more resistance against RSB than indica entries and have less protein content, while the indica entries showed more resistance against RLM insect and higher protein content. As so, the source plays role in host resistance .

It can be concluded that chemical components may play an important role in plant resistance, but is not the detrimental factor and other morphological, histological, and/or chemical factors share in plant resistance.

INTRODUCTION

Rice is considered as one of the most important cereal crops in Egypt. The Rice stem borer (RSB), *Chilo agamemnon* Bles. and the rice leaf miner (RLM), *Hydrellia prosternalis* Deem are considered the main insects that cause significant

losses in rice crop yearly.

Realizing the side effects of insecticides has caused increased interest in search for new measures of insect control. The use of integrated pest management (IPM) in insect control is the new approach in this respect. Selection of resistant varieties is an important item in IPM field. Chemical resistance strongly shares in varietal resistance and helps in selecting and presenting new resistant varieties in breeding programs.

Little attention has been given to the chemical resistance of the host plant in Egypt. This study is an attempt to correlate the contents of certain chemical components with the resistance of rice entries against RSB and RLM attack to present some chemical components that play a role in rice plant resistance.

MATERIALS AND METHODS

Four field experiments were conducted during 1993 and 1994 at Sakha Station, Kafr El-Sheikh Governorate. Different rice entries were planted for insect and chemical evaluations.

A. Field experiments

Through 1993 season, eleven rice entries, with known sources, were planted (3 rows x 6 m x 34 rep./entry) in two dates i.e. recommended (15 May) and late (15 June), for insect and protein content evaluation. Through 1994, nineteen entries were planted in the same dates and design for insect, silica and free amino acids content evaluation.

B. Insect evaluation

Infestation of two major rice insects was evaluated i.e., Rice Stem Borer (RSB) and Rice Leaf Miner (RLM). As for RSB, percentage of dead hearts (60 days after transplanting) as well as white heads (10 days before harvest) and then total damage percentage (dead hearts + white heads) of 25 hills/replicate were calculated. For RLM insect, percentage of damaged leaf area (DLA %) (as an average of 40 and 60 days after transplanting results) of 50 leaves/rep. was calculated. The average of infestation of each pest at the two sowing dates was calculated.

C. Chemical evaluation

Samples of the hills/rep. were air-dried for three days and put in an oven at

60°C for 48hr., finely ground and kept for determining total protein, free amino acids and crude silica contents.

Protein and silica contents were assessed according to Yoshida *et al.* (1976). Free amino acids were extracted according to Mengel and Helal (1968), and separated and determined using the paper chromatographic technique of Block *et al.* (1958). The chromatograms were sprayed with 0.2% ninhydrine in acetone w/v as Smith (1958) and then dried for 15 minutes at 60°C. The amino acids in the samples were identified according to the R_f values of known amino acids.

Correlation between infestation and chemical contents was done.

RESULTS AND DISCUSSION

A. Protein content, insect infestation and plant source correlation

As for RSB insect, Table 1 showed the relationship between protein content of the rice plant and the percentage of RSB damage as an average of the two sowing date results. Data showed that protein content differed according to the rice entry. It ranged from 17.61% in Giza-176 resistant, entry of 5% infestation to 27.39% in GZ 4255-8-1, susceptible entry of 7.9%. Telle Hamsa entry that had a high protein content (23.32 %) was severely infested (12.0 %), whereas GZ 3486-34-3 entry which had low protein content (18.21 %) was less infested (5.9 %). Statistical analysis showed a positive correlation between protein contents and RSB infestations ($r=0.45$). Therefore, it appeared that the high protein content in rice plant rendered it more susceptible to RSB infestation. These results are in agreement with Pathak (1967). Some entries which are not in agreement with this rule means that there are other factors which share in plant resistance.

Data summarized in table 2 indicate that entries of japonica source showed more resistance (4.6 %) with lower protein content (20.3%), while those of indica source showed more susceptibility (7.6%) with higher protein content (22%). Similar results have been achieved by Tantawi (1982), Kim *et al.* (1986) and Gu *et al.* (1989). This relation may be attributed to genetic bases.

Concerning RLM insect, table 3 showed the relationship between protein content and percentage of damaged leaf area (DLA %) with RLM insect as an average of the two sowing date results. The statistical analysis showed that there was an in-

significant negative correlation between the percentages of damage in different entries and their protein contents ($r=-0.076$).

This negative correlation can be noticed in GZ 4255-8-1 entry which had low infestation (0.27 % DLM) and the highest protein content (27.39 %), and Gz 4386-34-3 entry which had the highest infestation (0.58 % DLA) and low protein content (18.21%). The same negative correlation appeared in GZ 4565-S-10 and Giza 176 entries. On the other hand, few entries showed a relatively positive correlation like Giza 171 and Todorokiwase. Some entries had the same value of infestation but differed in protein contents (GZ 4255-6-4 and GZ 4255-8-1), while others had different infestation values in spite of the same protein content (Kagahikari and IR 40). These results indicate that protein plays a role in plant resistance but it is not the limiting factor.

Data summarized in table 4 indicated that entries of Indica sources are less infested (0.25 % DLA) with higher protein content (22 %), but those of Japonica source showed higher infestation (0.37 % DLA) and less protein content (20.3%) (Tantawi, 1982).

B. Silica content and insect infestation correlation

As for RSB insect, table 5 showed the relationship between silica content and the percentage of RSD damage as an average of two sowing date results. There was an insignificant negative correlation between damage and silica content ($r = -0.13$). The most susceptible entries Gz 4255-3-1 and Telle Hamsa which showed higher percentage of damage (17.1 and 15.4%) had lower silica contents (6.6 and 6.2 mg/100g dry weight, respectively), while the resistant entries GZ 13868-S-5-4 and IR 40 which showed lower percentage of damage (3.2 and 3.0 %) had higher silica contents (10.2 and 9.5 mg, respectively). However, other entries (Todorokiwase and Kagahikari), in spite of their low silica contents (5.5 and 6.2 mg), received low percentages of damage (2.6 and 2.7%, respectively).

For RLM insect, table 6 showed the correlation between silica contents and the percentage of damaged leaf area (DLA %) of RLM insect. The statistical analysis indicated that there was an insignificant negative correlation between silica content and RLM damage ($r=-0.23$). Some entries showed this correlation as IR 28 that received low infestation (0.30 % DLA) and highest amount of silica (10.4 mg/100 g). While Giza 171 entry, of the highest infestation (0.82 % DLA), contained a low amount of silica (6.0 mg). GZ 4122-23-4-2 and IR 40 rice entries had high silica contents (9.0 and 9.5 mg, respectively) and were subjected to low RLM damage

(0.35% for both), whereas Kagahikari and GZ 4255-3-1 entries had low silica (6.2 and 6.6 mg) and showed high RLM infestation (0.57 and 0.56%, respectively). Some other entries, Giza 172 and GZ 4127-1-3 showed low RLM infestation (0.18 and 0.24% DLA) in spite of their low silica contents (6.2 and 7.2 mg, respectively). On the other hand, GZ 1368-S-5-4 rice entry received high infestation (0.52%), though it had high silica content (10.2 mg/100g).

It was apparent that silica in some entries may play an important role as a chemical resistance factor against RSB and RLM insects. This phenomenon could have been due to the unsuitability of the plants with high silica contents for larval survival and development as reported by Ukwungwu (1984) and Ukwungwu and Odobiya (1985). Silica is not the only factor involved in rice insect resistance, but other resistance factors may share it.

C. Free amino acid (FAA) contents and insect infestation correlation

Little has been done about the possible role of free amino acids in susceptibility of rice plant. Therefore, this study is an attempt to correlate the contents of certain amino acids with the susceptibility of rice entries for RSB and RLM infestation.

Concerning RSB insect, Table 7 showed the relationship between certain FAA contents and RSB damage. Results indicated that there was an insignificant negative correlation between (ornithine + lysine) content and the percentage of damage ($r = -0.15$). Most of rice entries that had low (ornithine + lysine) content (5mg/100 g dry wt) received high infestation but those which had high content (10 mg) received low infestation. The contents of (histidine + arginine) were also insignificantly negatively correlated with the damage ($r = -0.27$). For example, the susceptible entries GZ 4255-3-1 and Telle Hamsa (17.1% and 15.4% damage) contained less amounts of these two amino acids (6.06 and 14.42 mg) than in the resistant entries Giza 176 (6.1 % damage) and TKY - 1024 (5.1 % damage) with (56.2 mg for both). The same correlation was found between (aspartic + serine) contents and insect damage ($r = -0.14$). The maximum amount of these amino acids (94.58 mg) was found in the resistant entry Giza 176 (6.1 % damage), while these amino acids decreased to 23.82 mg in the susceptible one, Telle Hamsa (15.4 % damage). Glycine contents and insect damage followed the same correlation ($r = -0.07$). TKY 1024 entry received low damage (5.1%) and had high glycine content (85.0 mg), while Telle Hamsa entry received higher damage (15.4 %) and had lower glycine content (35.1 mg). Glutamic acid showed an insignificant positive correlation in different tested entries

($r=0.13$). GZ 4122-23-4-2, TKY 1024 and Todorokiwase were the best examples for this correlation. As for the other amino acids, some were negatively correlated with the insect damage i.e., valine ($r=-0.31$), alanine ($r=-0.29$), methionine ($r=-0.29$) and phenylalanine ($r=-0.01$). Telle Hamsa and Todorokiwase are good examples for this correlation. Some other entries which were positively correlated with the insect damage i.e., proline ($r = 0.19$), leucine ($r=-0.14$) and tryptophane ($r = 0.08$), and GZ 4255-3-1, and Giza 180 entries are good examples for this correlation.

For RLM insect, table 8 showed the relationship between amino acid contents and the percentage of damaged leaf area (DLA %) of RLM insect. The statistical analysis showed that this correlation is either positive i.e., some amino acids increase damage as they increase, or negative i.e., some others decrease the damage as they increase. The amino acids that decrease the damage, in order, are (ornithine + lysine) $r = -0.44$, alanine $r = -0.29$, phenylalanine $r = -0.20$, glutamic acid $r = -0.18$ (histidine + arginine) $r = -0.14$, methionine $r = -0.09$, and glycine $r = -0.07$. The other amino acids, that increase the insect damage, in order, are tryptophan $r = 0.32$, proline $r = 0.20$, leucine $r = 0.18$, (aspartic + serine) $r = 0.05$ and valine $r = 0.04$. For more elucidation, (ornithine + lysine) were mentioned to have the highest negative effects $r = -0.44$. The results showed that the entries with higher contents of these two amino acids (10.00 mg), which are Giza 176; IR 40 and Giza 178, had lower insect damage (0.39, 0.35 and 0.18% DLA, respectively) and vice versa. Tryptophan was found to show the highest positive correlation $r = 0.32$. Data also showed that the maximum value of tryptophan (53.64 mg) was in the susceptible entry GZ 1386-S-5-4 (0.52 % DLA), while this decreased to 8.24 mg in the resistant one GZ 4127-2-1-3 (0.24 % DLA).

As a general result, most of the high concentrations of the negatively effected amino acids (alanine, histidine + arginine, glycine and aspartic + serine) increase the plant resistance against RSB infestation and so do (ornithine + lysine, histidine + arginine, glutamic and methonine) against RLM infestation. There is a possibility that these compounds possess inhibitory effect on insect feeding or growth and thus reduce the damage inflicted on the plant. While most of the high concentrations of the amino acids of the positive effects (leucine and proline) decrease the plant resistance against RSB infestation and so do (tryptophan, valin, leucine, proline and aspartic + serine) against RLM infestation. That may be due to certain activating effects of these amino acids on plant growth to become more attractive for insect attack (Kamano, 1971). Some entries had low infestation in spite of having low toxic

REFERENCES

- 1 . Block, R.J., E.L. Durrum and G. Zweig. 1958. A manual of paper chromatography and paper electrophoresis. Academic Press, Inc. Publishers, New York.
- 2 . Gu, Y., Y.F. Xiao and Y.M. Wang. 1989. Resistance of main rice varieties or lines to brown planthopper, white backed planthopper and stripped stem borer in Jiangsu. *Jiangsu. J. Agric. Sci.*, 5 (2) : 38-41.
- 3 . Ishii, S. and C. Hirano. 1955. Qualitative studies on the essential amino acids for the growth of larvae of the rice stem borer, *Chilo agamemnon* Butler, under aseptic conditions. *Bull. Natl. Inst. Agric. Sci., Tokyo (c)*, 5 : 35-48 .
- 4 . Kamano, S.I. 1971. Studies on artificial diets of the rice stem borer *Chilo suppressalis* Walker. *Bull. Natl. Inst. Agric. Sci., Jpn. Ser. (C)* 25 : 1-45.
- 5 . Kim, Y.A., J.O. Lee and H.G. Goh. 1986. Resistance of Tongil (japonica x indica) and japonica varieties to *Chilo suppressalis*. Res. Reports of the Rural Development Administration, Plant Environment, Mycology and Farm-Products, Utilization Korea Republic, 28 (1) : 90-93.
- 6 . Mengel, K. and H.M. Helal. 1968. The effect of varied nitrogen and potassium nutrition on the content of soluble amino compounds in aerial parts of oats. *Z.Pflanzenernaehr. Bodenk*, 120 (1) 12-20.
- 7 . Pathak, M.D. 1967. Insect pests of rice at the International Rice Research Institute, Los Banos, Laguna, Philippines, 68 p.
- 8 . Smith, I. 1958. Amino acids, amines and related compounds (Chapter 4), Chromatographic Techniques, Clinical and Biochemical Applications. William Heinemann, Medical Book Ltd., London Interscience Publishers, INC, New York.
- 9 . Tantawi, A.M. 1982. Rice stem borer, *Chilo agamemnon* Bles. infestation in relation to certain rice varieties. *Proc. Egypt. Nat. Dec.*, 1:215-222.
- 10 . Ukwungwu, M.N. 1984. Effect of silica content of rice plants on damage caused by the larvae of *Chilo zacconius*. *WARDA Tech. News* 1., 5 (1) : 20-21.
- 11 . Ukwungwu, M.N. and J.A. Odobi. 1984. Yield losses in resistant and susceptible varieties of rice in Nigeria due to *Chilo zacconius* and other stem borers. *Trop Pest Manage.*, 30 (3) : 291-295.
- 12 . Yoshida, S., D.A. Forno, J.H. Cock and K.A. Gomez. 1976. Laboratory manual for physiological studies of rice. Third Edition, International Rice Research Institute, Los Banos Laguna, Philippines, 83 p.

تأثير بعض المكونات الكيميائية فى نبات الأرز وموطنه الأصيل على درجة مقاومته لثاقبة ساق الأرز وصانعة الأنفاق

على محمود سليمان^١، وفاء محمد العطار^٢، رفعت ابو العلا^٣.

أحمد عزت عبد الوهاب^٤

١ معهد بحوث وقاية النباتات - مركز البحوث الزراعية - الدقى .

٢ كلية الزراعة - جامعة الزقازيق.

٣ كلية العلوم - جامعة القاهرة.

٤ معهد بحوث المحاصيل الحقلية - مركز البحوث الزراعية - الجيزة.

تم زراعة أصناف وسلالات مختلفة من الأرز معلومة الموطن، وتم تقييمها ضد حشرتى ثاقبة ساق الأرز وصانعة الأنفاق لمعرفة درجة إصابتها، كما تم تحليلها كيميائياً لتقدير كل من البروتين الخام والسلكا الخام وخمسة عشر حمض أمينى حر فى كل صنف وبالتحليل الإحصائى تم حساب الارتباط بين هذه المكونات والإصابة فى كل صنف، وتم الحصول على النتائج التالية :

- البروتين يزيد الإصابة بثاقبة ساق الأرز فى معظم الأصناف المختبرة بنسبة ارتباط (٣ = ٠.٤٥)، بينما تقل الإصابة بصانعة الأنفاق فى معظمها بنسبة ارتباط (٣ = ٠.٧٧) .

- السلكا تقلل الإصابة بكلتا الحشرتين فى معظم الأصناف المختبرة بنسبة ارتباط (٣ = ٠.١٣) .
للثاقبة ، = ٠.٢٣ . لصانعة الأنفاق) .

- الأحماض الأمينية الحرة أظهرت تأثيرات مختلفة حسب التركيز ونوع الحشرة. فبعضها مثل الألانين ، جليسين، هستيدين مع الأرجنين ، الأسبارتيك مع السيرين ، والفالين قللت الإصابة بثاقبة ساق الأرز فى معظم الأصناف وبدرجات ارتباط مختلفة ، وبعضها الآخر مثل الأورنيثين مع الليسين ، هستيدين مع الأرجنين ومثيونين قد قللت الإصابة بصانعة الأنفاق بدرجات ارتباط متفاوتة . وهناك أحماض أخرى أدت الى زيادة الإصابة باى من الحشرتين .

- الأصناف اليابانية الاصل اكثر مقاومة لثاقبة ساق الارز من الهندية ، بينما الأصناف الهندية كانت اكثر مقاومة لصانعة الأنفاق من اليابانية مع وجود نسبة اقل من البروتين فى الأصناف المقاومة.

وعلى ذلك يمكن القول بان بعض المكونات الكيميائية قد تلعب دورا فى المقاومة الصنفية لكنها ليست هى العامل المحدد لها بل يشاركها فى ذلك عوامل مورفولوجية وهستولوجية وجينية اخرى.