

Research article

Mitigating Physiological and Histological Changes Induced by a High-Salt Diet in Male Albino Rats Using *Origanum majorana* and *Olea europaea* Leaf Extracts

Mohamed Gaber Shalan *

Zoology and Entomology Department, Faculty of Science, Arish University, Egypt

HIGHLIGHTS

- High NaCl feeding induces hypertension.
- Protective effects of marjoram and olive leaf extracts.
- A high-salt diet disrupted liver and kidney functions.
- High-salt diet caused sperm abnormalities and histopathological changes in vital organs.
- Restoring balance with natural extracts.

ARTICLE INFO

ABSTRACT

Keywords:

Marjoram
Olive
Sodium Chloride
Oxidative stress,
Histological alterations

This research explored the possible protective impact of marjoram and olive leaf extracts against hypertension triggered by a salt-rich diet in rats. Twenty male albino rats (*Rattus norvegicus*) were separated into four equal groups; the first group was the reference control, and the second group was provided 250 mg/Kg body weight of *Origanum majorana* leaves extract and 250 mg/Kg body weight of *Olea europaea* leaves extract daily. The third group was given freely a high 8% NaCl diet. The fourth group was administered an 8% salt diet supplemented with 250 mg/Kg body weight of *O. majorana* leaves extract and 250 mg/Kg body weight of *O. europaea* leaves extract. Samples were taken after 4 weeks of treatments. Feeding rats, a high sodium chloride diet consisting of 8% NaCl induces hypertension in rats. The high salt diet enhanced marked elevation in body weights and reduced most relative organ weights. The high salt feeding elevated sodium levels in hepatic and kidney tissues, in addition to a rise in the liver's MDA level. Feeding on a high salt diet reduced superoxide dismutase and catalase activities. It promoted considerable disruption in liver and kidney functions. It also triggered a decrease in sperm count and an increase in abnormal sperm morphology. The high salt feeding induced histological changes in the liver, kidney, brain, and testes of rats. Combined supplementation with both *O. majorana* and *O. europaea* extracts alleviated salt-induced abnormalities in rats.

1. Introduction

High sodium chloride (NaCl) consumption is a major dietary factor contributing to multiple organs' widespread physiological and histopathological alterations [1-3]. The primary mechanism involves the disruption of cellular and systemic homeostasis [4], leading to oxidative stress, inflammation, and lipid peroxidation (LPO) [5]. Chronic high salt consumption is closely linked to the development of hypertension [6] by disruptions of fluid balance and vascular function [7] and promotes systemic organ damage [8].

* Corresponding author

E-mail address: mohammed.shalaan@sci.aru.edu.eg

<https://doi.org/10.21608/APJ.2025.347045.1000>

Received 23 December 2025; Received in revised form 21 January 2025, Accepted 21 January 2025

3062-4878/© 2025 The Author (s) Published by Arish University. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>)

Excessive sodium intake increases extracellular fluid volume by promoting water retention, which elevates blood pressure [9]. Additionally, high NaCl intake impairs the renin-angiotensin-aldosterone system (RAAS), resulting in vasoconstriction and elevated vascular resistance [10]. Sodium also influences vascular endothelial cells, causing oxidative stress, decreased nitric oxide (NO) availability, and impaired endothelial function contributing to cardiovascular complications [11].

High-salt diets exacerbate oxidative damage by promoting the generation of reactive oxygen species (ROS) and depleting the body's natural antioxidant reserves [12]. This oxidative imbalance drives LPO, damaging cellular membranes and contributing to systemic inflammation [13]. High NaCl consumption induces pathological changes in multiple organs. The liver experiences metabolic disruptions, oxidative damage, and inflammatory responses [14,15]. In the kidneys, high sodium intake accelerates fibrosis, impairs filtration, and promotes hypertension-related damage [16,17]. In the brain, sodium chloride contributes to neuroinflammation and oxidative damage, affecting cognitive functions [18,19]. The testes are also vulnerable, with oxidative stress impairing spermatogenesis and hormonal balance, potentially affecting reproductive health [20,21].

Given the systemic nature of these effects, there is increasing interest in natural, plant-derived interventions for mitigating the damage caused by high salt intake. Extracts from *Origanum majorana* (marjoram) and *Olea europaea* (olive) have shown potential in addressing oxidative stress and histological damage [22,23]. Marjoram is rich in flavonoids and polyphenols recognized for its strong radical scavenger and anti-inflammatory effects. At the same time, olive leaves are abundant in oleuropein and hydroxytyrosol, which improve vascular health and reduce oxidative stress [24,25]. Combined supplementation with olive and marjoram leaf extracts offers a promising strategy, as their bioactive compounds may synergistically mitigate oxidative stress, reduce LPO, and improve organ function.

Despite their benefits, research examining the combined use of olive and marjoram extracts in addressing high salt-induced hypertension and systemic organ damage are scarce. Combining these extracts offers a promising protective strategy, as their complementary bioactive compounds may exert synergistic effects in countering the physiological and histological damage induced by high salt feeding. Therefore, the current study investigated the combined efficacy of *O. majorana* and *O. europaea* leaf extracts in mitigating NaCl-induced alterations in rat organs.

3. Materials and methods

3.1. Chemicals

Pure sodium chloride was purchased from El Naser Pharmaceutical Chemicals Co., Egypt.

3.2. Animals

Twenty male albino rats (*Rattus norvegicus*) four weeks old, weighing (70 ± 5 g), were obtained from the Animal care facility, Faculty of Science, Suez Canal University, Egypt. The rats were housed in plastic housing and maintained on a 12-hour light-dark cycle with regulated temperature ($25 \pm 2^\circ\text{C}$) and relative humidity ($45 \pm 5\%$). They were provided with a standard balanced diet and water *ad libitum*, and allowed a 7-day acclimatization period to the laboratory conditions. The experimental procedures was evaluated and authorized by the Research Ethics Committee of the Faculty of Science at Arish University, under approval number "ARU/SF.13". Our techniques followed the guidelines outlined in the 8th edition of the guide for the care and use of laboratory animals, ensuring strict adherence to ethical standards.

3.3. Preparation of marjoram and olive leaf extracts

The leaves of marjoram (*O. majorana*) and Olive (*O. europaea*) were collected from the Arish area, North Sinai, Egypt. The plants were examined and classified at the Faculty of Science, Arish University, by a specialist professor in plant taxonomy. The plant leaves were dried by leaving them at room temperature for one week. The desiccated leaves were crushed into powder. Ten grams of desiccated powder of each plant were dissolved in 100 mL of 70% ethyl alcohol to prepare a 10% (w/v) stock solution [26,27]. The mixture is stirred, and placed in a cold and dark place for 48-72 hours to extract the effective substances. Then, the extract was filtered. A rotary evaporator was used to evaporate ethanol and get the dry extract, which was then stored in a tight and dark container, away from humidity and excess heat to preserve the extract's effectiveness.

3.4. Experimental design

After acclimatization, the rats were haphazardly assigned into four equal groups. The first group acted as the normal control, the second group received 250 mg/kg body weight of *O. majorana* leaves extract [28] and 250 mg/kg body weight of *O. europaea* leaves extract [29] daily. The third group was fed an unrestricted 8% NaCl diet [30]. The fourth group was given an 8% salt diet, supplemented with 250 mg/kg body weight of *O. majorana* leaves extract and 250 mg/kg body weight of *O. europaea* leaves extract.

3.5. Duration time

Upon completion of the experiment (28 days), all rats were euthanized utilizing ketamine (50 mg/kg body weight, intramuscular injection) and rapidly dissected. Tissues were removed, weighted, blotted on filter paper, and kept at –30°C till additional biochemical examinations. The body weights of rats were taken weekly.

3.6. Non-invasive blood pressure measurement

The systolic and diastolic blood pressures of the rats were evaluated non-invasively with a tail-cuff plethysmography system (CODA@Monitor, Kent Scientific, Torrington, CT, USA), incorporating a specialized retraining chamber designed for animal comfort and stability.

1. **Acclimatization:** Rats were acclimated to the restrainer and device for five consecutive days (10-15 minutes per day) to minimize stress-related variability. The chamber allowed the tail to extend freely for easy access, reducing handling stress.
2. **Chamber design:** The chamber was constructed to securely restrain the animal's body while leaving the tail accessible. The chamber size was customized to prevent unnecessary movement, ensuring consistent positioning during measurements.
3. **Setup:** Measurements were conducted in a temperature-controlled environment (30-35 °C) to ensure tail vasodilation and optimal blood flow.
4. **Measurement protocol:** The rat was placed into the chamber, and its tail extended through the opening. A pneumatic cuff with a sensor was placed around the proximal tail. The cuff was inflated to occlude blood flow and then deflated gradually, with oscillometric signals recorded to determine systolic and diastolic pressures.
5. **Data collection and validation:** Each rat underwent three consecutive measurements, and the average was used for analysis. Outliers caused by movement artifacts were excluded, and readings were repeated if necessary.

3.7. Tissue preparation for microscopic examination

Following the dissection of the animals, the liver, kidney, testes, and brain were extracted, dried on filter paper, and weighed. Representative samples were then placed in a 10% formalin solution and processed into paraffin for histological sectioning and staining.

3.8. Histological methods

Histological sections, 5 microns thick, were formulated and stained with hematoxylin and eosin. Microscopic examination of the samples was performed blindly using an Olympus BX43 microscope [31].

3.9. Biochemical analysis

Sodium was measured using spectrum kits, Cairo, Egypt, according to Henry et al. [32]. Oxidative stress parameters were assessed using BioDiagnostic kits (Cairo, Egypt). Catalase (CAT) activity was assessed using the Aebi method [33]. The concentration of malondialdehyde (MDA) was evaluated using the method of Ohkawa et al. [34]. Alanine aminotransferase (ALT) and aspartate aminotransferase (AST) activities were quantified using the modified version of the method of Schumann and Klauke [35]. Uric acid levels was quantified utilizing Spinreact kits (Spain) and the

technique of Fossati et al. [36]. Creatinine was determined with Diamond Diagnostics kits (Holliston, USA) using the Heinegard and Tiderstrom technique [37].

3.10. Statistical analysis

The results are presented as means \pm standard deviation (SD) for five rats per group. Data from the control and treated groups were analyzed using a one-way analysis of variance (ANOVA), followed by Tukey's test for multiple comparisons. A probability-value of <0.05 was considered statistically significant. Data analysis was conducted leveraging the statistical Package for Social Sciences (SPSS) software, version 22.0 for windows.

4. Results

4.1 Body weights

The results showed that salt-rich diet feeding induced considerable elevation ($P<0.05$) in body weights after two and four weeks of treatment, by 38.0% and 33.25%, respectively (Fig. 1). On the other hand, combined supplementation with both olive and marjoram extracts restores normal levels.

4.2 Organ weights

The results showed that high salt feeding induced significant ($P<0.05$) decrease in relative liver and testicular weights by 17.94% and 44.5% after 4 weeks of treatment, respectively. However relative kidney and brain weights remain unaffected after 4 weeks of treatment (Fig. 1). On the other hand, combined supplementation with both olive and marjoram extracts not only reversed these effects, but also elevated relative kidney, brain, and testis significantly ($p<0.5$) by 42.6, 65.76 and 46.07%, respectively, compared with normal controls after 4 weeks of treatment.

4.3 Blood pressure

The 8% salt feeding induced a considerable elevation in systolic blood pressure of rats after 2 and 4 weeks of treatment with magnitudes of 19.25 and 22.35% compared with normal controls, respectively (Fig. 2). It also enhanced significant ($P<0.05$) elevation in diastolic blood pressure by 24.71 and 28.94% compared with normal controls following 2 and 4 weeks of treatment, respectively. In contrast, combined supplementation with marjoram and olive extracts restores blood pressure to approach control.

4.4 Biochemistry analysis

After 4 weeks of high sodium chloride intake, there was a notable rise in MDA levels in hepatic tissue by 131% and sodium levels in liver and kidney tissues by 21.95% and 17.87%, respectively in comparison to the normal control group (Fig. 3). SOD and CAT activities were reduced significantly ($p<0.05$), by 21.87 and 27.38% compared to normal rats, upon 4 weeks of high NaCl administration, respectively. It enhanced considerable disruption in hepatic (AST & ALT) and renal (urea & creatinine) functions (Table 1). High salt feeding caused a substantial decrease in sperm count, and considerable elevation in abnormal sperm morphology; however, plasma testosterone and sperm motility are approximately not affected (Table 2). Nevertheless, supplementation with both marjoram and olive leaf extracts ameliorated these effects.

4.5 histopathology findings

The tissue architecture of the kidney, liver, rain, and testis in the control group displayed normal cellular organization, with well-defined structures and appropriate cellular integrity. Similarly, treatment with *O. majorana* and *O. europaea* leaves extracts resulted in tissue architectures of these organs that were more or less similar to the control group (Fig. 4-7), showing preserved cellular morphology and minimal disruption. The effects of high NaCl feeding on rats' histopathology were detected. A marked impairment of the renal tubules through glomeruli congestion, signs of tubular cell lining vacuolation, and intra-tubular hemorrhage could be seen after 4 weeks of salt ingestion (Fig. 4). Hepatocytes showed the formation of vacuoles in the cytoplasm and abnormal structure organization after chronic high salt ingestion (Fig. 5). On the other hand, brain cells displayed signs of cell atrophy and surrounded vacuolation after administering 8% salt diet for 28 days (Fig. 6). It was observed that the germinal testicular epithelium was adversely affected, as evidenced by the wide separation of the cells and a decrease in

spermatid count within the lumen in rats chronically given high salt diet (Fig. 7). Conversely, the dosage of *O. majorana* and *O. europaea* leaf extracts significantly improved tissue damage. These findings suggest that both plant extracts offer protective effects against potential damage caused by high NaCl exposure.




Fig. 1. The impact of high salt feeding on body weight (A) and relative organ weights (B) (g/100 g bw) of rats. Om: *O. majorana*, Oe: *O. europaea*; HS: high salt; bw: body weight. Data are presented as means \pm SD for five rats in each group. a: Significantly different from control group at $p < 0.05$ (Tukey's *post-hoc* test), b: significantly different from *O. majorana* and *O. europaea* extracts-treated group at $p < 0.05$ (Tukey's *post-hoc* test), c: significantly different from high salt group at $p < 0.05$ (Tukey's *post-hoc* test), d: significantly different from high salt + *O. majorana* + *O. europaea* extracts-treated group at $p < 0.05$ (Tukey's *post-hoc* test).




Fig. 2. Effect of high salt feeding on mean systolic and diastolic blood pressure in rats after 2 weeks of treatment (A) and 4 weeks of treatment (B). Om: *O. majorana*, Oe: *O. europaea*; HS: high salt; Data are presented as means \pm SD for five rats in each group. a: Significantly different from the control group at $p < 0.05$ (Tukey's *post-hoc* test), b: significantly different from *O. majorana* and *O. europaea* extracts-treated group at $p < 0.05$ (Tukey's *post-hoc* test), c: significantly different from high salt group at $p < 0.05$ (Tukey's *post-hoc* test), d: significantly different from high salt + *O. majorana* + *O. europaea* extracts-treated group at $p < 0.05$ (Tukey's *post-hoc* test).

5. Discussion

It has been demonstrated that high salt ingestion significantly elevates the body weight of rats, likely due to its effects on sodium levels, fluid retention, and appetite regulation. Increased sodium disrupts fluid balance, leading to water retention as the body attempts to maintain sodium homeostasis [38,39]. Additionally, high salt intake promotes thirst and water consumption, which further contributes to weight gain [40]. Evidence also suggests that salt enhances appetite, potentially important to increased food intake and fat accumulation [41]. Chronic salt consumption alters lipid metabolism and affects hormones such as leptin and insulin, disrupting appetite regulation

and promoting fat storage [42-44]. Furthermore, chronic high salt intake induces low-grade inflammation, which has been implicated in promoting fat storage and weight gain [45]. Alterations in the gut microbiota due to high salt intake may also impact digestion and nutrient absorption, influencing body weight [46]. At the cellular level, NaCl exposure can principal to intracellular fluid accumulation and tissue swelling, contributing to increased body weight [47]. Stress-related pathways, such as RAAS, are also activated by high salt consumption, promoting fat deposition, reducing energy expenditure, and increasing hypertension [43].




Fig. 3. Effects of high salt feeding for four weeks on plasma superoxide dismutase (A) and catalase (B) activities, and malondialdehyde (C) level, and sodium concentration in the liver (D) and kidney (E) of rats. Om: *O. majorana*, Oe: *O. europaea*; HS: high salt. Data are presented as means \pm SD for five rats in each group. a: Significantly different from control group at $p < 0.05$ (Tukey's *post-hoc* test), b: significantly different from *O. majorana* and *O. europaea* extracts-treated group at $p < 0.05$ (Tukey's *post-hoc* test), c: significantly different from high salt group at $p < 0.05$ (Tukey's *post-hoc* test), d: significantly different from high salt + *O. majorana* + *O. europaea* extracts-treated group at $p < 0.05$ (Tukey's *post-hoc* test).

Table 1. Effects of high dietary salt intake on liver and kidney function tests and the protective effect of *O. majorana* (Om) and *O. europaea* (Oe) supplementation in male albino rats.

28 days	ALT (U/L)	AST (U/L)	Uric acid (mg/dL)	Creatinine (mg/dL)
Control	30.3 \pm 4.74	85.5 \pm 12.9	1.76 \pm 0.28	0.48 \pm 0.07
Om + Oe	28.3 \pm 3.79 ^{c,d}	83.6 \pm 12.5 ^c	1.60 \pm 0.25 ^{c,d}	0.47 \pm 0.06 ^c
HS	48.2 \pm 6.44 ^{a,b,d}	114.5 \pm 16.9 ^{a,b}	3.12 \pm 0.46 ^{a,b,d}	0.65 \pm 0.09 ^{a,b}
Om + Oe + HS	38.7 \pm 5.38 ^{b,c}	103.2 \pm 14.0	2.48 \pm 0.35 ^{a,b,c}	0.53 \pm 0.08

Data are presented as mean \pm SD (n = 5). a: Significantly different from normal control group at $p < 0.05$ (Tukey's *post-hoc* test), b: significantly different from *O. majorana* and *O. europaea* extracts-treated group at $p < 0.05$ (Tukey's *post-hoc* test), c: significantly different from high salt group at $p < 0.05$ (Tukey's *post-hoc* test), d: significantly different from high salt + *O. majorana* + *O. europaea* extracts-treated group at $p < 0.05$ (Tukey's *post-hoc* test).

Table 2. Effects of high dietary salt intake on plasma testosterone, sperm parameters, and the protective effect of *O. majorana* (Om) and *O. europaea* (Oe) supplementation in male albino rats.

28 days	Testosterone (ng/dL)	Sperm count x 10 ⁶ /mL	Sperm motility (%)	Abnormal sperm morphology (%)
Control	26.3 ± 4.37	100.2 ± 15.0	93.0 ± 14.6	4.4 ± 0.54
Om + Oe	27.5 ± 4.39 ^c	103.4 ± 11.9 ^c	95.0 ± 13.8	3.20 ± 0.83 ^c
HS	19.8 ± 2.90 ^b	76.40 ± 10.8 ^{a,b}	74.2 ± 11.4	7.00 ± 2.54 ^{a,b}
Om + Oe + HS	23.1 ± 3.28	88.80 ± 10.7	80.0 ± 12.2	540 ± 0.54

Data are presented as mean ± SD (n = 5). a: Significantly different from normal control group at $p < 0.05$ (Tukey's *post-hoc* test), b: significantly different from *O. majorana* and *O. europaea* extracts-treated group at $p < 0.05$ (Tukey's *post-hoc* test), c: significantly different from high salt group at $p < 0.05$ (Tukey's *post-hoc* test).

A high-salt diet has reflective effects on blood pressure, mainly through increased plasma sodium concentrations that lead to fluid retention, increase in blood volume, and heightened cardiac output [48]. The vascular effects of salt include increased stiffness and reduced relaxation of blood vessels, leading to elevated peripheral resistance and hypertension [49,50]. Dysregulation of the RAAS under chronic salt exposure raises angiotensin II levels, causing vasoconstriction and elevated systolic and diastolic blood pressure [51]. Salt also disrupts endothelial function, decreasing nitric oxide synthesis and raising oxidative stress, which exacerbates vascular dysfunction [52].

Figure 4




Fig. 4. Histological sections of the rat kidney after 4 weeks of treatment, show a section in the cortex of the control (A & B) rat with a regular structure. The marjoram and olive extracts-treated group (C & D) is more or less similar to the control group. The high salt-fed group (E & F) shows marked affection for the renal tubules through glomeruli congestion (blue arrow), signs of tubular cell lining vacuolation (black arrow), and intratubular hemorrhage. The marjoram and olive extracts + high salt-fed group (G & H) shows minimal affection for the renal tubules with almost normal glomeruli and tubules (X 200: H & E section A, C, E & G; X 400: H & E section B, D, F & H).

Figure 5




Fig. 5. Histological sections of the rat liver after 4 weeks of treatment, show a section in the liver of the control (A & B) rat with the typical arrangement of the hepatocytes around the central veins. The marjoram and olive extracts-treated group (C & D) is more or less similar to the control group. The high salt-fed group (E & F) shows marked affection for the hepatocytes, which appear with vacuolated cytoplasm (black arrowhead) and distorted arrangement. The marjoram and olive extracts + high salt-fed group (G & H) shows minimal affection for liver parenchyma with more or less standard structure (X 200: H & E section A, C, E & G; X 400: H & E section B, D, F & H).

Oxidative stress is a major result of high salt consumption, as indicated by increased plasma malondialdehyde (MDA) levels, a marker of LPO [53]. The production of reactive oxygen species (ROS), including superoxide anions and hydroxyl radicals, leads to damage in cellular structures like lipids, proteins, and DNA [53,54]. MDA reacts with cellular components to form advanced glycation end products (AGEs), which further lead to tissue dysfunction [54]. The mechanisms underlying increased ROS production include monocyte activation, NADPH oxidase pathway dysregulation, mitochondrial dysfunction, and inflammation [55]. Pro-inflammatory cytokines, including TNF- α , IL-1, IL-6, and IL-33, further amplify ROS production and tissue damage [56]. These processes also reduce activities of antioxidant enzyme, such as CAT and superoxide SOD, which impair the body's ability to neutralize ROS [57,58]. Prolonged salt consumption leads to a suppression of antioxidant defense mechanisms, exacerbating oxidative damage over time [21,59].

Chronic salt intake stimulates the sympathetic nervous system, causing hypertension, which increases the risk of cardiovascular diseases such as hypertension, heart failure, stroke, and kidney disorder [60, 61]. It exerts detrimental effects on multiple organs. In the kidneys, it increases MDA levels, promoting inflammation, fibrosis, and glomerular damage [62]. This results in reduced filtration efficiency and elevated plasma urea and creatinine concentrations [63,64]. Prolonged exposure to high salt levels leads to glomerulosclerosis and reduced renal blood flow, impairing waste elimination and increasing intra-glomerular pressure reducing the efficiency of the kidneys in filtering waste products [63,65]. In the liver, high salt intake disrupts hepatocyte integrity through oxidative stress, LPO, and mitochondrial impairment, resulting in vacuolation and impaired metabolic functions [66,67]. Hepatocyte damage is further aggravated by inflammation, and immune response, which interferes with the liver's detoxification and protein synthesis abilities [68-70]. Chronic high salt intake disrupts cellular osmotic balance, causing hepatocyte swelling, membrane damage, and the release of AST and ALT enzymes [66,71]. It also induces mitochondrial damage and endoplasmic reticulum dysfunction, leading to protein misfolding, hepatocyte death, and enzyme leakage [72].

Figure 6




Figure 7




Fig. 6. Histological sections of the rat brain after 4 weeks of treatment, show a section in the cerebral cortex of the control (A & B) rat with a standard structure. The marjoram and olive extracts-treated group (C & D) is more or less similar to the control group. The high salt-fed group (E & F) shows marked affection for the brain cells with signs of cell atrophy and surrounded vacuolation (blue arrow). The marjoram and olive extracts + high salt-fed group (G & H) shows almost normal cerebral cortex nerve cells (X 200: H & E section A, C, E & G; X 400: H & E section B, D, F & H).

Fig. 7. Histological sections of rat testis after 4 weeks of treatment, show a section in the testis of a control (A & B) rat with the standard structure of the seminiferous tubules with regular germinal epithelium arrangement. The marjoram and olive extracts-treated group (C & D) is more or less similar to the control group. The high salt-fed group (E & F) shows marked affection for the germinal epithelium through wide separation of the cells (black arrow) and diminished spermatids in the lumen (black arrowhead). The marjoram and olive extracts + high salt-fed group

(G & H) shows minimal affection for the renal tubules with almost normal seminiferous tubules (X 200: H & E section A, C, E & G; X 400: H & E section B, D, F & H).

The brain is also vulnerable to high salt exposure, with evidence of neuronal atrophy and vacuolation in brain cells due to hypernatremia-induced osmotic stress and inflammation [73,74]. Pro-inflammatory cytokines, such as IL-4, IL-6, and TNF- α , lead to neuronal damage and impaired hippocampal neurogenesis [75,76]. Oxidative stress impairs mitochondrial function and synaptic integrity, resulting in impairments in learning, memory, and executive tasks [77-81]. A high-salt diet also compromises the blood-brain barrier, allowing harmful substances to infiltrate the brain and exacerbate neurodegenerative processes [82-84]. Loss of structural integrity in brain cells can manifest deficits in learning, memory, and executive functions [85,86].

In the male reproductive system, high salt intake reduces testosterone levels and impairs sperm quality, including motility and morphology [21,87]. Oxidative stress in testicular tissue disrupts Leydig and Sertoli cell function, primary to reduced testosterone production and abnormal spermatogenesis [88-92]. Increased ROS production damages sperm DNA and plasma membranes, resulting in reduced fertility potential [93]. Sodium-induced osmotic stress disrupts the germinal epithelium, further impairing sperm development [94,95].

The administration of *O. majorana* and *O. europaea* extracts demonstrated significant protective effects against the detrimental impacts of high salt intake. These extracts, rich in flavonoids, polyphenols, and terpenoids, exhibited strong antioxidant and anti-inflammatory properties [96-98]. They reduced oxidative stress by neutralizing ROS and enhancing antioxidant enzyme activities, protecting tissues from LPO and structural damage [99-101]. Both extracts also restored metabolic and hormonal functions, improving nutrient absorption and utilization [102,103].

In the kidneys, these extracts reduced sodium-induced inflammation and fibrosis, preserving nephron integrity and enhancing waste elimination [104-110]. In the liver, they protected hepatocytes from LPO and oxidative stress, maintaining cellular architecture and metabolic functions [111-116]. Neuroprotective effects included reduced neuroinflammation, oxidative damage, and vacuolation, improving cognitive function and preserving neuronal integrity [117-121]. In the reproductive system, *O. majorana* restored testosterone levels and protected the germinal epithelium, while *O. europaea* improved sperm quality and reduced morphological abnormalities through its antioxidant properties [122-124].

Overall, *O. majorana* and *O. europaea* extracts offer promising therapeutic potential to counteract the negative impacts of chronic high salt intake on body weight, organ function, and oxidative stress. Additional studies is necessary to explain their molecular processes and assess their prolonged efficacy in mitigating salt-induced damage.

5. Conclusion

In conclusion, the investigation provides compelling evidence that high salt intake contributes to a range of physiological and biochemical alterations, including weight gain, hypertension, oxidative stress, organ damage, and impaired reproductive health. The study also highlights the potential of *O. majorana* and *O. europaea* extracts as therapeutic agents that can mitigate these harmful effects. By enhancing antioxidant defense mechanisms and reducing inflammation, these extracts offer a promising strategy to counteracting the harmful consequences of chronic salt consumption on various organs and systems. Additional research is required to clarify the basic molecular mechanisms and to confirm the extended-term efficacy of these extracts in preventing salt-induced damage.

Credit authorship contribution statement

Mohamed Gaber Shalan: conducted experimental design, all testing procedures and measurements, statistical analysis, and wrote the manuscript.

Ethics statements

The experimental procedures were evaluated and authorized by the Research Ethics Committee of the Faculty of Science at Arish University under approval number "ARU/SF.13". Our techniques adhere to the 8th edition guide for the care and use of laboratory animals, ensuring rigorous ethical standards.

Funding

This study did not receive any specific funding from public, commercial, or non-profit funding agencies.

Data availability statement

The data will be made available upon request.

Declaration of competing interest

The authors declare no known competing financial interests or personal relationships that could have influenced the work presented in this paper.

List of abbreviations

ALT	Alanine aminotransferase
AST	Aspartate aminotransferase
CAT	Catalase
DNA	Deoxyribonucleic acid
EDTA	Ethylenediaminetetraacetic acid
FSH	Follicle-stimulating hormone
HS	High salt
IL	Interleukin
LH	Luteinizing hormone
MDA	Malondialdehyde
LPO	Lipid peroxidation
NaCl	Sodium chloride
NADPH	Nicotinamide adenine dinucleotide phosphate hydrogen
No	Nitric oxide
RAAS	Renin-angiotensin-aldosterone system
ROS	Reactive oxygen species
SD	Standard deviation

SOD	Superoxide dismutase
TNF- α	Tumor necrosis factor-alpha

References

- [1] Stocker SD, Kinsman BJ, Farquhar WB, Gyarmati G, Peti-Peterdi J, Sved A. FPhysiological Mechanisms of Dietary Salt Sensing in the Brain, Kidney, and Gastrointestinal Tract. *Hypertension* 2024;81(3):447–55, <https://doi.org/10.1161/HYPERTENSIONAHA.123.19488>.
- [2] Zostawa J, Adamczyk J, Sowa P, Adamczyk-Sowa M. The influence of sodium on pathophysiology of multiple sclerosis. *Neurol Sci.* 2017;38(3):389-98, <https://doi.org/10.1007/s10072-016-2802-8>.
- [3] Ghanem M, Zeineldin M, Eissa A, El Ebissy E, Mohammed R, Abdelraof Y. The effects of saline water consumption on the ultrasonographic and histopathological appearance of the kidney and liver in Barki sheep. *J Vet Med Sci.* 2018;17:596.
- [4] Bkaily G, Simon Y, Jazzar A, Najibeddine H, Normand A, Jacques D. High Na⁺ Salt Diet and Remodeling of Vascular Smooth Muscle and Endothelial Cells. *Biomedicines* 2021;9(8):883, <https://doi.org/10.3390/biomedicines9080883>.
- [5] Krajina I, Stupin A, Šola M, Mihalj M. Oxidative Stress Induced by High Salt Diet-Possible Implications for Development and Clinical Manifestation of Cutaneous Inflammation and Endothelial Dysfunction in Psoriasis vulgaris. *Antioxidants* 2022;11(7):1269, <https://doi.org/10.3390/antiox11071269>.
- [6] Grillo A, Salvi L, Coruzzi P, Salvi P, Parati G. Sodium Intake and Hypertension. *Nutrients* 2019;11(9):1970, <https://doi.org/10.3390/nu11091970>.
- [7] Patik JC, Lennon SL, Farquhar WB, Edwards DG. Mechanisms of Dietary Sodium-Induced Impairments in Endothelial Function and Potential Countermeasures. *Nutrients* 2021;13(1):270, <https://doi.org/10.3390/nu13010270>.
- [8] Liu, L., & Zuo, Z. Excessive dietary salt promotes neuroinflammation to worsen retinopathy in mice with streptozotocin-induced diabetes. *Biochimica et Biophysica Acta* 2022;1868(9):166426, <https://doi.org/10.1016/j.bbadis.2022.166426>.
- [9] Rakova N, Kitada K, Lerchl K, Dahlmann A, Birukov A, Daub S, Kopp C, Pedchenko T, Zhang Y, Beck L, Johannes B, Marton A, Müller DN, Rauh M, Luft FC, Titze J. Increased salt consumption induces body water conservation and decreases fluid intake. *J Clin Invest.* 2017;127(5):1932-43, <https://doi.org/10.1172/JCI88530>.
- [10] Gonzalez SR, Ferrão FM, de Souza AM, Lowe J, Morcillo LL, Inappropriate activity of local renin-angiotensin-aldosterone system during high salt intake: impact on the cardio-renal axis. *Brazilian Journal of Nephrology* 2018;40(2):170-8, <https://doi.org/10.1590/2175-8239-JBN-3661>.
- [11] Wang YJ, Yeh TL, Shih MC, Tu YK, Chien KL. Dietary Sodium Intake and Risk of Cardiovascular Disease: A Systematic Review and Dose-Response Meta-Analysis. *Nutrients* 2020;12(10):2934, <https://doi.org/10.3390/nu12102934>.
- [12] Dornas W C, de Lima W G, dos Santos, R.C, Guerra J F, de Souza MO, Silva M, Souza E, Silva L, Diniz MF, Silva, ME. High dietary salt decreases antioxidant defenses in the liver of fructose-fed insulin-resistant rats. *The Journal of Nutritional Biochemistry* 2013;24(12):2016–22, <https://doi.org/10.1016/j.jnutbio.2013.06.006>.
- [13] Basdeki ED, Kollias A, Mitrou P, Tsirimiagkou C, Georgakis MK, Chatzigeorgiou A, Argyris A, Karatzi K, Manios Y, Sfrikakis PP, Protogerou AD. Does Sodium Intake Induce Systemic Inflammatory Response? A Systematic Review and Meta-Analysis of Randomized Studies in Humans. *Nutrients* 2021;13(8):2632, <https://doi.org/10.3390/nu13082632>.
- [14] Kumar R, Marrapu S. Dietary salt in liver cirrhosis: With a pinch of salt! *World J Hepatol.* 2023;15(10):1084-90. <https://doi.org/10.4254/wjh.v15.i10.1084>.

- [15] Takahashi F, et al. The Association of Salt Intake and Non-alcoholic Fatty Liver Disease in People With Type 2 Diabetes: A Cross-Sectional Study. *Front Nutr.* 2022;9:943790, <https://doi.org/10.3389/fnut.2022.943790>.
- [16] O'Callaghan CA. Dietary salt intake in chronic kidney disease: recent studies and their practical implications. *Polish archives of internal medicine* 2024;134(5):16715. <https://doi.org/10.20452/pamw.16715>.
- [17] Nishimoto M, Griffin KA, Wynne, BM, Fujita T. Salt-Sensitive Hypertension and the Kidney. *Hypertension* 2024;81(6):1206–17. <https://doi.org/10.1161/HYPERTENSIONAHA.123.21369>.
- [18] Kajiwara S, Hasegawa Y, Fujimori K, Tomiyasu S, Kamenon K, Uchikawa H, Morioka M. Persistent brain exposure to high sodium induces stroke onset by upregulation of cerebral microbleeds and oxidative stress in hypertensive rats. *Hypertension Research* 2024;47(1):78–87. <https://doi.org/10.1038/s41440-023-01447-z>.
- [19] Liu W, xing S, Wei F, Yao Y, Zhang H, Li Y, Liu Z. Excessive Dietary Salt Intake Exacerbates Cognitive impairment progression and increases dementia risk in older adults. *J Am Med Dir Assoc.* 2023;24(1):125-9.e4. <https://doi.org/10.1016/j.jamda.2022.10.001>.
- [20] Lee EY, Cho KH. High-dose consumption of NaCl resulted in severe degradation of lipoproteins associated with hyperlipidemia, hyperglycemia, and infertility via impairment of testicular spermatogenesis. *Toxicol Res.* 2016;5(2):557-69. <https://doi.org/10.1039/c5tx00059a>.
- [21] Abdelnour SA, Abd El-Hack ME, Noreldin AE, Batiha GE, Beshbishy AM, Ohran H, Khafaga AF, Othman SI, Allam AA, Swelum AA. High salt diet affects the reproductive health in animals: an overview. *Animals* 2020;10(4):590. <https://doi.org/10.3390/ani10040590>.
- [22] Gheitasi I, Motaghi N, Sadeghi H, Sadeghi H, Moslemi Z, Eftekhari M, Shakerinasab N, Doustimotlagh AH. Antioxidant and anti-inflammatory effects of *Origanum majorana* L. methanolic extract on bile duct ligation in male rats. *Evid Based Complement Alternat Med.* 2021;2021(1):9927196. <https://doi.org/10.1155/2021/9927196>.
- [23] Ben Attia T, Ben Ali R, Nahdi A, Galai S, Ghali R, Rammeh S, Véronique El May M, Mhamdi A. *Olea europaea* L. leaf extract mitigates oxidative and histological damage in rat heart tissue exposed to combined noise and toluene: An experimental study. *Saudi Pharm J.* 2023;31(8):101683. <https://doi.org/10.1016/j.jsps.2023.06.016>.
- [24] Attia FA, El-Sallami I, Ibrahim O. Variability of essential oil content, composition and antioxidants of sweet Marjoram grown at different locations in Egypt. *Assiut Journal of Agricultural Sciences* 2023;54:85-101. <https://doi.org/10.21608/ajas.2023.170726.1194>.
- [25] Markhali FS, Teixeira JA, Rocha CMR. Olive tree leaves—a source of valuable active compounds. *Processes* 2020;8(9):1177. <https://doi.org/10.3390/pr8091177>.
- [26] Samet I, Han J, Jlaiel L, Sayadi S., Isoda H. Olive (*Olea europaea*) leaf extract induces apoptosis and monocyte/macrophage differentiation in human chronic myelogenous leukemia k562 cells: Insight into the underlying mechanism. *Oxid Med Cell Longev.* 2014;(2014):927619, <https://doi.org/10.1155/2014/927619>.
- [27] Zhou L, Wang S, Qu J, Zhou H, Fatma AKKA, Kang W, Xi X, Liu Z. Anti-inflammatory phytoconstituents of *Origanum majorana*. *Journal of Food Quality* 2022;(2022):3888075, <https://doi.org/10.1155/2022/3888075>.
- [28] Al-Howiriny T, Alsheikh A, Alqasoumi S, Al-Yahya M, ElTahir K, Rafatullah S. Protective effect of *Origanum majorana* L. 'Marjoram' on various models of gastric mucosal injury in rats. *The American Journal of Chinese Medicine* 2009;37(3):531-45, <https://doi.org/10.1142/S0192415x0900703X>.
- [29] Kamran MA, Yildirimhan HS, Senlik B. Exploring the anthelmintic activity of *Olea europaea* L (Olive) leaves extract and oleuropein in mice naturally infected with *Aspiculuris tetraptera*. *Helminthologia* 2023;60(3):240-5, <https://doi.org/10.2478/helm-2023-0025>.
- [30] Dornas WC, de Lima WG, dos Santos RC, Guerra JFC, ed Souza MO, Silva M, e Silva LS, Diniz MF, Silva ME. High dietary salt decreases antioxidant defenses in the liver of fructose-fed insulin-resistant rats. *The Journal of Nutritional Biochemistry* 2013;24(2):2016-22, <https://doi.org/10.1016/j.jnutbio.2013.06.006>.
- [31] Edrissi A. Routine Histological Techniques. Sudan, University of Elimam Elmahdi press, 2022.

- [32] Henry RJ, Cannon DC, Winkelman JW. Clinical chemistry principle and technics. 2nd Edition, Hagerstown, USA, Harper and Row, Medical Dept., 1974.
- [33] Aebi H. Catalase in vitro. *Methods Enzymol.* 1984;105:121-6, [https://doi.org/10.1016/s0076-6897\(84\)05016-3](https://doi.org/10.1016/s0076-6897(84)05016-3).
- [34] Ohkawa H, Ohishi N, Yagi K. Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. *Anal Biochem.* 1979;95(2):351-8, [https://doi.org/10.1016/0003-2697\(79\)90738-3](https://doi.org/10.1016/0003-2697(79)90738-3).
- [35] Schumann G, Klauke R. New IFCC reference procedures for the determination of catalytic activity concentrations of five enzymes in serum: preliminary upper reference limits obtained in hospitalized subjects. *Clin Chim Acta* 2003;327(1-2):69-79, [https://doi.org/10.1016/s0009-8981\(02000341-8](https://doi.org/10.1016/s0009-8981(02000341-8).
- [36] Fossati P, Prencipe L, Berti G. Use of 3,5-dichloro-2-hydroxybenzenesulfonic acid/4-aminophenazone chromogenic system in direct enzymic assay of uric acid in serum and urine. *Clin Chem.* 1980;26(2): 227-31, <https://doi.org/10.1093/LINCHEM/26.2.227>.
- [37] Heinegard D, Tiderstrom G. Determination of serum creatinine by a direct colorimetric method. *Clinica Chemica Acta* 1973;43:305-10, [https://doi.org/10.1016/0009-8981\(73\)90466-x](https://doi.org/10.1016/0009-8981(73)90466-x).
- [38] Rakova N, Kitada K, Lerchl K, Dahlmann A, Birukov A, Daub S, Kopp C, Pedchenko T, Zhang Y, Beck L, Johannes B, Marton A, Muller DN, Rauh M, Luft FC, Titze J. Increased salt consumption induces body water conservation and decreases fluid intake. *J Clin Invest.* 2017;127(5):1932-43, <https://doi.org/10.1172/JI188530>.
- [39] Ma Y, He FJ, MacGregor GA. Is high salt intake independent risk factor for obesity? *Hypertension* 2015;66(4):843-9, <https://doi.org/10.1161/HYPERTENSIONAHA.115.05948.Epub>.
- [40] Stachenfeld NS. Acute effects of sodium ingestion on thirst and cardiovascular function. *Curr Sports Med. Rep.* 2010;7(4 Suppl):S7-13, <https://doi.org/10.1249/JSR.0b013e31817f23fc>.
- [41] Bolhuis DB, Costanzo A, Newman LP, Keast RSJ. Salt promotes passive overconsumption of dietary fat in humans. *The Journal of Nutrition* 2016;146(4):838-45, <https://doi.org/10.3945/jn.115.226365>.
- [42] Wang Y, Liu X, Zhang C, Wang Z. High salt diet induces metabolic alterations in multiple biological processes of Dahl salt sensitive rats. *The Journal of Nutritional Biochemistry* 2018;56:133-41, <https://doi.org/10.1016/j.jntbio.2018.01.007>.
- [43] Wu Q, Burley G, Li L, Lin S, Shi Y. The role of dietary salt in metabolism and cardiovascular disease. *Diabetes Obesity and Metabolism* 2023;25(5):1147-61, <https://doi.org/10.1111/som.14980>.
- [44] Mosavat M, Mirsanjari M, Arabiat D, Smyth A., Whitehead L. The role of sleep curtailment on leptin levels in obesity and diabetes mellitus. *Obes. Facts* 2021;14(2):214-21, <https://doi.org/10.1159/000514095>.
- [45] Choi J, Heo Y. The association between dietary sodium intake and adiposity, inflammation, and hormone markers: A preliminary study. *J Nutr Health* 2017;50(6):578-84, <https://doi.org/10.4163/jnh.2017.50.6.578>.
- [46] Wang C, Huang Z, Yu K, Ding R, Ye K, Dai C, Xu X, Zhou G, Li C. High-salt diet has a certain impact on protein digestion and gut microbiota: A sequencing and proteome combined study. *Front Microbiol.* 2017;8:1838, <https://doi.org/10.3389/fmicb.2017.01838>.
- [47] Oppelaar JJ, Rorije NMG, Engberink RHGO, Chahid Y, van Vlies N, Verberne HJ, van der Born BH, Vogt L. Perturbed body fluid distribution and osmoregulation in response to high salt intake in patients with hereditary multiple exostoses. *Molecular Genetics and Metabolism Reports* 2021;29:100797, <https://doi.org/10.1016/j.ymgmr.2021.100797>.
- [48] He FJ, Macgregor GA. Salt intake, plasma sodium, and worldwide salt reduction. *Annals of Medicine* 2012;44(sup1):S127-37, <http://doi.org/10.3109/07853890.2012.660495>.
- [49] Ozkayar N, Dede F, Ates I, Akyel F, Yildirim T, Altun B. The relationship between dietary salt intake and ambulatory blood pressure variability in non-diabetic hypertensive patients. *Nefrologia* 2026;36(6):694-700, <https://doi.org/10.1016/j.nefro.2015.12.004>
- [50] Edwards GE, Farquhar WB. Vascular effects of dietary salt. *Curr Opin Nephrol Hypertens.* 2015;24(1):8-13, <https://doi.org/10.1097/MNH.0000000000000089>.
- [51] Pojoga LH, Yao TM, Sinha S, Ross RL, Lin JC, Raffetto JD, Adler GK, Williams GH, Khalil R. Effect of dietary sodium on vasoconstriction and eNOS-mediated vascular relaxation in Caveolin-1 deficient mice. *Am J Physiol Heart Circ Physiol.* 2008;294(3):H1258-65, <https://doi.org/10.1152/ajpheart.01o14.2007>.

- [52] Oberleithner H, Riethmuller C, Schillers H, Hausberg M. Plasma sodium stiffens vascular endothelium and reduces nitric oxide release. *PNAS* 2007;104(41):16281-86, <https://doi.org/10.1073/pnas.0707791104>.
- [53] Zvanarou S, Vagnerova R, Mackievic V, Usnich S, Smolich I, Sokolik A, Yu M, Huang X, Angelis KJ, Demidchik V. Salt stress triggers generation of oxygen free radicals and DNA breaks in *Physcomitrella patens* protonema. *Environmental and Experimental Botany*. 2020; 180:104236, <https://doi.org/10.1016/j.envexpot.2020.104236>.
- [54] Shalan MG. Mitigating lead acetate-induced histopathologic and physiologic disorders in rats receiving vitamin C and glutathione supplement. 2025;11(1):e41256, <https://doi.org/10.1016/j.heliyon.2024.e41256>.
- [55] Barbaro NR, Beusecum JV, Xiao L, de Carmo L, Pitzer A, Loperena R, Foss JD, Elijovich F, Laffer C, Montaniel KR, Galindo CL, Chen W, Ao M, Mernaugh RL, Alsouqi A, Ikizler T A, Fogo AB, Moreno H, Zhao S, Davies SS, Harrison DG, Kirao A. Sodium activates human monocytes via the NADPH oxidase and isolevuglandin formation. *Cardiovasc Res*. 2020; 117(5):1358-71, <https://doi.org/10.1093/cvr/cvaa207>.
- [56] Kany S, Vollrath JT, Relja B. Cytokines in inflammatory disease. *Int J Mol Sci*. 2019;20(23):6008, <https://doi.org/10.3390/ijms20236008>.
- [57] Edwards DG, Farquhar WB. Vascular effects of dietary salt. *Curr Opin Nephrol Hypertens*. 2015;24(1):8-13, <https://doi.org/10.1097/MNH.000000000000089>.
- [58] Shin MH, Rhie G, Kim YK, Park C, Cho KH, Kim KH, Eun HC, Chung JH. H₂O₂ accumulation by catalase reduction changes MAP kinase signaling in aged human skin in vivo. *Journal of investigative dermatology* 2005;125(2):221-29, <https://doi.org/10.1111/j.0022-202x.2005.23823.x>.
- [59] Lenda DM, Boegehold MA. Effect of a high salt diet in microvascular antioxidant enzymes. *J Vasc Res*. 2002;39(1):41-50, <https://doi.org/10.1159/000048992>.
- [60] Adejare A, Oloyo A, Dahud Y, Adeshina M, Agbaje A, Ejim C, Ismail-Badmus K, Jaja M. Renal denervation ameliorated salt-induced hypertension by improving cardiac work, cardiac enzyme and oxidative balance in Sprague-Dawley rats. *Int J Cardiol Cardiovasc Risk Prev*. 2024; 21:200290, <https://doi.org/10.1016/j.ijcrp.2024.200290>.
- [61] Mills KT, Chen J, Yang W, Appel LJ, Kusek JW, Alper A, Delafontaine P, Keane MG, Mohler E, Ojo A, Rahman M, Ricardo AC, Soliman EZ, Steigerwalt S, Townsend R, He J. Sodium excretion and the risk of cardiovascular disease in patients with chronic kidney disease. *JAMA*. 2016;315(20):2200-10, <https://doi.org/10.1001/jama.2016.4447>.
- [62] Tian N, Thrasher KD, Gundy PD, Hughson MD, Jr DM. Antioxidant treatment prevents renal damage and dysfunction and reduces arterial pressure in salt-sensitive hypertension. *Hypertension* 2005;45(5):934-9, <https://doi.org/10.1161/01.HYP.0000160404.08866.5a>.
- [63] Cianciaruso B, Bellizzi V, Minutolo R, Tavera A, Capuano A, Conte G, De Nicola L. Salt intake and renal outcome in patients with progressive renal disease. *Miner Electrolyte Metab*. 1998; 24(4):296-301, <https://doi.org/10.1159/000057385>.
- [64] Sugiura T, Takase H, Ohte N, Dohi Y. Dietary salt intake is a significant determinant of impaired kidney function in the general population. *Kidney and Blood Pressure Research* 2018;43(4):1245-54, <https://doi.org/10.1159/000492406>.
- [65] Boero R, Pignataro A, Qurello F. Salt intake and kidney disease. *J Nephrol*. 2002;15(3):225-9.
- [66] Wang G, Yeung CK, Wong WY, Zhang N, Wei YF, Zhang JL, Yan Y, Wong CY, Tang JJ, Chuai M, Lee KK, Wang LJ, Yang X. Liver fibrosis can be induced by high salt intake through excess reactive oxygen species (ROS) production. *J Agric Food Chem*. 2016; 64(7):1610-7, <https://doi.org/10.1159/000492406>.
- [67] Uetake Y, Ikeda H, Irie R, Tejima K, Matsui H, Ogura S, Wang H, Mu S, Hirohama D, Ando K, Sawamura T, Yatomi Y, Fujita T, Shimosawa T. High-salt in addition to high-fat diet may enhance inflammation and fibrosis in liver steatosis induced by oxidative stress and dyslipidemia in mice. *Lipids Health Dis*. 2015; 14:6, <https://doi.org/10.1186/s12944-015-0002-9>.
- [68] Zhang X, Liang Y, Jiang J, Lu C, Shi F, Cao Q, Zhang Y, Diao H. A high-salt diet exacerbates liver fibrosis through Enterococcus-dependent macrophage activation. *Microbiol Spectr*. 2023;11(2):e0340322, <https://doi.org/10.1128/spectrum.03403-22>.

- [69] Min B, Fairchild RL. Over-salting ruins the balance of the immune menu. *J Clin Invest*. 2015;125(11):4002-4, <https://doi.org/10.1172/JCI84690>.
- [70] Hernandez AL, Kitz A, Wu C, Lowther DE, Rodriguez DM, Vudattu N, Deng S, Herold KC, Kuchroo VK, Kleinewietfeld M, Hafler DA. Sodium chloride inhibits the suppressive function of FOXP3+ regulatory T cells. *J Clin Invest*. 2015;125(11):4212-22, <https://doi.org/10.1172/JCI81151>.
- [71] Huh JH, Lee KJ, Lim JS, Lee MY, Park HJ, Kim MY, Kim JW, Chung CH, Shin JY, Kim H, Kwon SO, Baik SK. High dietary sodium intake assessed by estimated 24-h urinary sodium excretion is associated with NAFLD and hepatic fibrosis. *PLoS One*. 2015;10(11):10(11):e0143222, <https://doi.org/10.1371/journal.pone.0143222>.
- [72] He X, Liu Y, Usa K, Tian Z, Jr AWC, Liang M. Ultrastructure of mitochondria and the endoplasmic reticulum in renal tubules of Dahl salt-sensitive rats. *Am J Physiol Renal Physiol*. 2014;306:F1190-7, <https://doi.org/10.1152/ajprenal.00073.2014>.
- [73] Noto JG, Bollu R, Sturzoiu T, Nanda S. Hyponatremia A systems-based approach. *International Journal of Academic Medicine* 2018;4(3):266-70, https://doi.org/10.4103/IJAM.IJAM_8_18.
- [74] Binger KJ, Gebhardt M, Heinig M, Rintisch C, Schroeder A, Neuhofer W, Hilgers K, Manzel A, Schwartz C, Kleinewietfeld M, Voelkl J, Schatz V, Linker RA, Lang F, Voehringer D, Wright MD, Huner N, Dechend R, Jantsch J, Titze J, Muller DN. High salt reduces the activation of IL-4- and IL-13-stimulated macrophages. *J Clin Invest*. 2015; 125(11):4223-38, <https://doi.org/10.1172/JCI80919>.
- [75] Deng Z, Wang Y, Zhou L, Shan Y, Tan S, Cai W, Liao S, Peng L, Lu Z. High salt-induced activation and expression of inflammatory cytokines in cultured astrocytes. *Cell Cycle* 2017;16(8):785-94, <https://doi.org/10.1080/15384101.2017.1301330>.
- [76] Liu YZ, Chen JK, Li ZP, Zhao T, Ni M, Li DJ, Jiang CL, Shen FM. High-salt diet enhances hippocampal oxidative stress and cognitive impairment in mice. *Neurobiol Learn Mem*. 2014;114:10-5, <https://doi.org/10.1016/j.nlm.2014.04.010>.
- [77] Ge Q, Wang Z, Wu Y, Huo Q, Qian Z, Tian Z, Ren W, Zhang Z, Han J. High salt diet impairs memory-related synaptic plasticity via increased oxidative stress and suppressed synaptic protein expression. *Mol Nutr Food Res*. 2017;61(10):1700134, <https://doi.org/10.1002/mnfr.201700134>.
- [78] Ying B, Yang G, Guo Z, Cai W, Chen S, Zhou X, Li Y. Chronic high-salt intake induces cardiomyocyte autophagic vacuolization during left ventricular maladaptive remodeling in spontaneously hypertensive rats. *Experimental and Therapeuti Medicine* 2023; 25(4):148-61, <https://doi.org/10.3892/etm.2023.11847>.
- [79] Zook F, Pan YE, Wipplinger A, Kerschbaum HH, Clements RJ, Ritter M, Stauber T, Model MA. Delayed vacuolation in mammalian cells caused by hypotonicity and ion loss. *Scientific Reports* 2024;14:29354, <https://doi.org/10.1038/s41598-79815-z>.
- [80] Rossitto G, Mary S, Chen JY, Boder P, Chew KS, Neves KB, Alves RL, Montezano AC, Welsh P, Petrie MC, Graham D, Touyz RM, Delles C. Tissue sodium excess is not hypertonic and reflects extracellular volume expansion. *Nat Commun*. 2020;11(1):4222, <https://doi.org/10.1038/s41467-020-17820-2>.
- [81] Sulyok E, Farkas B, Nagy B, Varnagy A, Kovacs K, Bodis J. Tissue sodium accumulation: pathophysiology and clinical implications. *Antioxidants* 2022;11(4):750, <https://doi.org/10.3390/antiox11040750>.
- [82] Yuan M, Wang Y, Wen J, Jing F, Zou Q, Pu Y, Pan T, Cai Z. Dietary salt disrupts tricarboxylic acid cycle and induces Tau hyperphosphorylation and synapse dysfunction during aging. *Aging Dis*. 13(5):1532-45, <https://doi.org/10.14336/AD.2022.0220>.
- [83] Fyfe I. High-salt diet promotes Alzheimer's disease-like changes. *Nature Reviews Neurology* 2020; 16:2-3, <https://doi.org/10.1038/s41582-019-0289-7>.
- [84] Na S, Janakiraman M, Leliavski A, Krishnamoorthy G. High-salt diet suppresses autoimmune demyelination by regulating the blood-brain barrier permeability. *PNAS* 2021;118(12):e2025944118, <https://doi.org/10.1073/pnas.2025944118>.
- [85] Siskova Z, Justus D, Kaneko H, Friedrichs D, Henneberg N, Beutel T, Pitsch J, Schoch S, Becker A, von der Kammer H, Remy S. Dendritic structural degeneration is functionally linked to cellular hyperexcitability in a

- mouse model of Alzheimer's disease. *Neuron* 2014;84(5): 1023-33, <https://doi.org/10.1016/j.neuron.2014.10.024>.
- [86] Vernooij MW, Ikram A, Vrooman HA, Wielopolski PA, Krestin GP, Hofman A, Niessen WJ, Van der Lugt A, Breteler MMB. White matter microstructural integrity and cognitive function in a general elderly population. *Arch Gen Psychiatry* 2009;66(5):545-53, <https://doi.org/10.1001/archgenpsychiatry.2009.5>
- [87] Mohamed RS, Mohamed RH, Wehrend A, Tahoun EA, Hussein HA; The effect of saline water consumption on sperm parameters, testicular histopathology, hormonal and antioxidants concentrations in Barki Rams. *BMC Veterinary Research* 2024;20:219, <https://doi.org/10.1186/s12917-024-04047-2>.
- [88] Lee E, Cho K. High-dose consumption of NaCl resulted in severe degradation of lipoproteins associated with hyperlipidemia, hyperglycemia, and infertility via impairment of testicular spermatogenesis. *Toxicol. Res.* 2016;5:557-69, <https://doi.org/10.1039/C5TX00059A>
- [89] Trafimovich Y. The impact of salt on testosterone levels: Unveiling the culprit. *MedShun* 2024 Jan 22. Available from: <https://medshun.com/article/does-salt-affect-testosteron>
- [90] Asadi N, Bahmani M, Kheradmand A, Rafeian-Kopaei M. The impact of oxidative stress on testicular function and the role of antioxidants in improving it: A review. *J Clin Diagn Res.* 2017;11(5):IE01-5, <https://doi.org/10.7860/JCDR/2017/23927.9886>
- [91] Dutta S, Sengupta P, Sama P, Roychoudhury S. Oxidative stress, testicular inflammatory pathways, and male reproduction. *Int J Mol Sci.* 2021;22(18):10043, <https://doi.org/10.3390/ijms221810043>.
- [92] Iranloye B, Oludare GO, Morakinyo AO, Esume NA, Ekeh LC. Reproductive parameters and oxidative stress status of male rats fed with low and high salt diet. *J Hum Reprod Sci.* 2013;6(4):267-72, <https://doi.org/10.4103/0974-1208.126308>.
- [93] Yumura Y, Saito K, Suzuki K, Ogawa T, Kanno H, Sato K, Iwasaki A, Hosaka M. The effect of high concentration of sodium and potassium ions on motility of human sperm preserved in electrolyte-free solution. *Nihon Hinyokika Gakkai Zasshi.* 2002;93(3):440-3, <https://doi.org/10.5980/jpnjurol1989.93.440>.
- [94] Iranloye B, Oludare GO, Esume NA, Ekeh LC. Reproductive parameters and oxidative status of male Sprague Dawley rats treated with low and high salt diet. *Metabolism and Endocrinology* 2013;Proc 37th IUPS, PCB350, Available from: <https://www.physco.org/abstracts/reproductive-parameters-and-oxidative-status-of-male-sprague-dawley-rats-treated-with-low-and-high-salt-diet/>
- [95] Adekunbi DA, Ogunsola AO, Oyelowo OT, Aluko EO, Popoola AA, Akinboboye OO. Consumption of high sucrose and/or high salt diet alters sperm function in male Sprague-Dawley rats. *Neuroendocrinology* 2014; Proc Physiol Soc 32, PC065, Available form: <https://www.physoc.org/abstrats/consumption-of-high-surose-and-or-high-salt-diet-alter-sperm-function-in-male-sprague-dawley-rats/>
- [96] Revelou PK, Konteles SJ, Batrinou A, Xagoraris M, Tarantilis PA, Strati IF. Origanum majorana L. as a flavoring agent: Impact on quality indices, stability, and volatile and phenolic profiles of extra virgin olive oil (EVOO). *Foods* 2024;13(19):3164, <https://doi.org/10.3390/foods13193164>.
- [97] Baatour O, Nasri-ayachi MB, Mahmoudi H, Tarchoun I, Nasri N, Zaghoudi M, Abidi W, Kaddour R, M'rah S, Hamdawi G. Salt effect on physiological, biochemical and anatomical structures of two origanum majorana varieties (Tunisian and Canadian). *African Journal of Biotechnology* 2012;11(27):7109-18, <https://doi.org/10.5897/AJB11.3493>.
- [98] Nediani C, Ruzzolini J, Romani A, Calorini L. oleuropein, a bioactive compound from *Olea europaea* L., as a potential preventive and therapeutic agent in non-communicable diseases. *Antioxidants* 2019;8(12):578, <https://doi.org/10.3390/antiox8120578>.
- [99] Wang S, Zhou L, Attia FAKK, Tang Q, Wang M, Liu Z, Waterhouse GIN, Liu L, Kang W. Origanum majorana L: A nutritional supplement with immunomodulatory effects. *Front. Nutr.* 2021;8:748031, <https://doi.org/10.3389/fnut.2021.748031>.
- [100] De Ciccio P, Maisto M, Tenore GC, Ianaro A. Olive leaf extract, from *Olea europaea* L., reduces palmitate-induced inflammation via regulation of murine macrophages polarization. *Nutrients* 2020;12(12):3663, <https://doi.org/10.3390/nu12123663>.

- [101] Mehanna S, Issa MY, Hassan NH, Hussien AM, Ibrahim MA, Hassanen EI. Origanum majorana essential oil improves the rat's sexual behavior and testicular oxidative damage induced by imidacloprid via modulating the steroidogenesis pathways. Saudi Pharm J. 2022;30(9):1315-26, <https://doi.org/10.1016/j.jsps.2022.06.016>.
- [102] Baatour O, Mahmoudi H, Tarchoun I, Nasri N, Traelsi N, Naddour R, Zaghoudi M, Hamdawi G, Ksouri R, Lahaal M, Marzouk B. Salt effect on phenolics and antioxidant activities of Tunisian and Canadian sweet marjoram (*Origanum majorana* L.) shoots. J Sci Food Agric. 2013;93(1):134-41, <https://doi.org/10.1002/jsfa.5740>.
- [103] Buciantini M, Leri M, Nardiello P, Casamenti F, Stefani M. Olive polyphenols: Antioxidant and anti-inflammatory properties. Antioxidants 2021;10(7):1044, <https://doi.org/10.3390/antiox10071044>.
- [104] Gheitasi I, Azizi A, Omidifar N, Coustimotlagh AH. Renoprotective effects of *Origanum majorana* methanolic L and carvacrol on ischemia/reperfusion-induced kidney injury in male rats. Evidence-Based Complementary and Alternative Medicine 2020;2020:9785932, <https://doi.org/10.1155/2020/9785932>.
- [105] AlHaithloul HAS, Alotaibi MF, Bin-Jumah M, Elgebaly H, Mahmoud AM. *Olea europaea* leaf extract up-regulates Nrf2/ARE/HO-1 signaling and attenuates cyclophosphamide-induced oxidative stress, inflammation and apoptosis in rat kidney. Biomedicine and Pharmacotherapy 2019; 111:676-85, <https://doi.org/10.1016/j.biopha.2018.12.112>.
- [106] Abdelsalam HM, Diab AA, Zahra MH, Eldehawy S, Hendawy BA. Therapeutic effects of *Origanum majorana* and Propolis against nephrotoxicity induced by adenine. Biol Eng Med. 2019;4:1-9, <https://doi.org/10.15761/BEM.1000176>.
- [107] Soussi R, Hfaiedh N, Sakly M, Ben Rhouma K. The aqueous extract of *Olea europaea* leaves protects from haemotoxicity and kidney damage induced by diclofenac in Swiss albino mice. RSC Adv. 2019;9:23352-61, <https://doi.org/10.1039/C9RA01670H>.
- [108] Ivanov M, Vajic U, Mihailovic-Stanojevic N, Miloradovic Z, Jovovic D, Grujic-Milanovic J, Karanovic D, Dekanski D. Highly potent antioxidant *Olea europaea* L. leaf extract affects carotid and renal hemodynamics in experimental hypertension: The role of oleuropein. EXCLI J. 2018;17:29-44, <https://doi.org/10.17179/excli2017-1002>.
- [109] Salla NS, Haggag MH, Rabeh NM, Ahmed OM. Effect of aqueous extract of SAGE and Marjoram on chronic kidney patients. Egypt J Appl Sci. 2020;35(11):186-201, <https://doi.org/10.21608/EJAS.2020.136650>.
- [110] Ibrahim D, Halboup A, Al Ashwal M, Shamsher A. Ameliorative effect of *Olea europaea* leaf extract on Cisplatin-induced nephrotoxicity in the rat model. Int J Nephrol. 2023;2023: 2074498, <https://doi.org/10.1155/2023/2074498>.
- [111] Rateb MO, Adelimagid AD, Ebd Elhamid OM, Said AM. Evaluation of the antidiabetic and hepatoprotective effect of *Origanum majorana* leaf extract and its nanoparticles in streptozotocin-induced diabetes in rats. Benha Veterinary Medical Journal 2024;46(2):23-7, <https://doi.org/10.21608/BVMJ.2024.288584.1825>.
- [112] Fki I, Sayadi S, Mohmoud A, Daoued I, Marrekchi R, Ghorbel H. Comparative study on beneficial effects of hydroxytyrosol- and Oleuropein-rich olive leaf extracts on high-fat diet-induced lipid metabolism disturbance and liver injury in rats. BioMed Research International 2020; 2020(1):1315202, <https://doi.org/10.1155/2020/1315202>.
- [113] Mossa ATH, Nawwar GAM. Free radical scavenging and antiacetylcholinesterase activities of *Origanum majorana* L essential oil. Human and Experimental Toxicology 2011;30(10):1501-13, <https://doi.org/10.1177/0960327110391686>.
- [114] Benavente-Garcia O, Castillo J, Lorente J, Ortuno A, Del Rio JA. Antioxidant activity of phenolics extracted from *Olea europaea* L. leaves. Food Chemistry 2000;68: 457-62, [https://doi.org/10.1016/s0308-8146\(99\)00221-6](https://doi.org/10.1016/s0308-8146(99)00221-6).
- [115] Wahby MM, Yacout G, Kandeel K, Awad D. LPS-induced oxidative inflammation and hyperlipidemia in male rats: The protective role of *Origanum majorana* extract I. Beni-Suef University Journal of Basic and Applied Sciences 2015;4(4): 291-98, <https://doi.org/10.1016/j.bjbas.2015.11.004>.

- [116] Rahmani AH, Almatroudi A, Allemailem KS, Alharbi HOA, Babiker AY, Althwab SA, Alsuhaymi N, Alsugoor MH, Khan AA, Al-Megrin WAI. Oleuropein, a phenolic component of *Olea europaea* L. ameliorates CCl₄-induced liver injury in rats through the regulation of oxidative stress and inflammation. *Eur Rev Med Pharmaol Sci.* 2024;28(4):1259-71, https://doi.org/10.26355/eurrev_202402_35447.
- [117] Wagdy R, Abdel-Kader RM, El-Khatib AH, Linscheid MW, Handoussa H, Hamdi N. *Origanum majorana* L. protects against neuroinflammation-mediated cognitive impairment: a phyto-pharmacological study. *BMC Complementary Medicine and Therapies* 2023;23:165, <https://doi.org/10.1186/s12906-023-03994-x>.
- [118] Aykac A, Terali K, Ozbeyli D, Ede S, Albayrak O, Baser KHC, Sener G. A multi-parameter evaluation of the neuroprotective and cognitive-enhancing effects of *Origanum onites* L. (Turkish oregano) essential oil on scopolamine-induced amnesic rats. *Metab Brain Dis.* 2022;37:1041-55, <https://doi.org/10.1007/s11011-022-00933-6>.
- [119] Sabogal-Guaqueta AM, Munoz-Manco JI, Ramirez-Pineda JR, Lamprea-Rodriguez M, Osorio E, Cardona-Gomez GP. The flavonoid quercetin ameliorates Alzheimer's disease pathology and protects cognitive and emotional function in aged triple transgenic Alzheimer's disease model mice. *Neuropharmacology* 2015;93:134-45, <https://doi.org/10.1016/j.neuropharm.2015.01.027>.
- [120] Mohagheghi F, Bigdeli MR, Rasouljan B, Hashemi P, Pour MR. The neuroprotective effect of olive leaf extract is related to improved blood-brain barrier permeability and brain edema in rat with serial focal cerebral ischemia. *Phytomedicine* 2011;18(2-3):170-5, <https://doi.org/10.1016/j.phymed.2010.06.007>.
- [121] Sissaoui Y, Boukhari Y, Mahi MA. The anti-inflammatory effect of polyphenolics olive leaf extracts *Olea europaea* L. *Phytotherapie* 2022; 20(1):17-22, <https://doi.org/10.3166/phyto-2021-0251>.
- [122] ElSawy NA, Alkushi AG, AlAsmari WAM, Sinna MM, Meader EA, ElMadouly MA, Sakran AMEA. Does oregano protect against testicular toxicity produced by ethylene glycol in adult male albino rat? *Int. J. Morphol.* 2019;37(1):358-62.
- [123] Hakemi SG, Sharififar F, Haghpanah T, Babae A, Hassan S, Eftekhari-Vaghefi. The effects of olive leaf extract on the testis, sperm quality and testicular germ cell apoptosis in male rats exposed to usulfan. *Int J Fertil Steril.* 2019;13(1):57-65, <https://doi.org/10.22074/ijfs.2019.5520>.
- [124] Sarishegi M, Gorgich EAC, Khajavi O. Olive leaves extract improved sperm quality and antioxidant status in the testis of rat exposed to rotenone. *Nephro-Urology Monthly* 2017;9(3):e47127, <https://doi.org/10.5812/numonthly.47127>.