

Molecular and physiological insights of salt tolerance in hulled barley (*Hordeum vulgare* L. var. *nudum*)

Noura S. Soud^a, Ashraf H. Fahmy^b, Ahmed M.A. El-Sharkawy^a, Mona H. Hussein^a

^aGenetic Engineering Research Center and Department of Genetics, Faculty of Agriculture, Cairo University, Giza, Egypt, ^bAgricultural Genetic Engineering Research Institute (AGERI), Agriculture Research Center, Ministry of Agriculture

Correspondence to Noura S. Soud, PhD, Genetic Engineering Research Center and Department of Genetics, Faculty of Agriculture, Cairo University, Giza 12613, Egypt. Tel: +20 101 809 1399; e-mail: Noura.soud@post.agr.cu.edu.eg

Received: 14 October 2023

Revised: 27 November 2023

Accepted: 27 November 2023

Published: 28 June 2024

Egyptian Pharmaceutical Journal 2024, 0:0–0

Background

The nutritional value of hulled barley makes it a promising resource for creating new healthy foods globally. However, improving the salt tolerance of certain barley cultivars remains a challenge, despite their inherent salt tolerance.

Objective

This study aims to investigate the physiological and molecular mechanisms associated with salt tolerance in barley, focusing on the expression of genes involved in regulating cellular ion homeostasis, detoxification, and water transport.

Materials and methods

Three barley cultivars were subjected to different levels of NaCl concentrations. Data on several growth parameters and gene expression were measured and recorded.

Results and conclusion

Increasing salinity affected shoot and root length, fresh and dry weight, depending on genotype. Giza-130 showed higher dry weight, followed by Giza-135, while Giza-136 showed the lowest value. Giza-130 exhibits the ability to regulate intracellular ion concentration through a higher expression level of the NHX1-gene, demonstrating its ability to effectively absorb water under salinity stress, due to its high expression level of the *hvpip* –aquaporin gene and effectively remove reactive oxygen species and reduces oxidative stress through the accumulation of higher concentrations of catalase, ascorbate peroxidase, glutathione S-transferase, and superoxide dismutase. In contrast, Giza-136 showed down-regulated gene expression and higher sensitivity to salt stress. Giza-130 was salt tolerant, followed by Giza-135 while Giza-136 was very sensitive. The genotype-specific regulation of gene expression not only highlights the important role of these genes in protecting plants against salt-induced oxidative stress but also improves our understanding of the salt stress tolerance of barley and plays an important role in improving salt tolerance in other crops.

Keywords:

barley, enzymes, expression, gene, oxidative, salt

Egypt Pharmaceut J 0:0–0

© 2024 Egyptian Pharmaceutical Journal

1687-4315

Criteria for inclusion in the authors'/contributors' list: MHS and AHF planned the experiments. NSS carried out field and laboratory sample preparations and experiments. All authors contributed to the interpretation of the results. NSS took the lead in performing the research, analyzing data, and writing the manuscript. All authors provided critical feedback and helped shape the research, analysis, and manuscript.

Introduction

Barley is a widely cultivated cereal crop in temperate regions and is the fourth most produced crop in the world after wheat, rice, and maize [1]. It accounts for ~8% of the world's agricultural land. Hulled barley (*Hordeum vulgare* L. var. *nudum*) is a variety of barley that is cultivated in East Asia, mainly on the Qinghai-Tibet Plateau at an average altitude of 4,500 m [2]. Despite its potential, barley is not widely used for

human consumption and is usually grown for cultural or regional reasons where wheat is less profitable [3]. However, it is now gaining importance as a health food in Europe, North America, and other nontraditional barley-growing regions [4]. Barley grains are rich in minerals, protein, and lysine, and have high β -glucan content, which helps inhibit cholesterol synthesis [5]. Hulled barley also has advantages for the food industry as it eliminates the need for processing steps. Therefore, hulled barley is considered a valuable resource for breeding new healthy foods around the world [6]. Salt stress, a significant abiotic stress, poses a widespread challenge to plant growth and

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

productivity. This is a result of salt accumulation in the soil, leading to reduced water and nutrient availability for plants [7,8]. Drought, salinity, heat, floods, air pollution, and climate change are some of the main causes of reduced crop growth and productivity [9]. Recent studies indicate that soil salinity is expected to increase further due to the effects of climate change [10]. Therefore, improving crop productivity through the development of salt-tolerant crop varieties becomes a strategy to mitigate salt stress. Salt stress causes an increase in the production of reactive oxygen species (ROS), which can damage cellular components [11,12].

Plants have evolved various adaptive mechanisms to cope with salt stress, including changes in gene expression. Previous research has identified several genes involved in plant responses to salt stress, including those involved in ROS scavenging, ion transport, and osmotic regulation [13]. Understanding the regulation and role of these genes in salt stress tolerance can provide insights into developing salt-tolerant crops and improving agricultural productivity in saline environments [14]. Many studies have focused on identifying and characterizing genes involved in plant responses to salt stress, which play important roles in maintaining cellular ion homeostasis, detoxifying reactive oxygen species, and regulating carbon metabolism under salt stress conditions, such as *hvip* [15], *nhx1* [16–18], *GSTTPS*, *sod*, *apx* [19,20].

This study aimed to investigate the impact of salt stress on plant biomass production in three cultivars of hulled barley and to explore the molecular mechanisms that underlie barley's response to stress. Specifically, we examined the expression of *hvip*-aquaporin, *nhx1* and genes encoding antioxidant enzymes (*cat*, *apx*, *sod* and *gst*) under salt stress conditions. A comprehensive understanding is necessary for devising plans to enhance plant salt tolerance and improve food security in areas with high salt concentrations.

Materials and methods

Plant materials and saline treatments

In this experiment, three cultivars of Egyptian hulled barley were grown (Giza-130, Giza-135, and Giza-136). Cultivation was carried out in 1 L plastic pots filled with a soil mixture consisting of equal parts of sandy soil and peat moss (ratio 1 : 1 v/v). To ensure optimal growth conditions, seedlings were irrigated with 400 ml of one-tenth concentration MS solution

[21] every day and placed in a greenhouse with controlled environmental conditions. To ensure adequate soil moisture, soil water tension was maintained below 60 kPa throughout the experiment. After 30 days of growth, plants were exposed to salt stress. This was achieved by gradually introducing NaCl (sodium chloride) into the nutrient solution at different concentrations: .0 mM (control), 50, 100, and 200 mM for 15 days. Throughout the experiment, the temperature in the greenhouse was kept constant at 35°C, and the photo-synthetically active radiation (PAR) was set at 2743 $\mu\text{mol m}^{-2} \text{s}^{-1}$. To ensure statistical validity and reliable results, each NaCl treatment and control group without NaCl treatment were repeated five times.

Measurement of growth

To evaluate the effects of salt stress on plant growth, shoot, and root length as well as plant fresh weight of salt treated and control plants were measured. Then, the sample was dried in a forced drying oven (Heraeus-0871, USA) at 80°C for three days. After the drying process, the dry weight of the sample is determined.

RNA extraction and real-time-PCR (qPCR) analysis

RNA extraction and quantitative polymerase chain reaction (qPCR) analysis were accomplished through the following methods. Frozen leaf tissue weighing 100 mg was used for total RNA isolation. The RNeasy Plant Mini kit (Qiagen, Germany, CAT NO.74903 and 74904) was used according to the manufacturer's instructions. Total RNA was then reverse transcribed into cDNA using the Revert Aid First Strand cDNA Synthesis Kit (Thermo Scientific, USA, Cat NO. K1621). The expression level of messenger RNA (mRNA) related to salt tolerance in barley shoot tissues was measured using real-time PCR (qPCR). Maxima SYBR Green/ROX qPCR Master Mix kit (Thermo Scientific, USA, Cat NO. K0221) was used for this purpose. Primer nucleotides were designed using Primer 3 software (Table 1) and the Actin gene was used as a housekeeping gene. The PCR reaction included an initial denaturation at 95°C for 10 min, followed by 40 cycles of denaturation at 95°C for 15 s, annealing at 60°C for 60 s, extension at 72°C for 60 s, and a final extension at 72°C for 4 min. qPCR reactions were performed in triplicate for all genes and a Stratagene MX3000P qPCR machine was used.

Data analysis

Data analysis involved measuring the comparative Ct value ($\Delta\Delta\text{Ct}$), obtained by subtracting the ΔCt value of the calibrator from the ΔCt value of the treated samples. Additionally, relative expression fold

Table 1 Real-time polymerase chain reaction primers used

Gene name	Sequences	
	Forward (5→3')	sReverse (5→3')
Actin	AGACCTTCAACACCCCTGCTATGT	CCAATCCAGACACTGTACTTCCTT
NHX1	TACGGTTTTCTGCCTCTGTCACA	ACAA CATCTGGTCATACTGCCG
HvPIP2:1	GCTAGCTTAGCAATGGCCAAGGAC	GTCGGACTGGTGCTTGTACC
Gst	AGCATCTCGTCAGAAACCCGT	TCCTTCAGGTTGCCCTCTCTT
Sod	CCGAAGATGAAATCCGCCAT	CGGCAATGATTGAATGTGG
Cat	CGACGACAAGATGCTGCAGT	TGGTTGTTCTTGAAGCCGC
Apx	CGGAGCTTTTGTAGTGGTGACA	CCGCAGCATATTTCTCCACAA

changes were calculated using the $2^{-\Delta\Delta CT}$ formula of Livak and Schmittgen [22]. To determine the significance (P value < 0.05) between mean genotype differences, an independent, unpaired Student's t test was applied.

Results impact of salt stress on plant growth the effects of salt stress on shoot and root length, plant fresh and dry weight were investigated in three barley cultivars, namely: Giza-130, Giza-135, and Giza-136 under salt stress conditions. Shoot length is an important indicator of the plant's ability to cope with salt stress. In the case of Giza-135 and Giza-136, the leaves turned yellow and eventually died 2 weeks after exposure to salt stress, unlike the control plants which remained green (Fig. 1). The results showed that shoot length decreased significantly in all three cultivars when increasing NaCl concentration. Giza-130 is more tolerant to salinity stress while Giza-136 exhibited the highest sensitivity to salt stress, with the greatest reduction in shoot length at the highest NaCl concentration of 200 mM (Table 2 and Fig. 1). Salt stress also has a significant impact on plant fresh weight. The results showed that plant fresh weight decreased significantly as NaCl concentration increased for the three cultivars. The Giza-136 showed the highest sensitivity to salt stress, with the greatest reduction in fresh weight at the highest NaCl concentration of 200 mM. Similarly, salt stress affected plant dry weight in all cultivars, with variations depending on genetic background (Table 2). The cultivar Giza-130 showed a less pronounced decrease in dry weight, followed by Giza-135, while the cultivar Giza-136 showed the most significant decrease in dry weight under salt stress conditions.

Expression of genes encoding antioxidant enzymes (SOD, GST, CAT and APX)

The changes in *sod* gene expression under salt stress in hulled barley cultivars was examined Fig. 2a. At 50 mM NaCl, Giza-135 accumulated higher levels of *sod* transcripts compared with other cultivars. As NaCl

Figure 1

The effect of salinity stress on plant growth of three hulled barley cultivars. A: Giza-130, B: Giza-135 and C: Giza-136.

concentration increased up to 100 mM, expression levels increased, with Giza-130 showing a 9-fold increase compared with a 7-fold increase in Giza-135. However, in Giza-136 it only reached a 2-fold increase at the same concentration. At 200 mM NaCl, *sod* expression levels were significantly decreased in all cultivars, with a significant decrease observed in Giza-136.

The expression of the *gst* gene was affected by salt stress, and an increase in expression was observed when the NaCl concentration increased to 100 mM. Giza-

Table 2 The effect of salt stress on biomass production in three barley cultivars

Cultivars	NaCl Treatment (mM)	Shoot length (cm)	Root length (cm)	Whole plant (cm)	FW (g)	DW (g)
Giza-130	0	25±1.1	14±1.2	39	5.72±0.2	5.12 (100%)
	50	26±2.0	18±1.5	44	6.21±0.1	5.49 (107%)
	100	27±1.4	17±1.3	44	5.46±0.2	4.80 (93%)
	200	25±1.2	17±1.4	42	4.41±0.3	3.81 (74%)
Giza-135	0	28±1.4	14±1.5	42	3.79±0.1	3.37 (100%)
	50	27±1.4	17±1.7	44	4.58±0.2	4.07 (120%)
	100	21±1.8	15±1.3	36	3.22±0.2	2.87 (85%)
	200	15±3.0	16±1.3	31	2.87±0.2	2.73 (69%)
Giza-136	0	25±2.1	15±1.1	40	5.78±0.3	4.97 (100%)
	50	27±2.0	17±1.2	44	5.42±0.1	4.72 (94%)
	100	22±1.1	15±1.4	37	4.60±0.1	4.04 (81%)
	200	14±1.1	14±1.3	28	2.60±0.1	2.13 (42%)


Data are mean of five replicates±SE.

130 showed a 10-fold increase in expression compared with 3-fold increase in Giza-135, while Giza-136 down-regulated *gst* expression at the same concentration. At 200 mM NaCl, Giza-130 maintained a higher expression level compared with Giza-135, and Giza-136 showed a negative expression level under salt stress (Fig. 2b).

In this study, *cat* gene expression was investigated in different cultivars, and the results indicated that Giza-

130 exhibited significantly higher levels of expression at 100 mM NaCl compared with other cultivars tested at the same salt concentration. However, expression levels decreased significantly with an increase in NaCl concentration in all cultivars, as shown in Fig. 2c.

The effect of salt stress on *apx* gene expression in different hulled barley cultivars was investigated, and the results presented in Fig. 2d. Increasing the salt concentration from 50 to 100 mM increased *apx* gene

Figure 2

The expression of genes encoding antioxidant enzymes in hulled barley under salinity stress conditions: A-*sod*, B-*gst*, C-*cat* and D-*apx*.

expression. It was observed that, Giza-130 showed the highest expression level (6-fold increase) compared with other cultivars at the same salt concentration. However, increasing the salt concentration to 200 mM led to a significant decrease in *apx* gene expression. Giza-130 showed higher expression levels (3-fold increase), followed by Giza-135 (2-fold increase), while the expression was completely inhibited in Giza-136 at 200 mM NaCl.

Expression of genes associated with water transport and ion homeostasis in barley under salt stress

In this study, the impact of salt stress on the *HvPIP* gene, which plays a crucial role in water transport activity and regulation of barley PIP aquaporins, was investigated in three different hulled barley cultivars. The results represented in Fig. 3 reveal that the folding of the *HvPIP* transcript is enhanced under salt stress conditions. At 50 mM NaCl, Giza-130 exhibited a 5-fold increase in expression, while Giza-135 and Giza-136 showed 4 and ~3-fold increases, respectively. Furthermore, increasing NaCl concentration from 50 mM to 100 mM resulted in increased expression of *HvPIP* gene, with Giza-130 exhibiting the highest increase, followed by Giza-135, and Giza-136 showing the lowest expression. However, at high salinity levels (200 mM), *HvPIP* expression levels decreased in all cultivars, with Giza-135 exhibiting the greatest decrease and Giza-130 maintaining relatively higher expression levels.

In this study, we examined *Nhx1* gene expression levels in three different cultivars of hulled barley under salt stress. According to the data shown in Fig. 4, *Nhx1* expression levels increased with increasing salt concentration. For example, increasing NaCl concentration from 50 mM to 100 mM increased the


expression levels from 1.5-fold to 6-fold in Giza-130 and from 2-fold to 7-fold in Giza-135. However, in Giza-136, the expression was down-regulated even at all salt concentration. A higher salt concentration of 200 mM decreased expression levels in all cultivars. Giza-136 showed a significant decrease compared with Giza-130, which maintained higher expression levels, followed by Giza-135 at the same salt concentration.

Discussion

Salt stress is a well-known abiotic as a major abiotic stress factor that negatively affects the growth, development, and yield of crops through ionic and osmotic stress [7,8,23]. In this study, seedlings of three hulls barley cultivars were exposed to saline stress for 15 days. Shoot length is an important indicator of a plant's ability to withstand salt stress. Increased NaCl concentrations significantly decreased shoot and root length in all three cultivars. Salinity stress also led to a reduction in plant fresh and dry weights, but the extent of the reduction varied among cultivars. Dry weight loss was minimal in Giza-130, followed by Giza-135, and Giza-136 showed the most significant reduction. These results show that Giza-130 has better salt tolerance than other cultivars, while Giza-136 is more sensitive.


Plants have developed various mechanisms, including regulation of gene expression, to cope with salt stress conditions. Several genes such as *hvpip*, *nhx1*, *cat*, *gst*, *sod*, and *apx* have been identified to play a role in plant response to salt stress [7,9]. *gst* is involved in detoxifying ROS in plants under stress conditions, including salt stress [24,25]. This study found that *gst* gene expression was upregulated under salt stress and Giza-130 showed the highest increase in

Figure 3



The effect of salinity stress on *hvpip* gene expression of hulled barley cultivars.

Figure 4

The effect of salinity stress on *nhx1* gene expression of hulled barley cultivars.

expression levels compared with Giza-135 and Giza-136. These findings suggest that the salt tolerance observed in Giza-130 is due to its ability to accumulate higher levels of *gst* expression in response to salt stress. The results also showed that *sod* gene expression was upregulated under salt stress, Giza-130 showed a higher increase in expression levels compared with Giza-135, in contrast, the expression in Giza-136 significantly decreased at 100 mM NaCl. These findings suggest that the response of barley cultivars to salt stress varies depending on their *sod* gene expression levels, which play a crucial role in protecting plants from oxidative stress caused by high salt concentrations, which in agreement with previous studies [26].

Cat and *apx* gene products are involved in detoxifying ROS in plant cells under salt stress conditions [19]. The results showed that *cat* gene expression was significantly higher in Giza-130 compared with other cultivars at 100 mM NaCl, but decreased significantly with an increase in salt concentration in all cultivars. *apx* gene expression was also upregulated at 100 mM, with Giza-130 showing the highest expression level (6-fold increase) compared with other cultivars at the same salt concentration. However, increasing the salt concentration to 200 mM led to a significant decrease in *apx* gene expression, with Giza-130 showing higher expression levels (3-fold increase), followed by Giza-135 (2-fold increase), while it was completely inhibited in Giza-136. These findings suggest that *cat* and *apx* play crucial roles in protecting plants from oxidative stress caused by high salt concentrations and that the response of Hulled barley cultivars to salt stress varies

depending on their *cat* and *apx* gene expression levels. These results agreed with previous studies [27].

The effect of salt stress on *hvpip* gene involved in water transport activity was investigated, in three Hulled barley cultivars. Under salt stress conditions, *hvpip* transcriptional folding was enhanced, Giza-130 showing the greatest increase in expression at 50 mM NaCl. Increasing NaCl concentration from 50 mM to 100 mM increased *hvpip* gene expression, with Giza-130 showing the greatest increase, consistent with previous studies [28,29].

Nhx1 gene is involved in the compartmentalization of sodium ions within plant cells, essential for maintaining cellular ion homeostasis under salt stress conditions. *nhx1* is upregulated in plants subjected to salt stress [30] and participate in K⁺ homeostasis, endosomal pH regulation, and salt tolerance in Arabidopsis [31]. The results showed that *nhx1* expression levels increased with increasing salt concentration in Giza-130 and Giza-135, but were down-regulated in Giza-136 at the same salt concentration. These findings suggest that the *nhx1* gene plays a role in sequestering Na⁺ in roots via vascular Na⁺/H⁺ antiporters, limiting the transport of toxic Na⁺ to shoots. Giza-130 maintained higher expression levels and showed better growth under high salinity conditions compared with Giza-136, which showed impaired *nhx1* gene expression at high salt concentration.

Conclusion

Genes involved in the regulation of cellular ion homeostasis, ROS detoxification, and water

transport in salt-stressed barley plants were investigated. The results of this study show that the expression levels of these genes vary between cultivars, with Giza-130 generally exhibiting higher expression levels and growing better under high salinity conditions than Giza-136. The results show that, these genes play an important role in protecting barley plants against oxidative stress caused by high salt concentrations and that different hulled barley cultivars have different abilities to cope with salt stress. These results could be useful for breeding programs aimed at developing salt-tolerant cultivars and improving crop productivity in saline environments.

Acknowledgments

The authors would like to express their sincere gratitude to the Genetic Engineering Research Center, Faculty of Agriculture, Cairo University for their invaluable support throughout the entire duration of this research work. Their guidance, expertise, and resources have been instrumental in the successful completion of this work. Also would like to extend their thanks to Dr. Ahmed Maher from the NRC for generously providing the barley grains for this study. His contribution has been essential in obtaining reliable results and achieving the objectives of this research.

Financial support and sponsorship

Nil.

Conflicts of interest

The authors declare that they have no conflict of interest.

References

- Girardo P, Benavente E, Manzano-Agugliaro F, Gimenez E. worldwide research trends on wheat and barley. A bibliometric comparative analysis. *Agronomy* 2019; 9:352.
- Zeng X, Guo Y, Xu Q, Mascher M, Guo G, Li S, *et al.* Origin and evolution of qingke barley in Tibet. *Nature Communications* 2018; 9:5433.
- Fischbeck G. Chapter 3 Diversification through breeding. In: Roland von Bothmer Tv H H K, Kazuhiro S, editors. *Developments in Plant Genetics and Breeding* 2003; 29–52
- Dickin E, Steele K, Edwards-Jones G, Wright D. Agronomic diversity of Hulled barley (*Hordeum vulgare* L.): a potential resource for breeding new food barley for Europe. *Euphytica* 2012; 184:85–99.
- Edney MJ, Tkachuk R, Macgregor AW. Nutrient composition of the hull-less barley cultivar, condor. *Journal of the Science of Food and Agriculture* 1992; 60:451–456.
- Sharma P, Gujral HS. Milling behavior of hulled barley and its thermal and pasting properties. *Journal of Food Engineering* 2010; 97:329–334.
- Cramer GR, Urano K, Delrot S, Pezzotti M, Shinozaki K. Effects of abiotic stress on plants. A systems biology perspective. *BMC Plant Biol* 2011; 11:1471–2229.
- Zhang HM, Zhu JH, Gong ZZ, Zhu JK. Abiotic stress responses in plants. *Nat. Rev. Genet.* 2022; 23:104–119.
- Rezvi HUA, Arif MTU, Azim MA, Tumpa TA, Tipu MMH, Najnine F, *et al.* Rice and food security. Climate change implications and the future prospects for nutritional security. *Food Energy Secur* 2022; 12:e430.
- Dmitrieva VA, Tyutereva EV, Voitsekhovskaja OV. Singlet oxygen in plants. Generation, detection, and signaling roles. *Int. J.Mol. Sci.* 2020; 21:3237.
- Hasanuzzaman M, Parvin K, Bardhan K, Nahar K, Anee TI, Masud AAC, *et al.* Biostimulants for the regulation of reactive oxygen species metabolism in plants under abiotic stress. *Cells* 2021; 10:2537.
- Salvaterra A, Mateluna P, Toro G, Solis S, Pimentel P. Genome-Wide Identification and Gene Expression Analysis of Sweet Cherry Aquaporins (*Prunus avium* L.) under Abiotic Stresses. *Genes* 2023; 14:940.
- Guo W, Xing Y, Luo X, Li F, Ren M, Liang Y. Reactive Oxygen Species: A Crosslink between Plant and Human Eukaryotic Cell Systems. *International Journal of Molecular Sciences* 2023; 24:13052.
- Bui QTN, Ki JS. 'Two novel superoxide dismutase genes (Cu Zn SOD and Mn SOD) in the toxic marine dinoflagellate *Alexandrium pacificum* and their differential responses to metal stressors'. *Chemosphere* 2023; 313:137532.
- Yamaguchi T, Aharon GS, Sottosanto JB, Blumwald E. Vacuolar Na⁺/H⁺ antiporter cation selectivity is regulated by calmodulin from within the vacuole in a Ca²⁺-and pH-dependent manner. *Proceedings of the National Academy of Sciences* 2005; 102:16107–16112.
- Katiyar-Agarwal S, Zhu J, Kim K, Agarwal M, Fu X, Huang A, *et al.* The plasma membrane Na⁺/H⁺ antiporter SOS1 interacts with RCD1 and functions in oxidative stress tolerance in Arabidopsis. *Proceedings of the National Academy of Sciences* 2006; 103:18816–18821.
- Munns R, Tester M. Mechanisms of salinity tolerance. *Annu. Rev. Plant Biol* 2008; 59:651–681.
- Zhu JK. Plant salt tolerance. *Trends in plant science* 2001; 6:66–71.
- Fathi A, Oveysi M, Nasri M, Tohidi-Moghadam HR, Kasraie P. Effect of Plant Hormones on Antioxidant Response and Essential Oil Production of Peppermint (*Mentha Piperita*) at Different Levels of Salinity Stress. *Gesunde Pflanzen* 2023; 75:1–12.
- Alshareef NO, Wang JY, Ali S, Al-Babili S, Tester M, Schmöckela SM. Overexpression of the NAC transcription factor JUNGBRUNNEN1 (JUB1) increases salinity tolerance in tomato. *Plant Physiology and Biochemistry* 2019; 113–121
- Murashige T, Skoog F. A revised medium for rapid growth and bioassays with tobacco tissue culture. *Physiologia Plantarum* 1962; 15:473–497.
- Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2- $\Delta\Delta$ CT method. *Methods* 2001; 25:402–408.
- Jain M, Ghanashyam C, Bhattacharjee A. Comprehensive expression analysis suggests overlapping and specific roles of rice glutathione S-transferase genes during development and stress responses. *BMC Genomics* 2010; 11:1–17.
- Wang R, Ma J, Zhang Q, Wu C, Zhao H, Wu Y, *et al.* Genome-wide identification and expression profiling of glutathione transferase gene family under multiple stresses and hormone treatments in wheat (*Triticum aestivum* L.). *BMC genomics* 2019; 20:1–15.
- Azeem M, Pirjan K, Qasim M, Mahmood A, Javed T, Muhammad H, *et al.* Salinity stress improves antioxidant potential by modulating physio-biochemical responses in *Moringa oleifera* Lam. *Scientific Reports* 2023; 13:2895.
- Kesawat MS, Satheesh N, Kherawat BS, Kumar A, Kim HU, Chung SM, *et al.* Regulation of reactive oxygen species during salt stress in plants and their crosstalk with other signaling molecules—Current perspectives and future directions. *Plants* 2023; 12:864.
- Horie T, Kaneko T, Sugimoto G, Sasano S, Panda SK, Shibasaki M, *et al.* Mechanisms of water transport mediated by PIP aquaporins and their regulation via phosphorylation events under salinity stress in barley roots. *Plant and Cell Physiology* 2011; 52:663–675.
- Bechtaoui N, Rabiou MK, Raklami A, Oufdou K, Hafidi M, Jemo M. Phosphate-dependent regulation of growth stresses management in plants. *Frontiers in Plant Science* 2021; 12:679916.
- Rodríguez-Rosales MP, Jiang X, Gálvez FJ, Aranda MN, Cubero B, Venema K. Overexpression of the tomato K⁺/H⁺ antiporter LeNHX2 confers salt tolerance by improving potassium compartmentalization. *New Phytologist* 2008; 179:366–377.
- Bassil E, Ohto MA, Esumi T, Tajima H, Zhu Z, Cagnac O, *et al.* The Arabidopsis intracellular Na⁺/H⁺ antiporters NHX5 and NHX6 are endosome associated and necessary for plant growth and development. *The Plant Cell* 2011; 23:224–239.
- Barragan V, Leidi EO, Andres Z, Rubio L, De Luca A, Fernandez JA, *et al.* Ion exchangers NHX1 and NHX2 mediate active potassium uptake into vacuoles to regulate cell turgor and stomatal function in Arabidopsis. *The Plant Cell* 2012; 24:1127–42.