

**GROWTH PERFORMANCE AND CARCASS  
RESPONSE OF RABBITS FED ON DIETS  
SUPPLEMENTED WITH DIFFERENT LEVELS OF  
SILVER NANO PARTICLES.**

**Hoda M. El-Hosseiny<sup>1</sup> ; G. F. Shaheen<sup>1\*</sup> ; Kout-Elkloub M. Moustafa<sup>2</sup> ;  
G.E. Younan<sup>3</sup> ; E. M. El- Kotamy<sup>1</sup> ; A. Ghoniem<sup>1</sup> ; M. M. El-Nahrawy<sup>1</sup> ;  
Mona E. Farag<sup>1</sup> ; A. A. Abdou<sup>1</sup> and M. E. Sayed-Ahmed<sup>1</sup>**

1 Department of Animal Nutrition Research. 2 Department of Poultry Nutrition  
Research 3 Department of rabbits, Turkey and waterfowl Breeding Research,  
Animal Production Research Institute (APRI), Agricultural Research center  
(ARC), Dokki, Giza, Egypt.

\*E-mail-G.F Shaheen, [gamal.shaheen@arc.sci.eg](mailto:gamal.shaheen@arc.sci.eg).

**ABSTRACT**

Fifty male New Zealand White rabbits, aged 6 weeks and average live body weight of  $710.74 \pm 10$  g, were randomly assigned to five groups (10 rabbits each) and fed basal diets containing varying levels of Silver Nano Particles (Ag NPs) for a 14-weeks experimental period. This study evaluated the efficacy of using different levels of Ag NPs as dietary supplementation in growing rabbits.

A completely randomized design was employed, with the first group G1 (control) receiving the basal diet without supplementation. Rabbits in the other groups G2, G3, G4, and G5 were fed the basal diet supplemented with Ag NPs at levels of 1.5, 2.0, 2.5, and 3.0 mg/kg of diet, respectively.

The obtained results indicated that the highest values of live body weight, daily weight gain, hot carcass weight, and dressing percentage were observed in rabbits fed diets supplemented with Ag NPs across all levels compared to the control group. However, nutrient digestibility and nutritive values were not significantly affected by the nanoparticle supplementation. Additionally, feed conversion ratios, net revenue, and economic efficiency values improved in the groups receiving Silver Nano Particles when compared to the control group.

Our findings suggest that the Silver Nano Particles in rabbit diets positively influenced performance and economic efficiency compared to the control group. The optimal results were obtained at 2 mg Ag NPs/kg of diet.

**Key Words:** Rabbits, Ag NPs, Productive performance and Economic efficiency.

## INTRODUCTION

Nanomaterials and nanoparticles are key components of nanotechnology, typically ranging in size from 1 to 100 nanometers in at least one dimension (Ahmadi and Kordestany, 2010). The size of these particles is crucial because it influences the action of nanoparticles (Mikhailova, 2020). Nanoparticles smaller than 300 nm can circulate within the body, while those smaller than 100 nm can more easily and quickly reach target cells, tissues, and organs (Choi and Hu, 2008; Prabhu and Poulouse, 2012). This small size also helps reduce the presence of residual compounds and decreases the withdrawal time of drugs in livestock tissues (Vadalasetty *et al.*, 2018). In medicine, nanoparticles can be used to target specific tissues while minimizing side effects and toxicity (Sadr *et al.*, 2023). Nanoparticles can be administered through drinking water, feed, injections, or inhalation (Salleh *et al.*, 2020). Recently, nanotechnology has been widely applied in animal nutrition to help explain metabolic and physiological mechanisms, improve fiber digestion, promote health, and enhance animal production overall (Rajendran *et al.*, 2014). When used as supplements, nano-minerals offer several advantages over traditional minerals, including higher bioavailability at lower doses, larger surface areas, higher surface activity, and stronger adsorption capabilities (Chaudhry and Castle, 2011 ; Shi *et al.*, 2011; Albanese *et al.*, 2012 and Rajendran *et al.*, 2014). Silver nanoparticles (Ag-NPs) play a significant role in animal production due to their disinfectant and antimicrobial properties. Ag-NPs had a potent inhibitory effect against *E. coli* growth in in vitro trail (Awaad *et al.*, 2021). While in in vivo trail, decreased the number of harmful bacteria represented as *E. coli* and had no effect on microflora represented as lactobacillus (Moustafa *et al.*, 2015). Also, it used as drugs to combat bacteria, fungi, and viruses (Sawosz *et al.*, 2007 ; Zhang *et al.*, 2016 and Ali, 2019). Moreover, Ag-NPs are known to have anticancer properties (Hamed *et al.*, 2017 and Salem *et al.*, 2021) and are seen as alternatives to antibiotics in the fight against drug-resistant bacteria (Hill and Li, 2017 and Zhang *et al.*, 2016). Silver nanoparticles help reduce ammonia and nitrogen oxide emissions (Xu *et al.*, 2013), aid in regulating rumen fermentation and overcoming reproductive challenges in animals (Gopi *et al.*, 2017). Moustafa *et al.*, (2015) investigated adding silver nanoparticles to broiler chicken diets (2, 4, 6, 8 and 10 ppm/kg) improved body weight, body weight gain and feed conversion ratio . In addition, cholesterol & total lipids were significantly decreased ,and increased total serum antioxidant capacity . Also, increase the European Production Efficiency Index (EPEI) and generally, the best level was 4 mg/kg diet. Ahmed *et al.*, (2023) found that adding Ag-NPs to chicken diets enhanced performance, improved nutrient utilization, and boosted economic efficiency. Other studies found that supplemented with Ag-NPs had no

significant effect on chicken performance when it added at a ratio of 0, 4, 8 or 12 ppm in drinking water by **Ahmadi and Rahimi (2011)**, at 0,10 and 25 ppm by **Pineda et al., (2010)**, at 50 ppm by **Vadalasetty et al., (2018)**. **Al-Sultan et al., (2022)** suggested that high doses of Ag-NPs (20 mg/kg) might reduce protein digestion and inhibit the absorption of sugars and amino acids, leading to lower body weight and organ mass in broilers. Similarly, **Tammam et al., (2023)** noted no significant impact of silver or graphene nanoparticle supplementation on carcass characteristics or immune organs in broilers. While **Loghman et al., (2012)**, observed liver damage at high doses (12 ppm) of Ag-NPs of broiler chicks. In rabbits, **Kotb et al., (2024)** reported that supplementing diets with Ag-NPs, especially at 1 mg/kg, improved body weight, weight gain, and feed efficiency. Lower doses (0.25 mg/kg) enhanced liver and kidney function, lipid profile, immunity, antioxidant capacity, and digestive enzyme activity while promoting beneficial gut bacteria with rabbits. **Abdelsalam et al., (2019)**, found no significant impact on daily gain or feed intake when rabbits were injected with Ag-NPs at doses of 0.5 mg and 1 mg/kg body weight. However, they noted that plasma cholesterol and triglyceride levels were lower in treated rabbits compared to the control ration, and the accumulation of silver in the blood and meat increased with higher doses. There is a lack of findings on experimental utility of Nano particles, especially Silver nitrate for rabbit production. So, the present study was aimed to evaluate the effect of using different levels of Ag NPs as dietary supplementation on nutrient digestibility, growth performance, carcass traits, some blood plasma biochemical, histological features of liver, kidney and economic efficiency of growing New Zealand White rabbits during the fattening period (6-14 weeks of age).

## MATERIAL AND METHODS


### Ethical Approval

This study was conducted at the Sakha Experimental Research Station in Kafr El-Sheikh, under the Animal Production Research Institute, Agriculture Research Center, Ministry of Agriculture, Egypt. The study followed all ethical guidelines for animal welfare, in line with the European Union's regulations (2010/63/EU) on the protection of animals used in experiments. The study was officially registered with the serial number (372429).

### Preparation and characterization:

Silver nanoparticles (Ag-NPs) were chemically prepared in the Poultry Nutrition Department's laboratories at the Animal Production Research Institute. The process involved reducing silver nitrate (AgNO<sub>3</sub>) using diluted aqueous solutions that contained cetyl trimethyl ammonium bromide and hydrazine hydrate as a dispersing agent to produce Ag-NPs. A colour change was observed during the mixing process, indicating

successful nanoparticle formation, as described by **Khan *et al.*, (2011)** and **Moustafa *et al.*, (2015)**. To confirm the synthesis of Ag-NPs, a UV-visible spectrophotometric analysis was performed. This analysis, conducted at the National Research Center in Dokki, Egypt, showed a peak wavelength of 424 nm (Fig.1), a key indicator that silver nitrate (AgNO<sub>3</sub>) had converted to nanoparticles (Ag-NPs). Additionally, the size and distribution of the particles were examined using a transmission electron microscope (TEM), revealing that the nanoparticles ranged in size from 5 to 25 nm and their size distribution was relatively wide (Fig.1). Flame atomic absorption spectrometry (Agilent Technologies 200 Series AA, Agilent Technologies, Santa Clara, CA) was used to measure the concentration of Ag-NPs at wavelength 328.1 nm. When silver particles reach nano-size, they exhibit a gray colour due to surface plasmon resonance (**Govindaraju *et al.*, 2009**), which was confirmed with an absorption peak at 424 nm. The analysis, conducted using the JEM 1400 microscope at Cairo University, confirmed the successful synthesis of Ag-NPs. The concentration of Ag-NPs was found to be 2.15 g/L. Different feed concentrations of Ag-NPs were then prepared at levels of 1.5, 2, 2.5, and 3 mg/kg of diet.



**Figure 1.** UV-visible spectrophotometer analysis and Transmission Electron Microscopy of Ag-NPs.

#### **Animals and management:**

Fifty weaned male New Zealand White rabbits, each around six weeks old with an average live body weight of  $710.74 \pm 10$  g, were randomly divided into five groups (10 rabbits per group). Each group was fed a

different level of Silver Nanoparticles (Ag-NPs) for the entire experimental period, which lasted at 14 weeks of age. The rabbits were housed individually in battery cages (30 × 60 × 40 cm) located in a naturally ventilated building. The environmental conditions included a temperature range of 22-25°C, relative humidity of 60-70%, and a lighting schedule of 14 hours of light and 10 hours of darkness. The experiment took place from November to December, and all rabbits were maintained under the same management, hygiene, and environmental conditions throughout the study.

#### Experimental diets:

Five experimental treatments were used, with a completely randomized design. The first group (G1), serving as the control and it was fed a basal diet without any Ag-NPs supplements to meet maintenance and production requirements of rabbits, as mentioned in **NRC guidelines (1977)**. The other four groups (G2, G3, G4, and G5) received the same basal diet with added Ag-NPs at levels of 1.5, 2.0, 2.5, and 3.0 mg/kg of diet, respectively. The rabbits were fed *ad-libitum* and fresh water was available at all times through an automatic system. The composition and calculated analysis of the basal diet are presented in Table 1.

**Table 1: The composition and calculated analysis of the basal diet.**

Ingredient	%	Calculated analysis	
Berseem hay	30.05	Crude protein (%)	17.75
Barley	24.60	Digestible energy kcal/kg	2500
Wheat bran	21.50	Crude fiber (%)	15.38
Soybean meal (44% CP)	17.50	Ether extract (%)	2.27
Molasses	3.00	NFE (%)	54.16
Di-calcium phosphate	1.60	Ash (%)	10.44
Limestone	0.95	Calcium (%)	1.24
Sodium chloride (NaCl)	0.30	Total phosphorus (%)	0.80
Vitamin & mineral mixture*	0.30	Lysine (%)	0.98
DL-Methionine	0.20	Methionine (%)	0.46
		Methionine + Cysteine (%)	0.76
		Sodium (%)	0.16

\*Each kilogram of diet contained : Vitamin A, 6000 IU; Vitamin D<sub>3</sub>, 900 IU; Vitamin E, 40 mg; Vitamin K<sub>3</sub>, 2 mg; Vitamin B<sub>1</sub>, 2 mg; Vitamin B<sub>2</sub>, 4 mg; Vitamin B<sub>6</sub>, 2 mg; Pantothenic acid, 10 mg; Vitamin B<sub>12</sub>, 0.01 mg; Niacin, 50 mg; Folic acid, 3 mg; Biotin, 0.05 mg; Choline, 250 mg; Fe, 50 mg; Mn, 8.5 mg; Cu, 5 mg; Co, 0.1 mg; Se, 0.1 mg; I, 0.2 mg and Zn, 50 mg.

#### Digestibility trials:

At the end of the experimental period, a digestibility trial was conducted to assess the rabbits' nutrient digestibility. This involved five separate digestion experiments, one for each group, using all rabbits from each group when they reached 14 weeks old. Each rabbit was housed individually in metabolic cages (60 × 50 × 40 cm) designed to separate feces from urine. Feces were collected daily and treated with a 2% boric acid solution to trap ammonia for five days. After collection,

the feces were dried at 60°C for 48 hours, ground into fine powder, mixed thoroughly to ensure uniformity, and stored for analysis according to AOAC (1995) guidelines.

**Growth performance:**

The rabbits live body weight (LBW) and feed intake were recorded monthly, weight gain (difference between final and initial weight/period), and feed conversion ratio (FCR) (dividing daily feed intake by daily weight gain) were calculated.

**Carcass traits:**

At the end of the study, three rabbits from each group were randomly taken and fasted for 16 hours before slaughter. Carcass traits were evaluated, including carcass weight, edible organs (liver, heart, kidney, spleen), total non-carcass fat %, fur, and dressing %, following the method described by Hassan *et al.*, (2016).

**Blood sampling:**

During slaughtering, blood samples (5 ml/each sample) were taken individually from three rabbits in each group (15 samples). Blood samples were collected in dry clean centrifuge tubes containing a few drops of heparin solution and centrifuged at 3000 rpm for 20 minutes to separate blood plasma and stored in a deep freezer at approximately -20°C until the time of analysis to estimate blood parameters (Kairalla *et al.*, 2022). Blood plasma samples were determined by spectrophotometer (Spectronic 21 DUSA) using commercial diagnostic kits supplied by Bio Merieux, France, following the same steps described by manufacturers. The blood plasma was measured for total protein, albumin, glucose, urea, creatinine, AST, ALT, cholesterol, HDL, LDL, triglycerides, T<sub>3</sub> and T<sub>4</sub>.

**Histological study:**

Representative samples were taken from the medial part of the liver and kidney of slaughtered male rabbits from each group (1 sample/rep.). Samples were placed in 10% buffered formalin and paraffin-embedded. Sections were stained with hematoxylin and eosin (H&E) stains and histologically examined using a light microscope (Optico XSZ-107B Binocular, China) at x200 magnifications (Bancroft *et al.*, 1996).

**Economic efficiency:**

Economic efficiency was calculated by comparing the income from weight gain to the cost of feed consumed at 14 weeks of age.

Net revenue (LE) = Total revenue – Total feed cost.

Economic efficiency % = (Net revenue (LE)/ Total feed cost (LE)) \*100.

Ingredient prices and the market price of live rabbits (100 Egyptian Pounds LE, per kilogram) were based on local market conditions in January 2022.

**Statistical Analysis:**

The data were analyzed using a one-way analysis of variance (ANOVA) based on a Completely Randomized Design, applying the General Linear Models procedure (SAS, 2009). The model used for statistical was as follows:

$$Y_{ij} = \mu + T_i + e_{ij}$$

Where:

$Y_{ij}$  = The observation,

$\mu$  = Overall mean,

$T_i$  = Effect of treatments (i: 1 to 5),

$e_{ij}$  = random error, assumed to be normally distributed.

Duncan's multiple range test (Duncan, 1955) was used to detect significant differences among treatment means.

**RESULTS****Nutrients digestibility and nutritive values:**

Results in Table 2 show that the rabbit fed  $G_3$ , supplemented with Ag NPs, had the highest nutrient digestibility percentages across all nutrients. Rabbits followed this in group  $G_4$ , then group  $G_2$  and finally the animals fed the control rations  $G_1$ . The lowest digestibility values were in the  $G_5$  group, fed the highest level of Ag NPs. However, the differences in nutrient digestibility among these groups were not statistically significant. Similarly, the nutritive values, expressed as Total Digestible Nutrients (TDN) and Digestible Crude Protein (DCP), were not significantly impacted by the supplementation of Ag NPs at any level when compared to the control group.

**Table 2: Nutrients digestibility and nutritive values of rabbit's diet with Nano Silver supplementation.**

Item	Experimental groups				
	$G_1$	$G_2$	$G_3$	$G_4$	$G_5$
<b>Digestibility (%)</b>					
DM	64.84±1.68	65.51±3.17	66.41±5.20	65.65±2.84	64.20±3.58
OM	71.83±0.94	72.66±3.74	73.68±3.79	72.96±2.24	71.56±3.54
CP	65.17±2.30	65.25±3.39	66.09±3.73	65.71±4.53	64.01±3.28
EE	76.17±4.18	76.39±5.80	76.84±5.50	76.57±6.84	75.90±2.49
CF	52.33±1.40	52.43±2.13	53.69±1.77	52.96±1.80	52.14±2.45
NFE	78.17±4.43	78.83±3.72	79.51±4.78	78.93±4.23	78.08±5.29
<b>Nutritive values (%)</b>					
TDN	65.85±1.14	66.23±2.70	66.96±2.38	66.46±1.40	65.57±3.16
DCP	11.57±0.24	11.58±0.84	11.73±0.75	11.66±0.87	11.63±0.91

$G_1$ : Control contained zero Ag NPs of diet.  $G_2$ : Contained 1.5 mg/kg Ag NPs of diet.  $G_3$ : Contained 2 mg/kg Ag NPs of diet.  $G_4$ : Contained 2.5 mg/kg Ag NPs of diet.  $G_5$ : Contained 3 mg/kg Ag NPs of diet. DM: Dry matter. OM: Organic matter. CP: Crude protein. EE: Ether extract. CF: Crude fiber. NFE: Nitrogen free extracts. TDN: Total digestible nutrients. DCP: Digestible crude protein.

**Growth performance:**

The Results presented in Table 3 indicated that rabbits fed diets supplemented with Ag NPs had significantly higher average live body weight

(LBW) ( $p \leq 0.05$ ) compared to those in the control group ( $G_1$ ) by the end of the 14-weeks. Whereas, the best average LBW was recorded with  $G_3$ , during the same period. Average daily gain (g/head) was increased ( $p \leq 0.05$ ) with Ag NPs supplementation within different levels and rabbit ages in comparison with those in the control group ( $G_1$ ). Rabbits receiving 2 mg/kg of Ag NPs ( $G_3$ ) showed the highest daily weight gain compared to other groups. The results also reveal that the average daily feed intake (g/head/day) was a slight increase in rabbits supplemented with Ag NPs compared to the control group during the periods from (6 to 10) and (10 to 14) weeks of age. Moreover, a significant improvement in feed conversion ratio ( $p \leq 0.05$ ) was observed in rabbits fed diets supplemented with silver nanoparticles compared to the control group during the experimental period. The best feed conversion ratios were observed in group  $G_3$ , which fed diets supplemented with 2 mg/kg of silver nanoparticles compared with the other treatments. Overall, rabbits fed 2 mg/kg of Ag NPs ( $G_3$ ) demonstrated improved feed intake and the best feed conversion ratio compared to all other treatments.

**Table 3 Rabbits performance values as affected by different levels of Nano silver supplementation at different ages**

Item	Experimental groups					±SEM
	$G_1$	$G_2$	$G_3$	$G_4$	$G_5$	
<b>Live Body Weight, g</b>						
Initial Weight	717	708.5	696.0	712.7	719.5	35.47
at 10 weeks	1311.0 <sup>b</sup>	1393 <sup>ab</sup>	1444.5 <sup>a</sup>	1439.5 <sup>a</sup>	1360.5 <sup>ab</sup>	27.99
at 14 weeks	1921.16 <sup>c</sup>	2111 <sup>ab</sup>	2213.5 <sup>a</sup>	2157 <sup>ab</sup>	2024 <sup>b</sup>	40.42
<b>Daily weight gain, g/head</b>						
6-10 weeks	21.19 <sup>c</sup>	24.45 <sup>ab</sup>	27.95 <sup>a</sup>	25.96 <sup>ab</sup>	23.03 <sup>b</sup>	0.99
10-14 weeks	21.79 <sup>d</sup>	25.64 <sup>ab</sup>	26.72 <sup>a</sup>	25.63 <sup>ab</sup>	23.77 <sup>c</sup>	1.45
6-14 weeks	21.49 <sup>c</sup>	25.05 <sup>ab</sup>	27.33 <sup>a</sup>	26.09 <sup>ab</sup>	23.50 <sup>b</sup>	0.75
<b>Daily feed intake, g/head/day</b>						
6-10 weeks	61.82	62.25	62.21	62.14	61.73	0.90
10-14 weeks	102.84	105.32	106.50	105.68	104.61	1.03
6-14 weeks	82.33	83.79	84.36	83.91	83.17	0.78
<b>Feed conversion ratio</b>						
6-10 weeks	2.92 <sup>a</sup>	2.55 <sup>cb</sup>	2.23 <sup>c</sup>	2.39 <sup>bc</sup>	2.68 <sup>b</sup>	0.11
10-14 weeks	4.72 <sup>a</sup>	4.11 <sup>ab</sup>	3.99 <sup>b</sup>	4.12 <sup>ab</sup>	4.40 <sup>a</sup>	0.24
6-14 weeks	3.83 <sup>a</sup>	3.35 <sup>bc</sup>	3.09 <sup>c</sup>	3.22 <sup>bc</sup>	3.54 <sup>b</sup>	0.09

<sup>a, b, c, d</sup> means in the same row with different superscripts are significantly different ( $P \leq 0.05$ ).

$G_1$ : control contained zero Ag NPs of diet.  $G_2$ : contained 1.5 mg/kg Ag NPs of diet.  $G_3$ : contained 2 mg/kg Ag NPs of diet.  $G_4$ : contained 2.5 mg/kg Ag NPs of diet.  $G_5$ : contained 3 mg/kg Ag NPs of diet. Feed conversion ratio= feed intake, g / daily weight gain, g

### Carcass traits:

The results in Table 4 indicate that both the hot carcass weight and dressing percentage were significantly ( $p \leq 0.05$ ) higher in rabbits supplemented with Ag NPs compared to the control group. The highest



values for these parameters were observed in Groups G3 and G4, supplemented with 2 and 2.5 mg Ag NPs per kg of diet, respectively. For the relative weights of the kidney, liver, heart, spleen, giblets, and fur, no significant differences were noted among the experimental groups, except for the spleen, which showed a significant difference ( $p \leq 0.05$ ) between groups. Conversely, the relative weights of the head and total non-carcass fat were the lowest in the control group (G1) compared to the Ag NP-treated groups, with significant differences ( $p \leq 0.05$ ).

**Table 4: Carcass parameters of growing rabbits as affected by different levels of Nano silver supplementation.**

Item	Experimental groups					±SE
	G <sub>1</sub>	G <sub>2</sub>	G <sub>3</sub>	G <sub>4</sub>	G <sub>5</sub>	
Per-slaughter (g)	1900 <sup>ab</sup>	1990 <sup>a</sup>	1890 <sup>b</sup>	1953 <sup>ab</sup>	1941 <sup>ab</sup>	50.26
Hot-carcass, %	42.04 <sup>c</sup>	50.42 <sup>ab</sup>	53.45 <sup>a</sup>	52.28 <sup>a</sup>	51.74 <sup>ab</sup>	1.34
Dressing, %	55.19 <sup>c</sup>	64.00 <sup>ab</sup>	65.47 <sup>a</sup>	65.29 <sup>a</sup>	61.37 <sup>b</sup>	1.27
Liver %	2.52	2.61	2.47	2.73	2.57	0.15
Kidney %	0.84	0.61	0.76	0.81	0.64	0.08
Heart %	0.37	0.39	0.34	0.34	0.28	0.03
Spleen %	0.06 <sup>ab</sup>	0.1 <sup>a</sup>	0.1 <sup>a</sup>	0.08 <sup>ab</sup>	0.04 <sup>b</sup>	0.02
Giblets weight %	3.37	3.61	3.57	3.88	3.49	3.89
Non-carcass fat weight %	0.58 <sup>b</sup>	1.05 <sup>ab</sup>	1.86 <sup>a</sup>	1.55 <sup>a</sup>	1.22 <sup>a</sup>	0.46
Head %	5.55 <sup>b</sup>	6.46 <sup>ab</sup>	6.66 <sup>a</sup>	6.66 <sup>a</sup>	6.80 <sup>a</sup>	0.16
Fur %	18.00	18.5	18.18	17.40	15.25	1.03

<sup>a, b</sup> means in the same row with different superscripts are significantly different ( $P \leq 0.05$ ).

G<sub>1</sub>: control contained zero Ag NPs of diet. G<sub>2</sub>: contained 1.5 mg/kg Ag NPs of diet.

G<sub>3</sub>: contained 2 mg/kg Ag NPs of diet. G<sub>4</sub>: contained 2.5 mg/kg Ag NPs of diet.

G<sub>5</sub>: contained 3 mg/kg Ag NPs of diet.

**Table 5: Some blood plasma parameters of male rabbits fed diets containing different levels of Nano silver particles.**

Item	Experimental groups					±SE
	G <sub>1</sub>	G <sub>2</sub>	G <sub>3</sub>	G <sub>4</sub>	G <sub>5</sub>	
Total protein (g/dl)	6.55 <sup>a</sup>	6.00 <sup>b</sup>	5.80 <sup>b</sup>	5.93 <sup>b</sup>	6.47 <sup>ab</sup>	0.22
Albumin (g/dl)	3.94	4.69	4.72	4.36	4.72	0.26
Globulin (g/dl)	2.51 <sup>a</sup>	1.31 <sup>b</sup>	1.08 <sup>b</sup>	1.57 <sup>ab</sup>	1.75 <sup>ab</sup>	0.35
Cholesterol (mg/dl)	131.00 <sup>a</sup>	108.0 <sup>ab</sup>	104.0 <sup>ab</sup>	101.67 <sup>ab</sup>	90.00 <sup>ab</sup>	5.34
HDL (mg/dl)	40.25 <sup>a</sup>	35.65 <sup>b</sup>	30.50 <sup>b</sup>	30.43 <sup>b</sup>	30.25 <sup>b</sup>	6.56
LDL (mg/dl)	49.35 <sup>bc</sup>	79.15 <sup>a</sup>	49.47 <sup>bc</sup>	38.30 <sup>c</sup>	41.00 <sup>c</sup>	7.64
Triglycerides (mg/dl)	123.33 <sup>a</sup>	114.30 <sup>ab</sup>	113.67 <sup>ab</sup>	113.00 <sup>ab</sup>	111.67 <sup>ab</sup>	12.20
Glucose (mg/dl)	98.0 <sup>b</sup>	104.00 <sup>ab</sup>	129.33 <sup>a</sup>	107.00 <sup>ab</sup>	121.67 <sup>a</sup>	7.78
Urea (mg/dl)	45.67 <sup>b</sup>	52.00 <sup>a</sup>	56.03 <sup>a</sup>	55.50 <sup>a</sup>	52.30 <sup>a</sup>	7.94
Creatinine (mg/dl)	1.46	1.36	1.31	1.31	1.35	0.14
AST (U/l)	43.67 <sup>a</sup>	29.33 <sup>ab</sup>	29.00 <sup>ab</sup>	27.33 <sup>b</sup>	25.00 <sup>b</sup>	4.20
ALT (U/l)	87.33	66.67	71.50	87.00	78.00	12.78
T <sub>3</sub> (ng/ml)	1.20	0.96	0.94	0.73	0.74	0.18
T <sub>4</sub> (ng/ml)	3.49	2.00	2.83	2.37	2.10	0.68

<sup>a, b, c</sup> means in the same row with different superscripts are significantly different ( $P \leq 0.05$ ).

G<sub>1</sub>: control contained zero Ag NPs of diet. G<sub>2</sub>: contained 1.5 mg/kg Ag NPs of diet. G<sub>3</sub>:

contained 2 mg/kg Ag NPs of diet. G<sub>4</sub>: contained 2.5 mg/kg Ag NPs of diet. G<sub>5</sub>:

contained 3 mg/kg Ag NPs of diet.


**Blood parameters:**

Results in Table 5 show that the total plasma protein, globulin, cholesterol and triglycerides concentrations were lowest ( $p \leq 0.05$ ) in the groups treated with Ag NPs compared to the control group. However, there were no significant effects of Ag NP supplementation on plasma albumin, creatinine, ALT,  $T_3$  and  $T_4$  levels. On the other hand, the plasma LDL (except in groups  $G_4$  and  $G_5$ ), glucose, and urea concentrations were numerically higher in the Ag NP-treated groups compared to the control group, with significant differences ( $p \leq 0.05$ ) among them. In contrast, HDL and AST concentrations significantly decreased ( $p \leq 0.05$ ) in the Ag NP-treated groups compared to the control.

**Histological Study:**


The histological analysis of the liver from slaughtered rabbits (Plate 1) indicated that the liver architecture in the Ag NP-treated groups ( $G_2$ ,  $G_3$ ,  $G_4$  and  $G_5$ ) was normal, with no pathological lesions observed, similar to the control group ( $G_1$ ). Likewise, the histological examination of the kidneys (Plate 2) showed typical renal cortex and medulla architecture across all treated groups, with no differences compared to the control group.

**Plate 1:** The histological examination of liver of slaughtered rabbits.



( $\times 200$ , H&E stains)

**Plate 2:** The histological examination of Kidneys of slaughtered rabbits.



(×200, H&E stains)

**Table 6: Economical efficiency for growing rabbits feeding diets containing different levels of Ag NPs.**

Item	Experimental groups				
	G <sub>1</sub>	G <sub>2</sub>	G <sub>3</sub>	G <sub>4</sub>	G <sub>5</sub>
Total body weight gain (kg)	1.208	1.403	1.518	1.444	1.305
Price of 1 kg body weight (L.E)	100	100	100	100	100
Selling price/rabbit (L.E) A	120.8	140.3	151.8	144.4	130.5
Total feed intake (kg)	4.627	4.692	4.724	4.700	4.657
Price of 1kg feed (LE)	14	14	14	14	14
Total feed cost/rabbit (LE) B	64.78	65.69	66.14	65.80	65.20
Net revenue (L.E) <sup>1</sup>	56.02	74.61	85.40	78.60	65.30
Economic efficiency <sup>2</sup>	0.87	1.14	1.29	1.20	1.00
Relative EEF, %	100	131.04	148.28	137.93	114.94

<sup>1</sup>Net revenue = A-B, <sup>2</sup> Economical efficiency= Net revenue / B

Based on prices of the Egyptian market during the experimental period (2022).

The price of one kg live body weight was 100 Egyptian Pound (LE).

G<sub>1</sub>: control contained zero Ag NPs of diet.

G<sub>2</sub>: contained 1.5 mg/kg Ag NPs of diet.

G<sub>3</sub>: contained 2 mg/kg Ag NPs of diet.

G<sub>4</sub>: contained 2.5 mg/kg Ag NPs of diet.

G<sub>5</sub>: contained 3 mg/kg Ag NPs of diet.

**Economic efficiency:**

Table 6 indicates that rabbits fed diets supplemented with Ag NPs at various levels exhibited notable profit margins compared to the control group at the marketing age of 16 weeks. The relative economic efficiencies (%) were as follows: 131.04 for G2, 148.28 for G3, 137.93 for G4, and 114.94 for G5. The highest relative economic efficiency was observed in Group G3, which was treated with a 2.5 mg/kg diet of Ag NPs, reaching 148.28%. This suggests that incorporating Ag NPs into the diets of growing male rabbits is economically feasible.

**DISCUSSION****Nutrient digestibility and nutritive value:**

Silver nanoparticles (Ag NPs) are still being researched in the livestock, rabbit, and poultry sectors. The present study was aimed to evaluate the impact of Ag NPs as dietary supplementation on different parameters, including nutrient digestibility, growth performance, blood consistency, carcass traits, histological examination, and economic efficiency in growing New Zealand White (NZW) rabbits. The findings revealed that the nutrient digestibility and nutritive values (%), including Total Digestible Nutrients (TDN) and Digestible Crude Protein (DCP), showed insignificant improvement across all groups treated with Ag NPs compared to the control group. This could be attributed to the potential of Ag NPs to enhance the population of beneficial microbes in the gut. Notably, the highest level of Ag NPs (3 mg/kg diet, G<sub>5</sub>) yielded the least favorable outcomes, indicating a dose-dependent effect. Research by **Albanese et al., (2012)** suggests that Ag NPs can stimulate digestive enzyme activity, thereby improving nutrient digestibility (**Kumar et al., 2020; Bolandi et al., 2021 and Kotb et al., 2024**). Previous studies (**Cho et al., 2005 and Sawosz et al., 2007**) highlighted the antimicrobial properties of silver nanoparticles, which can adversely affect both Gram-positive and Gram-negative bacteria, including antibiotic-resistant strains. **Sotiriou and Pratsinis (2010)**, noted that Ag NPs release silver ions (Silver + ions from their surface) in contact with water, which exhibit strong antimicrobial effects by disrupting cellular respiration and compromising bacterial cell membranes (**Vadalasetty et al., 2018**). Moreover, Ag NPs may alter the phosphotyrosine profile of bacterial peptides, hindering growth (**Shrivastava et al., 2007**). However, **Awaad et al., (2021)**, indicated that the effectiveness of Ag NPs is dose-dependent and is more pronounced against Gram-negative bacteria. On the other hand, the antimicrobial activity of metal nanoparticles can be attributed to cell membrane disruption, cell wall damage, and cytoplasmic leakage, ultimately leading to microbial death (**Gajbhiye et al., 2009 and Hassan et al., 2014; 2015**).

**Growth performance:**

The current study demonstrated that average live body weight (LBW) in rabbits supplemented with Ag NPs was significantly higher ( $p \leq 0.05$ ) compared to the control group (G1) at 14 weeks. The percentage increases in

LBW were 9.88% for G1, 15.22% for G2, 12.28% for G3, and 5.35% for G4. Additionally, Ag NPs supplementation in rabbit diet at different levels increased ( $p \leq 0.05$ ) average daily gain (g/head) of G1, G2, G3 and G4 groups as compared to the control group. The increasing rate was 16.57, 27.29, 21.41 and 9.35 %, respectively at 14 weeks. The highest LBW and average daily gain were observed in groups G<sub>3</sub> and G<sub>4</sub>, supplemented with 2 mg and 2.5 mg Ag NPs per kg of diet. Rabbits in groups G<sub>1</sub>, G<sub>2</sub>, G<sub>3</sub>, and G<sub>4</sub> also exhibited significant improvements ( $p \leq 0.05$ ) in total feed intake (g/head/week), average daily feed intake (g/head/day), and feed conversion ratio compared to the control group (G<sub>1</sub>) during the 6-14 week period. The best feed conversion ratio (g feed intake/g daily weight gain) was noted in Group G<sub>3</sub> (2 mg/kg diet). These improvements in growth performance may be attributed to the stimulation of digestive enzyme activity by Ag NPs (Kumar *et al.*, 2020; Bolandi *et al.*, 2021 and Kotb *et al.*, 2024), an increase in beneficial microbial populations, enhanced nutrient digestibility and absorption (Sadr *et al.*, 2023), and reduced ammonia and nitrogen emissions (Xu *et al.*, 2013). Moustafa *et al.*, (2015) and Saleh and El Magd (2018) noted that chicks receiving Ag NPs at 50 ppm exhibited increased body weight gain and muscle weight, along with improved feed conversion.

In contrast, Abdelsalam *et al.*, (2019) found that although final body weight increased with Ag NPs injection at 0.5 mg/kg body weight, while, body weight gain, feed intake, and feed conversion ratio showed no significant changes. El-Faham *et al.*, (2017), reported that Ag NPs in drinking water at 0.5 or 10 ppm did not significantly affect live body weight, daily weight gain, or feed conversion in rabbits and broilers. Hang and Tra (2013) also observed significant increases ( $p \leq 0.05$ ) in live body weight when rabbits received Ag NPs in drinking water.

#### **Carcass traits:**

The results presented in Table 4 indicate that carcass traits, particularly hot carcass weight and dressing percentage, were significantly affected ( $p \leq 0.05$ ) by the supplementation of Ag NPs compared to the control rations. The highest values were observed in the G3 and G4 groups, which were supplemented with Ag NPs at 2 and 2.5 mg/kg of diet, respectively. This improvement in carcass traits can be attributed to the enhanced digestion coefficients and nutritional value (as shown in Table 2), which positively impacted both live body weight and average daily gain, ultimately leading to increased hot carcass weight and dressing percentages. The relative organ weights of the kidneys, liver, heart, giblets, and fur were numerically similar to those in the control group, except for the spleen, which exhibited significant differences among the treated and control group. These findings align with Abdelsalam *et al.*, (2019), who reported similar relative weights for the liver, kidney, and heart in rabbits injected with Ag NPs at 0.5 or 1 mg/kg body weight. Conversely, Hassan (2018), found that Ag NPs supplementation in the

drinking water or feed of broilers significantly affected dressing percentage, liver, and heart weights.

Additionally, **Al-Sultan et al., (2022)**, demonstrated that broiler diets supplemented with Ag NPs at various concentrations (2.5, 5, 10, or 20 ppm) improved the relative weights of warm dressing, heart, liver, and gizzard. However, **Kotb et al., (2024)**, reported no significant effects on carcass traits such as dressing percentage and organ weights (except for liver percentage) due to Ag NPs supplementation in rabbit diets. Similarly, **El-Faham et al., (2017)** found no significant impact on the carcass traits of quail, broilers, and rabbits when Ag NPs were supplemented in drinking water at concentrations of 0.5 or 10 ppm. **Ahmadi et al., (2013) and Abdelsalam et al., (2019)**, also reported no significant effects of Ag NPs on liver, heart, kidney, or spleen percentages and dressing percentages or carcass traits. **Tammam et al., (2023)**, found no significant differences in carcass traits (including carcass percentage, giblets, total edible parts, and abdominal fat percentage) or organ parts (breast, thigh, drumstick, and wings percentages) when Ag NPs were used in broiler drinking water.

#### **Blood parameters:**

##### **Blood plasma metabolites**

The results presented in Table 5 ,regarding blood plasma metabolites, indicate that plasma total protein and globulin levels were significantly ( $p \leq 0.05$ ) decreased of the treated groups receiving Ag NPs compered with the control rations. Plasma albumin concentrations showed an insignificant increase compared to the control group. **Ashour et al., (2004)**, noted that albumin concentration is linked to an animal's ability to store and synthesize protein. Additionally, **Jones and Park (1979)**, reported that changes in albumin levels reflect liver function, as the liver is responsible for albumin synthesis, whereas globulin is formed by lymphoid tissue. Our results are partially consistent with those of **Sultan et al., (2022)**, who found that supplementation of 2.5 mg Ag NPs/kg of broiler diet significantly increased total protein levels, while albumin and globulin showed insignificant increases. In contrast, **Kotb et al., (2024)**, reported significant increases in total protein, albumin, and globulin levels in rabbits fed diets supplemented with Ag NPs at 0.25 and 0.75 mg. **Abdelsalam et al., (2019)**, stated that blood plasma content of total protein, albumin, and globulin were not significantly affected when rabbits were injected with Ag NPs at 0.5 or 1 mg/kg body weight. **Moustafa et al., (2015)**, demonstrated that broiler diets supplemented with Ag NPs at 2, 4, 6,8 and 10 ppm significantly affected blood globulin level, while total protein and albumin were not significantly impacted. **Syrvatka et al., (2014)** reported that silver nanoparticle supplementation did not alter any blood parameters.

##### **Blood Lipid Profile**

The results presented in Table 5 indicate that the concentrations of plasma total cholesterol and triglycerides were significantly decreased in rabbits

treated with Ag NPs compared to the control group. These findings align with those of **Abdelsalam et al., (2019)**, who reported that rabbits injected with Ag NPs at a dosage of 0.5 mg exhibited reduced plasma concentrations of total cholesterol and triglyceride levels. **Saleh and El-Magd (2018)**, also noted significant reductions in plasma triglycerides and total cholesterol when broiler chickens were fed a diet containing 50 ppm/kg of Ag NPs, without affecting total protein levels. Additionally, **Kotb et al., (2024)** found that serum total cholesterol and LDL concentrations significantly decreased in rabbits receiving 0.75 mg Ag NPs/kg diet compared to the control and other treatment groups. Conversely, **Al-Sultan et al., (2022)**, reported a significant increase in blood total cholesterol levels and an insignificant increase in triglyceride concentrations in broilers fed diets supplemented with Ag NPs at 2.5 mg/kg. Furthermore, **Ahmadi (2012)**, observed no significant changes in lipid profiles due to Ag NPs supplementation. Also, **Sawosz et al., (2009)**, reported that silver nanoparticles did not affect the activities concentrations of cholesterol.

Our study also revealed a significant ( $p \leq 0.05$ ) decrease in plasma HDL concentrations in rabbits fed diets containing Ag NPs at all levels compared to the control group. Conversely, LDL concentrations significantly ( $p \leq 0.05$ ) increased with a 1.5 mg/kg diet compared to the other treated and the control groups. This finding is consistent with **Ahmadi (2012)**, who noted that chickens fed diets containing small amounts (8 and 12 ppm/kg) of Ag NPs exhibited increased LDL cholesterol and decreased HDL cholesterol concentrations.

#### **Liver Function**

In terms of liver function, the data indicated that plasma ALT and AST activities decreased in rabbits treated with Ag NPs compared to the control group. The decrease in AST was significant ( $p \leq 0.05$ ), while the decrease in ALT was not significant. This transaminase activity suggests no damage to hepatic cells due to Ag NPs, in accordance with **Mezey (1976)**, who stated that any damage to hepatic cells or their membranes results in increased AST and ALT levels in the blood serum. **Dosoky et al., (2021)**, supported this indicating that decreased liver enzyme activity, such as ALT and AST, correlates with protein metabolism and varying levels of transaminases. Our findings are consistent with those of **Moustafa et al., (2015)** who demonstrated that broiler diets supplemented with Ag NPs at 2, 4, 6,8 and 10 ppm significantly affected AST level, while ALT were not significantly impacted. **Kotb et al., (2024)**, found significant reductions in serum AST and ALT levels with Ag NPs supplementation at up to 0.75 mg/kg diet. **Ahmadi (2012)**, also reported decreased AST and ALT activities in the blood plasma of chickens receiving Ag NPs at 20, 40, and 60 ppm/kg of feed. In contrast, other studies (**Sawosz et al., 2009; Andi et al., 2011; Saleh and El-Magd, 2018**), indicated that Ag NPs supplementation in broiler chickens had no significant effects on ALT and AST activity levels.



### **Kidney Function**

Regarding kidney function, the data in Table 5 revealed that Ag NPs supplementation in rabbit diets significantly ( $p \leq 0.05$ ) increased plasma urea values, while plasma creatinine levels were decreased (though insignificantly) compared to the control group. These results are partially consistent with those reported by **Kotb et al., (2024)**, who found that rabbits fed diets including Ag NPs at 0.50 and 0.75 mg/kg exhibited significantly lower creatinine concentrations. They also reported that urea concentration was significantly decreased at 0.25 mg Ag, while 0.50 and 1 mg Ag NPs led to increased serum urea levels. Conversely, **Sultan et al., (2022)** observed insignificantly higher urea concentrations in broilers fed Ag NPs at mg/kg, while **Dosoky (2021)**, noted decreased urea and creatinine concentrations in broiler chicks fed Ag NPs hydrocolloid. **Lohakare and Abdel-Wareth (2022)** found no effect on urea or creatinine levels in broiler diets.

### **Hormonal and Glucose Parameters**

The data in Table 5 also showed that plasma thyroid hormones (T3 and T4) concentrations were insignificantly decreased in rabbits treated with Ag NPs compared to the control group. However, glucose concentrations significantly ( $p \leq 0.05$ ) increased in rabbits treated with Ag NPs compared to the control group. These findings contrast with those of **Sawosz et al., (2009)**, who reported that silver nanoparticles did not affect the activities of glucose concentrations.. Overall, the blood plasma metabolites analyzed in this study were within the normal levels for healthy rabbits, as indicated by **Özkan et al., (2012)**.

### **Histological Examination:**

The histological examination of the liver and kidneys from slaughtered rabbits revealed normal hepatic lobule and renal cortex structures across all treated groups. This observation is attributed to the blood plasma parameters remaining within normal physiological ranges, indicating that the developing rabbits demonstrated remarkable adaptive abilities to maintain high productive performance levels without compromising liver and kidney function. **Hassan (2009)**, emphasized that the normal histological structure correlates with the AST and ALT activity results in the blood serum of rabbits, which were within normal values for growing rabbits (Table 5).

### **Economic efficiency:**

From an economic standpoint, feeding growing rabbits diets supplemented with Ag NPs led to reduced feed costs and increased net revenue and relative economic efficiency compared to the control group. The highest relative economic efficiency was observed in the G3 (148.28%) and G4 (137.93%) groups, which were treated with 2 and 2.5 mg Ag NPs/kg, respectively. These improvements can be attributed to enhanced growth performance, dressing percentage, and overall health care of the rabbits. **Saleh and El-Magd (2018)**, attributed growth improvements to Ag NPs' positive



effects on the balance of beneficial and harmful bacteria in the intestines and their impact on digestive enzyme activity, which improved nutrient absorption and promoted a healthier digestive tract. Other studies (**El-Sabry *et al.*, 2018 and Salem and Fouda., 2021**), have reported that nanoparticles increased survivability, production performance, and overall physical function across various animal species.

### CONCLUSION:

The present study concluded that supplementation of Silver Nanoparticles (Ag NPs) in rabbit diets at varying doses positively affected growth performance, carcass quality, health status, and economic efficiency, without adverse effects on blood metabolism or histological findings. Optimal results were obtained at 2 mg Ag NPs/kg of diet. There is a notable lack of studies exploring the experimental benefits of nanoparticles, particularly silver nitrate, as feed additives for rabbits. Thus, further research is warranted to determine a safe dosage that balances the health status and productivity of the growing rabbits while minimizing any potential harmful effects. Additionally, caution should be exercised regarding high doses of Ag NPs, as they can lead to toxicity due to accumulation in blood plasma and meat.

### REFERENCES

- Abdelsalam, M. ; I. Al-Homidan ; T. Ebeid ; O. Abou-Emera ; M. Mostafa ; M.A. El-Razik ; M. Shehab -El-Deen ; S. Abdel Ghani and M. Fathi (2019)**. Effect of silver nanoparticle administration on productive performance, blood parameters, antioxidative status, and silver residues in growing rabbits under hot climate. *Anim.* , 9(10): 845.
- Ahmadi, F. (2012)**. Impact of different levels of silver nanoparticles (AgNPs) on performance, oxidative enzymes and blood parameters in broiler chicks', *Pak. Vet. J.*, 26(3) 325-328.
- Ahmadi, F. ; M.M. Khah ; S. Javid ; A. Zarneshan ; L. Akradi and P. Salehifar (2013)**. The effect of dietary silver nanoparticles on performance, immune organs, and lipid serum of broiler chickens during starter period', *Int. J. Biosci.*, 3: 95–100.
- Ahmadi, F. and A.H. Kurdestany (2010)**. The impact of silver nano particles on growth performance, lymphoid organs and oxidative stress indicators in broiler chicks. *Glob. Vet.*, 5(6): 366–370.
- Ahmadi, F. and F. Rahimi (2011)**. The effect of different levels of Nano Silver on performance and retention of silver in edible tissues of broilers. *World Appl. Sci. J.*, 12: 1–4.
- Ahmed, M.M.N. ; Z.S.H. Ismail ; I. Elwerdany and A.A.A. Abdel-Wareth. (2023)**. Applications of biosynthesis of silver nanoparticles for

- sustainable poultry production under hot climatic conditions. A review. *Int. J. Agric. Sci.*, 5 (1): 137-151.
- Albanese, A. ; P.S. Tang and W.C. Chan (2012).** The effect of nano particle size, shape and surface chemistry on biological systems. *Annual Review of Biomed. Eng.* 14:1-6
- Ali, Z. S. (2019).** Synthesis, characterization of silver nano particles using *Nigella sativa* seeds and study their effects on the serum lipid profile and DNA damage on the rat's blood treated with hydrogen peroxide. *Iraqi J. Vet. Med.*,43(2):23-37.
- Al-Sultan S.I.;A.R.T. Hereba; K.M.A.Hassanein ; S.M.S. Abd-Allah; U.T. Mahmoud and S.M. Abdel-Raheem (2022).** The impact of dietary inclusion of silver nanoparticles on growth performance, intestinal morphology, caecal microflora, carcass traits and blood parameters of broiler chickens. *Ital. J. Anim. Sci.*, 21: 967–978.
- Andi, M.A. ; M. Hashem and F. Ahmadi (2011).** Effects of feed type with/without nano silver on cumulative performance, relative organ weight and some blood parameters of broilers. *Glob. Vet.*, 7: 605–609.
- AOAC. (1995).** Official Methods of Analysis, 15<sup>th</sup> Ed. Association of Official Analytical Chemists, Washington, DC, USA.
- Ashour, G. ; S.A. Ibrahim ; A.M. Ismail and K.H. El-Kholy (2004).** Physiological reactions and biological performance of rabbits to summer heat stress. 2<sup>th</sup> Sci. Conf. on Physio. Resp., to Environ. Condition, 28-31 July, 2004, El-Arish. Egypt, pp:165-186.
- Awaad, M.H.H. ; K.M. El-Moustafa ; Zoulfakar, S.A.M.S.Elhalawany, ; F.F. Mohammed ; R.M. El-Refay and E.A. Morsy (2021).** The role of silver nanoparticles in the reluctance of colisepticemia in broiler chickens. *J. Appl. Poult. Res.*, 30:100155
- Bancroft, J.D. ; A. Stevens and D.R. Turner. (1996).** Theory and Practice of Histological Techniques. 4th ed.Churchill, Livingstone, New York.
- Bolandi, N. ; S.R. Hashemi ; D. Davoodi ; B. Dastar ; S. Hassani and A. Ashayerizadeh (2021).** Performance, intestinal microbial population, immune and physiological responses of broiler chickens to diet with different levels of silver nanoparticles coated on zeolite.*Ital. J. Anim. Sci.*, 20: 497–504.
- Chaudhry, Q. and L. Castle (2011).** Food applications of nano technologies: An overview of opportunities and challenges for developing countries. *Food Sci. and Technol.*, 22(11): 595- 603.
- Cho, K.H. ; J.E. Park ; T. Osaka and S.G. Park (2005).** ‘The study of antimicrobial activity and preservative effects of Nano-Silver ingredient’. *Electrochim. Acta*, 51: 956–960.

- Choi, O. and Z. Hu (2008).** Size dependent and reactive oxygen species related nano silver toxicity to nitrifying bacteria. *Environ. Sci., Technol.*, 42(12):4583-4588.
- Dosoky, W.M. ; M.M.G. Fouda ; A.B. Alwan ; N.R. Abdelsalam ; A.E. Taha ; R.Y. Ghareeb ; M.R. El-Aassar and A.F. Khafaga (2021).** Dietary supplementation of silver-silica nanoparticles promotes histological, immunological, ultrastructural, and performance parameters of broiler chickens. *Sci. Rep.*, 11: 4166.
- Duncan, D.B. (1955).** Multiple range and multiple F-test. *Biometrics*. 11:1-42.
- El-Faham, A.I. ; M.H.S. El-Sanhoury and M.M.E. Mostafa (2017).** Effect of nano-silver particles supplementation in drinking water on performance and intestinal microflora population of growing poultry. *Egypt. J. Nut. and Feeds*. 20 (3): 519-528.
- El-Sabry, M.I. ; K.W. McMillin and C.M.Sabliov (2018).** Nanotechnology considerations for poultry and livestock production systems–A review. *Annals of Anim. Sci.*,18 (2): 319.
- Gajbhiye, M. ; J. Kesharwani ; A. Ingle ; A. Gade and M. Rai. (2009).** Fungus-mediated synthesis of silver nanoparticles and their activity against pathogenic fungi in combination with flu- conazole. *Nanomed*.5:382–386.
- Gopi, M. ; B. Pearlin ; R.D. Kumar ; M. Shanmathy and G. Prabaker (2017).** Role of nanoparticles in animal and poultry nutrition. Modes of action and applications in formulating feed additive food processing. *Int. J. Pharmacol.*, 13 (7): 723 – 731.
- Govindaraju, K. ; V. Kiruthiga ; V. Kumar and G. Sigar-avelu (2009).** Extracellular synthesis of silver nanoparticles by a marine alga *Sargassum wightii* Grevilli and their antibac- terial effects. *J. Nanosci. Nanotechnol.*, 9:5497–5501.
- Hamed, S. ; M. Emara ; R.M. Shawky ; R.A. El-Domany and T. Youssef (2017).** Silver nanoparticles antimicrobial activity, cytotoxicity, and synergism with Nacetyl cysteine. *J. Basic Microbiol.*, 57(8): 659–668
- Hang, D.T. and T.T.T. Tra (2013).** Effect on rabbit reproduction of adding silver-nano suspension to the drinking water. *Livest. Res. Rural. Dev.*, 25: 1–7.
- Hassan, A.A. ; H.K. Mahmoud ; H. Taha ; R.H. Sayed El-Ahl and H. H. Mahmoud (2015).** Herbal biosynthesis of zinc nanoparticles and evaluation of their antifungal and antibacterial effect for buffaloes skin affections. *Int. J. Curr. Res.*, 7:24338–24349.
- Hassan, A.A. ; N.H. Oraby ; A.E. Mohamed and H. H. Mahmoud (2014).** The possibility of using zinc oxide nano particles in controlling some fungal and bacterial strains isolated from buffaloes. *Egypt. J. Appl. Sci.* 29:58–83.

- Hassan, A.M. (2009).** Effect of boldenoneundecy Lenate treatment on buffalo male calf performance. M.Sc. Thesis, Faculty of Agric., Menoufiya University.
- Hassan, A.M. (2018).** Effect of nano silver on performance and some physiological parameters of broiler chicks under south condition. Int. J. Innoy, Appi. Res., 6: 1-8.
- Hassan, F. ; K.M. Mahrose and M.M. Basyony (2016).** Effects of grape seed extract as a natural antioxidant on growth performance, carcass characteristics and antioxidant status of rabbits during heat stress. Arch Anim Nutr., 70 (2):141-154.
- Hill, E.K. and J. Li (2017).** Current and future prospects for nanotechnology in animal production. J. Anim. Sci. Biotechnol., 8 (1): 1-13.
- Jones, E.A. and P.D. Bark (1979).** Chemical Diagnosis of Disease. Brown, S.S., F. L. Mitchell and D.S. Young (Ed.), Elsevier, Biomedical Press, Amsterdam, New York, Oxford, p.325 -363.
- Kairalla, M.A. ; M.I. Alshelmani and A.A.Aburas (2022).** Effect of diet supplemented with graded levels of garlic (*Allium sativum* L.) powder on growth performance, carcass characteristics, blood hematology, and biochemistry of broilers. Vet. J., 12(5): 595-601.
- Khan, Z. ; S.A. Al-Thabaiti ; A.Y. Obaid and A.O. Al- Youbi (2011).** Preparation and characterization of silver nanoparticles by chemical reduction method. Colloids Surf. B Biointerfaces 82:513–517.
- Kotb, O.A. ; A.I. Attia ; F.M. Reda ; S.A. Mahgoub ; M. Alagawany and M.S. El- Kholy (2024).** Impact of silver nanoparticles (Ag-NPs) as dietary supplement on growth performance, carcass traits, blood metabolites, digestive enzymes, and cecal microbiota of growing rabbits. Annals of Anim. Sci., 24(4): doi:10.2478/aoas-2024-0034.
- Kumar, I. ; J. Bhattacharya ; B.K. Das and P. Lahiri (2020).** Growth, serum biochemical, and histopathological responses of broilers administered with silver nanoparticles as a drinking water disinfectant, Biotechnol., 10(94): 1–12.
- Loghman, A. ; S.H. Iraj ; D.A. Naghi and M. Pejman (2012).** Histopathologic and apoptotic effect of nano silver in liver of broiler chickens. Afr. J. Biotechnol., 11(22): 6207-6211.
- Lohakare, J. and A.A.A. Abdel-Wareth (2022).** Effects of dietary supplementation of oregano bioactive lipid compounds and silver nanoparticles on broiler production, Sustainability,14, 13715.
- Mezey, E. (1976).** Dagnosis of Liver Disease By Laboratory Methods. In The Laboratory Of Clinical Medicine. Interpretation And Application. Ed. Holsted, J. Pub. W.B. Saundess Co., London, Toronto, pp. 417-429.

- Mikhailova, E.O. (2020).** Silver Nanoparticles: Mechanism of Action and Probable Bio- Application. *J. Funct. Biomater.*, 11: 84.
- Moustafa, K.E. ; M.; A.A. Ghazalah and A.A.A. Rehan (2015).** Effect of Dietary Nanosilver on Broiler Performance. *Int. J. of Poultry Sci.*, 14 (3): 177-182.
- NRC, (1977).** Nutrient Requirements of Rabbits. National Research Council, Washington DC.
- ÓZkan, C. ; A. Kaya and Y. Akgül (2012).** Normal values of haematological and some biochemical parameters in serum and urine of New Zealand white rabbits. *World Rabbit Sic.*, 20: 253 -259.
- Pineda, L. ; E. Chwalibog ; A. Sawosz ; J. Hotowy ; X. Hotowy ; F. Sawosz ; T. Niemiec and A. Ali (2010).** Effects Of Nano Silver On Energy Metabolism And Growth Performance Of Broiler Chicken. 3<sup>rd</sup> EAAP Internatuional Symposium on Energy and Protein Metabolism and Nutrition (ISEP). 6-10 September. Parma, Italy.
- Prabhu, S. and E.K. Poulose (2012).** Silver nanoparticles: mechanism of antimicrobial synthesis, medical applications, and toxicity effects. *Int. Nano Lett.*, 2: 32.
- Rajendran, D. ; A. Thulasi ; S. Jash ; S. Selvaraju and S.B.N. Rao (2014).** Synthesis and Application Of Nano Minerals In Livestock Industry. *Animal Nutrition And Reproductive Physiology (Recent concepts)*. Satish Serial Publishing House, Delhi. 517 – 530. Adugodi, Bangalore-560 030, Karnataka, India
- Sadr, S. ; N. Lotfalizadeh ; S. A. Ghafouri ; M. Delrobaei ; N. Komeili and A. Hajjafari (2023).** Nanotechnology innovations for increasing the productivity of poultry and the prospective of nanobiosensors. *Vet. Med. Sci.*, 11: 1–14.
- Saleh, A.A. and M.A. El-Magd (2018).** Beneficial effects of dietary silver nanoparticles and silver nitrate on broiler nutrition. *Environ. Sci. Pollut. Res.*, 25: 27031–27038.
- Salem, H.M. ; E. Ismael and M.E. Shaalan (2021).** Valuation of the effects of silver nanoparticles against experimentally induced necrotic enteritis in broiler chickens, *Int. J. Nanomed.*, 5: 6783–6796.
- Salem, S.S. and A. Fouda (2021).** Green synthesis of metallic nanoparticles and their prospective biotechnological applications: An overview. *Biol. Trace. Elem. Res.*, 199: 344–370
- Salleh, A. ; R. Naomi ; N.D. Utami ; A.W. Mohammad ; E. Mahmoudi ; N. Mustafa and M.B. Fauzi (2020).** The potential of silver nanoparticles for antiviral and antibacterial applications: A mechanism of action. *Nanomaterials*, 10: 1566.

- SAS (2009).** User,s Guide . Statistics. Version, 9.2, 2002-2009, SAS, Inst. Inc., Cary, NC, USA.
- Sawosz, E. ; M. Binek ; M. Grodzik ; M. Zielinska ; P. Sysa ; M. Szmidt ; T. Niemiec and A. Chwalibog (2007).** Influence of hydrocolloidal silver nanoparticles on gastrointestinal microflora and morphology of enterocytes of quails. Arch. Anim. Nutr., 61: 444–451.
- Sawosz, E. ; M. Grodzik ; M. Zieliska ; T. Niemiec ; B. Olszaska and A. Chwalibog (2009).** Nanoparticles of silver do not affect growth, development and DNA oxidative damage in chicken embryos, Arch. Geflugelk., 73: 208–213.
- Shi, L.; X.Wenjuan; Y. Wenbin; Z.Chunxiang ;R. Youshe; L.Qiang ;W. Qian and S. Lei (2011).** Effect of elemental nano-selenium on feed digestibility, rumen fermentation, and purine derivatives in sheep. Anim. Feed Sci. and Technol., 163:136-142.
- Shrivastava, S. ; T. Bera ; A. Roy ; G. Singh ; P. Ramachandrarao and D. Dash (2007).** Characterization of enhanced antibacterial effects of novel silver nanoparticles, Nanotechnol., 18: 225103.
- Sotiriou, G.A. and S.E. Pratsinis (2010).** Antibacterial activity of nano silver ions and particles. Environ. Sci. Technol., 44(14): 5649–5654.
- Syrvatka, V. ; I. Rozgoni ; Y. Slyvchuk ; G. Milovanova ; I. Hevkan and I. Matyukha (2014).** Effects of silver nanoparticles in solution and liposomal form on some blood parameters in female rabbits during fertilization and early embryonic development. J. Microbiol. Biotechnol. Food Sci., 3: 274–278.
- Tammam, A.M. ;S.A. Ibrahim ; A.A. Hemid ; F. Abdel-Azeem ; A.I. El-Faham ; N.G.M. Ali and W. Salem (2023).** Comparative effects of nano particles (silver vs graphene) in drinking water for broiler chickens. Egyptian J. Nut. and Feeds,26 (2):1-8.
- Vadalasetty, K.P. ; C. Lauridsen ; R.M. Engberg ; R. Vadalasetty ; M. Kutwin ; A. Chwalibog and E. Sawosz (2018).** Influence of silver nanoparticles on growth and health of broiler chickens after infection with *Campylobacter jejuni*. BMC Vet. Res., 14: 14.
- Xu, Y. ; H. Tang ; J.H. Liu ; H. Wang and Y. Liu (2013).** Evaluation of the adjuvant effect of silver nano particles both in vitro and in vivo. Toxicol. Letters, 219:42-48.
- Zhang, X.F. ; W. Shen and S. Gurunathan (2016).** Silver nanoparticle-mediated cellular responses in various cell lines, an in vitro model. Int. J. Mol. Sci., 17(10).ID:1603 doi: org/10.3390/ijms17101603.

## أداء النمو وأستجابة الذبيحة للأرانب التي تتغذى على علائق مضاف إليها مستويات مختلفة من جزيئات نانو فضة

هدى محمد الحسيني<sup>1</sup>، جمال فاروق شاهين<sup>1</sup>، قوت القلوب محمد مصطفى<sup>2</sup>، جورج يونان<sup>3</sup>، عصام القتامي<sup>1</sup>، عبد الغني غنيم<sup>1</sup>، مصطفى محمد النحراوي<sup>1</sup>، منى أحمد فراج<sup>1</sup>، على أحمد عبده<sup>1</sup>، محمد السيد سيداحمد<sup>1</sup>

1- قسم بحوث تغذية الحيوان ، 2- قسم بحوث تغذية الدواجن، 3- قسم بحوث تربية الأرانب و الرومي والطيور المائية معهد بحوث الإنتاج الحيواني ، مركز البحوث الزراعية ، الدقي، الجيزة، مصر .

أجريت الدراسة بأستخدام عدد 50 ذكر من أرانب النيوزيلندي الابيض، متوسط أعمارهم 6 أسابيع ووزنهم الحي  $710.74 \pm 10$  جرام، تم توزيعهم بشكل عشوائي على 5 مجموعات (10 أرانب لكل مجموعة) وتم تغذيتهم بأعلاف أساسية تحتوي على مستويات متفاوتة من جزيئات الفضة النانوية (Ag NPs) لفترة تجريبية مدتها 14 أسبوعاً. أجرى هذا البحث بهدف تقييم فعالية استخدام مستويات مختلفة من جزيئات الفضة النانوية كمكملات غذائية للأرانب النامية. حيث غذيت المجموعة الأولى G1 (المجموعة الضابطة) على علف الأساسي بدون مكملات. و تم تغذية الأرانب في المجموعات الأخرى G2 و G3 و G4 و G5 على العلف الأساسي (المجموعة الضابطة) مضاف إليه جزيئات الفضة النانوية بمستويات 1.5 و 2 و 2.5 و 3 مجم / كجم من العلف على التوالي. وكانت أهم نتائج هذه التجربة ما يلي:

وجد أن أعلى قيم لوزن الجسم الحي و معدل زيادة اليومية ووزن الذبيحة ونسبة التصافي كانت في الأرانب التي تغذت على مكملات جزيئات الفضة النانوية في جميع المستويات مقارنة بالمجموعة الضابطة. ومع ذلك، لم تتأثر معاملات الهضم والقيم الغذائية بشكل كبير بمكملات جسيمات النانو. كذلك تحسنت الكفاءة الغذائية و الاقتصادية في المجموعات التي تلقت جزيئات الفضة النانوية بالمقارنة بالمجموعة الضابطة.

من هذه الدراسة ينصح بأضافة جزيئات الفضة النانوية في علائق الأرانب كأضافات غذائية حيث أثرت بشكل إيجابي على الأداء الأنتاجي والكفاءة الاقتصادية مقارنة بالمجموعة الضابطة. وقد وجد أن اضافة 2 مجم من جزيئات الفضة النانوية / كجم من العلف أعطى أفضل النتائج.