

Arthroscopic-assisted mini-open rotator cuff repair outcome measures

Mohamed R.A. Saleh

Department of Orthopedics, Faculty of Medicine,
Helwan University, Helwan, Egypt

Correspondence to Mohamed R.A. Saleh, MD,
Tel: +20 100 665 4001;
e-mails: docmohamedrabie@gmail.com,
mohamed.rabie@med.helwan.edu.com

Received: 22 May 2022

Revised: 01 June 2022

Accepted: 03 June 2022

Published: 14 April 2023

The Egyptian Orthopaedic Journal 2023,
57:225–237

Background

Mini-open repair seems to be equivalent to all-arthroscopic repair in multiple nonrandomized comparative studies, and a randomized trial is underway to formally address this question.

Objectives

These include the ability to perform diagnostic arthroscopy, preservation of the origin of the deltoid, rapid hospital discharge, less postoperative pain, and accelerated rehabilitation.

Patients and Methods

A case series study was performed on 20 patients from May 2017 to May 2019. The technique is not technically demanding than all-arthroscopic repair while still having the advantages of arthroscopic repair. A total of 20 patients, 11 (55%) men and nine (45%) women, were included in this study with a mean age of 58.00 Å} 9.17 years.

Results

There was high significant gradual improvement in the modified University of California, Los Angeles (UCLA) shoulder scale among the studied patients at 1, 3, and 6 months postoperative follow-up compared with preoperative. Postoperative complications were found in six patients. Three (15%) cases had stiffness, and the other three (15%) cases had superficial wound infections. There were significant correlations between the modified UCLA shoulder scale with age, operative time, and postoperative complications, while there were no significant correlations with sex or side.

Conclusion

Our study concluded that surgery for rotator cuff tears improves self-reported patient outcomes and has a lasting and durable result at an average of 15 years after surgery. There was a high statistically significant gradual improvement in the modified UCLA shoulder scale among the studied patients at 1, 3, and 6 months postoperative follow-up compared with preoperative. Further studies including a larger sample are required to enhance the current findings.

Keywords:

arthroscopic, cuff repair, postoperative outcomes, University of California, Los Angeles

Egypt Orthop J 2023, 57:225–237

© 2023 The Egyptian Orthopaedic Journal

1110-1148

Introduction

Rotator cuff injuries are considered the most common tendon injuries in adults, overall. Rotator cuff injury is a common cause of shoulder pain and disability and becomes more common with advancing age [1]. Most symptomatic rotator cuff disease is seen between fifth and sixth decades. Rotator cuff tears are associated with pain and weakness and can result in significant disability [2]. It is also known that asymptomatic rotator cuff tears exist in a large percentage of patients, and asymptomatic tears increase with increasing age [3]. The cause of a rotator cuff tear is most likely related to a combination of several factors including impingement against the subacromial arch, age-related degeneration, overuse, and post traumatic [4]. A rotator cuff tear can range from small to large. One or more of your tendons in your rotator cuff could tear if you continue the activity and do not have the injury

treated. Then the tear can become worse. It is essential you receive the right treatment so your rotator cuff can function optimally [5]. There are different types of rotator cuff tears. Partial rotator cuff tears: this is a damaged rotator cuff tendon, but it is not torn all the way through [6]. This is also called a partial thickness tear. Complete rotator cuff tear: this is when you have soft tissue that tears into two different pieces. Often, the tendons tear away from the upper arm bone. A full thickness tear does not heal by itself since your muscles pull the tear's edges apart. But it is possible for a partial or full thickness tear to stabilize and leave your shoulder with reasonable function

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.


and comfort [7]. Acute tear: injury causes these tears if you lift a heavy object or fall on outstretched hands, you could have an acute tear. Shoulder injuries, such as a dislocated shoulder or broken collarbone, can also cause an acute rotator cuff tear [8]. Degenerative tear: genetics and specific health conditions like high cholesterol and diabetes can also cause degeneration of the cuff. Because of this, the patient's dominant side is more likely to get a rotator cuff tear because he tends to use it more, and it experiences repetitive stress. Degeneration also naturally with advancing age, increases the chances of injury over time [9]. Cofield classified rotator cuff tears are classified according to tear size to small less than 1 cm, medium 1–3 cm, large 3–5 cm, and massive more than 5 cm. Full thickness rotator cuff tear: Ellman and Gartsman [10] classified rotator cuff tear according to the tear morphology into: crescent reverse L, L-shaped, trapezoidal, and massive tear full-thickness rotator cuff tears (Fig. 1).

The rationale for repairing the rotator cuff is derived from multiple published studies demonstrating functional improvement and pain reduction after rotator cuff repair and rehabilitation [11]. Although complete healing of the tendon does not occur in all cases, rotator cuff repair is a beneficial procedure

for relieving pain, improving strength, and the for improving the range of motion. The earliest report of rotator cuff repair comes from Codman [12]. Since then, many studies have demonstrated good outcomes with improved pain and function following formal open repair of the rotator cuff with subacromial decompression and acromioplasty [13]. The method by which the cuff is repaired has changed during the past two decades, with a movement toward minimally invasive techniques (arthroscopic-assisted mini-open and arthroscopic repair) [14]. The arthroscopic-assisted mini-open or deltoid-splitting approach to the rotator cuff is a well-characterized procedure with excellent outcomes and is a successful method of rotator cuff repair [14].

Frisella and Cuomo [15], including rotator cuff tears. The ability to visualize the anatomy of the shoulder through the arthroscope inevitably led to strategies to treat rotator cuff tears by a less invasive technique. Before arthroscopy, rotator cuff tears were treated by open repair with approaches that violated the deltoid insertion on the acromion [16]. The deltoid was detached from the acromion in order to perform an acromioplasty and repaired to the acromion at the end of the procedure [17]. This approach carried the risk of

Figure 1



Classified rotator cuff tear according to the tear morphology.[10]

deltoid avulsion [17]. Diagnosis and characterization of tears by arthroscopy led to the description of the arthroscopically assisted, mini-open, or deltoid-splitting repair technique of rotator cuff repair [15]. Mini-open repair remains a viable alternative to arthroscopic repair and has advantages over both arthroscopic and open repair [15]. The mini-open rotator cuff repair represents a bridge between open and arthroscopic cuff repair. It has advantages and disadvantages when compared with other methods of cuff repair [18]. In addition, the deltoid origin is minimally disturbed, allowing for fast rehabilitation and decreasing the possibility of avulsion of deltoid as one of the complications [15]. The primary advantage of arthroscopic-assisted mini-open repair of the rotator cuff tear is the avoidance of passing and tying complex techniques of arthroscopic suture. It creates less surgical trauma, facilitating early hospital discharge and decreasing postoperative pain [19]. Thus, this study aimed to evaluate outcome measures and postoperative complications of arthroscopic-assisted mini-open rotator cuff repair.

Patients and methods

A case series study included 20 cases studied from June 2019 till June 2020 with a follow-up period of 12 months after obtaining approval from the local ethics committee. Women who agreed to participate gave their signed informed consent after explanation of the trial benefits and hazards. All procedures were carried out in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Declaration of Helsinki and its later amendments or comparable ethical standards. The trial was registered with local ethics committee of the Faculty of Medicine, Helwan University.

All patients were placed supine on the operating table. Regional anesthesia was administered, typically an interscalene block or general anesthesia. Once adequate anesthesia was given, an examination under anesthesia was done to check the full range of motion. Stiffness may develop in patients with rotator cuff tears, making it important to document adequate range of motion before the start of the procedure. If the patient has a stiff shoulder, a manipulation under anesthesia is performed to release adhesions. The patient is then ready for positioning in the beach chair position.

In this position, the buttocks were the most dependent position, ensuring that the patient stabled and not slipped down the table. The surgeon had adequate access to the posterior shoulder to the medial border of the scapula and the anterior shoulder to the mid-clavicle. The head was held in place with a head holder

Figure 2



Position of the patient in semi sitting position.

(Fig. 2). The shoulder was prepared and draped as for arthroscopy. A complete diagnostic arthroscopy was done and the presence of a cuff tear was confirmed and associated pathology addressed. The lateral arthroscopy portal incision was then extended and the deltoid was split, exposing the cuff tear. The shoulder is draped with care taken to ensure exposure of the widest area especially posteriorly. A standard posterior portal is created. The correct position of this portal in the medial–lateral direction may be found by feeling the notch in the spine of the scapula usually about 2 cm medial to the posterolateral corner of the acromion. The portal is then placed about 2 cm inferior to this point (Fig. 3). A blunt trocar is used to penetrate the posterior capsule and a diagnostic arthroscopy is begun (Fig. 4).

A thorough diagnostic arthroscopy is then performed. The glenohumeral joint is examined for lesions and cartilage loss. The labrum is inspected and examined along with the bicep's tendon (Fig. 5). The defect in the cuff may be marked with a percutaneously placed spinal needle, especially for a partial-thickness tear. A suture is advanced through the needle into the glenohumeral joint, allowing easier identification during the subacromial portion of the procedure and during the open repair. The arthroscope is then removed from the glenohumeral joint (Fig. 6). Once in the subacromial space, an anterolateral portal is created 2 cm posterior

Figure 3




A standard posterior portal is created, about 2 cm medial to the posterolateral corner of the acromion.

Figure 4




A blunt trocar is used to penetrate the posterior capsule.

Figure 5



The glenohumeral joint with intact labrum.

Figure 6



The arthroscopic view of articular rotator cuff tear.

and inferior to the anterolateral border of the acromion. The position of this portal may be modified to center it over the rotator cuff tear, which may be facilitated by the previously placed marking needle (Fig. 7).


There has been some controversy about the necessity of performing a subacromial decompression in the presence of a rotator cuff tear. In this study we routinely perform a subacromial decompression, and perform an acromioplasty before cuff repair. Debridement is done to the edges of the tear in the subacromial space. Burssectomy, especially laterally, will facilitate visualization during the procedure. The tendon edges are lightly debrided. At this point, needle may be placed through the edge of the cuff and attention turned to exposure of the rotator cuff. A 3- to

Figure 7



Marking needle in the rotator cuff tear.

Figure 8




Skin incision of mini open procedure.

4-cm skin incision was made from the anterolateral edge of the acromion distally, and dissection was made to the raphe between the anterior and middle deltoid (Fig. 8).

A stay suture was placed distally to prevent propagation of the deltoid split and potential injury to the axillary nerve (Fig. 9). Once the deltoid is split, the subacromial space is entered. Blunt self-retaining retractors may be helpful to hold the fibers of the deltoid apart, but care should be taken to avoid excess pressure and deltoid necrosis (Fig. 10). After assessing the adequacy of the acromioplasty by direct digital palpation, we placed a deltoid retractor for direct visualization of the rotator cuff and humeral head (Fig. 11). As the torn tendon was tagged by traction sutures after removing the hypertrophic bursal tissue around the split site to improve

Figure 9



Stay suture was placed distally to prevent propagation of the deltoid split.

Figure 10



Blunt split of the deltoid.

visualization (Fig. 12), we confirmed involvement and configuration of the torn tendon by rotating the arm and attempted anatomical reduction on the footprint of the greater tuberosity. After preparing the footprint using a ring curette or rasp (Fig. 13), the torn tendon was repaired by a single-row technique using suture

Figure 11



Retracted tendon of rotator cuff.


Figure 12



Removing the hypertrophic bursa.


anchors (Fig. 14). If pathology of the long head of the biceps tendon was found, tenodesis was performed under direct visualization. Appropriate rotation of the arm is the key to positioning the cuff tear underneath the deltoid split. By varying the position of the arm, different parts of the tendon can be brought into view.

Figure 13



Preparing the footprint of greater tuberosity using a ring curette or rasp.

Figure 14




The torn tendon was repaired by single row technique.

If the tear is massive, one or multiple traction sutures using simple stitches can be placed to help mobilize the cuff and allow easier repair (Fig. 15).

Traction on the tendon allows easy release and better excursion of the tendon. Extra-articular adhesions are released, allowing full mobilization of the tendon. The

Figure 15



Traction sutures.

Figure 16




Cuff sutures.

goal is to repair the tendon to bone with no tension while the arm is at the side. Intra-articular adhesions deep into the cuff as well as the coracohumeral ligament may need to be addressed case by case and they can be transected as necessary. Once adequate mobilization is obtained, the size and shape of the tear are again evaluated. U-shaped tears can be repaired with a combination of side-to-side sutures and bone fixation, while crescent-shaped tears are generally repaired directly to the bone. Once side-to-side sutures are placed, a smaller cuff edge will be attached to the bone (Fig. 16). Bony fixation can be accomplished either through trans-osseous tunnels or anchors. Anchors are placed in the 'footprint' of the cuff or anatomic insertion, and their position is chosen to allow an even repair of the tendon edge without excessive tension on one portion of the cuff (Fig. 17). Once the cuff has been repaired, the shoulder range of motion is checked to demonstrate the safe range for rehabilitation (Fig. 18). The wound is thoroughly irrigated and the deltoid fascia is meticulously repaired. A subcutaneous and subcuticular closure is performed and dressings are applied.

All the 20 patients are discharged on the same day of surgery; they received the same postoperative analgesia and followed up for constant–Murley sore preoperative,

Figure 17



Anchors are placed in the "footprint" of the cuff or anatomic insertion.

Figure 18



The cuff has been repaired.

1-month postoperative and 3 months postoperative. The cuff is repaired with trans-osseous sutures securely down to the bone. The wound is thoroughly irrigated and the deltoid fascia is meticulously repaired. A subcutaneous and subcuticular closure is performed and dressings are applied.

Postoperative protocol

The patient is discharged from the hospital on the day of surgery. The patient is placed in a sling and is allowed out of the sling only for physical therapy and exercises. Passive range-of-motion exercises begin; forward elevation, external rotation, and pendulum exercises are started. Internal rotation is not allowed until healing of the cuff is completed. Elbow and hand exercises are also begun. The patient performs pendulum, elbow, and wrist exercises at home several times a day, while passive motion exercises are performed either at home or with physical therapy many times a week. The goal of early rehabilitation is to minimize stiffness without putting tension on the cuff repair. At 6 weeks postoperatively, the sling is removed and active-assisted range-of-motion exercises are added.

Strengthening exercises are begun at 6–8 weeks postoperatively, depending on the size of the tear. A strengthening and stretching program was continued until 6–8 months postoperatively.

Modified University of California, Los Angeles shoulder scale

The patients were assessed postoperatively with the University of California, Los Angeles (UCLA) score at an average of 2 years, 7 years, and 15 years after surgery. The questionnaire was adapted for self-assessment, and we have previously described the reliability of the UCLA score as a self-assessment tool following rotator cuff surgery. In this scoring system, a maximum of 35 points is possible. A maximum 10 points is allocated for pain, 10 for function, 5 for active range of motion, 5 for strength of forward flexion, and 5 for overall satisfaction with the operation. Satisfaction with the operation is scored as 5 for satisfied or better and as 0 for dissatisfied or worse. This satisfaction score was used in the final analysis of overall satisfaction. The outcome results are categorically reported as excellent (34–35 points), good (28–33 points), fair (21–27 points), or poor (>20 points).

Outcome of the study

Three cases suffered from stiffness that was dealt with by manipulation under excellent results. In three cases of superficial wound infection, regular dressing was done for 3 weeks.

Statistical analysis

Results were tabulated and statistically analyzed using SPSS V.25 program and two types of statistics were done: descriptive statistics that includes description of data was in the form of mean±SD for quantitative data, and frequency and proportion for qualitative data and analytical statistics that includes standard Student's *t* test, Fisher's exact test, and χ^2 test and Pearson correlation (*r*). *P* value less than 0.05 was considered statistically significant.

Results

A total of 20 patients, 11 (55%) males and nine (45%) females, were included in this study with a mean age of 58.00 ± 9.17 years. Most of the patients (80%) had their right side affected and the affected shoulder was dominant. The mean operative time was 69.95 ± 14.32 min (Table 1).

There was high statistically significant gradual improvement in modified UCLA shoulder scale among the studied patients at 1, 3, and 6 months postoperative follow-up compared with preoperative ($P < 0.001$) (Table 2).

Postoperative complications were found in six patients. Three (15%) cases had stiffness, and the other three (15%) cases had superficial wound infection (Table 3).

Age, sex, operative time, modified UCLA shoulder scale preoperative and postoperative, and postoperative complications did not show any significant relations with dominant sides of affected shoulder ($P>0.05$) (Table 4).

Age, operative time, modified UCLA shoulder scale preoperative and postoperative and postoperative complications did not show any significant relations with sex ($P>0.05$) (Table 5).

There were significant correlations between modified UCLA shoulder scale with age, operative time, and postoperative complications ($P<0.05$), while there were no significant correlations with sex or side ($P<0.05$) (Table 6).

Discussion

Rotator cuff tears are a common pathology associated with degenerative changes in the shoulder joint. They cause significant disability, pain, and poor health status and their prevalence is increasing within an aging population. Many studies have documented excellent outcomes following rotator cuff repair surgery; however, complications can occur [20]. In the current study on a total of 20 patients, 11 (55%) males and nine (45%) females were included with a mean age of 58.00 ± 9.17 years. Most of the patients

Table 1 Descriptive data of the studied patients ($N=20$)

Descriptive data	Median	Minimum–maximum	Mean \pm SD
Age (years)	42.00	26.00–68.00	58.00 \pm 9.17
	<i>n</i>	%	
Sex			
Males	11	55	
Females	9	45	
Side			
Right	16	80.0	
Left	4	20.0	
Dominance of affected shoulder			
Dominant	16	80.0	
Nondominant	4	20.0	
Operative time (min)	50.00	50.00–100.00	69.95 \pm 14.32

Table 2 Modified University of California, Los Angeles shoulder scale of the studied patients ($N=20$)

UCLA shoulder scale	Preoperative	Postoperative follow-up [<i>n</i> (%)]			Total
		1 month	3 months	6 months	
Poor (< 21)	15 (75.0)	1 (5.0)	0	0	16 (20.0)
Fair (22–27)	5 (25.0)	17 (85.0)	10 (50.0)	0	32 (40.0)
Good (28–33)	0	2 (10.0)	10 (50.0)	14 (70.0)	26 (32.5)
Excellent (34–35)	0	0	0	6 (30.0)	6 (7.5)
<i>P</i> value [#]	–	<0.001**	<0.001**	<0.001**	–

UCLA, the University of California, Los Angeles score.

[#]*P* value compared 1, 3, 6-month follow-up versus preoperative using χ^2 test.

**No significance.

(80%) had their right side affected and the affected shoulder was dominant. The mean operative time was 69.95 ± 14.32 min. In agreement with us Shinnars *et al.* [21] found that the average age of the study group was 51 years (range, 31–72 years) at the time of surgery. There were 29 men and 11 women. The dominant shoulder was involved in 28 (68%) of the 41 cases. Also, Eid *et al.* [22] studied eight (66.7%) males and four (33.3%) females with an average age of 52.3 ± 2.6 years (range, 47–62 years), all (100%). The right shoulder was involved in 12 (100%) patients; also, the dominant side was involved in 12 (100%) shoulders. In addition, Kelly *et al.* [23] reported a response rate of 81%. Of those who completed follow-up, 122 (84.7%) patients were right handed and 22 (15.3%) patients were left handed. Overall, 92 (63.9%) patients had surgery on their dominant side, while 52 (36.1%) patients had surgery on their nondominant side. No participant had bilateral rotator cuff repair within the study period.

In this study, there was high statistically significant gradual improvement in modified UCLA shoulder scale among the studied patients at 1, 3, and 6 months postoperative follow-up compared with preoperative. In agreement, Eid *et al.* [22] revealed that using the UCLA scoring system, the final assessment (at a mean of 27.4 months postoperatively; range, 25–42 months) revealed poor results in one (8.3%), good results in seven (58.3%), and excellent results in four (33.4%) patients. As a result, the final overall results were satisfactory (good and excellent) in 11 (91.7%) and unsatisfactory (poor) in one (8.3%) patient. Also, the mean value of overall UCLA score, pain score, function score, active forward flexion score, and strength of active forward flexion score significantly improved from 8.8 ± 1.2 , 2.1 ± 0.5 , 1.8 ± 2.1 , 2.4 ± 1.1 , and 2.3 ± 2.3 preoperatively to 32.4 ± 2.4 , 9.3 ± 2.8 , 9.1 ± 4.1 , 4.6 ± 3.2 , and 4.4 ± 1.6 postoperatively ($P<0.05$), respectively. Also, Levy *et al.*

Table 3 Postoperative complications among the studied patients ($N=20$)

Postoperative complications	<i>n</i>	%
Stiffness 3 cases	3	15.0
Superficial wound infection	3	15.0

Table 4 Relation between dominance sides of affected shoulder with demographic data, University of California, Los Angeles shoulder scale preoperation and postoperation and postoperative complications

	Dominance of affected shoulder [n (%)]				FET	P value	95% CI
	Dominant side (right) (N=16)		Nondominant side (left) (N=4)				
Age (year)							
Mean±SD	57.75±9.73		59.0±7.62		U=0.277	0.792	-12.42 to 9.92
Sex							
Male	8	50.0	3	75.0	0.768	0.381	-
Female	8	50.0	1	25.0			
Operative time (min)							
Mean±SD	69.813±14.33		70.50±16.53		U=0.076	0.943	-25.22 to 23.84
Modified UCLA shoulder scale preoperative							
Poor (<21)	13	81.25	2	50.0	1.58	0.208	-
Fair (22-27)	3	18.75	2	50.0			
After 1 month							
Poor (< 21)	1	6.25	0	0.0			
Fair (22-27)	14	87.5	3	75.0	1.28	0.256	-
Good (23-33)	1	6.25	1	25.0			
After 3 months							
Fair (22-27)	8	50.0	2	50.0	0.00	1.00	-
Good (23-33)	8	50.0	2	50.0			
After 6 months							
Good (23-33)	10	62.5	4	100.0	2.036	0.154	-
Excellent (34-35)	6	37.5	0	0.0			
Postoperative complications							
Stiffness	2	12.5	1	25.0	0.00	1.00	-
Superficial wound infection	2	12.5	1	25.0			

CI, confidence intervals; FET, Fisher exact test; U, Mann-Whitney test; UCLA, the University of California, Los Angeles score.

Table 5 Relation between sex and demographic data, University of California, Los Angeles shoulder scale preoperation and postoperation and postoperative complications

	Sex [n (%)]				χ^2	P value	95% CI
	Males (N=11)		Females (N=9)				
Age (year)							
Mean±SD	58.91±5.59		56.89±12.57		t=0.447	0.664	-7.97 to 12.01
Operative time (min)							
Mean±SD	71.64±17.57		67.89±10.13		t=0.602	0.555	-9.42 to 16.92
Modified UCLA shoulder scale preoperative							
Poor (< 21)	9	81.82	6	66.67	0.606	0.436	-
Fair (22-27)	2	18.18	3	33.33			
After 1 month							
Poor (< 21)	1	9.09	0	0.0			
Fair (22-27)	8	72.73	9	100.0	2.88	0.236	-
Good (23-33)	2	18.18	0	0.0			
After 3 months							
Fair (22-27)	5	45.45	5	55.56	0.202	0.653	-
Good (23-33)	6	54.55	4	44.44			
After 6 months							
Good (23-33)	9	81.82	5	55.56	1.626	0.202	-
Excellent (34-35)	2	18.18	4	44.44			
Postoperative complications							
Stiffness	2	18.18	1	11.11	0.667	0.414	-
Superficial wound infection	1	9.09	2	22.22			

χ^2 , χ^2 test; CI, confidence intervals; t, independent t test; UCLA, the University of California, Los Angeles score.

Table 6 Modified University of California, Los Angeles shoulder scale in relation to clinical data of the studied patients

	Modified UCLA shoulder scale	
	<i>r</i>	<i>P</i>
Sex	0.174	0.463
Age (year)	0.392*	0.008
Side	0.289	0.217
Operative time (min)	0.652**	0.002
Postoperative complications	0.707**	0.001

r, correlation coefficient; UCLA, the University of California, Los Angeles score.

*Significant.

**No significance.

[24] reported that 80% of 25 patients ($N=20$) who underwent arthroscopic-assisted, mini-open rotator cuff repair and were monitored for a minimum of 1 year had good or excellent objective clinical results and 96% ($N=24$) were subjectively satisfied. Shinnors *et al.* [21] suggested that an arthroscopic-assisted mini-open technique of rotator cuff repair is an excellent alternative to standard open techniques.

On the contrary, Kang *et al.* [18] reported, in a retrospective study of 63 patients treated with mini-open rotator cuff repair and 65 treated with arthroscopic rotator cuff repair, no statistically significant improvements at 6 months in SF-36 general health, role-emotional, and mental health. Also, Pearsall *et al.* [25] reported that although there was a significant improvement in clinical outcome from preoperative (UCLA score) to the latest follow-up, the SF-36 was not significantly different postoperatively. These different findings may be related to the size of the tear which seems to be a determining factor in the functional outcome. Small and medium tears did better than large tears.

Regarding postoperative complications, the present study found postoperative complications in six patients. Three (15%) cases had stiffness, and the other three (15%) cases had superficial wound infection. In this line, Eid *et al.* [22] reported the postoperative complications included scar at the site of deltoid-split approach in one (8.3%) patient, superficial infection in one (8.3%) patient managed by antibiotics and regular dressings, and finally, postoperative shoulder stiffness in one (8.3%) patient. Also, Blevins *et al.* [26] found one of the three patients who required further surgery required a revision cuff repair (following a fall onto the operated shoulder), and two required revision arthroscopic subacromial decompression of soft tissue. All three did well following the second procedure. One additional patient complained of impingement symptoms but declined further surgery. This patient responded moderately well to two subacromial steroid injections.

Age, operative time, modified UCLA shoulder scale preoperative and postoperative, and postoperative complications did not show any significant relations with sex or dominance sides of affected shoulder. In agreement, Kelly *et al.* [23] found a mean age of 63 ± 10.1 years in the dominant group and 62 ± 8.6 years in the nondominant group. There were 48 females and 44 males in the dominant group, with 27 females and 25 males in the nondominant group. The mean overall outcome score was marginally higher in the dominant surgery group with a mean of 89.8 ± 14.2 compared with a mean of 87.4 ± 17.5 in the nondominant group. Multivariate analysis including age, sex, tear location, tear retraction, assessment to surgery time, and surgery to follow up time as individual input variables revealed this difference to be nonsignificant ($P=0.4$). They found no difference in patient-reported outcome measures between dominant and nondominant hand rotator cuff repair at a 3-year follow-up. Also, Eid *et al.* [22] found no statistically significant difference in the final results; among the different age groups (as categorized into decades) ($P>0.05$), between the male and female patient groups ($P>0.05$), among the patient groups of different durations of preoperative complaint (as divided into 6-month intervals) ($P>0.05$), and also among the patient groups of different mechanisms of injury ($P>0.05$).

On the other hand, Kelly *et al.* [23] showed that hand dominance was significantly associated with the side of rotator cuff tear ($P=0.005$). Also, Sahni and Narang [27] found that factors such as age, sex, and time from tear to surgery are more consistently cited as having an effect on outcomes. The different results could be explained by the small number of patients included in the different groups of age, sex, duration of preoperative complaint, and mechanism of injury.

The present findings showed significant correlations between modified UCLA shoulder scale with age, operative time, and postoperative complications, while there were no significant correlations with sex or side. This is consistent with Kelly *et al.* [23] as univariate analysis found no effect of age, sex, tear location, retraction, assessment to surgery time, or assessment to follow-up time on outcome. Sex did have a significant effect on outcome score ($P=0.03$). Also, Wolf *et al.* [28] and Montgomery *et al.* [29] compared results of 50 patients (average age, 58) with open repairs with those of 38 patients (average age, 66) with arthroscopic decompression alone at an average 2–5-year follow-up times and found 78 versus 61% satisfactory results. No correlation was identified among the size of tear, patient age or activity level, and results achieved with arthroscopic decompression. As well, Ogilvie-Harris and Demaziere [30] prospectively studied 45 patients with arthroscopic subacromial decompression versus open rotator cuff repair

and found pain relief with both, but better functional scores with cuff repair, although recovery was longer.

A possible explanation is that as these patients become older, their functional and physical demands decrease, leading to a result perceived by the patient as a satisfactory outcome. Older patients may be able to compensate satisfactorily in the presence of a recurrent or persistent defect in the rotator cuff.

Limitations

A limitation of this study is the short follow-up period of our patients. A second weakness of the study is the lack of imaging to determine cuff integrity, which could provide a better understanding of the natural history of rotator cuff repairs.

Summary

Arthroscopic-assisted mini-open rotator cuff repair is a successful procedure with multiple published studies demonstrating a high proportion of good-to-excellent results using well-validated outcome measures. The technique is not technically demanding than all-arthroscopic repair while still having the advantages of arthroscopic repair. These include the ability to perform diagnostic arthroscopy, preservation of the origin of the deltoid, less postoperative pain, rapid hospital discharge, and accelerated rehabilitation. Mini-open repair seems to be equivalent to all-arthroscopic repair in multiple nonrandomized comparative studies, and a randomized trial is underway to formally address this question. Arthroscopic-assisted mini-open repair remains a viable option for the surgeon who wishes to use the classic surgical suture passing and tying while also taking advantage of arthroscopic examination and treatment of the shoulder joint. The arthroscopic-assisted mini-open approach remains a useful tool for rotator cuff repair.

Conclusions

Our results suggest that the arthroscopically assisted mini-open technique for rotator cuff repair is an excellent approach for the repair of a torn rotator cuff. Surgery for rotator cuff tears improves self-reported patient outcomes and has a lasting and durable result at an average of 15 years after surgery. There was high statistically significant gradual improvement in modified UCLA shoulder scale among the studied patients at 1, 3, and 6 months postoperative follow-up compared with preoperative. Further studies with a large sample are required to enhance the current findings.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

References

- 1 Malik S, Aslam S, Soomro SK, Malik SZ, Rauf D. Effectiveness of graded motor imagery in pain and disability after rotator cuff injury. *Rehabilit J* 2018; 2:75–82.
- 2 Fahy K, Galvin R, Lewis J, McCreesh K. Large to massive rotator cuff tendon tears: a protocol for a systematic review investigating the effectiveness of exercise therapy on pain, disability and quality of life. *HRB Open Res* 2021; 4:75.
- 3 Barcia AM, Makovicka JL, Spenciner DB, Chamberlain AM, Jacofsky MC, Gabriel SM, *et al.* Scapular motion in the presence of rotator cuff tears: a systematic review. *J Shoulder Elbow Surg* 2021; 30:1679–1692.
- 4 Nyffeler RW, Schenk N, Bissig P. Can a simple fall cause a rotator cuff tear? Literature review and biomechanical considerations. *Int Orthop* 2021; 45:1573–1582.
- 5 Cunningham G, Nicodème-Paulin E, Smith MM, Holzer N, Cass B, Young AA. The greater tuberosity angle: a new predictor for rotator cuff tear. *J Shoulder Elbow Surg* 2018; 27:1415–1421.
- 6 Mihata T, Lee TQ, Hasegawa A, Kawakami T, Fukunishi K, Fujisawa Y, *et al.* Arthroscopic superior capsule reconstruction can eliminate pseudoparalysis in patients with irreparable rotator cuff tears. *Am J Sports Med* 2018; 46:2707–2716.
- 7 Zoga AC, Kamel SI, Hynes JP, Kavanagh EC, O'Connor PJ, Forster BB. The Evolving roles of MRI and ultrasound in first-line imaging of rotator cuff injuries. *Am J Roentgenol* 2021; 217:1–11.
- 8 Quillen DA, Wuchner M, Hatch RL. Acute shoulder injuries. *Am Fam Physician* 2004; 70:1947–1954.
- 9 Narvani AA, Imam MA, Godenèche A, Calvo E, Corbett S, Wallace AL, Itoi E. Degenerative rotator cuff tear, repair or not repair? A review of current evidence. *Ann R Coll Surg Engl* 2020; 102:248–255.
- 10 Ellman H, Kay SP, Wirth M. Arthroscopic treatment of full-thickness rotator cuff tears: 2- to 7-year follow-up stud. *Arthroscopy* 1993; 9:195–200.
- 11 Jo CH, Kim JE, Yoon KS, Lee JH, Kang SB, Lee JH, *et al.* Does platelet-rich plasma accelerate recovery after rotator cuff repair? A prospective cohort studies. *Am J Sports Med* 2011; 39:2082–2090.
- 12 Codman EA. Complete rupture of the supraspinatus tendon: operative treatment with report of two successful cases. *Boston Med Surg J* 1911; 164:708–710.
- 13 Strauss EJ, Salata MJ, Kercher J, Barker JU, McGill K, BachJrBR, *et al.* The arthroscopic management of partial-thickness rotator cuff tears: a systematic review of the literature. *Arthroscopy* 2011; 27:568–580.
- 14 Chahal J, Mall N, MacDonald PB, Van Thiel G, Cole BJ, Romeo AA, Verma NN. The role of subacromial decompression in patients undergoing arthroscopic repair of full-thickness tears of the rotator cuff: a systematic review and meta-analysis. *Arthroscopy* 2012; 28:720–727.
- 15 Frisella WA, Cuomo F. Mini-open rotator cuff repair. In: DeLee & Drez's, editor. *Minimally invasive surgery in orthopedics*. New York, NY: Springer; 2010. pp. 27–33.
- 16 Fealy S, Adler RS, Drakos MC, Kelly AM, Allen AA, Cordasco FA, *et al.* Patterns of vascular and anatomical response after rotator cuff repair. *Am J Sports Med* 2006; 34:120–127.
- 17 Yamaguchi K, Levine WN, Marra G, Galatz LM, Klepps S, Flatow E. Transitioning to arthroscopic rotator cuff repair: the pros and cons. *J Bone Joint Surg* 2003; 85:144–155.
- 18 Kang L, Henn RF, Tashjian RZ, Green A. Early outcome of arthroscopic rotator cuff repair: a matched comparison with mini-open rotator cuff repair. *Arthroscopy* 2007; 23:573–582.
- 19 Severud EL, Ruotolo C, Abbott DD, Nottage WM. All-arthroscopic versus mini-open rotator cuff repair: a long-term retrospective outcome comparison. *Arthroscopy* 2003; 19:234–238.
- 20 Kukkonen J, Joukainen A, Lehtinen J, Mattila KT, Tuominen EK, Kauko T, Äärämaa V. Treatment of nontraumatic rotator cuff tears: a randomized controlled trial with two years of clinical and imaging follow-up. *J Bone Joint Surg* 2015; 97:1729–1737.
- 21 Shinnars TJ, Noordsij PG, Orwin JF. Arthroscopically assisted mini-open rotator cuff repair. *Arthroscopy* 2002; 18:21–26.
- 22 Eid TA, Hannout YS, El-Sayed AS, Morsi ES. Arthroscopically assisted repair of massive full-thickness rotator cuff tears: an analysis of 2-year postoperative follow-up. *Egypt Orthop J* 2018; 53:77.

- 23 Kelly MA, Mc Donald CK, Boland A, Groarke PJ, Kaar K. The effect of hand dominance on functional outcome following single row rotator cuff repair. *Open Orthop J* 2017; 11:562.
- 24 Levy HJ, Uribe JW, Delaney LG. Arthroscopic assisted rotator cuff repair: preliminary results. *Arthroscopy* 1990; 6:55–60.
- 25 Pearsall AW, Ibrahim KA, Madanagopal SG. The results of arthroscopic versus mini-open repair for rotator cuff tears at mid-term follow-up. *J Orthop Surg Res* 2007; 2:1–8.
- 26 Blevins FT, Warren RF, Cavo C, Altchek DW, Dines D, Palletta G, Wickiewicz TL. Arthroscopic assisted rotator cuff repair: results using a mini-open deltoid splitting approach. *Arthroscopy* 1996; 12:50–59.
- 27 Sahni V, Narang AM. Risk factors for poor outcome following surgical treatment for rotator cuff tear. *J Orthop Surg* 2016; 24: 265–268.
- 28 Wolf EM, Pennington WT, Agrawal V. Arthroscopic rotator cuff repair: 4-to 10-year results. *Arthroscopy* 2004; 20:5–12.
- 29 Montgomery TJ, Yerger B, Savoie III FH. Management of rotator cuff tears: a comparison of arthroscopic debridement and surgical repair. *J Shoulder Elbow Surg* 1994; 3:70–78.
- 30 Ogilvie-Harris DJ, Demaziere A. Arthroscopic debridement versus open repair for rotator cuff tears. A prospective cohort studies. *J Bone Joint Surg Br* 1993; 75:416–420.