



Menoufia Veterinary Medical
Journal
(MVMJ)

Menoufia Veterinary Medical Journal

<https://vmmj.journals.ekb.eg>

MVMJ. Vol. 1 (1)



Heavy metals (Hg, Pb and Cd) assessment in Nile Fish *Oreochromis niloticus* and *Clarias lazera*

Elkady¹, S. Elbagory¹, A. Bazh², E. Shawish³, R.

¹ Food Hygiene Department, Faculty of Veterinary Med., Menoufia University, ² Parasitology Department, Faculty of Veterinary Med., Menoufia University, ³ Food Hygiene Department, Faculty of Veterinary Med., Sadat University,

ABSTRACT

Key words:

Heavy metal residues,
Oreochromis niloticus,
Clarias lazera, Acetic
acid.

*Correspondence

Vetsally99@gmail.co
m

Article History

Received: 11 Aug
2024.

Accepted: 9 sep 2024

Heavy metals are one of the most important pollutants that affect the aquatic environment and fish. The present study was carried out on 40 random samples of Nile fish represented by *Oreochromis niloticus* and *Clarias lazera* (20 of each) were collected from several fish markets in Shibin El-kom, El-Menoufia, Egypt. These samples were promptly assigned to the laboratory for analysis of their mercury (Hg), lead (Pb), and cadmium (Cd) content. The mean contents of metal, expressed in mg/kg wet weight, varied from 0.46 ± 0.05 in *Oreochromis niloticus* and 0.69 ± 0.07 in *Clarias lazera* for Hg, 0.29 ± 0.03 in *Oreochromis niloticus* and 0.44 ± 0.05 in *Clarias lazera* for Pb and 0.13 ± 0.01 in *Oreochromis niloticus* and 0.21 ± 0.01 in *Clarias lazera* for Cd. The permissible limit by E.O.S in mg/kg for Hg is 0.50, for Pb is 0.30 and for Cd is 0.10. Treatment of fish fillet by different acetic acid concentrations (1%), (3%), (5%) for 5 min; resulted in noticeable reduction of heavy metals by percentage in treated group fish compared with the untreated group, mercury reduced by 8.1, 17.4 and 32.6, respectively, while lead reduced by 12.3, 26.2 and 38.5, respectively, and finally cadmium reduced by 17.3, 28.9 and 46.2, respectively. The public health importance of heavy metals was discussed.

1. INTRODUCTION

Fish play a crucial role in aquatic ecosystems and serve as a vital source of animal protein production in Egypt, helping to meet the growing demand for protein [1] and in Africa it is one of the cheapest sources of animal protein [2]. Fish protein is highly preferred in the human diet due to its low cholesterol levels. Fish and its products are rich in high-quality proteins, essential vitamins, minerals and fatty acids, making them crucial sources of nutrition. [3]. The risks associated with heavy metal residues and their effects on public health are a critical concern for those in the field of food hygiene. The main source of toxic heavy metals that enter the aquatic system are anthropogenic impacts including industrial discharge, non-point source runoff, domestic sewage and

atmospheric precipitation [4]. However, metals can also naturally occur in small quantities and enter aquatic systems via ore-bearing rocks, wind-blown dust, forest fires, and vegetation [5]. Metals that are absorbed readily and eliminated slowly in fish tissues tend to accumulate at higher levels [6]. High concentrations of mercury have been identified in certain fish species [7]. The health risks of mercury may encompass lung damage, neurological and psychological disorders, kidney damage and congenital abnormalities [8]. Lead (Pb) can enter water systems from various sources, including industrial and smelter emissions, the breakdown of old lead plumbing, pesticides containing lead, precipitation, lead dust deposition, street runoff and municipal wastewater [9]. When ingested, lead can accumulate in the human body, substituting

calcium in bones, which can lead to numerous health issues. These issues may include reduced intellectual abilities, kidney failure, miscarriage, high blood pressure, lung cancer and even death. [10]. The kidneys are significantly affected by prolonged exposure to cadmium, along with liver toxicity, skeletal damage, and neurological effects [11]. Long-term exposure to cadmium is linked to numerous diseases, such as heart disease, anemia, weakened bones, reduced immune system response, and disorders of the kidneys and liver [12]. Acetic acid is a weak organic acid that can lower the pH of fish tissue and form complexes with metal ions, thus reducing their solubility and bio-accessibility [13]. The reduction of metal loads increases as the acid concentration rises and the immersion period [14]. The decrease in heavy metals in fish treated with acetic acid may result from the formation of insoluble acetate salts of these metals [15]. So, the aim of the study is to determine the heavy metals content in *Oreochromis niloticus* and *Clarias lazera* and comparing with the stipulated permissible limits by [16]. Additionally, the beneficial effect of acetic acid in reducing the levels of these heavy metals in edible fish tissue was studied.

2. MATERIALS AND METHODS

3. 2.1 Sampling:

A total of 40 random samples of *Oreochromis niloticus* and *Clarias lazera* (20 of each) were compiled from various fish markets in Shibin Elkom. The collected samples were then directly transferred to the laboratory to determine their contents of mercury, lead, and cadmium as heavy metal residues.

2.2. Detection of heavy metals:

All collected samples were analyzed to determine the levels of heavy metals (mercury, lead, and cadmium) based on wet weight (mg/Kg).

2.2.1 Washing procedures according to AOAC(17).

2.2.2. Digestion technique [18].

2.2.3. Preparation of blank and standard solutions [19].

3: - Detection and analysis:

By using AAS (ATOMIC ABSORPTION SPECTROPHOTOMETER, VARIAN, Australia, model AA240 FS) which was adjusted at 217.0 nm. for lead, 228.8 nm. for cadmium and 253.7 nm. for mercury

4:- Quantitative detection of heavy metals:

Examined samples were calculated on wet weight according to the following equation:- $C = R \times (D/W)$. Where, C = Concentrations of element (mg/ kg) wet weight, R = reading of digital scale of AAS, D = Dilution of prepared sample, W= Weight of the sample.

5:- Experimental part :-

Effect of different concentrations of acetic acid (1%), (3%) and (5%) for 5 min to control heavy metal residues in fish fillet (Atta et al., 2015).

6:-Statistical Analysis

The results obtained from the examined fish samples were statistically analyzed using the methods (ANOVA) suggested by [41]

3.RESULTS

The results presented in table (1) showed that the mean mercury concentration in the examined samples was 0.53 ± 0.08 mg/kg for *Oreochromis niloticus* and 0.70 ± 0.09 mg/kg for *Clarias lazera*. Additionally, 40% of *Oreochromis niloticus* and 65% of *Clarias lazera* samples exceeded the maximum permissible limit (0.5 mg/kg), according to [16]. Table (2) indicated that the mean lead concentration was 0.40 ± 0.06 mg/kg for *Oreochromis niloticus* and 0.52 ± 0.06 mg/kg for *Clarias lazera*. Based on the permissible limit (0.30 mg/kg) stipulated by [16], 50% of *Oreochromis niloticus* and 70% of *Clarias lazera* samples were found unacceptable. Table (3) revealed that the mean cadmium concentration in the examined samples was 0.08 ± 0.02 mg/kg for *Oreochromis niloticus* and 0.15 ± 0.03 mg/kg for *Clarias lazera*. Furthermore, 30% of *Oreochromis niloticus* and 50% of *Clarias lazera* samples were deemed unfit for human consumption due to their exceeding content on the maximum permissible limit (0.1 mg/kg), according to [16]. After immersing fish fillet in different concentrations of acetic acid for 5 minutes, table (4), revealed that mercury level reduced from 0.86 ± 0.06 in controlled samples to 0.79 ± 0.04 (8.1%) by using 1% acetic acid, while with 3% acetic acid it reduced to 0.71 ± 0.04 (17.4%), and with 5% acetic acid it reduced to 0.58 ± 0.03 (32.6%). Meanwhile, lead concentrations reduced from 0.65 ± 0.05 in controlled samples to 0.57 ± 0.04 (12.3%) by using 1% acetic acid, while with 3% acetic acid it reduced to 0.48 ± 0.04 (26.2%) , and with 5% acetic acid it reduced to 0.40 ± 0.03 (38.5%), the cadmium level reduced from 0.52 ± 0.04 in controlled samples to 0.43 ± 0.03 (17.3%) by using 1% acetic acid, while with 3%

acetic acid it reduced to 0.37 ± 0.02 (28.9%), and (46.2%).
 finally with 5% acetic acid it reduced to 0.28 ± 0.02

Table (1): - Statistical analytical results of mercury concentration (mg/kg) in the examined samples of Nile fish and their acceptability according to their mercury content (n=20).

Fish species	Mean $\pm$ SE*	MRL mg/kg*	Accepted samples		Unaccepted samples	
			No	%	No	%
<i>Oreochromis niloticus</i>	0.46 ± 0.05	0.50	13	65	7	35
<i>Clarias lazera</i>	0.69 ± 0.07	0.50	11	55	9	45

S.E* = standard error of mean

* Maximum Residual Limit stipulated by Egyptian Organization for Standardization [16].

Table (2): - Statistical analytical results of lead concentration (mg/kg) in the examined samples of Nile fish and their acceptability according to their lead content (n=20).

Fish species	Mean $\pm$ SE*	MRL mg/kg*	Accepted Samples		Unaccepted Samples	
			No	%	No	%
<i>Oreochromis niloticus</i>	0.29 ± 0.03	0.30	15	75	5	25
<i>Clarias lazera</i>	0.44 ± 0.05	0.30	13	65	7	35

S.E* = standard error of mean

* Maximum Residual Limit stipulated by Egyptian Organization for Standardization [16].

Table (3): - Statistical analytical results of cadmium concentration (mg/kg) in the examined samples of Nile fish and their acceptability according to their cadmium content (n=20).

Fish species	Mean $\pm$ SE*	MRL mg/kg*	Accepted Samples		Unaccepted Samples	
			No	%	No	%
<i>Oreochromis niloticus</i>	0.13 ± 0.01	0.10	16	80	4	20
<i>Clarias lazera</i>	0.21 ± 0.01	0.10	14	70	6	30

S.E* = standard error of mean

* Maximum Residual Limit stipulated by Egyptian Organization for Standardization [16].

Table (4): - Effect of immersing fish fillet in different concentrations of acetic acid for 5 minutes on mercury, lead and cadmium residues (n=5).

Treatment	Mercury levels mg/Kg	Reduction %	Lead levels mg/Kg	Reduction %	Cadmium levels mg/Kg	Reduction %
Untreated control	0.86 ± 0.06	-----	0.65 ± 0.05	-----	0.52 ± 0.04	-----
1% acetic acid	0.79 ± 0.04	8.1	0.57 ± 0.04	12.3	0.43 ± 0.03	17.3
3% acetic acid	0.71 ± 0.04	17.4	0.48 ± 0.04	26.2	0.37 ± 0.02	28.9
5% acetic acid	0.58 ± 0.03	32.6	0.40 ± 0.03	38.5	0.28 ± 0.02	46.2

4. DISCUSSION

Continuous exposure to high levels of these metals presents potential health hazards to livestock and wildlife, including fish, birds, mammals and even humans. Their tendency to bioaccumulate in various fish tissues leads to toxic effects [20,21]. It has been established that mercury, lead and cadmium are among the most toxic metals [22]. The results of this study indicated that the *Clarias lazera* samples had higher levels of mercury, lead, and cadmium compared to the *Oreochromis niloticus* samples. The obtained results of mercury level in table (1) were slightly agree with those recorded by [23] (0.49 ± 0.05) for small size *Oreochromis niloticus* and in small size *Clarias lazera* were (0.72 ± 0.04); also with [24], (0.46 ± 0.03) in *Oreochromis niloticus*, (0.52 ± 0.04) in *Clarias lazera*; and with [25], (0.57 ± 0.08) in *Oreochromis niloticus*; While higher results were recorded by [23], for large size *Oreochromis niloticus* were (0.81 ± 0.05) and for large size *Clarias lazera* were (0.98 ± 0.08); and obtained by [26], (1.29 ± 0.12) in *O. niloticus*; also recorded by [27], (0.89 ± 0.01) in *O. niloticus*; and by [28], (1.13 ± 0.01) in *O. niloticus*, (1.44 ± 0.03) in *C. lazera*; and recorded by [29], (0.94 ± 0.01) in *O. niloticus*, (1.28 ± 0.02) in *C. lazera*, but lower results were obtained by [30], (0.013 ± 0.001) in *Oreochromis niloticus* and (0.040 ± 0.001) in *Clarias lazera*; and finally recorded by (Nasser et al., 2016), (0.16 ± 0.02) in Menoufia governorate and (0.08 ± 0.04) in Kafr El- Shaikh for *O. niloticus*.

Mercury compounds pose significant health risks due to their high toxicity, fat solubility, and ability to accumulate in erythrocytes and the central nervous system (CNS) [32]. Exposure to mercury can manifest in various symptoms, including oral inflammation (redness of lips, tongue, and throat), tooth loss, and skin abnormalities such as swelling, redness, and pink-red fingertips. Neurological effects, such as irritability, are also commonly associated with mercury poisoning [33]. While mercury occurs naturally in the environment at low levels, industrial activities have been the primary source of mercury pollution [34].

Regarding to the lead level, Table (2), lower results were obtained by [23], (0.11 ± 0.01) mg/kg for small size *Oreochromis niloticus* and (0.16 ± 0.03) mg/kg for large size one and (0.15 ± 0.02) mg/kg for small size *Clarias lazera* and (0.27 ± 0.04) mg/kg for large size one; and by [24], (0.08 ± 0.02) in *Oreochromis niloticus*, (0.14 ± 0.02) in *Clarias lazera*, also [31], reported in Kafr El- Shaikh (0.08

0.03) in *O. niloticus*, while in Menoufia reported results were nearly similar (0.40 ± 0.05) in *O. niloticus*; also slightly same results were also obtained by [25], (0.36 ± 0.06) in *Tilapia niloticus*, while higher results were recorded by [35], (0.53 ± 0.06) in *O. niloticus*; and by [36], (0.49 ± 0.01) in *O. niloticus*; and finally by [37], (0.78 ± 0.01) and (0.61 ± 0.01) mg/kg in examined *C. lazera* and *O. niloticus*.

Lead is a toxic substance that accumulates in the body due to its slow elimination process. Common symptoms of lead poisoning include abdominal pain, colic, anemia, and encephalopathy. Additionally, lead is regarded as an immunosuppressive agent in humans [38]. On the other hand, lead induces renal tubular dysfunction, as evidenced by proteinuria or aminoaciduria in humans. Additionally, lead inhibits heme synthesis, resulting in weakened cell walls of red blood cells, which leads to anemia [39].

During the 20th century, cadmium production, consumption and emissions to the environment have increased dramatically due to its industrial uses, such as in batteries, electroplating, plastic stabilizers, and pigments. This increase has consequently led to the contamination of aquatic habitats [8]. Cadmium is a toxic element that can cause insomnia, nausea, testicular atrophy, prostate cancer and kidney failure [40]. The obtained results of cadmium levels in Table (3) were similar to those reported by [23], with (0.11 ± 0.01) (mg/kg) for small-sized *Clarias lazera* and lower results for larger-sized ones at 0.20 ± 0.03 (mg/kg). In *Oreochromis niloticus*, the cadmium concentration was (0.08 ± 0.01) (mg/kg) for small-sized fish, while larger-sized fish had a lower concentration of 0.12 ± 0.02 (mg/kg), and slightly the same as the result reported by [24], (0.14 ± 0.02) for *Clarias lazera* and (0.10 ± 0.01) for *Oreochromis niloticus*, but the results were lower than [28], (0.43 ± 0.01) and (0.26 ± 0.01) (mg/kg) in examined *C. lazera*, *O. niloticus*, respectively, and [36], (0.25 ± 0.01) in *Clarias lazera* and (0.17 ± 0.01) in *O. niloticus*.

Treatment of fish with different concentrations of acetic acid reduced the level of heavy metals content [15]. The obtained results of mercury after immersion in acetic acid in table (4) slightly agree with [15], as mercury reduced by 75% after immersion in 5% acetic acid for 15 min, and with [29] as the mercury level reduced by (52.8%) after immersion in 5% acetic acid for 30 min. Moreover, the results of lead level in table (4) slightly agree

with [15], as lead level reduced by 76% after immersion in acetic acid 5% for 15 min, also agree with [14], as lead level reduced by (31.5%) after immersion in acetic acid 5% for 3 min. Finally, the results of cadmium in table (4) agree with [15], as cadmium level reduced by (41.7%) after immersion in acetic acid 5% for 15 min, also [14], cadmium level reduced by (25%), and with [29], as cadmium level reduced by (43.5%).

5. CONCLUSIONS

Examined fish samples, somewhat, were contaminated with heavy metal residues exceeding the safe permissible limits for human consumption. Addition of acetic acid to fish aid in reducing the level of heavy metals content, so it is recommended to try to minimize the level of heavy metals in river Nile and the aquatic creatures to protect the human and fish.

6-REFERENCES

1. Clauca, I. J. and Ward, A. R. (1996): Post-harvest fisheries development: a guide to handling, preservation, processing and quality. 443 pp.
2. Metin, S.I., Erkan, N.; Varlik, C.; Ozden, C.; Baygar, T., Kalafatoglu, H., Gun, H. (2000): Influence of modified atmosphere packaging on shelf life of fish-salami. *Fleish Witchcraft Inc.* 4:49 - 51.
3. Langston, W. J.; Burt, G. R. And Pope, N. D. (1999): Bioavailability of metals in sediments of the dogger bank (Central North Sea): A mesocosm study. *Estuarine, Coastal and Shelf Science.* 48: 519-540.
4. Fernandez-Leborans, G. and Olalla-Herrero, Y. (2000): Toxicity and bioaccumulation of lead and cadmium in marine protozoa communities. *Ecotoxicol. and Environ. Saf.* 47: 266-276.
5. Kalay M, Canli M. (2000): Elimination of essential (Cu, Zn) and non-essential (Cd, Pb) metals from tissues of a freshwater fish *Oreochromis zilli*. *Turk J Zool* 24: 429-436.
6. Abrefah, R., Adotey, D. K., Mensimah, E., Ampomah-Amoako, E., Sogbadji, R., & Opat, N. (2011): Biomonitoring of Occupational Exposure to Total Arsenic and Total Mercury in Urine of Goldmine Workers in Southwestern Ghana. *Environmental Research, Engineering and Management*, 2(56), 43–48.
7. Järup, L. (2003): Hazards of heavy metal contamination. *British Medical Bulletin*, 68, 167–182.
8. Sepe A, Ciaralli L, Ciprotti M, Giordano R, Fumari E, Costantini S, (2003): Determination of cadmium, chromium, lead and vanadium in six fish species from the Adriatic Sea. *Food Addit. Contam.* 20: 543-552. 19.
9. Awuah, G.K. (2016): Assessment of heavy metal concentrations in sediment, water and fish from the ankobra and tano river basins in Ghana. Thesis (MSc). school of public health. University of Ghana.
10. Misra, B. R.; Crance, K. A.; Bare, R. M.; Waalkes, M. P. (1998): Lack of correlation between the inducibility of metallothionein mRNA and metallothionein protein in Cadmium-exposed rodents. *Toxicology*, 7/7: 99-109.
11. Codex Alimentarius Commission Procedural Manual. (2001): Maximum residual limits (MRLs) for pesticides. CAC / MRL, Rev. 1.
12. Chan WS, Routh J, Luo C, Dario M, Miao Y, Luo D, Wei L. (2021): Metal accumulations in aquatic organisms and health risks in an acid mine-affected site in South China. *Environmental Geochemistry and Health.* 43,11.:4415–4440.
13. Atta, M.; Salem, M.; Elsayed, E. and Badr, M. (2015): Effect of acetic acid and cooking process on chemical composition and metal contents of polluted grey mullet (*Mugil cephalus*). *J. Food and Dairy Sci.*, 6 (2): 159-175.
14. Elnimr, T. (2011): Evaluation of some heavy metals in *Pangasius hypophthalmus* and *Tilapia nilotica* and the role of acetic acid in lowering their levels. *International Journal of Fisheries and Aquaculture.* 3(8): 151–157.
15. Egyptian Organization for Standardization "EOS" (2010): Maximum Levels for certain contaminants in foodstuffs. No 7136/2010. Egyptian Standards, Ministry of Industry, Egypt.
16. Association of Official Analytical Chemists "AOAC" (2006): Official Methods of Analysis. 31st Ed., W. Horwitz (Editor), Academic Press, Washington, D. C., USA.
17. Staniskiene, B.; Matusevicius, P.; Budreckiene, P. and Skibniewska, K.A. (2006): Distribution of heavy metals in tissues of freshwater fish in Lithuania. *Polish J. Environ Studies*, 15(4): 585-591.
18. Andreji, J.; Stranai, Z.; Massonyl, P. and Valent, M. (2005): Concentration of selected metal in muscle of various fish species. *J. Environ. Sci. Heal.* 40 (4):899-912.

19. Javed, M. (2012): Tissue-specific bioaccumulation of metals in fish during chronic waterborne and dietary exposures. *Pak. Vet. J.* 32, 567–570.
20. Hounkpatin, A.S.Y., Johnson, R.C., Guédénon, P., Domingo, E., Alimba, C.G., Boko, M., Etorh, P.A., (2012): Protective Effects of Vitamin C on Haematological Parameters in Intoxicated Wistar rats with Cadmium, Mercury and Combined Cadmium and Mercury. *Int. Res. J. Biolo. Sci.* 1 (8), 76–81.
21. Fernandes, C., Fontaínhas-Fernandes, A., Cabral, D., Salgado, M., (2008): Heavy metals in water, sediment and tissues of *Liza saliens* from Esmoriz-Paramos lagoon. *Port. Environ. Monit. Assess.* 136, 267–275.
22. Sohsah - Madiha, A.M. (2009): Studies on some heavy metal residues in freshwater fish with special reference to water environmental pollution. PhD, Thesis, (Meat Hygiene), Fac. Vet. Med., Benha Univ., Egypt.
23. Shaltout, A. F ; Hashim F. M ; El-nahas, S. (2015): Detection of some heavy metals in fish (*Tilapia nilotica* and *Clarias lazera*) at Menofia governorate. *Benha. Vet. Med. J.* 29(2):124-128.
24. Abd El Maksod, Hania, E.; Saad, M. S; Fatahalla, Gehan ; Samir, Maha. M. (2023): Chemical Residues in Some Farmed Fish Species Marketed in Sharkia Governorate, Egypt. *Benha. Vet. Med. J.* 45:141-145.
25. Helmy, Nariman, A.H.; Hassan, M.A; Hassanien, Faten. S; Maarouf, A.A. (2018): Detection of Heavy Metals Residues in Fish and Shellfish. *Benha. Vet. Med. J.* 34(2): 255-264
26. Shokr, Loaloo, A. (2019): Studies on Heavy Metal Residues Contaminating Nile and Marine Fishes. PhD. V.Sc. Thesis (Meat Hygiene), Food control Dep., Fac. Vet. Med., Benha Univ., Egypt.
27. AbouZeid, O.A.; El Bayoumi, Z.H; Edris, A.M; Shawish, R.R. (2023): Studies on some Heavy Metal Residues in Fish. PhD. V.Sc. Thesis (Meat Hygiene), Food co
28. Elhefnawy, M.A .; Edris, A.M ; El-sheikh, Nabila. I . (2023): Novel approaches with acetic acid, liquid smoke, and *Lactobacillus rhamnosus* for mitigating some heavy metals in Nile fish. *Benha. Vet. Med. J.* 45:221-225.
29. El-Zahaby-Dina, I.M. (2007): Micro-biological and chemical studies on some fish and fish products in Menoufia Governorate. M.V.Sc., Thesis. (Meat Hygiene), Fac. Vet. Med., Menoufia Univ., Sadat Branch, Egypt.
30. Nasser F. S. Ali; Hussein M. H. Mohamed; Fathi A. M. El-Nawawi. (2016): Heavy metals residues of public health significance in some freshwater fish farmed in Kafr-El Sheikh and Menofia governorates. *Egypt. J. Food. safety.* 3(2):37-46.
31. Castoldi, A. F., Coccini, T. And Manzo, L. (2003): Neurotoxic And Molecular Effects of Methyl Mercury in Humans. *Reviews On Environmental Health*, 18:19 – 31
32. Mert, W. (1987): Trace Element in Human and Animal Nutrition. 5th Ed. Vol. I And It London: Academic Press.
33. Clarkson, T.W. (2002): The three modern faces of mercury. *Environmental Health Prespect*, 110(1):11-32.
34. Helmy, Nariman, A.H.; Hassan, M.A; Hassanien, Faten. S; Maarouf, A.A. (2018): Detection of Heavy Metals Residues in Fish and Shellfish. *Benha. Vet. Med. J.* 34(2): 255-264.
35. Shokr, Loaloo, A. (2019): Studies on Heavy Metal Residues Contaminating Nile and Marine Fishes. PhD. V.Sc. Thesis (Meat Hygiene), Food control Dep., Fac. Vet. Med., Benha Univ., Egypt.
36. AbouZeid, O.A.; El Bayoumi, Z.H; Edris, A.M; Shawish, R.R. (2023): Studies on some Heavy Metal Residues in Fish. PhD. V.Sc. Thesis (Meat Hygiene), Food control Dep., Fac. Vet. Med., Benha Univ., Egypt.
37. Chisolm, J. (1973): Management of Increased Lead Absorption and Lead Poisoning. *Eng. J. Med.*, 289:1016 - 1017.
38. Jones, T.C., Hunt, R.D. (1983): *Veterinary Pathology*. 5th Ed. Lea and Febiger Philadelphia (USA).
39. Kikuchi, Y., Nomiyama, T., Kumagai, N.; Uemura, T., Omar, K. (2002): Cadmium concentration in current Japanese food beverages. *J. Occupational Health.* 44:240-247.
40. Feldman, D.; Ganon, J.; Haffman, R. and Simpson, J. (2003): *The solution for data analysis and presentation graphics*. 2nd Ed., Abacus Lancrpts, Inc., Berkeley, USA.
41. Konsowa, A. H. (2007): Ecological studies on fish farms of El-Fayoum depression. *E. J. Aquat. Res.* 33(1): 290-299.