

Journal of Food and Dairy Sciences


Journal homepage & Available online at: www.jfds.journals.ekb.eg

The Effect of Water Replacement with Different Proportions of Permeate on the Properties and Quality of Processed Cheese

Refaey, M. M. M.* ; M. S. Gomaa ; A. El-M. El -Shamy and Samar A. Zalma



Dairy Department Faculty of Agriculture ,Mansoura University



In this research, the effect of replacing water used in the manufacture of processed cheese with different percentages of permeate on the properties cheese. The results were as follows: an increase in the total solids of the cheese was observed with the increase in the percentage was (49.73%). Also, an increase in both protein and fat (8.82-19.64%) with the increase in the percentage of replacement was observed. Also, the percentage of fat in the solid material decreased (40.35%). An increase in the pH values (5.31) was observed with the increase in the percentage of replacement and a decrease with storage. The same reverse effect was observed for acidity in all samples. Regarding microbiological properties and bacterial count, it was observed that with increasing the addition percentage, the total microbial count increased. The lowest numbers were in the control sample, which was 2×10^3 , while the numbers increased with storage, reaching 11×10^3 at the end of storage. There was also a non-noticeable increase in the numbers of fungi and yeasts with increasing the replacement percentage, during the storage period. The lowest numbers were in the control sample, whether in fresh cheese or after 90 days ($1-9 \times 10^3$), while chloroform and staphylococcus bacteria were not present, whether in fresh cheese or at the end of storage. Regarding rheological properties, the hardness increased, and the rest of the rheological decreased with increasing the replacement percentage. The sensory properties of the cheese, whether, were better with increasing the replacement.

Keywords: processed cheese , permeate ,properties ,effect

INTRODUCTION

There are numerous definitions describing processed cheese. In additions, rocessed cheese continues to gain appeal because of its broad composition and wide applications (El Dakhkhny and Dabour, 2016). Among them, the definition proposed by (Guinee *et al.*, 2004) is the most frequently used and the simplest one.

The processed cheese as “the cheese which can be produced by blending natural cheese of different ages and degrees of maturity in the presence of emulsifying salts and other dairy and nondairy ingredients followed by heating and continuous mixing to form a homogeneous product Over the past century”, the manufacturing of processed cheese has grown substantially, resulting in the creation of a wide range of products all over the world. Processed cheese was typically divided into two categories: spreadable processed cheese and processed cheese, depending on the finished product's physical attributes (Smith, 1990). Processed cheese is defined by (Gulzar *et al.*, 2020) as a uniform blend of natural cheese types, vegetable oils, butter oil, milk solids, emulsifying salts, and additional dairy or nondairy ingredients with a longer shelf life. Processed cheese was created in response to the need to increase the shelf life of natural cheese, recycle cheese that was flawed, and create a cheese with unique flavor, texture, and useful qualities (Falih *et al.*, 2024). In Europe, processed cheese manufacture began in the early 1900s. Emulsifying salts (ES), like sodium phosphates and trisodium citrate (TSC), are frequently employed in the production of processed cheese to regulate melting, texture, and the creation of free oil (Meyer, 1973; Berger *et al.*, 1998 and Fox *et al.*, 2000).

Since these compounds are not surface-active chemicals, they are not true emulsifiers in the literal sense of the word, unlike mono- or diglycerides. Enhancing the emulsifying ability of cheese proteins is ES's crucial processing function (Caric *et al.*, 1985). The development of ultrafiltration (UF) technology over the past 20 years has now enabled the manufacturing of low-lactose whey protein concentrates (WPC) at a comparatively low cost. Producing WPC with various physicochemical and functional characteristics for use in various food systems is now feasible thanks to this breakthrough technology. There are various types of whey that can be used in the production of processed cheese, including "sweet" whey (pH > 5.8) from cheddar, The production of fresh acid cheeses like Danbo, Queso Blanco, and others, as well as "medium" acid whey (pH 5.0 to 5.8) and "acid" whey (pH < 5.0) were obtained from the manufacturing of fresh acid cheeses like mozzarella and other similar cheeses or rennet casein.

The aim of this research is studying the effect of replacing water used in processed cheese manufacturing with different ratios of permeate (25%, 50%, 75%, and 100%) on the chemical, microbiological, rheological, and sensory characteristics of the resultant processed cheese.

MATERIALS AND METHODS

Materials:

- 1-**Ras cheese:** which made at Dairy Research Unit, Faculty of Agriculture Mansoura University was used in the experimental search.
- 2-**Kareish cheese:** made in Dairy Unit Faculty of Agriculture Mansoura University.
- 3-**Skim milk powder:** made in Poland by "Arla "

* Corresponding author.

E-mail address: mahmoud_refae_2004@mans.edu.eg

DOI: 10.21608/jfds.2025.398213.1204

4- **Palm oil**: imported from Indonesia. Two types of palm oil, from kernel and the other from the fruit.

5-Emulsifying salts are:

a- EGY Phos S20 (Egy Dairy Company - East-Egypt). b- Joha NO with corrector (made in Germany, both were bought from EGY Phos Company found in AL-Aasher of Ramadan/Sharkia Governorate

6- **Permeate**: obtained from Elshehab company in Mansoura city.

7- **Commercial grade salt** (NaCl), and vegetable oil, were obtained from the local market.

Methods:

Manufacture of processed cheese spread:

Control processed cheese and other processed cheese treatments were prepared by using ingredients illustrated in Table (1) by using the method described by was El-Assar (1991) as follows:

Control processed cheese was prepared by grinding Ras cheese and kareish cheese, and mixing with skim milk powder, palm oil, emulsifying salts, and water. Other processed cheese treatments were prepared by replacing water with permeate at different ratios 25% (P1), 50% (P2), 75% (P3), and 100% (P4), followed by cooking at 90°C for 15 min, packaging in sterilized plastic containers, and cooling to 7°C. Finally keeping under refrigeration for 90 days as shown in Figure(1)

1-Chemical analysis:

Control processed cheese and other treatments containing permeate in replacement of water at different ratios were chemically analyzed when they were fresh, and after 30, 60, and 90 days of refrigerated storage to determine their content from total solids (T.S%), fat content (%), and protein content (%) (AOAC, 2019). Titratable acidity (%) was determined by titration with NaOH (1/9N), and pH values were determined by using a pH meter (Hanna Instruments Microprocessor pH-Meter 211, Woonsocket, RI, USA)

2-Organoleptic evaluation:

Control processed cheese and other treatments containing permeate in replacement of water at different ratios were sensory evaluated when they were fresh, and after 30, 60, and 90 days of refrigerated storage. A total score of 100 points was assigned as follows: 40points for flavor and taste, 30 points for body and texture, and 30 points for color and appearance as described by Hassan et al. (2004).

3-Microbial analysis

Control processed cheese and other treatments containing permeate in replacement of water at different ratios were microbiologically analyzed when they were fresh, and after 30, 60, and 90 days of refrigerated storage by using the standard plate count technique, where the examined samples were serially diluted in a sterile saline (0.85% sodium chloride). For the detection of total bacterial count selected dilutions were plated on Nutrient agar plates, and incubated under aerobic conditions at 37°C for 24-48 hours. The presence of Coliform and Staphylococci bacteria was determined by enumeration on MacConkey agar, and Staphylococcus agar 110, respectively with similar incubation conditions. Counts of Molds and yeasts were assessed by using potato dextrose agar (adjusted to pH 3.5), followed by incubation aerobically at 30°C for 24-48 hours according to (Standard Methods for the Examination of Dairy Products, 1978).

Table 1. The composition Processed cheese base ingredients (%).

	%	Chemical analysis			
		Fat/TS	TS%	Fat %	Protein %
Ras cheese	10_20%	54.88	62.35	34.22	25.21
Kareish cheese	20-30	0.02	25.00	0.5	21.2
Skim milk powder	8-12%	0.01	96.0	0.10	34.00
Palm oil	20-30%	100	100	100	0.0
Emulsifying salt	1-3%	0.00	99.30	0.00	0.00
Water	20-25%	0.0	0.00	0.00	000
Permeate		7.13	5.61	0.4	0.49
Total	100				




Fig. 1. Steps for making processed cheese

4- Textural Analyses

Using a texture analyzer (CNS/FARNELLFRA, Borehamwood, Hertfordshire, England), the texture characteristics of cheese samples were assessed. Fresh cheese samples were used for control and experimental purposes, and 60-day aged cheese samples that were stronger were measured right away. The dimensions of the cheese sample were 20 mm in height and 30 mm in circumference. The speed was 1 mm/s, and the penetration distance was 10 mm. At least twenty minutes before testing, the samples were left to stand at room temperature. The perplex cone TA15-450C probe was utilized. Using an LFRA texture analyzer and a computer interface, the texture profile parameters were computed after data was gathered on a computer. The texture profile characteristics described by Bourne (1978) were determined and obtained as follows:

- The hardness is measured by the compressive force (g) obtained at maximum compressive during the initial bite.
- As a cohesiveness metric, the ratio of the positive force

area under the curve during the first compression (a_2/a_1) to that during the second compression (bite).

- iii) As a measure of springiness, the height (mm) to which the sample recovered during the clamping period between the conclusion of the first bite and the beginning of the second bite.
- iV) Gumminess is measured as the product of hardness and cohesiveness (g). V) A measure of chewiness, calculated as the product of gumminess and springiness (g.mm). Vi) The modulus, which is the force slope that indicates the rigidity of the sample

RESULTS AND DISCUSSION

pH Values

Results from (Table 2) showed increased pH was observed with increasing permeate replacement ratio in all treatments, but with storage a decrease in pH values occurred. this is might by due to the existence of these organic preservatives, which prevented the growth of microorganisms while they were being stored. The pH of processed cheese is known to be closer to 5.7; block processed cheese has a lower pH, whereas processed cheese spread has a higher pH. Lactic acid was used to modify the pH values of cheese (Codex STAN-A-8(b)-1978). The pH and acidity levels of processed cheese were not included in (Egyptian Standards E.S 1132/2013).

Table 2. chemical composition of processes cheese made from different ratio of permeate.

Treatments	Storage periods	pH	Acidity	T.S %	TP %	Fat %	Fat/DM
Control	Fresh	5.22	0.23	44.69	8.76	19.22	43.01
	30	5.22	0.24	44.88	8.78	19.23	42.85
	60	5.21	0.24	44.91	8.79	19.27	42.91
	90	5.20	0.25	45.12	8.81	19.31	42.80
P1 (25 %)	Fresh	5.23	0.21	46.68	8.77	19.13	40.98
	30	5.22	0.22	47.12	8.78	19.20	40.75
	60	5.22	0.24	47.53	8.78	19.31	40.63
	90	5.20	0.25	47.73	8.79	19.32	40.48
P2 (50%)	Fresh	5.24	0.22	47.01	8.78	18.91	40.23
	30	5.23	0.23	47.67	8.81	19.00	39.86
	60	5.22	0.25	47.85	8.82	19.15	40.02
	90	5.21	0.26	48.03	8.83	19.17	39.91
P3 (75 %)	Fresh	5.30	0.23	48.35	8.81	19.1	39.50
	30	5.28	0.23	48.55	8.82	19.19	39.53
	60	5.23	0.25	48.71	8.83	19.23	39.48
	90	5.22	0.26	49.81	8.84	19.44	39.03
P4 (100%)	Fresh	5.31	0.24	48.68	8.82	19.64	40.35
	30	5.30	0.24	49.12	8.83	19.66	40.02
	60	5.28	0.25	49.53	8.84	19.69	39.75
	90	5.26	0.26	49.73	8.85	19.82	39.86

Titrateable acidity

In the same trend the acidity increased at 90 days storage periods which P1(25%), P2 (50%) P3 (75%) and P4 (100%) recorded highest values of acidity in comparing to other storage periods. The range of acidity in both fresh and preserved cheese was 0.20 to 0.30 percent. The values that were obtained are accurately correlated with the pH values. While the current results were consistent with Timothy *et al.* (2009), the acquired results were not in agreement with those reported by Metwaly *et al.* ,(2007).

Total Solids content

Total solids percentage T.S% increased with replacement ratio recorded highest values at P4 (100%) 49.73% in comparing other treatments either the control fresh processed cheese or after at 90 days storage periods. Egyptian Standards (E.S. 1132/2013) address manufactured cheese that contains vegetable oils. There should be at least

45% total solids. Five samples of each of the six processed cheese brands produced and marketed in India were examined by (Boghra *et al.*, 2006); the moisture content ranged from 44.9 to 48.68%.

Total protein content

The treatment P4 (100%) recorded the highest total protein (T.P%) 8.85at 90 days storage periods in comparing to other treatments and control either fresh cheese or during storage periods. Because the cheese's moisture level has decreased, the protein content of the control cheese seems to be higher. The permitted percentage of protein was not included in the Egyptian Standards (E.S1132/2013). Similarly, the legality of protein was not mentioned in Codex STAN-A8 (b) (1978). In addition to skim milk powder (SMP), a significant proportion of Ras and Kariesh cheese is used to produce our fresh processed cheese, which has a high protein content (18.00-24.34%).

Total Fat content

The treatment P4 (100%) recorded highest fat content 19.64 to 19.82 at fresh cheese or after 90 days of storage periods. For all treatments, processed cheese had a higher fat content. The fat content marginally rose as the storage period went on. The cheese's increased total solids are the cause of the apparent rise. The findings were consistent with those of Metwaly *et al.*, (2007).

Fat/Dry matter

On the other hands the P3 (75%) recorded the lowest Fat/DM 39.03 at 90 days storage periods which P3 at 90 days of storage periods recorded highest values in comparing to other treatments. Egyptian Standards (E.S1132/2013) listed Fat/DM not less than 65% for high cream cheese and not less than 35% for whole fat cheese as well 25-35 for half cream cheese, low fat cheese 10-25% fat cheese.

The obtained results are similar to those reported by (Awaad *et al.*, 2020); (Lara *et al.*, 2021) and (Bandyopadhyaya *et al.*, 2024).

Data presented in Table 3 showed the effect of replacement water with permeate cheese on microbiological properties processes cheese.

Table 3. Microbiological properties of processes cheese made from different ratio of permeate.

Treatments	Storage periods	T.C *10 ³	M&Y *10 ³	E. coli *10 ³	Staph *10 ³
Control	Fresh	2	ND	ND	ND
	30	3	1	ND	ND
	60	7	2	ND	ND
	90	11	4	ND	ND
P1 (25 %)	Fresh	3	ND	ND	ND
	30	5	6	ND	ND
	60	9	7	ND	ND
	90	12	9	ND	ND
P2 (50%)	Fresh	5	ND	ND	ND
	30	8	3	ND	ND
	60	11	4	ND	ND
	90	13	6	ND	ND
P3 (75 %)	Fresh	6	ND	ND	ND
	30	8	3	ND	ND
	60	11	5	ND	ND
	90	15	7	ND	ND
P4 (100%)	Fresh	7	ND	ND	ND
	30	10	4	ND	ND
	60	12	7	ND	ND
	90	17	9	ND	ND

The treatment P4 (100%) recorded highest values of T.C*10³ and M&Y *10³ 17 and 13 respectively and Microbiological analysis was conducted during the storage period, starting with fresh treatments, and the analysis process continued at 30 days and 60 days until the end of the

storage period at 90 days. It was noted in the following table that as the storage period increased, the total count content and the content of Molds and Yeasts increased for all treatments. It was also noted that all treatments were not detect of coliform bacteria and staphylococcus aureus during the storage period. The obtained results are similar to those reported by (Awaad *et al.*, 2020); (Lara *et al.*, 2021) and Bandyopadhaya *et al.*, 2024).

(Table 4) presented the effect of replacement water with permeate cheese on rheological properties processes cheese. From collected data the treatment of P4 (100 %) recorded the highest values of Hardness (N) 104.04, while Cohesiveness (ratio), Springiness (mm), and Gumminess (N) decrease 35.5, 0.26, 13.57 and 15.83 respectively at fresh cheese. The obtained results are similar to those reported by (Awaad *et al.*, 2020; Lara *et al.*, 2021 and Bandyopadhaya *et al.*, 2024).

Table 4. Rheological properties of processes cheese made from different ratio of permeate.

Treatments	Storage periods	Control	P1 (25%)	P2 (50%)	P3 (75%)	P4 (100%)
Hardness (N)	Fresh	91.4	94.00	97.60	101.10	104.04
	90	73.4	77.00	86.80	90.90	96.00
Adhesiveness (mj)	Fresh	60.81	57.90	41.9	36.3	35.5
	90	50.8	45.5	30.4	27.20	23.9
Cohesiveness (ratio)	Fresh	0.37	0.31	0.27	0.26	0.22
	90	0.26	0.29	0.26	0.24	0.19
Springiness (mm)	Fresh	7.75	13.15	12.45	13.22	13.57
	90	10.98	14.04	15.92	14.93	15.83
Gumminess (N)	Fresh	23.4	20.0	19.0	18.4	15.30
	90	42.4	36.04	33.1	29.5	28.0
Chewiness (mj)	Fresh	481	264	336	270	207
	90	365	230	255	233	171

On the other hand's during storage periods decrease in Hardness, Adhesiveness, cohesiveness, chewiness and increased springiness, gumminess days of storage periods.

Hardness (N):

It is defined as the force needed to achieve a specific deformation and is correlated with the strength of the cheese matrix. It is also a measure of the force needed to compress the cheese treatment. The hardness value increased for all treatments after the processed cheese was refrigerated for ninety days. This is due to the addition of emulsifying that led to the formation of additional iso peptide bonds, and the production of a gelatinous network with smaller aggregates and smaller pore sizes (Imm *et al.*, 2000).

Springiness (mm):

Defined as the ratio of which a distorted material returns to its main form when removing the distorting strength Szczesniak *et al.*, (1963). On the other hand, the Springiness of the control treatment increased during the first 90 days, and the results showed an increase in the Springiness value for all treatments at the end of storage compared to the control treatment.

Cohesiveness (%)

Defined as the force of internal bonds forming the body of the product Bourne (1978) and Szczesniak *et al.*, (1963). Cohesiveness values decreased gradually by the gradually addition of retentates amount in fresh processed cheese (Table,4). On the other hand, the Cohesiveness of the control treatment decreased during the

first 90 days, and the results showed a decrease in the Cohesiveness value for all treatments at the end of storage compared to the control treatment. A decrease in the Cohesiveness value for treatment at the end of storage. The obtained results are in accordance with Ossman. E. (2018).

Gumminess (N):

Defined as the power required to disintegrate a semi-solid food for swallowing Szczesniak *et al.*, (1963). Table (4) showed that addition of retentates increase the processed cheese values for some treatments, the gumminess increased from for fresh and 90 days storage respectively. The results showed an increase in the gumminess value for all treatments at the end of storage and an increase in the gumminess value for all treatment in fresh. The results obtained are consistent with Othman. E. (2018).

Chewiness (J):

The energy needed to chew a solid food item is called chewiness, and it is related to other textural characteristics like cohesion, hardness, and springiness (Prakasan *et al.*, 2015 and Salinas-valdes *et al.*, 2015). Chewiness values decreased gradually in fresh processed cheese (Table,5).

Table 5. The effect of replacement water with permeate cheese on organoleptic properties processes cheese.

Treatments	Storage periods	Flavor (40)	Int app (30)	Outer app (30) b	Total
Control	Fresh	36	27.5	26.5	90
	30	36	26	26	88
	60	35	25.5	25.5	86
	90	35	25.0	25	85
P1 (25 %)	Fresh	36	25.0	24	85
	30	35	25.0	24	84
	60	35	24	24	83
	90	33	23	23	79
P2 (50%)	Fresh	38	27	26	91
	30	37	27	25	89
	60	36	26	25	87
	90	34	24	25	83
P3 (75 %)	Fresh	38	28	26	92
	30	37	27	26	90
	60	36	26.5	25.5	88
	90	35	25	25	85
P4 (100%)	Fresh	38	28.5	27.5	94
	30	38	28	27	93
	60	36	27	26	89
	90	36	26	26	88

On the other hand, values decreased by storage time, the results showed a decrease in the Chewiness value of all treatments. The obtained results are in accordance with Salinas-Valdés *et al.*, (2015). The obtained results are similar to those reported by (Awaad *et al.*, 2020; Lara *et al.*, 2021 and Bandyopadhaya *et al.*, 2024).

Data presented in Table 4 showed the effect of replacement water with permeate cheese on organoleptic properties processes cheese. Results indicated that long storage periods affect the flavor which lowest values at 90 days storage periods and the best flavor recorded at 30 days of storage periods in comparing no storage (fresh). The best flavor recorded 38 at 30 days storage periods in treatment of P4 (100%). The organoleptic properties processes cheese affected by long storage periods. The obtained results are similar to those reported by (Awaad *et al.*, 2020; Lara *et al.*, 2021 and Bandyopadhaya *et al.*, 2024).

REFERENCE

- American Public Health Association (1980). Standard Methods for The Examination of Water and Waste Water. 15th Ed., New York, USA.
- Awaad, S. S., Moawad, A. A., Ayah, B., & Sallam, S. S. (2020). Impact of raw materials and processing techniques on the microbiological quality of Egyptian Domiati cheese analogue. th Arab. Mans. Conf. of Food , Dairy Sci, Tec(12)- 10215-10225
- A.O.A.C (1990). Official Method of Analysis 15th. Association of Official Analysis Chemists Inc., U.S.A

- Bandyopadhyaya, S.; Sarkar, U and Bera, D. (2024). Food Stability Simulation: Accelerated Shelf-Life Mechanism. Food Coatings and Preservation Technologies, 259-300.
- Bourne, M.C. (1978) Texture Profile Analysis. Food Technology, 32, 62-66, 72.
- Berger, W.; Klostermeyer, H.; Merkenich, K. and Uhlmann, G. (1998). Processed cheese manufacture, a JOHA guide. BK Giulini Chemie GmbH & Co. OHG, Ladenburg, Germany.
- Boghra, V.R; Borkhatrya, V. N; Sharma, R.S (2006)." Market survey on chemical Cheddar-style cheese on the biochemical and rheological properties of processed cheese Dairy Sci. Technol. 89 317–333.
- Caric, M.; Gantar, M. and Kalab, M. (1985). Effects of emulsifying agents on the microstructure and other characteristics of process cheese- a review. Food structure, 4(2),13.297-312.<https://digitalcommons.usu.edu/foodmicrostructure/vol4/iss2/13>
- Difo (1966). Manual, of Dehydrate Culture Media and Reagents Procedures. Nith. Ed Difco Laboratories, Detroit, Michigan, USA.
- Difco Manual of Dehydrate Culture Media and Reagents (1974). Pub Difco Laboratory Incorporated. Detroit, I., Michigan, 48, 201, USA.
- EL- Assar, M.A.E.G (1991). Studies on Processed Cheese. Ph.D. Thesis, Faculty of Agric. Cairo University. composition and aflatoxin MI in processed cheese " Indian Journal of Dairy
- El Dakhakhny, E. and Dabour, N. (2016). Processed cheese: Basics and possibility for the development of healthier products. Alexandria Journal of Food Science and Technology, 13(2), 45-62.
- Falih, M. A.; Altemimi, A. B.; ALKaisy, Q. H.; Awlqadr, F. H.; Abedelmaksoud, T. G.; Amjadi, S. and Hesarinejad, M. A. (2024). Enhancing safety and quality in the global cheese industry: A review of innovative preservation techniques. *Heliyon*. 10.1016/j.heliyon.2024.e40459
- Fox, P.F.; Guinee, T.R.; Cogan, T.M. and Mc Sweeney, P.L.M. (2000). Processed cheese and substitute or imitation cheese products. In: Fundamentals of Cheese Science. Aspen Publishers, Inc, Gaithersburg, Maryland, USA. pp.: 429-451. 10.1007/978-1-4899-7681-9
- Guinee, T. P.; Carić, M. and Kalab, M. (2004). Pasteurized processed cheese and substitute/imitation cheese products. In *Cheese: chemistry, physics and microbiology* (Vol. 2, pp. 349-394). Academic Press [https://doi.org/10.1016/S1874-558X\(04\)80052-6](https://doi.org/10.1016/S1874-558X(04)80052-6)
- Gulzar, N.; Sameen, A.; Muhammad Aadil, R.; Sahar, A.; Rafiq, S.; Huma, N.; Nadeem, M.; Arshad, R. and Muqadas Saleem, I. (2020). Descriptive sensory analysis of pizza cheese made from mozzarella and semi-ripened cheddar cheese under microwave and conventional cooking. Food, 9(2), 214.
- Lara-Castellanos, M. J.; Azuara, E.; Jimenez-Fernandez, V. M.; Luna-Solano, G. and Jimenez, M. (2021). Effect of casein replacement by modified casein on physicochemical, textural, sensorial properties and microbiological stability of fresh cheese. International Dairy Journal, 112, 104864.
- Ling, E.R (1963). A text book of dairy chemistry. Vol.1,2,3rd. ED. Chapman and Hall, Ltd.37Essex Str., London W.C.Z.Ltd.,(1sted.),11-29. <https://www.cabidigitallibrary.org/doi/full/10.5555/19740410281>
- Metwaly . M.Abo- Srea Ahmed; EL Tahra. M. A. Ammar and Abdel-Wahab El Shazly Khalil (2007). improvement the keeping quality and flavour of processed cheese analogue. th Arab. Mans. Conf. of Food , Dairy Sci, Tec(12)- 10215-10225
- Meyer, A. (1973). Processed Cheese Manufacture. London, U.K.: Food Trade Press Ltd.,(1sted.),11-29. <https://www.cabidigitallibrary.org/doi/full/10.5555/19740410281>
- Othman N.A.; EL-Shabrawy, S.A. and Awad, R.A. (2005). Preference evaluation and properties of reduced sodium processed cheese spreads. Annals Agric. Sci. Ain Shams Univ., Cairo, 50(1), 209 – 225.<https://www.cabidigitallibrary.org/doi/full/10.5555/20053149454>
- Osman, A. and E. M. Abd El-Wahed (2019). the effects of crude phenolic-rich extract from mushroom (*Agaricus bisporus*) on shelf-life extension and quality attributes of the feta like cheese. J. Food and Dairy Sci., Mansoura Univ., 10 (4): 85 - 91, 2019.
- Oxoid Manual of Culture Media (1962). Pub. By the Oxoid Division, Oxo Ltd, south Wark bridge Rd., London, S.E.I. Salinase-valdes et al., (2015) The Effect of Adding Microbial Transglutaminase Enzyme to Cheese Milk on the Quality of Halloumi-Like Cheese Made From Recombined Skimmed Milk Powder Egypt. J. Chem. Vol. 66, No. 7 pp. 577 - 591
- Smith, B.L. (1990). Codex Alimentarius: Abridged Version, Food and Agriculture Organization of the United Nations, Rome.pp:12.1012.16.
- Szczesniak, A., Brandt, M. Freidman, H (1963) . "Development of standard rating scales for mechanical parameters and correlation between the objective and sensory texture measurements" Food Tech., 22, pp. 50–54:(1963)
- Timothy P.; Guinee, B. T. and O'KENNEDY (2009) The effect of calcium content of Cheddar-style cheese on the biochemical and rheological properties of processedcheese Dairy Sci. Technol. 89 317–333

تأثير استبدال الماء بنسب مختلفة من الراشح الناتج من الترشيح الفائق على خواص وجودة الجبن المطبوخ

محمود مصطفى محمد رفاعي ، محمد شلبي جمعة ، احمد المتولى الشامى و سمر احمد عبد الجليل زلمه

قسم الالبان كلية الزراعة جامعة المنصورة

الملخص

في هذا البحث تم دراسة تأثير استبدال الماء المستخدم في صناعة الجبن المطبوخ بنسب مختلفة من البريميت permeate على الخواص الكيميائية والميكروبيولوجية والريولوجية والحسية للجبن المطبوخ وكانت النتائج كالتالي: تم ملاحظة زيادة المواد الصلبة الكلية للجبن المطبوخ مع زيادة نسبة الاستبدال من الراشح حتى وصلت نسبة الاستبدال إلى 100٪ وكانت (49.73٪) وكذلك زيادة كل من البروتين والدهن (8.82-19.64٪) مع زيادة نسبة الاستبدال وكذلك انخفضت نسب الدهن / المادة الصلبة (40.35٪) وتم ملاحظة زيادة قيم الأس الهيدروجيني (5.31) مع زيادة نسبة الاستبدال وانخفاضه مع التخزين ونفس التأثير العكسي بالنسبة للحموضة في كل العينات. بالنسبة للخواص الميكروبيولوجية والعد البكتيري لوحظ مع زيادة نسبة الاضافة زاد العد الكلي للميكروبات وكانت أقل الأعداد في العينة الكنترول وكانت 10×2^3 بينما زادت الأعداد مع التخزين حيث وصلت في نهاية التخزين إلى 10×11^3 كذلك حدث زيادة غير ملحوظة لأعداد الفطريات والخمائر مع زيادة نسبة الاستبدال وكذلك خلال فترة التخزين وكانت أقل الأعداد في العينة الكنترول سواء في الجبن الطازج أو بعد 90 يوم (10×9) بينما بالنسبة للكوليفورم والبكتريا العفوية كانت غير موجودة سواء في الجبن الطازج أو في نهاية التخزين. بالنسبة للخواص الريولوجية زادت الصلابة للجبن وانخفضت باقي الخواص الريولوجية مع زيادة نسبة الاستبدال وكانت الخواص الحسية للجبن سواء الطعم أو القوام والمظهر أفضل مع زيادة نسبة الاستبدال حتى 100 ٪