

IJMA



INTERNATIONAL JOURNAL OF MEDICAL ARTS

Volume 7, Issue 10 (October 2025)



<http://ijma.journals.ekb.eg/>

P-ISSN: 2636-4174

E-ISSN: 2682-3780



Available online on Journal Website
<https://ijma.journals.ekb.eg/>
 Main Subject [Pediatrics]



Original Article

Maternal and Fetal factors and Short-Term Neonatal Outcomes Associated with Late Preterm Delivery

Ibrahim Amr El-Deeb*, Hany Elkhalegy; Mohammed Mostafa Abo-Elmaaty

Department of Pediatrics, Damietta Faculty of Medicine, Al-Azhar University, Damietta, Egypt.

Abstract

Article information

Received: 14-03-2025

Accepted: 28-08-2025

DOI: [10.21608/ijma.2025.365354.2153](https://doi.org/10.21608/ijma.2025.365354.2153).

*Corresponding author

Email: amrieldeeb@gmail.com

Citation: El-Deeb IA, Elkhalegy H, Abo-Elmaaty MM. Maternal and Fetal factors and Short-Term Neonatal Outcomes Associated with Late Preterm Delivery. IJMA 2025 October; 7[10]: 6206-6212. doi: [10.21608/ijma.2025.365354.2153](https://doi.org/10.21608/ijma.2025.365354.2153).

Background: The primary cause of infant mortality and morbidity is still preterm births [PTB]. In our nation, little research has been done on the prevalence of late PTB and related perinatal outcomes for patients receiving treatment in the intensive care unit.

The aim of the work: The aim of the current study was to determine predictors of unfavorable neonatal outcomes and to estimate the impact of gestational age on short-term neonatal morbidity and mortality in late preterm infants [LPIs] treated in intensive care.

Patients and Methods: This cross sectional included 322 women whose deliveries were performed at Al-Azhar University Hospital in Damietta, Egypt from August 2023 to August 2024. The following items were collected through interview meeting with the mother and her infant; sociodemographic [Age, Schooling and Residency], clinical-obstetric and perinatal characteristics

Results: According to the neurological morbidities, seizures represent 1.6%, hypoxic ischemic encephalopathy [HIE] represent 2.2%, and intra ventricular hemorrhage [IVH] represent 4.3%. The mortality rate in our study was 5.3%, all of them were from the age group 34 weeks [P =0.001]. Logistic regression analysis revealed that gestational age, gender, and maternal DM were the only predictors for the respiratory morbidity [P =0.001 for all]. A statistically significant association was found between the mortality rate and respiratory morbidity, neurological morbidity, infections, blood transfusion, hypoglycemia [P =0.001 for all]

Conclusion: It is critical to determine the factors that predict unfavorable short-term outcomes in late preterm infants for evaluating clinical practices and guidelines aimed at reducing infant morbidity and mortality.

Keywords: Fetal Risk Factors; Late Preterm Infants; Maternal Risk Factors; Preterm Births; Seizures.



This is an open-access article registered under the Creative Commons, ShareAlike 4.0 International license [CC BY-SA 4.0] [<https://creativecommons.org/licenses/by-sa/4.0/legalcode>].

INTRODUCTION

According to the World Health Organization [WHO], the definition of prematurity encompasses all newborns born before 37 weeks of gestational age [GA]. Late premature infants are those born between 34 and 36 weeks and 6 days, and extreme premature infants are born before 28 weeks of gestational age [1].

Prematurity occurs in more than one in ten births, and it is the leading cause of morbidity and mortality among newborns, with about 1.1 million deaths per year worldwide. Of these deaths, most can be prevented with basic care and low-cost interventions [2].

The main maternal complications that lead to late preterm birth are premature labor [PL] and premature rupture of membranes [PROM]. Other obstetric conditions contribute to prematurity, such as urinary tract infection, hyper-tensive disease, gestational diabetes, and twin pregnancies. Among the main fetal factors are restricted intrauterine growth and the non-reassuring fetal status [3].

The risk of neonatal complications is inversely proportional to GA. Each week the fetus remains in the womb, its development improves and the frequency and severity of neonatal complications are improved. Early complications of late prematurity include respiratory distress syndrome, apnea, hypothermia, hypoglycemia, hyperbilirubinemia, feeding difficulties, central nervous system immaturity, and infections, with a risk of at least seven times more complications in this group compared to term infants [TI] [4,5].

Morbidity and risk of hospitalization during the first year of life are also higher compared to term infants [6].

Long-term studies indicate that, when compared to preterm infants, late preterm infants are at an increased risk for neurodevelopmental disorders and learning disabilities, with neurocognitive changes that may persist into adulthood [7].

Because extreme preterm infants receive greater attention from health professionals for their high risk of complications, LPI infants are often neglected and treated the same as TI, which increases their risk for complications [8]. Therefore, the term “late premature” rather than “near term” is recommended in order to avoid the misinterpretation that preterm infants that are born close to term, share the same risks as a TI [9]. Knowledge of the factors that lead to the birth of late preterm infants can provide subsidies for health professionals and managers to prevent them, and reduce neo-natal morbidity and mortality related to prematurity.

The purpose of this study is to determine predictors of poor neonatal outcomes and to estimate the impact of gestational age on short-term neonatal morbidity and mortality in LPIs treated in the intensive care unit.

PATIENTS AND METHODS

This cross-sectional study included 322 women whose deliveries were performed at Al-Azhar University Hospital in Damietta from August 2023 to August 2024. Our study was guided by the Helsinki declaration principles.

Ethical approval was obtained from the Institutional review board of Damietta Faculty of Medicine, Al-Azhar University. An Informed

written consent was obtained from the parents of the infants before the recruitment.

We included newborns met the gestational age criteria for late preterm birth. However, we excluded the following: 1] age less than 34 weeks or more than 37 weeks; 2] Cases of fetal deaths; 3] Cases with major malformations

Collected data

- Sex, gestational age, parity, birth weight, Apgar score [i.e., vitality assessment at birth within the first five minutes after birth], and length of hospitalization in the intensive care unit were among the data analyzed.
- Among the maternal risk factors we identified were the mother's age, parity, pregnancy complications, maternal morbidity, and pregnancy treatment.
- The data on short-term outcomes were separated into four groups: neurological, infectious, respiratory, and other outcomes.
- The presence of respiratory distress syndrome, transient tachypnea of the newborn, persistent pulmonary hypertension in the neonate [PPHN], neonatal pneumothorax [NP], or the requirement for ventilatory support, such as mechanical ventilation and continuous positive airway pressure, were all considered respiratory morbidity.
- Meningitis, pneumonia, and Culture-proven sepsis were categorized as infectious morbidities.
- Intraventricular hemorrhage [IVH] and neonatal convulsions were categorized as central nervous system morbidities.
- Hypoglycemia, anemia needing a blood transfusion, and jaundice necessitating phototherapy were among the other morbidities.
- The Department of Pediatrics defined the ranges for clinical diagnoses made by a pediatrician in compliance with diagnostic protocols, which served as the basis for infant outcomes.
- For neurological and respiratory pathologies, radiological diagnostic techniques were employed in addition to clinical evaluation.
- Laboratory and microbiological results were utilized in the evaluation of infection and entities categorized as additional morbidities.

Statistical analysis

Statistical analysis was performed with SPSS statistical software, version 25 [IBM, Chicago, Illinois, USA]. The normality of the data was tested by the Kolmogorov-Smirnov test. Qualitative data were presented as numbers and percentages and were compared by the Chi square test, while quantitative data were presented as mean and standard deviations and were compared by the independent t test. Binary logistic regression analysis was done to determine the predictors of late preterm labor. As a result, the p-value was considered significant at the level of <0.05

RESULTS

A total number of 322 late preterm infants were included in our study. The gestational age ranged from 34 to 36 weeks, with a mean of 35 ± 0.8 weeks. The male gender represents 51.6% versus 48.4% females. The birth weight ranged from 1.2 to 3.4 kg, with a mean of 2.4 ± 0.5 kg. The mean head circumference was 32.5 ± 1.4 cm with a range of 42 – 50. The mean length was 45.2 ± 2.4 cm with a range of 42 – 50 cm. In terms of the BW – GA, 10.9% were SGA, 81.1% were AGA, and 8.1% were LGA [Table 1].

According to the maternal demographic data, the mean maternal age was 27.8 ± 6.3 years with a range of 19 – 40 years. According to their residency, 59.6% were from urban areas, and 40.4% were from rural areas. As regards the maternal work or employment, 28.3% were worker. According to the smoking status, 6.8% were smokers. In terms of the parity, 69.9% were multipara and 30.1% were nullipara [Table 2].

As regards the maternal obstetric data, the delivery indications were as follow; 18.3% maternal diseases, 44.7% obstetric complications, 37% were isolated spontaneous delivery. According to the antenatal care [ANC], 86.6% had underwent proper ANC [Table 4] According to the pregnancy morbidities, the most common morbidities were Anemia [44.7%], UTI [31.4%], and Uterine bleeding [30.1%]. As regards the type of delivery, 82.6% underwent CS, and 17.4% underwent NVD. According to the pregnancy outcomes, 96.6% delivered single infant and 3.4% delivered twins [Table 2].

In our study, 3.1% of infants whose gestational age was 34 weeks had major congenital anomalies, however those in the age categories 35 or 36 had no congenital anomalies [$P=0.001$]. According to the NICU admission, 35.1% of the total included infants required NICU admission, most of them were in the age category 34 weeks [$P=0.001$] [Table 3].

In terms of respiratory morbidities, the most common types were TTN [19.6%], RDS [9.9%], PPHN [5%], and apnea [4.3%]. Most of these morbidities were reported in the 34 weeks age category [$P=0.001$].

Mechanical ventilation was required only in 9.9% of the total included infants all of them were 34 weeks gestational age [$P=0.001$]. Twenty-seven percentage [27%] of the infants required CPAP, 53.7% from the age category 34 weeks, 21% were from the age category 35 weeks, and 8.3% were from the age category 36 weeks [Table 4].

According to the neurological morbidities, Seizures represent 1.6%, HIE represent 2.2%, and IVH grade represents 4.3% with a statistically significant association between the age group and the incidence of neurological morbidities in which most of them were reported in the age group 34 weeks [$P=0.001$ for all] [Table 5].

As regards the infection, Sepsis was reported in 36 infants [33.3%] in the age group 34 weeks, 4 infants [3.8%] in the age group 35 weeks, and in 4 cases [3.7%] in the age category of 36 weeks [$P=0.001$]. Pneumonia was reported in 3.7% of the infants, all of them were from the age group 34 weeks. NEC was reported in 3.7% of the infants, all of them were from the age group 34 weeks [Table 6].

Blood transfusion was reported in 8.1% of the infants. Hypoglycemia was reported in 5.3% of the infants. Thirty-two cases [9.9%] needed rehospitalization, 21 infants were from the age group 34 weeks, and 11 infants were from the age category 35 weeks [$p=0.001$]. The mortality rate in our study was 5.3%, all of them were from the age group 34 weeks [$P=0.001$] [Table 6].

Logistic regression analysis revealed that, gestational age, Gender, and maternal DM were the only predictors for the respiratory morbidity [$P=0.001$ for all] [Table 7]. Logistic regression analysis revealed that, gestational age, Oligo/poly hydramnios, maternal UTI, Preeclampsia, and maternal DM were the only predictors for the infection [$P=0.001$, 0.002, 0.001, 0.05, 0.05 respectively] [Table 8].

A statistically significant association was found between the mortality rate and Respiratory morbidity, Neurological morbidity, Infections, Blood transfusion, Hypoglycemia [$P=0.001$ for all] [Table 9].

Table [1]: Demographic and baseline clinical data of the studied patients.

Measures		Values
Gestational age [weeks]	mean $\pm$ SD	35 ± 0.8
	Min. – Max.	34 - 36
Sex [n,%]	Male	166 [51.6%]
	Female	156 [48.4%]
Birth weigh [kg]	mean $\pm$ SD	2.4 ± 0.5
	Min. – Max.	1.2 – 3.4
Head circumference [cm]	mean $\pm$ SD	32.5 ± 1.4
	Min. – Max.	42 – 50
Length [cm]	mean $\pm$ SD	45.2 ± 2.4
	Min. – Max.	42 - 50
Birth weight – Gestational age [n,%]	Small for gestational age	35 [10.9%]
	Average for gestational age	261 [81.1%]
	Large for gestational age	26 [8.1%]

Table [2]: Maternal data associated with preterm.

	Measures	Values
Maternal age [year]	mean±SD	27.8 ± 6.3
	Min. – Max.	19 - 40
Residency	Urban	192 [59.6%]
	Rural	130 [40.4%]
Intensive Physical Work	yes	91 [28.3%]
	No	231 [71.7%]
Smoking	yes	22 [6.8%]
	No	300 [93.2%]
Parity	Multiparity	255 [69.9%]
	Nulliparity	97 [30.1%]
Indication for delivery	Maternal diseases	59 [18.3%]
	Obstetric complications	144 [44.7%]
	Isolated spontaneous	119 [37%]
Antenatal care	Proper	279 [86.6%]
	Not	43 [13.4%]
Pregnancy morbidity	Oligo/poly hydramnios	35 [10.9%]
	Uterine bleeding	97 [30.1%]
	Fetal distress	12 [3.7%]
	Premature rupture of membrane	39 [12.1%]
	Hypertension	40 [12.4%]
	Diabetes mellitus	61 [18.9%]
	Urinary tract infection	101 [31.4%]
	Preeclampsia	22 [6.8%]
	Anemia < 10	144 [44.7%]
Type of delivery	Cesarean section	266 [82.6%]
	Normal vaginal delivery	56 [17.4%]
Pregnancy outcome	Single	311 [96.6%]
	Twins	11 [3.4%]

Table [3]: Preterm outcomes.

Variables	Age				P value
	Total [n=322]	34 weeks [n= 108]	35 weeks [n= 105]	36 weeks [n= 109]	
Multiple Congenital abnormalities					
No	312 [96.9%]	98 [90.7%]	105 [100%]	109 [100%]	0.001*
Major	10 [3.1%]	10 [9.3%]	0 [0%]	0 [0%]	
Apgar score 1 min					
Mean ± SD	7.3 ± 1.2	6.8 ± 1.5	7.6 ± 0.7	7.6 ± 1	0.001 ^{ab}
Range	4 – 9	4 – 9	6 – 9	4 – 9	
Apgar score 5 min.					
Mean ± SD	8.6 ± 0.6	8.5 ± 0.8	8.8 ± 0.5	8.7 ± 0.5	0.04 ^{ab}
Range	6 – 10	7 – 9	8 - 10	6 – 9	
NICU admission					
Yes	113 [35.1%]	66 [61.1%]	28 [26.7%]	19 [17.4%]	0.001*
No	209 [64.9%]	42 [38.9%]	77 [73.3%]	90 [82.6%]	
Admission with jaundice					
Yes	54 [16.8%]	26 [24.1%]	16 [15.2%]	12 [11%]	0.03*
No	268 [83.2%]	82 [75.9%]	89 [84.8%]	97 [89%]	
Jaundice needed extensive intervention					
Yes	18 [5.6%]	15 [13.9%]	3 [2.9%]	0 [0%]	0.001*
No	304 [94.4%]	93 [86.1%]	102 [97.1%]	109 [100%]	

Table [4]: Respiratory morbidity of the preterm.

Variables	Age				P value
	Total [n=322]	34 weeks [n= 108]	35 weeks [n= 105]	36 weeks [n= 109]	
Transient tachypnea of newborn	63 [19.6%]	32 [29.6%]	22 [21%]	9 [8.3%]	0.001*
Persistent pulmonary hypertension of the newborn	16 [5%]	16 [14.8%]	0 [0%]	0 [0%]	0.001*
Apnea	14 [4.3%]	14 [13%]	0 [0%]	0 [0%]	0.001*

Respiratory distress syndrome	32 [9.9%]	32 [29.6%]	0 [0%]	0 [0%]	0.001*
Mechanical ventilation	32 [9.9%]	32 [29.6%]	0 [0%]	0 [0%]	0.001*
Continuous positive airway pressure	87 [27%]	58 [53.7%]	22 [21%]	7 [6.4%]	0.001*

Table [5]: Neurological morbidity of the preterm.

Variables	Age				P value
	Total [n=322]	34 weeks [n= 108]	35 weeks [n= 105]	36 weeks [n= 109]	
Seizures	5 [1.6%]	5 [4.6%]	0 [0%]	0 [0%]	0.007*
Hypoxic ischemic encephalopathy	7 [2.2%]	7 [6.5%]	0 [0%]	0 [0%]	0.001*
Hypoxic ischemic encephalopathy grade					
0	315 [97.8%]	101 [93.5%]	105 [100%]	109 [100%]	0.007*
1	2 [0.6%]	2 [1.9%]	0 [0%]	0 [0%]	
2	5 [1.6%]	5 [4.6%]	0 [0%]	0 [0%]	
3	0 [0%]	0 [0%]	0 [0%]	0 [0%]	
Intraventricular hemorrhage	14 [4.3%]	14 [13%]	0 [0%]	0 [0%]	0.001*
Intraventricular hemorrhage grade					
0	308 [95.7%]	94 [87%]	105 [100%]	109 [100%]	0.001*
1	2 [0.6%]	2 [1.9%]	0 [0%]	0 [0%]	
2	7 [2.2%]	7 [6.5%]	0 [0%]	0 [0%]	
3	5 [1.6%]	5 [4.6%]	0 [0%]	0 [0%]	

Table [6]: Infections of the preterm.

Variables	Age				P value
	Total [n=322]	34 weeks [n= 108]	35 weeks [n= 105]	36 weeks [n= 109]	
Sepsis	44 [13.7%]	36 [33.3%]	4 [3.8%]	4 [3.7%]	0.001*
Pneumonia	14 [4.3%]	14 [13%]	0 [0%]	0 [0%]	0.001*
Necrotizing enterocolitis	12 [3.7%]	12 [11.1%]	0 [0%]	0 [0%]	0.001*
Blood transfusion	26 [8.1%]	26 [24.1%]	0 [0%]	0 [0%]	0.001*
Hypoglycemia	17 [5.3%]	17 [15.7%]	0 [0%]	0 [0%]	0.001*
Rehospitalization	32 [9.9%]	21 [19.4%]	11 [10.5%]	0 [0%]	0.001*
Live	305 [94.7%]	91 [84.3%]	105 [100%]	109 [100%]	0.001*

Table [7]: Factors predicting Respiratory morbidity for late preterm infants

Variables	Respiratory morbidity			
	B	Exp [B] / OR	95% CI	P value
Gestational age	-1.8	0.16	95% [0.07-0.3]	0.001*
Birth weight	-0.7	0.4	95% [0.1-1.2]	0.2
Female gender	-1.7	0.17	95% [0.06-0.5]	0.001*
Rural Residency	0.7	2	95% [0.8-4.9]	0.1
Multiparity	0.47	1.6	95% [0.5-4.9]	0.4
Oligo/poly hydramnios	-0.02	1.1	95% [0.3-3.3]	0.9
Uterine bleeding	0.1	1.1	95% [0.4-2.9]	0.7
Premature rupture of membrane	0.94	2.5	95% [0.7-9.2]	0.1
Diabetes mellitus	-1.5	0.2	95% [0.06-0.6]	0.009*
Urinary tract infection	-0.01	0.9	95% [0.3-2.9]	
Cesarean section	-0.7	0.4	95% [0.1-1.8]	0.2

Table [8]: Factors predicting Infections for late preterm infants

Variables	Infections			
	B	Exp [B] / OR	95% CI	P value
Gestational age	-0.3	0.04	95% [0.01-0.4]	0.001*
Birth weight	0.18	1.2	95% [0.2-6.8]	0.8
Female gender	-0.6	0.5	95% [0.1-1.6]	0.2
Rural Residency	0.6	1.9	95% [0.5-5.7]	0.2
Multiparity	-0.2	0.7	95% [0.2-2.7]	0.6
Oligo/poly hydramnios	3.8	46.9	95% [3.9-564.7]	0.002*
Uterine bleeding	0.7	2	95% [0.7-5.9]	0.16
Premature rupture of membrane	-0.7	0.4	95% [0.1-2.1]	0.33
Diabetes mellitus	-1.4	0.2	95% [0.05-0.9]	0.05*
Urinary tract infection	-2.2	0.1	95% [0.02-0.4]	0.001*
Preeclampsia	2.5	12.7	95% [0.9-164]	0.05*
Cesarean section	2.1	8.9	95% [0.9-81.9]	0.05*
Antenatal care	-2.6	0.07	95% [0.01-0.3]	0.001*

Table [9]: Factors predicting mortality for late preterm infants

Variables	Mortality		P value
	Died [n=17]	Live [n=305]	
Respiratory morbidity	10 [58.8%]	68 [22.3%]	0.001*
Neurological morbidity	17 [100%]	4 [1.3%]	0.001*
Infections	17 [100%]	32 [10.5%]	0.001*
Blood transfusion	17 [100%]	9 [3%]	0.001*
Hypoglycemia	17 [100%]	0 [0%]	0.001*

DISCUSSION

Three hundred twenty-two LPI who were delivered at our university hospital were included in our study; this high incidence may be the result of lesser quality prenatal care. The lack of antenatal ultrasound equipment to detect fetal prenatal conditions and the ability to treat these diagnoses, the inability to identify mothers at risk of preterm birth early enough to provide timely care, the lack of multiple micronutrient supplementation, the poor referral systems, the poor health infrastructures, the lack of health supplies, and structural factors in the health system are the causes of this. The delivery and experience of high-quality maternity care at medical facilities are hampered by these technical, interpersonal, resource, and infrastructure issues.

The incidence was reported in multiple studies, in Mansoura University Hospital, Egypt, a study was done by reported an incidence of 24.3% [10]. In South-West Ethiopia the incidence was 25.9% [11].

The sociodemographic factors are considered an important risk factor for LPIs incidence. The mean age of the mothers was 27.8 ± 6.3 years. Most of them [59.6%] were from urban areas. Maternal factors are also considered an important risk factor in which, 13.4% of the mothers didn't have proper ANC, 10.9% had Oligo or poly hydramnios, 30.1% had uterine bleeding, 12.1% had PROM, 12.4% were hypertensive, 18.9% were diabetics, 31.4% had UTI, 6.8% had preeclampsia, and 44.7% were anemic [Hb<10 gm], which is in agreement with Vanin et al. [3] who discovered that 91.3% of pregnant women had a urinary tract infection and 7.5% had a bacterial infection at delivery.

Another important risk factor for the LPIs is the mother employment in which 28.3% are working intensive physical work. in a study by Etil et al. [12] found that mothers who were unemployed being 0.657 times less likely to give birth to premature children than those who had jobs.

Long work hours and temporary contracts were also linked to higher risks of preterm delivery at Mansoura University Hospital in Egypt [AOR=2.36, CI: 1.18–7.78] and [AOR=1.98, CI: 1.72–8.74] respectively [10]. Additionally, the type of work a pregnant woman does may have a different effect depending on whether it is manual or labor-intensive. In a similar vein, a high workload may cause stress for an expectant mother, which may result in pregnancy complications and early birth.

In our study most of the neonates were delivered by CS [82.6%], Tsai et al. [14], as well as Mekic et al. [13], reported a higher CS rate. However, Champion et al. [15] and Aliaga et al. [16] discovered that most late preterm infants [59.6%] were delivered vaginally, particularly when labor was induced.

Poor birth outcomes were also linked to parity [17]. Multipara mothers made up 69.9% of the mothers in our study. In line with a recent study involving 837,226 singleton births in the Netherlands, late-preterm birth was significantly and independently linked to a higher risk of unfavorable neonatal outcomes for nulliparous mothers. According to the aforementioned study, nulliparous mothers had a significantly higher risk of PTB than mothers who had given birth at least once [RR: 1.95, 95% CI: 1.89–2.00 for PTB] [18].

Identifying the risk factors for hospitalization and morbidity is crucial for creating treatment and prevention plans. For late-

preterm infants who may be at higher risk, careful observation and follow-up are necessary. The clinical diagnosis of respiratory distress syndrome [RDS] is made based on the presence of tachypnea, retractions, grunting, increased work of breathing, and other typical X-ray findings. One of the most prevalent respiratory conditions affecting LPIs is still RDS. Lung epithelium immaturity, the epithelial Na⁺ channel [ENaC] transition immaturity, and decreased surfactant production all contribute to the increased risk of TTN in LPI. All newborns are susceptible to NP, but premature newborns, especially those in need of mechanical ventilation, are at an even greater risk. High pulmonary vascular resistance and postpartum hypoxemia are hallmarks of persistent pulmonary hypertension of the newborn [PPHN] ^[121].

In our study most of the respiratory morbidities were in the newborns of gestational age 34 weeks in which TTN represent 19.6%, PPHN represent 5%, Apnea represent 4.3%, and RDS represent 9.9%. Newborns who need mechanical ventilation represent 9.9% and those who need CPAP represent 27%. Numerous studies show that late-preterm infants have high rates of TTN ^[19,20]. A higher rate of respiratory distress may result from lung immaturity-related factors like underdeveloped surfactants and impaired fetal alveolar fluid resorption, even though spontaneous preterm labor promotes fetal lung maturation and enhances pulmonary fluid clearance. The cessation of maternal antibody transmission and immature immune cells makes late-preterm newborns comparatively more vulnerable to lower respiratory tract infections ^[121].

Additionally, late-preterm newborns need more respiratory support than full-term newborns due to physiological and metabolic immaturity. Reducing the incidence of late-preterm births and using antenatal steroids and other appropriate prenatal care can help avoid respiratory issues and save money. The need for respiratory support in these infants can be further reduced by treating infections early and effectively. Four to twelve percent of late preterm infants have apnea of prematurity. According to Engle et al., it is linked to premature breathing control and potential neurodevelopmental deficits ^[9].

According to a study by **Pisani et al.** ^[122] infants born at 34 weeks GA had a > 10-fold higher rate of respiratory morbidity than those born at term, and there was a strong age-related trend in respiratory morbidity regardless of delivery mode.

Up to one-third of newborn deaths globally are caused by sepsis each year. According to the World Health Organization, neonatal sepsis is a serious worldwide health issue, with low- and middle-income nations bearing the brunt of the burden ^[23]. 13.7% of the patients in our study had sepsis, and the literature reports infection percentages ranging from 4.9% to 20.6% ^[24].

Both short-term and long-term neurological morbidity are more likely to occur in LPIs. Preterm newborns are more likely to experience neonatal convulsions and seizures that are solely detected by electrography ^[122]. Preterm infants are primarily

caused by hypoxic-ischemic encephalopathy and intracranial hemorrhage ^[25].

In our study, convulsions occurred in 1.6% of the patients; all of them were from the 34-week age group. IVH represents 4.3% of the total patients, and in 14 of the patients with an age of 34 weeks the percentage rises to 10.9% at 34 weeks GA when we only look at the data on LPIs admitted to the NICU. McIntire et al. reported rates of IVH grade 1 and 2 in LPIs of 0.5% at 34 weeks GA, 0.2% at 35 weeks GA, and 0.06% at 36 weeks GA. According to **Teune et al.** ^[26], who examined 22 studies, intracranial hemorrhage was more common in LPIs. LPIs had a very low rate of IVH grade 3 or 4, but it was still higher than the rate for term neonates [0.01% vs. 0.004%].

The nadir hematocrit is inversely proportional to GA and is lower in LPIs than in term neonates. Some LPIs tolerate normocytic, normochromic anemia well, while others need blood ^[27]. Given the morbidities of preterm infants needing NICU admission, it was not surprising that 8.1% of LPIs in our study had anemia requiring blood transfusions. As gestational age increased from 34 to 36 weeks, the rate of anemia declined.

As regards the mortality rate, it was 5.3%, all of them were from the age group 34 weeks. All of them had neurological morbidity, infection, hypoglycemic and needed blood transfusion. Only 10 cases had respiratory morbidity. Some studies showed higher mortality rates in late-preterm infants [0.3-1.3%] than in term infants [0.0-0.08%] ^[14]. Their results are in agreement with our findings [1.4% vs. 0.0%].

According to a prior study, the rehospitalization rate for late-preterm infants is two to three times higher than that of term infants. This was supported by the fact that 32 cases [9.9%] required rehospitalization. ^[28]

One of the **main limitations** of the present study is the absence of control group as the presence of a control group will allow for the identification of factors associated with early delivery, not just a description of the characteristics of late preterm infants. Another limitation is that more than 90% of the population studied had low income, which did not allow social comparisons, as other studies have done.

The strength of the study is that all infants were delivered and cared for at the same academic institution, and it provides insight into the maternal risk factors

Conclusion: Identifying the predictors of adverse short-term outcomes in late preterm infants is of crucial importance for informing and evaluating clinical practices and guidelines aimed at reducing infant morbidity and mortality

Financial and non-financial activities and relationships of interest: None

REFERENCES

1. Bick D. Born too soon: the global issue of preterm birth. *Midwifery*. 2012 Aug;28[4]:341-2. doi: 10.1016/j.midw.2012.06.010.
2. Quinn JA, Munoz FM, Gonik B, Frau L, Cutland C, Mallett-Moore T, et al; Brighton Collaboration Preterm Birth Working Group. Preterm birth: Case definition & guidelines for data collection, analysis, and presentation of immunisation safety data. *Vaccine*. 2016 Dec 1;34[49]:6047-6056. doi: 10.1016/j.vaccine.2016.03.045.
3. Vanin LK, Zatti H, Soncini T, Nunes RD, Siqueira LBS. MATERNAL AND FETAL RISK FACTORS ASSOCIATED WITH LATE PRETERM INFANTS. *Rev Paul Pediatr*. 2019 Nov 25;38:e2018136. doi: 10.1590/1984-0462/2020/38/2018136.
4. Araújo BF, Zatti H, Madi JM, Coelho MB, Olmi FB, Canabarro CT. Analysis of neonatal morbidity and mortality in late-preterm newborn infants. *J Pediatr [Rio J]*. 2012 May;88[3]:259-66. doi: 10.2223/jped.2196.
5. Leone A, Ersfeld P, Adams M, Schiffer PM, Bucher HU, Arlettaz R. Neonatal morbidity in singleton late preterm infants compared with full-term infants. *Acta Paediatr*. 2012 Jan;101[1]:e6-10. doi: 10.1111/j.1651-2227.2011.02459.x.
6. Natarajan G, Shankaran S. Short- and Long-Term Outcomes of Moderate and Late Preterm Infants. *Am J Perinatol*. 2016 Feb;33[3]:305-17. doi: 10.1055/s-0035-1571150.
7. McGowan JE, Alderdice FA, Holmes VA, Johnston L. Early childhood development of late-preterm infants: a systematic review. *Pediatrics*. 2011 Jun;127[6]:1111-24. doi: 10.1542/peds.2010-2257.
8. Ruth CA, Roos N, Hildes-Ripstein E, Brownell M. 'The influence of gestational age and socioeconomic status on neonatal outcomes in late preterm and early term gestation: a population based study'. *BMC Pregnancy Childbirth*. 2012 Jun 29;12:62. doi: 10.1186/1471-2393-12-62.
9. Engle WA, Tomashek KM, Wallman C; Committee on Fetus and Newborn, American Academy of Pediatrics. "Late-preterm" infants: a population at risk. *Pediatrics*. 2007 Dec;120[6]:1390-401. doi: 10.1542/peds.2007-2952. Erratum in: *Pediatrics*. 2008 Feb;121[2]:451. PMID: 18055691.
10. ElGilany A, ElKhawaga G, Ghanem A. Incidence and occupational risk factors of preterm delivery among working mothers: a single center study in Egypt. *TAF Prev Med Bull*. 2016;15[3]:199. Doi: 10.5455/pmb.1-1441637597
11. Bekele I, Demeke T, Dugna K. Prevalence of Preterm Birth and its Associated Factors among Mothers Delivered in Jimma University Specialized Teaching and Referral Hospital, Jimma Zone, Oromia Regional State, South West Ethiopia. *J Women's Heal Care*. 2017;06[1]:356. DOI: 10.4172/2167-0420.100035
12. Etal T, Opio B, Odur B, Lwanga C, Atuhaire L. Risk factors associated with preterm birth among mothers delivered at Lira Regional Referral Hospital. *BMC Pregnancy Childbirth*. 2023 Nov 23;23[1]:814. doi: 10.1186/s12884-023-06120-4. Erratum in: *BMC Pregnancy Childbirth*. 2023 Dec 11;23[1]:845. doi: 10.1186/s12884-023-06163-7.
13. Mekic N, Selimovic A, Cosickic A, Mehmedovic M, Hadzic D, Zulic E, Mustafic S, Serak A. Predictors of adverse short-term outcomes in late preterm infants. *BMC Pediatr*. 2023 Jun 17;23[1]:298. doi: 10.1186/s12887-023-04112-z.
14. Tsai ML, Lien R, Chiang MC, Hsu JF, Fu RH, Chu SM, Yang CY, Yang PH. Prevalence and morbidity of late preterm infants: current status in a medical center of Northern Taiwan. *Pediatr Neonatol*. 2012 Jun;53[3]:171-7. doi: 10.1016/j.pedneo.2012.04.003.
15. Champion V, Durrmeyer X, Dassieu G. [Short-term respiratory outcome of late preterm newborn in a center of level III]. *Arch Pediatr*. 2010 Jan;17[1]:19-25. French. doi: 10.1016/j.arcped.2009.10.002.
16. Aliaga S, Boggess K, Ivester TS, Price WA. Influence of neonatal practice variation on outcomes of late preterm birth. *Am J Perinatol*. 2014 Sep;31[8]:659-66. doi: 10.1055/s-0033-1356484.
17. Schimmel MS, Bromiker R, Hammerman C, Chertman L, Ioscovich A, Granovsky-Grisaru S, Samueloff A, Elstein D. The effects of maternal age and parity on maternal and neonatal outcome. *Arch Gynecol Obstet*. 2015 Apr;291[4]:793-8. doi: 10.1007/s00404-014-3469-0.
18. Koullali B, van Zijl MD, Kazemier BM, Oudijk MA, Mol BWJ, Pajkrt E, Ravelli ACJ. The association between parity and spontaneous preterm birth: a population based study. *BMC Pregnancy Childbirth*. 2020 Apr 21;20[1]:233. doi: 10.1186/s12884-020-02940-w.
19. Tan JH, Poon WB, Lian WB, Ho SK. A Comparison of the Short-term Morbidity and Mortality Between Late Preterm and Term Newborns. *Ann Acad Med Singap*. 2014 Jul;43[7]:346-54. PMID: 25142470.
20. Visrathan NK, Agarwal P, Sriram B, Rajadurai VS. Neonatal Outcome of the Late Preterm Infant [34 to 36 Weeks]: The Singapore Story. *Ann Acad Med Singap*. 2015 Jul;44[7]:235-43. PMID: 26377057.
21. Holt PG, Jones CA. The development of the immune system during pregnancy and early life. *Allergy*. 2000 Aug;55[8]:688-97. doi: 10.1034/j.1398-9995.2000.00118.x. PMID: 10955693.
22. Pisani F, Prezioso G, Spagnoli C. Neonatal seizures in preterm infants: A systematic review of mortality risk and neurological outcomes from studies in the 2000's. *Seizure*. 2020 Feb; 75:7-17. doi: 10.1016/j.seizure.2019.12.005.
23. Popescu CR, Cavanagh MMM, Tembo B, Chieme M, Lufesi N, Goldfarb DM, Kissoon N, Lavoie PM. Neonatal sepsis in low-income countries: epidemiology, diagnosis and prevention. *Expert Rev Anti Infect Ther*. 2020 May;18[5]:443-452. doi: 10.1080/14787210.2020.1732818.
24. McIntire DD, Leveno KJ. Neonatal mortality and morbidity rates in late preterm births compared with births at term. *Obstet Gynecol*. 2008 Jan;111[1]:35-41. doi: 10.1097/01.AOG.0000297311.33046.73.
25. Glass HC, Shellhaas RA, Tsuchida TN, Chang T, Wusthoff CJ, Chu CJ, Cilio MR, Bonifacio SL, Massey SL, Abend NS, Soul JS; Neonatal Seizure Registry study group. Seizures in Preterm Neonates: A Multicenter Observational Cohort Study. *Pediatr Neurol*. 2017 Jul; 72:19-24. doi: 10.1016/j.pediatrneurol.2017.04.016.
26. Teune MJ, Bakhuizen S, Gyamfi Bannerman C, Opmeer BC, van Kaam AH, van Wassenae AG, Morris JM, Mol BW. A systematic review of severe morbidity in infants born late preterm. *Am J Obstet Gynecol*. 2011 Oct;205[4]:374.e1-9. doi: 10.1016/j.ajog.2011.07.015.
27. Olivier F, Nadeau S, Caouette G, Piedboeuf B. Association between Apnea of Prematurity and Respiratory Distress Syndrome in Late Preterm Infants: An Observational Study. *Front Pediatr*. 2016 Sep 26;4:105. doi: 10.3389/fped.2016.00105.
28. Tomashek KM, Shapiro-Mendoza CK, Weiss J, Kotelchuck M, Barfield W, Evans S, Naninni A, Declercq E. Early discharge among late preterm and term newborns and risk of neonatal morbidity. *Semin Perinatol*. 2006 Apr;30[2]:61-8. doi: 10.1053/j.semperi.2006.02.003.

IJMA



INTERNATIONAL JOURNAL OF MEDICAL ARTS

Volume 7, Issue 10 (October 2025)



<http://ijma.journals.ekb.eg/>

P-ISSN: 2636-4174

E-ISSN: 2682-3780