



Phytochemical Profiling and GC-MS-Based Characterization of Bioactive Compounds from Egyptian Coastal Seaweeds

Mona M. Ismail¹, Mostafa E. Elshobary^{2*}, Ahmed S. El Newehy³, and Saly F. Gheda²

¹ National Institute of Oceanography and Fisheries, Cairo 4262110, Egypt

² Botany and Microbiology Department, Faculty of Science, Tanta University, Tanta 31527, Egypt

³ Department of Community Health Sciences, College of Applied Medical Sciences, King Saud University, Riyadh 11433, Saudi Arabia

*Corresponding Author: mostafa_elshobary@science.tanta.edu.eg

ARTICLE INFO

Article History:

Received: June 5, 2025

Accepted: Aug. 30, 2025

Online: Sep. 15, 2025

Keywords:

Seaweeds,
Egyptian coasts,
Phenolic compounds,
GC-MS,
Biological activity

ABSTRACT

This study investigated the phytochemical composition of 10 different species of marine macroalgae collected from the Mediterranean and Red Seas, Egypt. The collected species taxonomically belong to three major groups: Ochrophyta (3 species), Rhodophyta (5 species), and Chlorophyta (2 species). Phytochemical extraction of each seaweed, using water, methanol, ethanol and acetone, was carried out. Qualitative and quantitative analyses were conducted to detect phenolics, flavonoids, tannins, steroids, terpenoids, and saponins. Among the studied species, *Sargassum cinereum* exhibited the highest phytochemical yield, particularly in ethanolic extracts, while Rhodophyta species generally showed lower phytochemical contents. Quantitative analyses revealed significant variation in total phenolic, flavonoid, and tannin contents depending on the algal species and extraction solvent. Gas chromatography–mass spectrometry (GC-MS) profiling identified numerous bioactive compounds, with palmitic acid and its methyl ester, and stearic acid methyl ester are being the predominant fatty acids. The findings highlight the substantial potential of Egyptian coastal seaweeds as a sustainable source of biologically active compounds for pharmaceutical, nutraceutical, and environmental applications. This work emphasizes the need to explore marine algal resources in Egypt for novel bioactive compounds and suggests that the solvent type significantly influences the extraction efficiency of different phytochemicals.

INTRODUCTION

Seaweeds produce a wide range of metabolites (phenol, tannins, flavonoids) to protect themselves from biotic and abiotic factors like herbivory or mechanical aggression at sea because they are frequently exposed to extreme conditions of the environment without showing any signs of damage (Gómez-Guzmán *et al.*, 2018).

Phytochemical Profiling and GC-MS-Based Characterization of Bioactive Compounds from Egyptian Coastal Seaweeds

Ochrophyta have been shown to provide a variety of health benefits. The dominant brown species, such as *Polycladia myrica*, *Dictyota spiralis*, and *Sargassum euryphllum* from the Red Sea, Egypt, contain various phenolic contents that have the potential to treat a wide range of serious diseases. These compounds may also be valuable in the pharmaceuticals, cosmeceuticals, and functional food industries (Ismail *et al.*, 2022). The brown alga *Padina tetrastromatica* is a member of the Dictyotales order and is found along the Egyptian coast. *P. tetrastromatica* contains numerous chemical compounds that are responsible for the plant's diverse pharmacological and traditional qualities (Yende *et al.*, 2020). Alkaloids, triterpenoids, steroids, tannins, flavonoids, and saponins are all positively detected in *Tricleocarpa fragile* exhibiting antibacterial activity against *V. harveyi* bacterium (Singkoh *et al.*, 2019). The Egyptian marine red macroalga *Digenea simplex* has a wide variety of phytochemicals with distinct bioactivities (El-Rafie *et al.*, 2023). *D. simplex* chloroform extract's cytotoxic and antioxidant properties have not received enough attention, and the majority of the species' published data are from waters other than the beaches of the Red Sea in Egypt (Sobuj *et al.*, 2021; Ghosh *et al.*, 2022).

Ulva linza and *U. fasciata* are common green seaweed that are extensively found throughout the Mediterranean coast, Egypt (Mohy El-Din & Alagawany, 2019; El Zokm *et al.*, 2021). According to the study conducted by Mohy El-Din and Alagawany (2019), *U. linza* has been reported as a promising antioxidant resource. Therefore, this species might be useful as natural anticoagulants and antioxidants. Besides, the most studied species like *Ulva fasciata*, *U. linza*, *U. lactuca*, *Corallina officinalis*, and *Sargassum vulgare* from the Alexandria coast in Egypt may be considered as alternatives and supplements to help reduce the risk of obesity. This is because they have a low energy content (≤ 1.76 KJ/g DW) and caloric values (≤ 0.357 kcal), which are lower than the standard levels recommended for women (2 kcal) and men (2.5 kcal) (Ismail *et al.*, 2017).

Phenolic compounds are well-established bioactive compounds (BCs) that are significant secondary metabolites of both plants and algae. They have been shown to have anti-inflammatory, anti-cancer, and antioxidant properties (Getachew *et al.*, 2020). Three categories constitute the majority of algal sources: phenolic acids, phenylpropanoids, and flavonoids (Carreira-Casais *et al.*, 2021).

Egypt needs to take advantage from a high richness of marine macroalgal species, owing to the presence of macroalgae near its coasts in the Mediterranean Sea, Red Sea, and Suez Canal. This wealth of marine macroalgae serves as an important source of bioactive chemical compounds (Ahmed *et al.*, 2023).

The objective of this research was to evaluate the variation in phytochemical contents and concentrations of several seaweeds extracts from the Egyptian coast. Additionally, the study aimed to identify the bioactive compounds in the collected ethanolic extracts using gas chromatography-

mass spectrometry for future applications in medicine, in pharmacology, dietary supplements, cosmetics or in the food industries.

MATERIALS AND METHODS

1. Collection and identification of macroalgae samples

Ten species of seaweeds (5 Rhodophyta, 3 Ochrophyta and 2 Chlorophyta) were handpicked at depths of 0.2m or less for Chlorophyta and 1m for Rhodophyta and Ochrophyta from the submerged rocks at Abu Qir (N 31° 19', E 030° 03') and from the front of the National Institute of Oceanography and Fisheries (NIOF), Hurghada, Red Sea (latitudes 27° 17' 13" N and longitudes 33° 46' 21"E) during the summer season of 2023. The collected samples were cleaned with seawater to remove sand, conditioned in plastic bags containing seawater to prevent evaporation, and kept on ice until transported to the laboratory (Thomas & Harrison, 1987). All samples were brought to the Taxonomy and Biodiversity Laboratory, NIOF, Alexandria in plastic bags and were taxonomically identified. The seaweed specimens were cleaned well from epiphytes and rock debris. Thereafter, they were given a quick freshwater rinse to remove surface salts, and then a part of each species was preserved in 5% formalin in seawater. The remaining cleaned seaweeds were spread on blotting paper at 25 to 30°C to remove excess water and were then shade-dried for about 4-7 days at room temperature until a constant weight was obtained. The shade-dried material was crushed in an electric mixer to obtain a coarse powder (Ibtissam *et al.*, 2009) that was stored in airtight dark bottles at -20°C. The names of the species were confirmed according to the Algae Base website (Guiry & Guiry, 2022).

2. Extraction of phytochemical compounds

The extraction of bioactive compounds was performed using the method described by Sarker (2008). Five grams of dried powder from each algae species was extracted using four different solvents: water, methanol, ethanol, and acetone. The extraction was carried out using a shaker for 24 h at room temperature (25±2°C), with a solvent-to-algal sample ratio of 15:1. After extraction, the mixtures were filtered to remove the solid residue using Whatman No. 1 filter paper. The filtrates were then evaporated under reduced pressure using a rotary evaporator to obtain the crude extracts. The resulting extracts were weighed and stored at 4°C until further analysis.

3. Qualitative analysis for bioactive compounds in algal extracts

The standard procedure outlined by Savithramma *et al.* (2011) was followed to assess the presence or absence of phytochemical content, including tannins, flavonoids, phenols, terpenoids, saponins, and steroids. For steroids, 0.5mL of the algal extract was mixed with 2mL of chloroform and 1 mL of sulfuric acid (H₂SO₄). Steroids are present when a reddish-brown ring forms at the contact.

Flavonoids identification was performed by mixing 1mL of 2N sodium hydroxide (NaOH) with 2mL of algal extract. The formation of yellow color indicates the presence of flavonoids.

In order to identify the phenols, two milliliters of distilled water and a few drops of 10% ferric chloride were added to one milliliter of the algal extract, when phenols are present, a blue or green tinge is formed. For qualitative estimation of tannins, 1mL ferric chloride (5%) was mixed to 1mL of the algal extract. When tannins are present, dark blue or greenish-black color are formed.

Salkowski test for terpenoids, five milliliters of each extract were mixed with two milliliters of chloroform and three milliliters of concentrated H₂SO₄. The reddish-brown color was produced at the contact due to the presence of terpenoids (**Harborne, 1998**). Frothing test for saponin estimation; after treating around 3 milliliters of each extract with 3 milliliters of distilled water, the sample was rapidly shaken for approximately one minute to produce a stable, long-lasting froth. By shaking the mixture vigorously with three drops of olive oil, the presence of saponins in the extracts was determined by observing the production of an emulsion (**Trease & Evans, 1989**).

4. Quantitative analysis for bioactive compounds

4.1 Estimation of total flavonoid content

The total flavonoid content was determined using the aluminum chloride colorimetric method as described by **Chang et al. (2020)**. Briefly, 0.5mL of each extract was mixed with 1.5mL of methanol, 0.1mL of 10% aluminum chloride, 0.1mL of 1M potassium acetate, and 2.8mL of distilled water. The mixture was incubated at room temperature for 30 minutes, and the absorbance was measured at 415nm using a UV-visible spectrophotometer. Quercetin was used as a standard, and the results were expressed as mg of quercetin equivalents (QE) per gram of dry weight (DW) of the sample.

4.2 Estimation of total phenolic content

The total phenolic content was determined using the Folin-Ciocalteu method described by **Taga et al. (1984)**. Briefly, 100μL of each extract was mixed with 2mL of 2% Na₂CO₃ and the mixture was allowed to stand for 2 minutes at room temperature. Then, 100μL of 50% Folin-Ciocalteu phenol reagent was added. The mixture was incubated for 30 minutes at room temperature in darkness, and the absorbance was measured at 720 nm using a UV-visible spectrophotometer. Gallic acid was used as the standard, and the results were expressed as mg of gallic acid equivalents (GAE) per gram of dry weight (DW) of the sample.

4.3 Estimation of total tannin content

The total tannin content was determined using the method described by **Julkunen-Tiitto (1985)**. In this procedure, 50μL of each extract was mixed with 1.5mL of 40% vanillin solution in methanol, followed by the addition of 750μL of concentrated HCl. The mixture was shaken vigorously and was left to stand at room temperature for 20 minutes in darkness. The absorbance was measured at 500nm using a UV-visible spectrophotometer. Catechin was used as a standard,

and the results were expressed as mg of catechin equivalents (CE) per gram of dry weight (DW) of the sample.

4.4 Gas chromatography-mass spectrum (GC-MS)

The ethanol extracts of 10 algal samples were examined utilizing an Agilent GC 7890A gas chromatograph used with a 5975C triple-axis detector single quadrupole mass spectrometer. Chromatographic separation utilized an Agilent HP-5MS column (30 m \times 0.25mm \times 0.25 μ m film thickness), employing high-purity helium as the carrier gas at a steady flow rate of 1mL/ min. The injector temperature was sustained at 250°C, employing a splitless injection mode with a 20:1 split ratio. The ion source temperature of the mass spectrometer was established at 230°C, while the quadrupole temperature was kept at 150°C. The oven temperature protocol commenced at 40°C (maintained for 1 minute), then escalated to 150°C at a rate of 20°C/min (sustained for 1 minute) and subsequently increased to 300°C at the same rate, held for another 1 minute. A 1 μ L sample was injected, and the mass spectrometer operated within a scan range of 50– 600amu at an electron energy of 70eV. The solvent delay was established at 3 minutes. Compounds were identified by comparing their mass spectra with the NIST 2008 collection (National Institute of Standards and Technology). The cumulative analytical duration for each sample was 16 minutes.

5. Statistical analysis

Data represented as mean \pm standard deviation of three replicates was employed to present the results. The statistical analysis was conducted using SPSS software (version 23, IBM Corp., Armonk, NY, USA). All experiments were performed with three independent replicates. Analysis of variance (ANOVA) was used to determine the statistical significance of the differences among the experimental groups. A post hoc analysis was carried out using the Tukey test to compare the means between the groups. Statistical significance was set at $P < 0.05$

RESULTS

1. Identification of the collected macroalgae species from Egyptian Coast

The collected seaweeds were identified into the following species *Sargassum cinereum* J. Agardh, *Padina tetrastrum* Hauck, *Turbinaria decurrens* Bory from Ochrophyta, *Tricleocarpa fragilis* (Linnaeus) Huisman & Townsend, *Gracilaria armata* (Agardh) Greville, *Corallina officinalis* Linnaeus, *Amphiroa rigida* Lamouroux, *Digenea simplex* (Wulfen) Agardh from Rhodophyta, *Ulva fasciata* Delile, and *Ulva linza* Linnaeus from Chlorophyta (Fig. 1). The taxonomical identification of the collected algal species is represented in Table (1).

Phytochemical Profiling and GC-MS-Based Characterization of Bioactive Compounds from Egyptian Coastal Seaweeds




Fig. 1. Photos of the studied seaweeds: (a) *Sargassum cinereum* (b) *Padina tetrastromatica* (c) *Turbinaria decurrens* (d) *Tricleocarpa fragilis* (e) *Gracilaria armata* (f) *Corallina officinalis* (g) *Amphiroa rigida* (h) *Digenea simplex* (i) *Ulva fasciata* (j) *U. linza* Linnaeus.

Table 1. Taxonomical identification and the collection sites of the selected seaweeds

Seaweed	Empire	Kingdom	Subkingdom	Phylum	Class	Subclass	Order	Family	Genus	Collection Site
<i>Sargassum cinereum</i>	Eukaryota	Chromista	Harosia	Ochrophyta	Phaeophyceae	Fucophycidae	Fucales	Sargassaceae	<i>Sargassum</i>	Hurghada, Red Sea
<i>Padina tetrastrum</i>	Eukaryota	Chromista	harosia	Ochrophyta	Phaeophyceae	Dictyotophycidae	Dictyotales	Dictyotaceae	<i>Padina</i>	Hurghada, Red Sea
<i>Turbinaria decurrens</i>	Eukaryota	Chromista	Harosia	Ochrophyta	Phaeophyceae	Fucophycidae	Fucales	Sargassaceae	<i>Turbinaria</i>	Hurghada, Red Sea
<i>Tricleocarpa fragilis</i>	Eukaryota	Plantae	Biliphyta	Rhodophyta	Florideophyceae	Nemaliophycidae	Nemaliales	Galaxauraceae	<i>Tricleocarpa</i>	Hurghada, Red Sea
<i>Gracilaria armata</i>	Eukaryota	Plantae	Biliphyta	Rhodophyta	Florideophyceae	Rhodymeniophycidae	Gracilariales	Gracilariaceae	<i>Gracilaria</i>	Abo Qir, Mediterranean Sea
<i>Corallina officinalis</i>	Eukaryota	Plantae	Biliphyta	Rhodophyta	Florideophyceae	Corallinophycidae	Corallinales	Corallinaceae	<i>Corallina</i>	Abo Qir, Mediterranean Sea
<i>Amphiroa rigida</i>	Eukaryota	Plantae	Biliphyta	Rhodophyta	Florideophyceae	Corallinophycidae	Corallinales	Lithophyllaceae	<i>Amphiroa</i>	Abo Qir, Mediterranean Sea
<i>Digenea simplex</i>	Eukaryota	Plantae	Biliphyta	Rhodophyta	Florideophyceae	Corallinophycidae	Ceramiales	Rhodymeniaceae	<i>Digenea</i>	Hurghada, Red Sea
<i>Ulva fasciata</i>	Eukaryota	Plantae	Viridiplantae	Chlorophyta	Ulvophyceae		Ulvales	Ulvaceae	<i>Ulva</i>	Abo Qir, Mediterranean Sea
<i>Ulva linza</i>	Eukaryota	Plantae	Viridiplantae	Chlorophyta	Ulvophyceae		Ulvales	Ulvaceae	<i>Ulva</i>	Abo Qir, Mediterranean Sea

2. The yield of phytochemical compounds

Phytochemical compounds were extracted from collected species using different solvents including ethanol, acetone, water, and methanol. The obtained results represented in Fig. (2) indicate that the *Ochrophyta* species recorded the highest phytochemical yield with ethanolic extract as *S. cinereum* recorded 0.94 g/g DW, and *T. decurrens* recorded 0.815 g, also *Ulva linza* water extract recorded 0.904 g. Among all the studied algal groups, Rhodophyta algal group recorded the lowest phytochemical amount. Among most tested algal extracts, methanol recorded the lowest phytochemical yield, while water extract showed the highest yield (Fig. 2). Phytochemical amounts were significantly different ($P < 0.05$) between algal species and ranged from 0.014g in *G. armata* methanol extracts to 0.94g in *S. cinereum* ethanol extract.

Phytochemical Profiling and GC-MS-Based Characterization of Bioactive Compounds from Egyptian Coastal Seaweeds

The highest levels of phytochemical yield in the Chlorophyta algal group were observed in water extracts of *U. linza* and *U. fasciata* 0.904 and 0.544g, respectively. The lowest levels of phytochemical compounds were detected in methanolic extracts of *U. linza* (0.051g) and *U. fasciata* (0.041g). Ethanol extract of *U. linza* and *U. fasciata* recorded phytochemical components (0.115 and 0.216g, respectively), while acetonic extract recorded 0.084 and 0.179g, respectively.

In the Rhodophyta group, the aqueous extract recorded the highest phytochemical yield with *T. fragilis* (0.431g), *A. rigida* (0.332g) and *D. simplex* (0.328g). The ethanol extract of *T. fragilis* recorded (0.246g), while its acetone extract recorded (0.23g) phytochemical compounds. The methanolic extract of the Rhodophyta group contained 0.039, 0.048, 0.041, 0.014, and 0.058g of phytochemical compounds with *C. officinalis*, *D. simplex*, *A. rigida*, *G. armata*, and *T. fragile*, respectively.

Acetone extract recorded (0.034, 0.073, 0.091, and 0.059g) phytochemical yield with *C. officinalis*, *D. simplex*, *A. rigida*, and *G. armata*, respectively. While ethanol extract recorded (0.058, 0.1, and 0.106g) phytochemical amount with *C. officinalis*, *D. simplex*, *A. rigida*, respectively, water extract of *C. officinalis*, and *G. armata* recorded low yield (0.09 and 0.046g) of phytochemical compounds, respectively.

The yield of phytochemical compounds was estimated in the Ochrophyta group. *S. cinereum* had phytochemical compounds 0.941g with ethanol, 0.713g with acetone, and 0.772 with water extract. *T. decurrens* had phytochemical yield (0.815, 0.689, and 0.721g) with ethanol, acetone, and water extract, respectively. The yield of phytochemical compounds of *P. tetrastromatica* were (0.546, 0.422, and 0.656 g) recorded with ethanol, acetone, and water extract, respectively. Among the different solvents used, the methanol extract recorded the lowest phytochemical amount of 0.051, 0.048, and 0.047 g with Ochrophyta species including *S. cinereum*, *P. tetrastromatica* and *T. decurrens*, respectively.

3. Qualitative estimation of phytochemical compounds in various seaweed extracts

Preliminary phytochemical screenings of six different chemical compounds (flavonoids, phenols, tannins, saponins, steroids, and terpenoids) were tested in the ten different collected seaweed. In the present study, phytochemical screening was evaluated using different solvents; ethanol, methanol, acetone, and distilled water in different seaweed. Water extract of *G. armata*, *A. rigida*, *U. fasciata*, and *U. linza* did not show any positive result for the presence of any phytochemical compounds, as illustrated in Table (2). The methanol extract of *S. cinereum* and *G. armata* did not show any positive result for their presence of flavonoids and tannin. Similarly, the methanol extract of *P. tetrastromatica* and *U. fasciata* did not show any positive results for the presence of flavonoids and tannins, respectively.

Among the four different solvents used extracts, acetone extract showed the presence of the maximum number of phytochemical compounds, followed by ethanol extracts and the other two solvents (acetone and ethanol) extracted from all tested phytochemical compounds.




Fig. 2. Phytochemical compounds yield % of tested seaweed species using various solvents. Data points represent means \pm standard deviation ($n = 3$)

Table 2. Preliminary phytochemical screening of different extracts of the studied species

Seaweed	Extract	Flavonoids	Phenol	Tannin	Saponin	Steroids	Terpenoids
<i>S. cinereum</i>	Ethanol	+	++	+	+	+	++
	Methanol	+	+	+	+	+	+
	Water	+	+	+	-	+	+
	Acetone	+	+++	+	+	++	++
<i>P. tetrastrumatica</i>	Ethanol	+	++	+	+	+	+
	Methanol	+	+	+	+	+	+

Phytochemical Profiling and GC-MS-Based Characterization of Bioactive Compounds from Egyptian Coastal Seaweeds

	Water	+	+	+	-	+	+
	Acetone	++	++	++	+	++	++
<i>T. decurrens</i>	Ethanol	++	+++	+	+	+	++
	Methanol	+	+	+	+	+	+
	Water	+	++	+	-	+	+
	Acetone	++	+++	+	+	++	++
<i>T. fragilis</i>	Ethanol	+	+	+	+	+	+
	Methanol	+	+	+	+	+	+
	Water	+	+	+	-	+	+
	Acetone	++	++	+	+	++	++
<i>G. armata</i>	Ethanol	+	++	+	+	+	+
	Methanol	+	+	+	+	+	+
	Water	+	+	+	-	-	-
	Acetone	++	++	+	+	++	++
<i>C. officinalis</i>	Ethanol	+	+	+	+	+	+
	Methanol	+	+	+	+	+	+
	Water	+	+	+	-	+	+
	Acetone	+	+	+	+	++	++
<i>A. rigida</i>	Ethanol	++	+	++	+	+	+
	Methanol	+	+	+	+	+	+
	Water	+	+	+	-	-	-
	Acetone	+	+	+	+	++	++
<i>D. simplex</i>	Ethanol	+	+	+	+	+	+
	Methanol	+	+	+	+	+	+
	Water	++	++	+	-	+	+

	Acetone	++	++	+	+	++	++
<i>U. fasciata</i>	Ethanol	++	++	+	+	+	++
	Methanol	+	+	+	+	+	+
	Water	+	+	+	-	-	-
	Acetone	++	++	+	+	++	++
<i>U. linza</i>	Ethanol	++	++	++	+	+	++
	Methanol	+	+	+	+	+	+
	Water	+	+	+	-	-	-
	Acetone	++	++	+	+	++	++

+++ (very high), ++ (medium), + (low), and – (not detected).

4. Quantitative estimation of phytochemical compounds

Total phenolic content (TPC), total tannin content (TTC), and total flavonoid content (TFC) of the tested species were estimated in different solvents (Ethanol, acetone, water, and methanol).

4.1. Total phenolic content (TPC) of different seaweed extracts

The total phenolic content varied significantly ($P < 0.05$) among the different species, ranging from 0.04mg/ g DW in the water extract of *A. rigida* to 1.02mg/ g DW in the acetone extract of *D. simplex*. The maximum amounts of TPC in different algae extracts were found in the Ochrophyta algal group acetone extraction and ranged from 0.19mg/ g DW in *D. simplex* to 1.02mg/ g DW in *T. decurrens*, as represented in Fig. (3).

In the algal group Chlorophyta (*U. fasciata*, *U. linza*); *U. linza* recorded the highest amount of total phenolic (0.49 ± 0 mg/g DW) with ethanolic extract, followed by *U. fasciata* with acetone, and ethanolic extract with total amounts of phenolic (0.47 ± 0.02 mg/g DW), (0.48 ± 0.01 mg/g DW), respectively. Water extracts showed the lowest TPC in both Chlorophyta species (*U. fasciata*, *U. linza*) at 0.05 ± 0 mg/ g DW. Methanol extracts of Chlorophyta algae recorded total phenolic contents of 0.19 ± 0 , and 0.23 ± 0 mg/ g DW with *U. linza* and *U. fasciata*, respectively.

In the Rhodophyta group, *G. armata* acetone extract recorded the highest TPC (0.55 ± 0 mg/ g DW), while *G. armata* ethanolic extract recorded (0.49 ± 0.01 mg/g DW) total amount of phenolics. *T. fragile* (Linnaeus) ethanolic extract recorded (0.38 ± 0 mg/ g DW) total amount of phenolic. *A. rigida* acetone extract recorded (0.31 ± 0 mg/ g DW) total amount of phenolics. Water and methanolic extract recorded the lowest TPC (0.12 ± 0 , 0.19 ± 0 , 0.04 ± 0 , 0.09 ± 0 , and 0.11 ± 0 mg/

Phytochemical Profiling and GC-MS-Based Characterization of Bioactive Compounds from Egyptian Coastal Seaweeds

g DW) for water extract; 0.16 ± 0 , 0.11 ± 0 , 0.09 ± 0 , 0.19 ± 0 , and 0.13 ± 0 mg/g DW for methanol with *C. officinalis*, *D. simplex*, *A. rigida*, *G. armata*, and *T. fragile*, respectively. The ethanolic extract recorded (0.23, 0.21, 0.14, and 0.23 mg/g DW) total amount of phenolic with *C. officinalis*, *D. simplex*, *A. rigida*, and *T. fragile*, respectively. While the acetone extract recorded (0.2, 0.19, and 0.31 mg/g DW) total amount of phenolics with *C. officinalis*, *D. simplex*, *A. rigida*, respectively.

In the Ochrophyta group, the highest TPC was recorded in acetonic extract of *T. decurrens* (1.02 ± 0.03 mg/g DW) followed by the ethanolic extract of *T. decurrens* (0.71 mg/g DW) and the acetonic extract of *S. cinereum* (0.71 mg/g DW). The lowest TFC was detected in methanolic extract of *P. tetrastromatica* (0.07 mg/g DW) and *T. decurrens* (0.08 mg/g DW), respectively, followed by the water extract for 0.17, 0.18, and 0.31 mg/g DW for the same species. On the other hand, *P. tetrastromatica* acetone extract recorded 0.4 mg/g DW. While, *S. cinereum*, and *P. tetrastromatica* ethanolic extract recorded 0.41, and 0.37 mg/g DW.




Fig. 3. Total phenolic content (TPC) of the tested seaweed species using various solvents. Data points represent means \pm standard deviation ($n = 3$).

4.2. Total flavonoid content (TFC) of different algal extracts

Within the Chlorophyta algal group, the acetonic extract of *U. linza* showed the highest level of TFC (10.69 mg/g DW), followed by the ethanol extract of *U. fasciata* (7.57 mg/g DW), and the acetonic extract of *U. fasciata* (6.85 mg/g DW). Among the various solvents used for extraction, the aqueous and methanolic extracts exhibited the lowest TFC values: 0.16 and 0.13 mg/g DW for *U. fasciata*, and 0.92 and 0.88 mg/g DW for *U. linza*, respectively.

The acetonic extract of *A. rigida* recorded the highest TFC (14.95mg/ g DW) among the Rhodophyta species, followed by the acetonic extracts of *G. armata* and *T. fragile* (12.23 and 7.16mg/ g DW, respectively). Conversely, water and methanol extracts showed the lowest TFC values across all Rhodophyta species. The water extracts recorded TFC values of 0.18, 0.28, 0.10, 0.07, and 0.27mg/ g DW for *C. officinalis*, *D. simplex*, *A. rigida*, *G. armata*, and *T. fragile*, respectively. Methanolic extracts showed TFC values of 0.94, 0.87, 0.89, 0.50, and 0.49mg/ g DW for the same species, respectively.

Using ethanolic extracts, the TFC values were 0.59, 2.94, 2.82, 2.24, and 2.22 mg/g DW for *C. officinalis*, *D. simplex*, *A. rigida*, *G. armata*, and *T. fragile*, respectively.

The acetonic extract of *S. cinereum* showed the highest TFC (18.21mg/ g DW), followed by *P. tetrastromatica* (16.87 mg/g DW), and *T. decurrens* (8.71mg/ g DW). As observed with both the Chlorophyta and Rhodophyta groups, the lowest TFC values were obtained from water and methanolic extracts. Water extracts of *S. cinereum*, *P. tetrastromatica*, and *T. decurrens* recorded TFC values of 0.13, 0.19, and 0.45mg/ g DW, respectively. Methanolic extracts showed TFC values of 0.19, 0.73, and 0.11mg/ g DW for the same species, respectively. When using ethanolic extracts, TFC values were 3.97, 2.45, and 3.17mg/ g DW for *S. cinereum*, *P. tetrastromatica*, and *T. decurrens*, respectively.




Fig. 4. Total flavonoid content (TFC) (mg/g DW) of the tested seaweed species using various solvents. Data points represent means \pm standard deviation ($n = 3$).

4.3. Total tannin content (TTC) among different seaweeds extracts

The results of the quantitative tannin content (TTC) showed that the different types of seaweed had significantly different levels of tannins ($P < 0.050$). Tannin levels varied significantly among solvent extracts, ranging from 0.04 in *T. decurrens* to 0.56mg/ g DW in *U. fasciata*. Aqueous extract of *U. fasciata* exhibited the highest tannin level (0.56mg/ g DW), followed by the ethanolic extract of *A. rigida* (0.41mg/ g DW), as shown in Fig. (5). Tannin levels in the selected species vary significantly depending on the variety of seaweeds and the solvents used.

Phytochemical Profiling and GC-MS-Based Characterization of Bioactive Compounds from Egyptian Coastal Seaweeds

The maximum tannin level in the Chlorophyta algal group was detected in *U. fasciata* water extract (0.56mg/ g DW) followed by acetonic extract of *U. linza* (0.46 mg/g DW), while the ethanol extract of *U. fasciata*, and *U. linza* recorded close tannin levels of (0.13mg/ g DW) and (0.19mg/ g DW), respectively. Additionally, methanol extract of *U. fasciata*, and *U. linza* recorded 0.06 and 0.1mg/ g DW.

Among the different groups of Rhodophyta algae, the ethanolic extract of *A. rigida* exhibited the highest tannin level value (0.41mg/ g DW), followed by the water extract of *A. rigida* (0.32mg/ g DW), and the acetonic extract of *T. fragile* (0.3mg/ g DW). The lowest tannin level was recorded with aqueous extract of *G. armata* (0.04mg/ g DW). *C. officinalis* and *D. simplex* exhibited the lowest tannin level with all solvents. *C. officinalis* recorded values of 0.09, 0.08, 0.08 and 0.05mg/ g DW with ethanol, acetone, water, and methanol, respectively, while *D. simplex* recorded 0.05, 0.09, 0.08 and 0.08mg/ g DW with ethanol, acetone, water, and methanol, respectively. *G. armata* ethanolic and acetonic extract recorded close tannin level values with 0.2±0.0 and 0.25mg/ g DW, respectively. In addition, water and methanolic extract of *G. armata* recorded similar tannin level values of 0.04 and 0.05mg/ g DW, respectively. Whereas *T. fragile* exhibited tannin level values of 0.1, 0.05, and 0.06mg/ g DW with ethanol, water, and methanol extract, respectively.

The aqueous extract of *P. tetrastromatica* exhibited the highest tannin level value (0.48mg/ g DW) among the different Ochrophyta groups, while *T. decurrens* methanolic, acetonic, and ethanolic extracts had tannin level values close to each other at 0.29, 0.27, and 0.26mg/ g DW.

In the Ochrophyta group, *P. tetrastromatica* acetonic extract had a tannin level of 0.23 mg/g DW. *S. cinereum* exhibited tannin levels of 0.15, 0.14, 0.12, and 0.08mg/ g DW with ethanolic, acetonic, water, and methanolic extracts, respectively. In this context, *P. tetrastromatica* ethanolic and methanolic extracts had tannins levels of 0.16 and 0.06mg/ g DW, respectively.




Fig. 5. Total tannin content (TTC) (mg/g DW) of tested seaweed species using various solvents. Data points represent means \pm standard deviation (n = 3).

4.4 Gas chromatography–mass spectrometry analysis

Gas chromatography–mass spectrometry (GC–MS) is an authentic reliable tool commonly employed for the identification of chemical constituents. In the current investigations, the maximum number of bioactive compounds were found in ethanolic extract of *S. cinereum* (15 compounds) followed by 12 compounds in both *P. tetrastromatica* and *T. decurrens* extract. The investigation revealed various peaks of compounds, which were tentatively determined by comparing their height in percentage, peak retention time and other data reported in the literature and the National Institute of Standards and Technology library (NIST), as listed in Table (3) and Fig. (S1). The major compounds in *S. cinereum* were palmitic acid (37.43); Palmitic acid, methyl ester (17.75). The major compounds in the ethanolic extract of *P. tetrastromatica* were stearic acid, methyl ester (26.05); Palmitic acid (23.3); palmitic acid, methyl ester (20.61), and cholesterol (14.44). GC/MS analysis of *T. decurrens* extract identified different compounds, with the palmitic acid (56.09) being the major one; 13-docosenamide (Z) (11.06), cis-vaccenic acid (9.98) and palmitic acid, methyl ester (7.35).

GC–MS analysis of Rhodophyta showed that eighteen main constituents were identified in *D. simplex*; seventeen bioactive compounds were found in *G. armata*; fifteen peaks were present in the extract of *A. rigida*; fourteen bio compounds-were elucidated in *T. fragile*; and *C. officinalis* contained 11 constituents. The major constituents of *C. officinalis* were palmitic acid, methyl ester (20.01); palmitic acid (12.52664262); bicyclo [3.1.1] heptane, 2,6,6-trimethyl-, (1. alpha.,2. beta.,5. alpha.)- (10.84341219) and 2-butenic acid, 3-(methylamino)-, ethyl ester (8.922992623).

Phytochemical Profiling and GC-MS-Based Characterization of Bioactive Compounds from Egyptian Coastal Seaweeds

While the main constituents of *D. simplex* were palmitic acid (49.4). palmitic acid (35.46) and palmitic acid, methyl ester (11.75) represented the high area in *G. armata* extract. The GC-MS chromatogram analysis of the *A. rigida* extract revealed that the major compounds were palmitic acid (19.17%) and palmitic acid, methyl ester (31.995%). Similarly, the GC-MS analysis of *T. fragile* identified palmitic acid (22.17%), bicyclo[3.1.1]heptane, 2,6,6-trimethyl-, (1.α,2.β,5.α) (15.25%), palmitic acid, methyl ester (9.36%), and tetradecanoic acid (14.75%) as the predominant compounds, as shown in Table (4) and Fig. (S2).

Among the Chlorophyta algal group, the GC-MS analysis of *U. linza* revealed the presence of 11 phytoconstituents while *U. fasciata* contained nine constituents. The major compounds found in *U. fasciata* were palmitic acid (55.94) and palmitic acid methyl ester (19.16), while the main compounds in *U. linza* extract were palmitic acid (38.18); palmitic acid methyl ester (28.41) and 9-octadecenoic acid (Z)-, methyl ester (10.85), as illustrated in Table (5) and Fig. (S3).

Table 3. The bioactive compounds identified using GC chromatography and their biological activities in the ethanolic extracts of brown seaweed investigated in the present study

Compound Name	Common name	MW (g/mol)	(R T)	<i>S. cinereum</i>	<i>P. tetrastromatica</i>	<i>T. decurrens</i>	Biological Activity
				Area %			
n-Hexadecanoic acid (palmitic acid)	Palmitic acid	256.43	12.3	37.43	23.3	56.09	Antiandrogenic, Haemolytic, 5-Alpha reductase inhibitor (Kumar <i>et al.</i> , 2019).
Hexadecanoic acid, methyl ester	Methyl palmitate	270.45	12.0	17.75	20.61	7.35	Antimicrobial activity (Davoodbasha <i>et al.</i> , 2018; Shaaban <i>et al.</i> , 2021)
Octadecanoic acid, methyl ester	Methyl stearate	298.5	12.9	N D	26.05	3.06	Antiviral activity (Linton <i>et al.</i> , 2013)
13-Docosenamide, (Z)-	(Z)- Erucamide	337.59	15.9	N D	N D	11.06	Antinociceptive and anti-inflammatory activities (Shareef <i>et al.</i> , 2016)
cis-Vaccenic acid	cis-Vaccenic acid	282.47	13.1	6.14	2.86	9.98	Antibacterial and hypolipidemic effects (Hamazaki <i>et al.</i> , 2016)
11,13-Dimethyl-12-tetradecen-1-ol acetate	Tetradecyl acetate derivative	282.45	11.5	N D	N D	1.54	-
Tetradecanoic acid	Myristic acid	228.38	11.1	3.51	1.48	3.16	Anti-protozoa &Antimicrobial (Kumar <i>et al.</i> , 2019)

26-Nor-5-cholesten-3-beta.-ol-25-one	Nor-cholestenone	384.62	15.6	N	N	1.93	Antimicrobial, Diuretic, Anti-inflammatory, Anti-asthma (Cardeal <i>et al.</i> , 2006)
cis, cis,cis-7,10,13-Hexadecatrienal	Hexadecatrienal	234.34	13.6	N	N	1.4	-
trans-13-Docosenamide	trans-Erucamide	337.59	14.2	N	N	1.31	-
2,2-Dimethylpropanoic acid, 2,6-dimethylnon-1-en-3-yn-5-yl ester	Dimethylpropanoic acid ester	250.37	14.4	N	N	1.51	-
Cholesterol	Cholesterol	386.65	15.4	N	14.44	N	Anticancer activity, anticardiac activity, anti-inflammatory activity, antimicrobial activity, anti-psychotic activity, antioxidant activity, drug-loaded activity (Zhang <i>et al.</i> , 2021)
Gamolenic Acid	Gamma-Linolenic acid	278.43	13.6	N	4.64	N	anti-inflammatory, antithrombotic, antiproliferative, and lipid-lowering effects (Ponphaiboon <i>et al.</i> , 2024)
Bicyclo[3.1.1]heptane, 2,6,6-trimethyl-, (1.alpha.,2.beta.,5.alpha.)-	Camphene derivative	138.25	11.5	1.39	1.45	N	Antimicrobial (Srinivasan & Kumaravel, 2016)
9-Octadecyne	Octadecyne	250.48	11.7	4.51	1.42	N	-
Methyl tetradecanoate	Methyl myristate	242.4	10.8	N	1.39	1.56	-
13-Octadecenal, (Z)-	(Z)-13-Octadecenal	266.46	14.6	N	1.64	N	Antibacterial and antifungal activities (Hameed <i>et al.</i> , 2016)
2-Norbornaneacetic acid	Norbornaneacetic acid	154.21	9.2	N	0.7	N	-
9-Octadecenoic acid (Z)-, methyl ester	Methyl oleate	296.49	12.9	7.07	N	N	Anti-inflammatory, antiandrogenic cancer preventive, dermatitogenic hypocholesterolemic, 5-alpha reductase inhibitor, anemiagenic, insectifuge (Krishnamoorthy & Subramaniam, 2014)
,2-Benzenedicarboxylic acid, diisooctyl ester	Diisooctyl phthalate	390.56	14.8	5.77	N	N	antifungal activity (RAHMAN AND ANWAR 2006)
Phenobarbital	Phenobarbital	232.24	12.7	4.18	N	N	Reducing the spread of seizure activity from a seizure focus (Czapinski <i>et al.</i> , 2005).
2-Pentadecanone, 6,10,14-trimethyl-	Hexahydrofarnone	268.48	11.6	1.35	N	N	Antibacterial, anti-inflammatory, and anticancer properties, and is used in pain relief research (Avoseh <i>et al.</i> , 2021; Jiangseubchatveera <i>et al.</i> , 2015; G. Wei <i>et al.</i> , 2016)
Arachidonic acid	Arachidonic acid	304.47	13.6	4.29	N	N	anti-bacterial and anti-biofilm activities (Chamlagain <i>et al.</i> , 2024)
Tridecanoic acid, 12-methyl-, methyl ester	Methyl 12-methyltridecanoate	242.4	10.9	1.57	N	N	-

Phytochemical Profiling and GC-MS-Based Characterization of Bioactive Compounds from Egyptian Coastal Seaweeds

Oleic Acid	Oleic acid	282.47	13.4	0.83	N D	N D	hypocholesterolemic, antioxidant and lubricating activity(Selvamangai & Bhaskar, 2012)
Dodecanoic acid	Lauric acid	200.32	9.8	1.034	N D	N D	Antibacterial, antioxidant and anti-apoptotic activity (Renugadevi et al., 2021)
Hexadecanal, 2-methyl-	2-Methylhexadecanal	240.43	5.4	1.17	N D	N D	-

Table 4. The bioactive compounds identified using GC chromatography and their biological activities in the ethanolic extracts of red seaweed investigated in the present study

Compound Name	Common name	MW (g/mo l)	RT	<i>C. officinalis</i>	<i>D. simplex</i>	<i>G. armata</i>	<i>A. rigida</i>	<i>T. fragilis</i>	Biological Activity
				Area %					
Hexadecane acid, methyl ester	Methyl palmitate	270.25	12.20	20.01	ND	11.75	19.17	9.35	Antimicrobial activity (Davoodbasha <i>et al.</i> , 2018; Shaaban <i>et al.</i> , 2021)
Bicyclo[3.1.1]heptane, 2,6,6-trimethyl-, (1.alpha.,2.beta.,5.alp ha.)-	Camphane (bornane)	138.14	11.72	10.84	4.25	4.27	ND	15.54	Antimicrobial (Srinivasan & Kumaravel, 2016)
2-Butenoic acid, 3-(methylamino)-, ethyl ester	Ethyl 3-(methylamin o)crotonate	143.09	13.21	8.92	ND	ND	ND	ND	-
n-Hexadecanoic acid	Palmitic acid	256.24	12.59	12.52	49.45	35.45	31.94	22.16	Antiandrogenic, Haemolytic, 5-Alpha reductase inhibitor (Kumar <i>et al.</i> , 2019).
9-Eicosyne	9-Icoyne	278.29	11.96	4.85	ND	ND	ND	ND	Anti-microbial and cytotoxic properties (Kumar <i>et al.</i> , 2019)
5-Eicosene, (E)-	(E)-5-Icocene	280.31	13.63	13.36	ND	ND	ND	ND	Antimicrobial, cytotoxic, antihyperglycemic , antioxidant, insecticidal (Banakar & Jayaraj, 2018)
6-Octadecenoic acid, (Z)-	Petroselinic acid	282.25	13.39	9.13	ND	ND	2.53	ND	anti-inflammatory <i>acti</i>

									<i>vity (Diab et al., 2021) ,</i>
Tetradecanoic acid	Myristic acid	228.20	11.35	5.94	4.36	4.51	5.35	14.75	Anti-protozoa &Antimicrobial (Kumar et al., 2019)
Benzo[h]quinoline, 2,4-dimethyl-	2,4-Dimethylbenzo[h]quinoline	207.10	14.95	6.13	ND	ND	ND	ND	-
Benzene, 1,3-dimethyl-	m-Xylene	106.07	4.41	5.39	ND	ND	ND	ND	-
2-Pentanethiol	Amyl mercaptan	104.06	6.04	2.868	ND	ND	ND	ND	-
Pentadecanoic acid, 14-methyl-, methyl ester	Methyl 14-methylpentadecanoate	270.25	12.49	ND	6.20	ND	ND	ND	antifungal and antimicrobial activity (Akpuaka et al., 2013).
3,7,11,15-Tetramethyl-2-hexadecen-1-ol	Phytol	296.30	11.72	ND	4.54	ND	ND	ND	metabolism-modulating, cytotoxic, antioxidant, autophagy- and apoptosis-inducing, antinociceptive, anti-inflammatory, immune-modulating, and antimicrobial effects (Islam et al., 2018).
9-Octadecenamide, (Z)-	Oleamide	281.27	13.17	ND	6.69	ND	ND	ND	
1,2-Benzenedicarboxylic acid, diisooctyl ester	Diisooctyl phthalate (DIOP)	390.27	11.33	ND	1.68	ND	ND	ND	cytotoxic activity (Krishnan et al., 2014a), antibacterial (Rahman et al., 2007)
9-Octadecyne	9-Octadecyn-1-yne	250.26	15.07	ND	2.48	ND	ND	ND	-
2H-Pyran-2-one, 5,6-dihydro-6-pentyl-	6-Pentyl-5,6-dihydro-2H-pyran-2-one	168.11	11.96	ND	4.10	1.859	ND	ND	-
Oleic Acid	Oleic acid	282.25	9.49	ND	3.84	ND	ND	ND	Anticancer, anti-aging and Anti-inflammatory , Anti-tumor (Carrillo Pérez et al., 2012)

Phytochemical Profiling and GC-MS-Based Characterization of Bioactive Compounds from Egyptian Coastal Seaweeds

Tetradecanamide	Myristamide	227.2 2	13.3 5	ND	3.2 3	ND	ND	ND	-
2-Butanone, 4-(dimethylamino)-3-methyl-	4-(Dimethylamino)-3-methyl-2-butanone	129.1 1	13.5 6	ND	1.9 3	ND	ND	ND	-
tert-Hexadecanethiol	tert-Hexadecyl mercaptan	258.2 3	11.0 1	ND	1.6 6	ND	ND	ND	-
Dodecanoic acid	Lauric acid	200.1 7	13.8 9	ND	1.2 3	ND	1.90	1.6 0	antibacterial, antioxidant and anti-apoptotic activity (Renugadevi <i>et al.</i> , 2021)
. beta.-D-Glucopyranose, 1-thio-,1-[N-hydroxy-5-(methylthio)pentanimide]	(Complex, not widely named)	341.0 9	10.0 4	ND	1.2 7	ND	ND	ND	-
Xylitol	Xylitol	152.0 6	5.85	ND	1.0 2	ND	ND	ND	Antibacterial (Cocco <i>et al.</i> , 2020)
1,3-Propanediamine, N,N'-diethyl-	N,N'-Diethyl-1,3-propanediamine	130.1 4	6.08	ND	1.0 04	ND	ND	ND	-
1H-Pyrazole, 1,5-dimethyl-	1,5-Dimethylpyrazole	96.06	6.95	ND	1.0 03	ND	ND	ND	-
2-Furanmethanol	Furfuryl alcohol	98.03	4.05	ND	ND	0.74 2	ND	ND	antiviral (Premathilaka & Silva, 2016)
2(4H)-Benzofuranone, 5,6,7,7a-tetrahydro-4,4,7a-trimethyl-, (R)-	(R)-4,4,7a-Trimethyl-5,6,7,7a-tetrahydrobenzofuran-2-one	180.1 1	9.74	ND	ND	0.81	ND	ND	Antifungal (Parker <i>et al.</i> , 1997)
Propylphosphonic acid, fluoroanhydride, decyl ester	(No widely used common name; phosphonic acid derivative)	266.1 8	10.0 24	ND	ND	0.85	ND	ND	-
Cyclohexane, 1,2,4,5-tetraethyl-, (1.alpha.,2.alpha.,4.alpha.,5.alpha.)-	Tetraethylcyclohexane	196.2 1	10.8 87	ND	ND	1.19	ND	ND	-
1-Hexadecyne	Hexadecyne	222.2 3	11.7 7	ND	ND	1.29	ND	ND	Antibacterial & antioxidant

									activities (Mou <i>et al.</i>, 2013)
1-Octadecene	Octadecene	252.28	12.53	ND	ND	1.40	ND	ND	-
Phenobarbital	Phenobarbital	232.08	12.75	ND	ND	15.06	ND	ND	increasing synaptic inhibition. reducing the spread of seizure activity from a seizure focus (Czapinski <i>et al.</i>, 2005).
9-Octadecenoic acid (Z)-, methyl ester	Methyl oleate	296.27	12.92	ND	ND	6.89	7.25	5.40	Anti-inflammatory, antiandrogenic cancer preventive, dermatitogenic hypocholesterolemic, 5-alpha reductase inhibitor, anemiagenic, insectifuge (Krishnamoorthy & Subramaniam, 2014)
cis-13-Octadecenoic acid	Cis-vaccenic acid	282.25	13.19	ND	ND	6.43	ND	ND	-
9-Tricosene, (Z)-	Z-9-Tricosene (Muscalure)	322.36	13.42	ND	ND	2.71	ND	ND	-
2-Butyl-5-methyl-3-(2-methylprop-2-enyl)cyclohexanone	Menthone derivative	222.19	14.09	ND	ND	1.79	ND	ND	-
1,2-Benzenedicarboxylic acid, mono(2-ethylhexyl) ester	Mono(2-ethylhexyl) phthalate (MEHP)	278.15	12.32	ND	ND	2.93	8.17	ND	Cytotoxic activity (Krishnan <i>et al.</i>, 2014b)
2-Pentadecanone, 6,10,14-trimethyl-	Hexahydrofarnesone	268.27	11.582	ND	ND	ND	4.64	ND	Antibacterial, anti-inflammatory, and anticancer properties, and is used in pain relief research (Avoseh <i>et al.</i>, 2021 ; Jiangseubchatveera <i>et al.</i>, 2015 ; G. Wei <i>et al.</i>, 2016)
Phthalic acid, cyclohexylmethyl propyl ester	(No widely used common name;	304.16	11.77	ND	ND	ND	1.63	ND	-

Phytochemical Profiling and GC-MS-Based Characterization of Bioactive Compounds from Egyptian Coastal Seaweeds

	phthalate ester)								
Dodecane, 1-chloro-	1-Chlorododecane	204.16	9.07	ND	ND	ND	2.30	ND	-
Methyl tetradecanoate	Methyl myristate	242.22	10.85	ND	ND	ND	3.19	ND	-
Tetradecane, 1-chloro-	1-Chlorotetradecane	232.19	10.56	ND	ND	ND	1.41	ND	-
trans-13-Octadecenoic acid	Elaidic acid	282.25	13.20	ND	ND	ND	4.51	ND	Arachidonic acid inhibitor, inhibits the production of uric acid, anti-inflammatory, cancer preventive, anemiagenic, insectifuge, and flavor (Abdullah et al., 2020)
Ethyl 2-(2-chloroacetamido)-3,3,3-trifluoro-2-(4-fluoroanilino) propionate	(No common name; fluorinated propionate derivative)	200.17	15.46	ND	ND	ND	1.90	ND	-
Icosapent	Eicosapentaenoic acid ethyl ester (EPA ethyl ester)	302.225	13.72	ND	ND	ND	2.93	ND	-
9,17-Octadecadienal, (Z)-	(Z)-9,17-Octadecadienal	264.24	8.36	ND	ND	ND	1.05	ND	Antimicrobial property (Rajeswari et al., 2012)
9-Octadecenoic acid, (E)-	Elaidic acid	296.27	13.20	ND	ND	ND	ND	4.37	-
Noracymethadol	(No common name; noracymethadol opioid)	339.22	4.78	ND	ND	ND	ND	4.72	-
2,2-Dimethyl-N-tert-butylaziridine-1-carboxamide	(No common name; aziridine derivative)	170.14	6.92	ND	ND	ND	ND	2.52	-
Z-(13,14-Epoxy) tetradec-11-en-1-ol acetate	Epoxy myristyl acetate	268.20	13.83	ND	ND	ND	ND	2.58	-
2-Mercaptopropanoic acid	Thiolactic acid	106.00	6.20	ND	ND	ND	ND	3.63	-

1-Propanamine, 3-propoxy-	3-Propoxypropylamine	117.11	5.53	ND	ND	ND	ND	3.16	-
2-Tridecanone	Methylnonyl ketone	198.19	5.39	ND	ND	ND	ND	1.52	-
2-Heptanone, 4-methyl-	Methylisoamyl ketone	128.12	7.47	ND	ND	ND	ND	1.43	-

Table 5. The bioactive compounds identified using GC chromatography and their biological activities in the ethanolic extracts of green seaweeds investigated in the present study

Compound Name	Common name	M W (g/mol)	(R T)	U. fascia ta	U. linz a	Biological Activity
				Area %		
n-Hexadecanoic acid	Palmitic acid	256.24	12.49	55.94	38.18	Antiandrogenic, Haemolytic, 5-Alpha reductase inhibitor (Kumar et al., 2019).
Hexadecanoic acid, methyl ester	Methyl palmitate	270.25	12.20	19.16	28.41	Antimicrobial activity (Davoodbasha et al., 2018; Shaaban et al., 2021)
2-Pentadecanone, 6,10,14-trimethyl-	Hexahydrofarnesone	268.27	11.78	4.38	ND	Antibacterial, anti-inflammatory, and anticancer properties, and is used in pain relief research (Avoseh et al., 2021; Jiangseubchatveera et al., 2015; G. Wei et al., 2016).
(R)-(-)-(Z)-14-Methyl-8-hexadecen-1-ol	(R)-(Z)-14-Methylhexadecenol	254.26	11.96	2.98	ND	-
Hexanedioic acid, bis(2-ethylhexyl) ester	DEHA (Di(2-ethylhexyl) adipate)	370.30	14.43	5.34	ND	-
Phenobarbital	Phenobarbital	232.08	12.90	5.65	ND	Phenobarbital acts on GABAA receptors, increasing synaptic inhibition. This has the effect of elevating seizure threshold and reducing the spread of seizure activity from a seizure focus (Czapinski et al., 2005).
Ethyl tetradecyl ether	Ethyl myristyl ether	242.26	13.57	2.92	ND	-
Oleic Acid	Oleic acid	282.25	13.35	2.36	2.26	hypocholesterolemic, antioxidant, Anti-inflammatory and lubricating activity(Selvamangai & Bhaskar, 2012)
12-Octadecenoic acid, methyl ester	Methyl petroselinate	296.27	13.13	1.27	ND	Antimicrobial (L. S. Wei et al., 2011)
9-Octadecenoic acid (Z)-, methyl ester	Methyl oleate	296.27	12.93	ND	10.85	Anti-inflammatory, antiandrogenic cancer preventive, dermatitigenic hypocholesterolemic,5-alpha reductase inhibitor, anemiagenic,

Phytochemical Profiling and GC-MS-Based Characterization of Bioactive Compounds from Egyptian Coastal Seaweeds

						insectifuge(Krishnamoorthy & Subramaniam, 2014)
cis-Vaccenic acid	cis-Vaccenic acid	282.25	13.158	ND	5.56	antibacterial and hypolipidemic effects (Hamazaki <i>et al.</i> , 2016)
Bicyclo[3.1.1]heptane, 2,6,6-trimethyl-, (1.alpha.,2.beta.,5.alpha.)-	Camphene derivative	138.14	11.52	ND	4.70	Antimicrobial (Srinivasan & Kumaravel, 2016)
Di-n-octyl phthalate	Dioctyl phthalate	390.27	14.88	ND	3.25	-
Tetradecanoic acid	Myristic acid	228.20	11.13	ND	2.25	Anti-protozoa &Antimicrobial (Kumar <i>et al.</i> , 2019)
3,7,11,15-Tetramethyl-2-hexadecen-1-ol	Phytol	296.30	11.76	ND	1.63	-
Z-10-Pentadecen-1-ol acetate	Z-10-Pentadecenyl acetate	268.24	13.36	ND	1.60	-
Pentadecanoic acid, methyl ester	Methyl pentadecanoate	256.24	11.45	ND	1.32	-

DISCUSSION

The pronounced influence of the extraction solvent on crude yield and phytochemical recovery reflects trends reported for the tested brown species. *Sargassum cinereum* and *Turbinaria decurrens* exhibited the highest extractable material with ethanol, consistent with **Ismail *et al.* (2019)**, who observed that moderately polar ethanol preferentially solubilizes phenolic phlorotannins in Phaeophyceae. Additionally, **Sujatha *et al.* (2019)** demonstrated that ethanolic fractions of *Sargassum swartzii* contain richer bioactive profiles. On the other hand, **Mahendran *et al.* (2021)** showed that methanolic extraction of the red seaweeds *Gracilaria edulis* and *Hypnea valentiae* maximizes total phenolic content, highlighting the importance of matching solvent polarity to target metabolite classes. The variation in extraction yield of bioactive compounds from algal species may be related to solvent type, age, algal species, and differences in habitat (**Gazali *et al.*, 2023**).

Qualitative screening revealed the consistent presence of phenols, flavonoids, tannins, steroids, terpenoids, and saponins across most species, with alkaloids detected more variably. This echoes **Ganesan *et al.* (2008)**, who found that Ochrophyta often display the widest suite of secondary metabolites. The absence of certain compounds in high-polarity (water, methanol) extracts of Rhodophyta and Chlorophyta aligns with the “polarity window” hypothesis of **Herawati and Pudjiastuti (2021)**, whereby intermediate solvents such as acetone and ethanol co-extract both polar and moderately nonpolar constituents simultaneously. Additionally, qualitative phytochemical screening by **Al-Hashdy *et al.* (2025)** detected phenolic and tannin compounds in the methanolic extract of *Sargassum vulgare*, whereas these were absent in its nonpolar n-hexane

fraction, further underscoring the critical role of solvent polarity in recovering flavonoids and condensed tannins from brown macroalgae.

Quantitatively, total phenolic, flavonoid, and tannin contents varied significantly ($P < 0.05$) among species and solvents. The highest total phenolic content (1.02 ± 0.03 mg GAE/g DW) was found in the acetone extract of *Turbinaria decurrens*. Regarding total phenolic content, **Nguyen and Tran (2025)** reported that the ethanolic extract of *Dictyota implexa* contained 85.95 ± 1.21 mg GAE/g extract, a value substantially higher than many brown algal extracts and closely mirroring the elevated phenolic levels we observed in our ethanol fractions. This concordance not only validates the reliability of ethanol as an extraction solvent for phenolic compounds in Phaeophyceae but also suggests that the intrinsic biochemical composition of *D. implexa* and related species shares a consistent propensity for phenolic accumulation under similar collection and processing conditions. Therefore, our data, in conjunction with Nguyen and Tran's findings, underscore the critical influence of solvent polarity and algal physiology on maximizing phenolic recovery across diverse brown macroalgae. **Nagah *et al.* (2023)** reported that the cold-water extract of *Sargassum swartzii* (SCWE) contained $5,749.04 \pm 118.2$ μ g GAE/mL, compared with $4,559.09 \pm 79.8$ μ g GAE/mL in the autoclaved water extract and only 662.21 ± 26.3 μ g GAE/mL in the ethanolic extract, underscoring how solvent polarity dramatically influences phenolic recovery.

The highest flavonoid content was recorded in acetone extracts of *Ulva linza* (10.7 ± 0.2 mg QE/g DW) and *Sargassum cinereum* (18.2 ± 0.04 mg QE/g DW), confirming similar solvent biases noted by **Koivikko *et al.* (2005)** and **Lee *et al.* (2022)** for green and brown seaweeds. In this respect, **Subbiah *et al.* (2025)** reported that *Phyllospora comosa* exhibited a peak flavonoid content of 0.73 mg QE/g at 0 h during colonic fermentation, further emphasizing the strong influence of solvent polarity and species-specific chemistry on flavonoid recovery in marine algae. The cold-water extract of *Sargassum swartzii* (SCWE) contained 352.22 ± 7.7 μ g QE/mL, compared with only 39.00 ± 1.7 μ g QE/mL in its ethanolic extract, nearly a tenfold difference, underscoring the profound impact of solvent polarity on flavonoid recovery.

Tannin distributions, peaking in aqueous *Ulva fasciata* (0.56 ± 0.01 mg CE/g DW), further illustrate how solvent polarity governs the leaching of condensed tannins (**Koivikko *et al.*, 2005**).

Comparative GC-MS profiling of ethanolic extracts from ten Egyptian seaweeds revealed a phytochemical suite dominated by fatty acids, esters, alcohols, and hydrocarbons. Four core metabolites—palmitic acid, palmitic-acid methyl ester, cis-vaccenic acid, and 9-octadecenoic-acid methyl ester—recurred at high abundance, with the palmitic-acid pair alone contributing $\geq 30\%$ of the total peak area in *T. decurrens*, *S. cinereum*, *P. tetrastromatica*, *U. fasciata*, and *G. armata*. Our results are in good accordance with **Al-Hashdy *et al.* (2025)**, who stated that the prevalence of palmitic acid derivatives mirrors earlier GC-MS reports for the brown alga *S. cinereum* and the red alga *Corallina officinalis* (**Hassan & Shobier, 2018**). Saturated medium-chain lipids such as lauric and myristic acids, both prominent in many marine algal extracts, have been shown to scavenge free radicals, attenuate lipid peroxidation, suppress a wide range of Gram-positive skin

pathogens, and competitively inhibit type-1 and type-2 5- α -reductase, the key enzyme in androgen activation (Nakatsuji *et al.*, 2009). Red-algal extracts displayed lineage-specific markers: *Dichotomaria simplex* accumulated phenobarbital; *Acanthophora rigida* and *Tetraclada fragile* contained 2-pentadecanone, 6,10,14-trimethyl- (hexahydrofarnesyl acetone); and *Corallina officinalis* was enriched in tetradecanoic (myristic) acid metabolites. These compounds have been linked to antinociceptive/anti-inflammatory (hexahydrofarnesyl acetone) and NF- κ B-mediated anti-inflammatory (myristic acid) activities (Huang *et al.*, 2023). Such chemovariation likely reflects interacting environmental drivers (climate, seasonality, geography) and intrinsic genetic diversity that modulate secondary-metabolite biosynthesis and, consequently, medicinal quality (Briskin, 2000). Collectively, the dominance of bioactive fatty acids, esters, and alkenes, supplemented by species-specific compounds, underscores the therapeutic promise of Egyptian seaweed extracts and rationalizes their strong antioxidant and cytotoxic activities observed *in vitro*.

CONCLUSION

Phytochemical screening of various seaweed extracts revealed the presence of phenolic compounds with varying concentrations depending on the algal species and the solvent used. The study clearly indicates that *Sargassum cinereum* is a promising source of phytochemical compounds. This research also supports the health benefits of common Egyptian seaweed extracts in medicinal applications by emphasizing their significance as possible sources of natural phenolic compounds with a variety of bioactivities. According to the results obtained, it can be said that the solvent used had a significant effect on the total phytochemical content and concentrations. However, further studies are needed to isolate and purify the bioactive compounds from the most promising and dominant species responsible for the biological activity. This will allow them to be utilized as natural ingredients in the nutritional and pharmaceutical industries.

REFERENCES

- Ahmed, H.; Heneidak, S.; Rasmey, A.H. and El Shoubaky, G. (2023). Evaluation of antioxidant activities of some abundant marine macroalgae, egypt. *Bulletin of Pharmaceutical Sciences Assiut University*, 46(2), 913–925.
- Al-Hashdy, D.F.; El-Shaibany, A.; Abdelkhalek, A. S.; Raweh, S. M.; Elaasser, M. M. and Raslan, A. E. (2025). Phytochemical profiling, toxicity, and pharmacological investigations of the brown seaweed *Sargassum vulgare* collected from the Red Sea coast combined with molecular docking analysis and SwissADMET prediction. *South African Journal of Botany*, 182, 165–184. <https://doi.org/10.1016/j.sajb.2025.05.006>
- Briskin, D.P. (2000). Medicinal Plants and Phytomedicines. Linking Plant Biochemistry and Physiology to Human Health. *Plant Physiology*, 124(2), 507–514. <https://doi.org/10.1104/pp.124.2.507>
- Carreira-Casais, A.; Otero, P.; Garcia-Perez, P.; Garcia-Oliveira, P.; Pereira, A.G.; Carpena, M.; Soria-Lopez, A.; Simal-Gandara, J. and Prieto, M.A. (2021). Benefits and

- drawbacks of ultrasound-assisted extraction for the recovery of bioactive compounds from marine algae. *International Journal of Environmental Research and Public Health*, 18(17), 9153.
- Chang, C.C.; Yang, M.H.; Wen, H.M. and Chern, J.C.** (2020). Estimation of total flavonoid content in propolis by two complementary colometric methods. *Journal of Food and Drug Analysis*, 10(3). <https://doi.org/10.38212/2224-6614.2748>
- Cotas, J.; Leandro, A.; Pacheco, D.; Gonçalves, A.M. and Pereira, L.** (2020). A comprehensive review of the nutraceutical and therapeutic applications of red seaweeds (Rhodophyta). *Life*, 10(3), 19.
- El Zokm, G.M.; Ismail, M.M. and El-Said, G. F.** (2021). Halogen content relative to the chemical and biochemical composition of fifteen marine macro and micro algae: Nutritional value, energy supply, antioxidant potency, and health risk assessment. *Environmental Science and Pollution Research*, 28(12), 14893–14908. <https://doi.org/10.1007/s11356-020-11596-0>
- El-Rafie, H.M.; Hasan, E.A. and Zahran, M.K.** (2023). Enhancement of cytotoxic and antioxidant activities of *Digenea simplex* chloroform extract using the nanosuspension technique. *Bioprocess and Biosystems Engineering*, 46(2), 279–296.
- El-Shahir, A.A.; Alzamel, N. M.; Abuzaid, A.O.; Loutfy, N. and Alwaleed, E.A.** (2024). Antifungal Properties of *Sargassum cinereum* and *Padina boergeresenii* Extracts Against Fungi Associated with Strawberry Fruits Concerning Mycotoxin Production. *Plants*, 13(22), 3115. <https://doi.org/10.3390/plants13223115>
- Ganesan, P.; Kumar, C.S. and Bhaskar, N.** (2008). Antioxidant properties of methanol extract and its solvent fractions obtained from selected Indian red seaweeds. *Bioresource technology*, 99(8), 2717-2723.
- Gazali, M.; Effendi, I.; Wahyuni, S.; Kurniawan, R.; Matin, A. and Syafitri, R.** (2023). Boosting the growth of tilapia fish using marine macroalgae (*Sargassum* sp.) extracts. *Aquaculture, Aquarium, Conservation and Legislation*, 16(3), 1626-1635.
- Getachew, A.T.; Jacobsen, C. and Holdt, S.L.** (2020). Emerging technologies for the extraction of marine phenolics: Opportunities and challenges. *Marine Drugs*, 18(8), 389.
- Ghosh, S.; Sarkar, T.; Pati, S.; Kari, Z.A.; Edinur, H.A. and Chakraborty, R.** (2022). Novel bioactive compounds from marine sources as a tool for functional food development. *Frontiers in Marine Science*, 9, 832957.
- Guiry, M.D. and Guiry, G.M.** (2022). In. *AlgaeBase* (National University of Ireland, Galway: World-wide electronic publication). *AlgaeBase*, <https://www.algaebase.org/>.

- Harborne, J.** (1998). *Phytochemical Methods* London Chapman and Halls. 91.
- Hassan, S. and Shobier, A.** (2018). GC/MS identification and applications of bioactive seaweed extracts from Mediterranean coast of Egypt. *Egyptian Journal of Aquatic Biology and Fisheries*, 22(5 (Special Issue)), 1–21. <https://doi.org/10.21608/ejabf.2018.17952>
- Herawati, D. and Pudjiastuti, P.** (2021). Effect of different solvents on the phytochemical compounds of Sargassum sp. from Yogyakarta and East Nusa Tenggara. In *Journal of Physics: Conference Series* 1783(1), 012001
- Huang, Q.; Chen, C.; Zhang, Z. and Xue, Q.** (2023). Anti-inflammatory effects of myristic acid mediated by the NF- κ B pathway in lipopolysaccharide-induced BV-2 microglial cells. *Molecular Omics*, 19(9), 726–734. <https://doi.org/10.1039/D3MO00063J>
- Ibtissam, C.; Hassane, R.; Jose, M.; Francisco, D.S.J.; Antonio, G.V.J.; Hassan, B. and Mohamed, K.** (2009). Screening of antibacterial activity in marine green and brown macroalgae from the coast of Morocco. *African Journal of Biotechnology*, 8(7). <https://www.ajol.info/index.php/ajb/article/view/60101/48355>
- Ismail, M.M.; Mostafa, M.H., and El Sikaily, A.M.** (2022). Phytochemical variation, antioxidant and antidiabetic capacity of extracts of common brown seaweeds from the Red Sea, Egypt. *Egyptian Journal of Aquatic Biology and Fisheries*, 26(5), 445–462. <https://doi.org/10.21608/ejabf.2022.262605>
- Ismail, M. M.; El Zokm G.M., and El-Sayed, A.A.M.** (2017). Variation in biochemical constituents and master elements in common seaweeds from Alexandria Coast, Egypt, with special reference to their antioxidant activity and potential food uses: Prospective equations | Request PDF. ResearchGate. <https://doi.org/10.1007/s10661-017-6366-8>
- Julkunen-Tiitto, R.** (1985). Phenolic constituents in the leaves of northern willows: Methods for the analysis of certain phenolics (world). ACS Publications; *American Chemical Society*. <https://doi.org/10.1021/jf00062a013>
- Mahendran, S.; Maheswari, P.; Sasikala, V.; Jaya Rubika, J. and Pandiarajan, J.** (2021). In vitro antioxidant study of polyphenol from red seaweeds dichotomously branched gracilaria Gracilaria edulis and robust sea moss Hypnea valentiae. *Toxicology reports*, 8, 1404-1411.
- Mohy El-Din, S.M. and Alagawany, N.I.** (2019). Phytochemical Constituents and Anticoagulation Property of Marine Algae Gelidium crinale, Sargassum hornschurchii and Ulva linza. *Thalassas: An International Journal of Marine Sciences*, 35(2), 381–397. <https://doi.org/10.1007/s41208-019-00142-6>

- Nagah, A.; Ali, M. and Aly, S.H.** (2023). Physicochemical properties and antioxidant activities of crude fucoidan extracted from *Sargassum cinereum*. <http://jifi.farmasi.univpancasila.ac.id/index.php/jifi/article/view/823>
- Nakatsuji, T.; Kao, M. C.; Fang, J.Y.; Zouboulis, C.C.; Zhang, L.; Gallo, R.L. and Huang, C.M.** (2009). Antimicrobial Property of Lauric Acid Against *Propionibacterium Acnes*: Its Therapeutic Potential for Inflammatory Acne Vulgaris. *Journal of Investigative Dermatology*, 129(10), 2480–2488. <https://doi.org/10.1038/jid.2009.93>
- Trang Thuy, N.N., and Men, T.T.** (2025). Phytochemical and Bioactive Analysis of Extracted Brown Macroalgae (*Dictyota implexa*) Collected in Vietnam. *Biochemistry Research International*, 2025(1), 9461117. <https://doi.org/10.1155/bri/9461117>
- Sarker, S.D.** (2008). *Natural Products Isolation*. Springer Science and Business Media.
- Sobuj, M.K.A.; Islam, M. A.; Islam, M.S.; Islam, M.M.; Mahmud, Y. and Rafiquzzaman, S.** (2021). Effect of solvents on bioactive compounds and antioxidant activity of *Padina tetrastrum* and *Gracilaria tenuistipitata* seaweeds collected from Bangladesh. *Scientific Reports*, 11(1), 19082.
- Subbiah, V.; Ebrahimi, F.; Agar, O.T.; Legione, A.R.; Dunshea, F.R.; Barrow, C.J. and Suleria, H.A.R.** (2025). Bioactive phenolic compounds from Australian seaweeds and their impact on gut health. *Scientific Reports*, 15(1), 20954. <https://doi.org/10.1038/s41598-025-04689-8>.
- Sujatha, R.; Siva, D. and Nawas, P.M.A.** (2019). Screening of phytochemical profile and antibacterial activity of various solvent extracts of marine algae *Sargassum swartzii*. *World Scientific News*, (115), 27-40.
- Taga, M.S.; Miller, E.E. and Pratt, D.E.** (1984). Chia seeds as a source of natural lipid antioxidants. *Journal of the American Oil Chemists' Society*, 61(5), 928–931. <https://doi.org/10.1007/BF02542169>
- Thomas, T.E. and Harrison, P.J.** (1987). Rapid ammonium uptake and nitrogen interactions in five intertidal seaweeds grown under field conditions. *Journal of Experimental Marine Biology and Ecology*, 107(1), 1–8. [https://doi.org/10.1016/0022-0981\(87\)90118-3](https://doi.org/10.1016/0022-0981(87)90118-3)
- Trease, G. and Evans, W.** (1989). *Pharmacognosy*. 11th edn. Brailliar Tiridel Can.