



EGYPTIAN ACADEMIC JOURNAL OF
BIOLOGICAL SCIENCES
TOXICOLOGY & PEST CONTROL

F



ISSN
2090-0791

WWW.EAJBS.EG.NET

Vol. 17 No. 2 (2025)

www.eajbs.eg.net



Physiological and Biochemical Disturbances in the Insect Pests Infected with Entomopathogenic Nematodes: A comprehensive review

Karem Ghoneim^{*1} and Reda F.A. Bakr²

¹Department of Zoology and Entomology, Faculty of Science, Al-Azhar University, Cairo, Egypt.

²Department of Entomology, Faculty of Science, Ain Shams University, Abbassia- Cairo, Egypt.

*E-mail: karemghoneim@gmail.com

REVIEW INFO

Review History

Received:10/8/2024

Accepted:17/9/2025

Available:21/9/2025

Keywords:

Antioxidant enzymes, detoxification enzymes, digestive enzymes, carbohydrate, immunity-related enzymes, intermediary metabolism, lipid, protein %48

ABSTRACT

The excessive and indiscriminate uses of synthetic chemical insecticides have led to many serious problems of human health, ecosystem and economics. Therefore, it is necessary to search for safe and eco-friendly alternatives. Among these promising alternatives is the use of entomopathogenic nematodes (EPNs) for pest control. The objective of the present review was to give an insight into some of the physiological and biochemical disturbances in insects due to the infection with EPNs. Therefore, the current article discussed (1) the impaired intermediary metabolism including the main body metabolites (proteins, lipids and carbohydrates), and food consumption and utilization efficiencies; (2) the drastic effects of EPNs on some developmental and reproductive criteria of the insects; (3) the disrupted activities of major digestive enzymes including carbohydrate, protein and lipid hydrolyzing enzymes; (4) the disturbing effects of EPNs on the detoxification enzymes and the major humoral immunity-related enzymes. The disruptive effects of the EPNs on insects have suggested the potential of these entomopathogens for pest control. To the best of our knowledge, this is the first comprehensive review for understanding the principal physiological and biochemical mechanisms involved in the insects to combat the EPN oxidative stress. The current review will no doubt suggest new research avenues to many investigators, and it will make a valuable contribution to the published literature. However, some points of research need more investigation in future, such as the link between humoral immunity defenses and cellular immunity defenses to understand the defense integration in insects against EPNs.

Contents:

Introduction	
1. Impaired intermediary metabolism of insect pests by EPNs.....	
1.1. Disrupted main body metabolites in insects by EPNs	
1.1.1. Disruptive effects of EPNs on protein content in insects	
1.1.2. Disruptive effects of EPNs on lipid content in insects	
1.1.3. Disruptive effects of EPNs on carbohydrate content in insects	
1.2. Disrupted food consumption and utilization in insects by EPNs.....	

2. Drastic impacts of EPNs on developmental and reproductive physiology of insect pests	
3. Enzymatic disturbance in insect pests by the infection with EPNs	
3.1. Disturbed activities of the major digestive enzymes in insects.....	
3.1.1. Activity disturbance of carbohydrate hydrolyzing enzymes.....	
3.1.2. Activity disturbance of protein hydrolyzing enzymes	
3.1.3. Activity disturbance of lipid hydrolyzing enzymes	
3.1.4. Activity disturbance of some digestive-related enzymes	
3.2. Disturbed activities of the major detoxifying enzymes in insects by the infection with EPNs	
3.2.1. Disturbed activities of phosphatases in insects.....	
3.2.2. Disturbed activities of transaminases in insects.....	
3.2.3. Disturbed activity of Acetylcholinesterase in insects.....	
3.2.4. Disturbed activity of carboxylesterase in insects.....	
3.2.5. Disturbed activities of esterases in insects	
3.2.6. Disturbed activity of glutathione S-transferase in insects	
3.3. EPN infection interferes with activities of the major humoral immunity-related enzymes in insects	
3.3.1. Humeral immune defences in insects: a synopsis	
3.3.2. Effects of EPNs on the activity of peroxidase	
3.3.3. Effects of EPNs on the activity of catalase	
3.3.4. Effects of EPNs on the activity of superoxide dismutases	
3.3.5. Effects of EPNs on the activity of tyrosinase	
3.3.6. Effects of EPNs on the activity of immune response-mediated phenoloxidase in insects	
Summary points	
Conclusions and Prospectives	
Declaration.....	
References	

INTRODUCTION

Although the integrated pest management (IPM) strategies are increasingly developed in different parts of the world (Veres *et al.*, 2020), the control of insect pests still relies mainly on application of the synthetic insecticides (Jeschke *et al.*, 2011; Meslin *et al.*, 2021). The intensive and indiscriminate uses of these chemicals have led to negative impacts on the ecosystems including water, air and soil pollution (Tiryaki and Temur, 2010; Gunstone *et al.*, 2021) and drastically affect the natural enemies (like parasites and predators) leading to outbreaks of the pest populations (Demok *et al.*, 2019) as well as adverse effects on domestic animals (Vattikonda and Sangam, 2017; Shahzad *et al.*, 2020). Acute or chronic illnesses can develop in people who have been exposed to insecticides, either directly or indirectly (Mostafalou and Abdollahi, 2012). Additionally, conventional pesticides have serious impacts on both target and non-target organisms including birds, fish, amphibians, earthworms, and pollinators (Gill and Garg, 2014). In Egypt, as for example, the control of cutworms depends mainly on the application of conventional insecticides (Vattikonda and Sangam, 2017; Ismail, 2021). Some authors reported that the chemical control for these pests is often ineffective and remains inadequate because of the larval hiding behavior during the daylight hours causing hidden damage in fields (Capinera, 2001; Takeda, 2008; Kumar *et al.*, 2022), besides the fast development of resistance and cross resistance to almost all marketed insecticides (Yu Dong *et al.*, 2012; Mahmoud *et al.*, 2016; Shaurub *et al.*, 2018).

Therefore, various research institutions in the world have searched for new control agents as alternatives to synthetic insecticides (Laznik and Trdan, 2012; Derbalah *et al.*, 2014; Glare *et al.*, 2016). These alternative agents should be eco-environmentally safe (Liao *et al.*, 2017; Kunbharet *et al.*, 2018), effective at low concentrations (Walkowiak *et al.*, 2015) and biodegradable into harmless compounds (Tiryaki and Temur, 2010; Li *et al.*, 2017). One of these alternatives is the biocontrol of insect pests using natural enemies (parasitoids, predators and pathogens). Biocontrol agents are highly promising (Amutha *et al.*, 2021) because they are safe for humans and the environment, as well as they have little or no effect on other non-targeted organisms (Jagodič *et al.*, 2019).

Among the biocontrol agents, entomopathogenic nematodes (EPNs) have broad potential to kill the soil-dwelling insect pests as well as the above-ground insects which have soil-dwelling stages (Laznik and Trdan, 2011; Lacey and Georgis, 2012; James *et al.*, 2018; Du Preez *et al.*, 2021a, b; Kumar *et al.*, 2022). Moreover, various EPNs species are now applied against a wide range of foliar pests occurring in different habitats (Brusselman *et al.*, 2012; Beck *et al.*, 2013; Mahmoud, 2014; Gözel and Gozel, 2016). For example, EPNs *Steinernema carpocapsae* and *Heterorhabditis indica* possess significant potential for controlling several insect pests (Sunanda *et al.*, 2014; Khan *et al.*, 2018; Viteri *et al.*, 2018).

Distribution, virulence, and usage of EPNs in IPM programs have been studied globally (Çağlayan *et al.*, 2021; Ali *et al.*, 2022) because they are harmless to non-target organisms, human health and the environment (Odendaal *et al.*, 2016; Gulcuet *et al.*, 2017; Kumar *et al.*, 2022; Peçen and Kepenekci, 2022). Also, EPNs have high reproductive potential, the ability to kill hosts quickly, high virulence, broad host range, and safety to plants and vertebrates (Kaya and Gaugler, 1993). Most biocontrol agents require days or weeks to kill the insect pests, yet EPNs, working with their symbiotic bacteria, kill insects in 24-72 h (Adams and Nguyen, 2002). In addition, EPNs can be easily mass produced and applied using common irrigation and pesticide equipment (Askary and Abd-Elgawad, 2021; Yağci *et al.*, 2021a,b).

As biocontrol agents, EPNs can be used individually or in combination with other entomopathogenic bacteria or fungi in order to improve their efficacy for controlling the insect pests (Krishnayya and Grewal, 2002; Laznik *et al.*, 2012; Amizadeh *et al.*, 2019). Also, EPNs were found compatible with different agrochemicals (for review, see Ghoneim and Hamadah, 2024). EPNs have a high tolerance to the variations in environmental conditions (Toledo *et al.*, 2010). For some details, see Vashisth *et al.* (2013), Sujatha and Jeyasankar (2018), Jagodič *et al.* (2019), Askary and Abd-Elgawad (2021), Kumar *et al.* (2022), Shaurub (2023) and Ghoneim and Hassan, 2024).

It is important to point out that the EPNs parasitism leads to the suppression of the immune system of the insect host (Lewis and Clarke, 2012; Shapiro-Ilan and Brown, 2013; Lacey *et al.*, 2015; Kaliaskaret *et al.*, 2022). Moreover, EPNs have an association with certain symbiotic bacteria which are carried in the intestine of infective juveniles (IJs) of EPNs (Arthurs *et al.*, 2004; Lewis and Clarke, 2012; Chaston *et al.*, 2013). Among 23 families of EPNs, Steinernematidae and Heterorhabditidae are the two most potential families living naturally in the soil (Kumar *et al.*, 2015) and used as biocontrol agents against many insect pests (Lacey and Georgis, 2012; Abd-Elgawad, 2020; Koppenhöfer *et al.*, 2020; Yüksel *et al.*, 2022). Their association with symbiotic bacteria is found to be the primary cause of insect mortality (Leonaret *et al.*, 2022). Symbiotic bacteria of the genera *Xenorhabdus* and *Photorhabdus* associate with the EPN families, Steinernematidae and Heterorhabditidae, respectively, which produce natural products with insecticidal potential (Vicente-Díez *et al.*, 2021) for the suppression of the immune system of the insect host (for more detail, see Lewis and Clarke, 2012; Shapiro-Ilan and Brown, 2013; Lacey *et al.*, 2015; Koppenhöfer *et al.*, 2020; Bhat *et al.*, 2020; Kaliaskaret *et al.*, 2022; Ghoneim, 2024).

In this context, the pathogenicity of EPNs begins immediately after entering the insect's haemocoel where they release their symbiotic bacteria and rapidly multiply causing lethal septicemia to the insect host (Nickle and Welch, 1984). Also, the success of entomopathogens for the insect control depends on their stress potential and capability to impair certain physiological processes and biochemical constituents of the insect hosts. Over the past two decades, there has been increasing evidence that some EPNs influenced certain metabolic processes (Shaurubet *et al.*, 2020; Ghoneim *et al.*, 2022) and some important enzymatic activities (Abdel-Razek *et al.*, 2004; Shaurubet *et al.*, 2015; Ibrahim *et al.*, 2015; Shairraet *et al.*, 2016; Vidhya *et al.*, 2016; Ghoneim *et al.*, 2023a).

On the other hand, haemolymph is the only extracellular fluid in the insect body that is usually circulated by an open system within the body cavity. It performs several functions, such as the transportation of food materials to the cells and metabolic waste products away from those cells. It also transports hormones for regulation of larval moulting, growth, metamorphosis, metabolism and other physiological processes of insects (Hietakangas and Cohen, 2009). In insects, the use of haemolymph as a medium for controlling insect pests has been investigated because the changes occurring in the haemolymph are quickly transferred to other portions of the insect's body (Rodriguez-Ortega *et al.*, 2003; Pugazhvendan and Soundararajan, 2009). Therefore, physiological and biochemical disturbances in the haemolymph are expected, since the haemolymph is the main site of action of invading EPNs (Ghoneim *et al.*, 2022, 2023a).

Some attempts have been made to understand the principal physiological and biochemical mechanisms involved in the insect infection. Therefore, objective of the **present review** was to give an insight into the physiological and biochemical disturbances in the insect hosts, post-infection with EPNs, including impaired intermediary metabolism, disrupted developmental and reproductive processes, disturbed digestive enzymes, detoxifying enzymes and the major humoral immunity-related enzymes.

1. Impaired Intermediary Metabolism of Insect Pests by EPNs:

1.1. Disrupted Main Body Metabolites in Insects by EPNs:

In insects, different biological and physiological processes need adequate energy (Chapman, 1998; Fagan *et al.*, 2002). The content of macromolecules (such as protein, lipid and carbohydrate) is a valuable indicator of the level of metabolism, after treatment with exogenous materials (Zhu *et al.*, 2012). It is important to mention that protein synthesis is crucial for insect development and reproduction. Carbohydrates are the main source of energy during insect metamorphosis. Energy reserves, such as proteins, lipids, and glycogen in the haemolymph, are also an important indicator of the level of metabolism in insects (Chowanskiet *et al.*, 2015; Ferreira *et al.*, 2014; Ismail, 2018). These energy reserves are closely related to different physiological and biochemical processes in insects (Nawaz *et al.*, 2017).

1.1.1. Disruptive Effects of EPNs on the Protein Content in Insects:

Proteins are important organic constituents of animal tissues, including insects, and they play an important role in energy production (Taşkın and Aksoylar, 2011). The protein synthesis in insects is a prerequisite process for the development and reproduction (Hahn and Denlinger, 2007; Taşkın and Aksoylar, 2011). As reported by several authors (Suarez *et al.*, 2005; Bernstein and Jervis, 2008; Sugumaran, 2010), proteins perform a wide variety of physiological and metabolic functions and play a key role in the production of microsomal detoxifying enzymes.

Based on the available literature, the total protein content in the haemolymph of 4th instar of the mosquito *Culex pipiens* (Diptera: Culicidae) was reduced after infection with the EPN *Romanormisculicivora* (Schmidt and Platzer, 1980). The haemolymph protein content of the Egyptian cotton leafworm *Spodoptera littoralis* (Lepidoptera: Noctuidae) larvae was markedly reduced 30 hrs post-infection with some EPNs (El-Bishry, 1989). Also,

the total protein content of *S. littoralis* larvae was significantly reduced after infection with the EPNs *Steinernema riobrave* and *Heterorhabditis bacteriophora* (Ahmed et al., 2014). Four EPNs *H. bacteriophora* AS1, *H. bacteriophora* HP88, *Steinernema carpocapsae* ALL, and *S. riobrave* ML29 caused a remarkable decline in the total protein content of larvae of the Mediterranean fruit fly *Ceratitis capitata* (Diptera: Tephritidae) (Shaurub et al., 2015).

In Egypt, also, Hassan et al. (2016) studied the disturbance of protein content in 6th instar larvae of the black cutworm *Agrotis ipsilon* (Lepidoptera: Noctuidae) at different time intervals after infection with *S. glaseri* and *H. bacteriophora*. They determined a significant reduction in total protein content 24 hr post-infection. Two years later, infection of 5th instar nymphs of the desert locust *Schistocerca gregaria* (Orthoptera: Acrididae) with 1000 and 2000 IJs of *H. bacteriophora* resulted in reduction of the total protein content in nymphs (Gaber et al., 2018). Shaurub et al. (2020) incubated the newly moulted 4th instar larvae of *S. littoralis* with LD₅₀ of EPNs *Steinernema riobrave* and *H. bacteriophora* for 24 h. They determined decreasing protein content in the infected larvae. Also, Gomaa et al. (2020) evaluated the efficacy of two EPN isolates (*H. bacteriophora* and *S. carpocapsae*) and the Entomopathogenic fungus *Beauveria bassiana*, separately and in combination, on the 3rd instar larvae of *S. littoralis*. According to their results, total protein content was reduced post-infection with all treatments. In a recent study conducted in Egypt by Ghoneim et al. (2022), a predominant reduction of the protein content in haemolymph of last instar larvae of *A. ipsilon* was determined after infection with *S. carpocapsae* and *H. bacteriophora*. Moreover, *H. bacteriophora* exhibited stronger reducing potency against the protein content than *S. carpocapsae*, at 48 h post-infection (See Table 1).

Table 1. Total protein content in haemolymph (g/dL) of last (6th) instar larvae of *Agrotis ipsilon* as influenced by treatment of the newly moulted 5th instar larvae with LC₅₀ values of the tested entomopathogenic nematodes.

Nematode species		Time interval		
		6 hr	24 hr	48 hr
<i>S. carpocapsae</i>	mean±SD	6.10±0.12 a	5.78±0.33 a	4.84±0.12 b
	Change (%)	-13.84	-9.26	-21.94
<i>H. bacteriophora</i>	mean±SD	5.75±0.59 a	5.47±0.55 a	4.56±0.37 b
	Change (%)	-18.78	-14.12	-26.45
Control	mean±SD	7.08±0.84	6.37±1.21	6.20±0.49

Mean ± SD followed with letter: a: insignificant (P >0.05), b: significant (P<0.05), c: highly significant (P<0.01), d: very highly significant (P<0.001) (After Ghoneim et al., 2022).

In contrast, the current literature contains, also, few reported results of increasing protein content in larvae of some insects after infection with certain ENPs, such as *C. capitata* larvae at 4 and 18 h post-infection with *S. feltiae* Filipjev (Ghally et al., 1988). Also, total protein content significantly increased in the full-grown larvae of pink bollworm *Pectinophora gossypiella* (Lepidoptera: Gelechiidae) after treatment with *S. riobrave* but slightly increased after infection with *H. bacteriophora* (Shairra et al., 2016). Infection of the 5th instar nymphs of *S. gregaria* with the concentrations 1000 and 2000 IJs of *H. bacteriophora* resulted in disturbance of total protein contents in nymphs (Gaber et al., 2018).

Reduction of proteins in haemolymph of insects after infection with some EPNs may be attributed to the proteolytic activity in the haemolymph of the infected larvae. This activity was suggested to be the main cause of the host quick death (El-Bishry, 1989). According to some authors (von Brando, 1973; Lee and Atkinson, 1976), the considerable reduction of protein content may occur because many EPNs secrete chemicals (including toxins and digestive enzymes) to facilitate penetration and migration through the host tissues, and for feeding and avoiding the host immune responses. More than two decades later, some authors

(Istkhari and Chaubey, 2019; Wee et al., 2000) suggested that the production of proteases by EPN-symbiotic bacterial cells, followed by the breakdown of insect's protein and serving as nutritional resources for EPN/bacterium development. Also, reduction of proteins after infection with EPNs may be due to the stimulation of protein catabolism in the host fat body-the major organ for metabolism, nutrient storage, and synthesis of vitellogenin, a yolk protein precursor (Kamruzzaman et al., 2020) - to acquire a dietary supply of amino nitrogen from haemolymph (Gordon et al., 1973).

Early, Schmidt and Platzner (1980) reported that the protein degradation, when the mosquito *Cx. pipiens* were infected with EPN *R. culicivora* may be attributed to the production of some proteases from EPNs leading to this degradation in haemolymph. Also, the protein reduction in haemolymph might be due to the conversion of some proteins to fat, resulting in low protein content in the infected larvae (Abdel-Razek et al., 2004). In addition, breakdown of the proteins into free amino acids would ultimately lead to decrease in protein content. Moreover, the decreased protein content may be expected to suppress the immune response of infected larvae, including encapsulation, prophenoloxidase activity, phenoloxidase activity, total haemolymph proteins, and hemocyte density (Wilson et al., 2019).

1.1.2. Disruptive Effects of EPNs on the Lipid Content in Insects:

Lipids represent a principal source of energy for insects. They are transferred from their synthesis site *via* the haemolymph towards the target organs for use, in particular chitin synthesis, oogenesis, vitellogenesis, embryogenesis and continuous muscular activity (Dapporto et al., 2008; Zhou and Miesfeld, 2009). In addition, to the sites of lipid storage in the body, egg lipids play a very important role for achieving energy needed for the developing embryo (Boz and Gülel, 2012). Quantity of lipids available for the reserves seems to be the result of a balance between the obtained food and the requests for reserves by processes, such as maintenance, growth and reproduction, and this balance is disturbed by any xenobiotic stress (Canavoso et al., 2001). Also, impaired synthesis of lipids has been resulted in adversely influenced physiology and subsequently disrupted vital functions of growth and reproduction (Ghoneim et al., 2022).

Total lipids in *C. capitata* larvae declined after 18 h post-infection with *S. feltiae* (Ghally et al., 1988). The total lipid content of the *S. littoralis* larvae was significantly decreased post-infection with *S. riobrave* or *H. bacteriophora* (Ahmed et al., 2014). It was remarkably decreased in the 4th instar larvae of *S. littoralis* after incubation with LD₅₀ values of *S. riobrave* or *H. bacteriophora* for 24 hr (Shaurub et al., 2020). In Egypt, Ghoneim et al. (2022) obtained similar results of lipid reduction. Depending on their results, the lipid content in haemolymph of the EPN-infected *A. ipsilon* larvae was gradually reduced with age. The greatest reduction of lipids was determined in larvae at 48 h (26.55 & 21.73% lipid reduction, by *S. carpocapsae* or *H. bacteriophora*, respectively). As clearly seen, *S. carpocapsae* exerted stronger reducing action on the lipid content of infected larvae than *H. bacteriophora*. On the other hand, lipid content in the fat body of the red palm weevil *Rhynchophorus ferrugineus* (Coleoptera: Curculionidae) larvae increased after infection with *S. carpocapsae* or *H. bacteriophora* (Abdel-Razek et al., 2004). In Egypt, also, the *P. gossypiella* full-grown larvae were infected with LC₅₀ of *H. bacteriophora* and *S. riobrave* by Shairra et al. (2016) who recorded an increase of free fatty acids.

To interpret the reduction of total lipid content in last instar larvae of different insects after infection with some EPNs, it may be important to point out that the lipid turnover in insects is regulated by neuroendocrine-controlled feed-back loops (Kim et al., 2002; Etebariet al., 2007). Therefore, the reduced lipid content might be due to the disrupting effects and stress of these EPNs on neurosecretion or other hormones in larvae. Also, the declined lipid level in larvae might be due to shift in energy metabolism towards lipid catabolism as a result

of physiological stress caused by the tested EPNs. In other words, these EPNs induced stress on larvae to use lipids and glucose for cell repair and increasing protein catabolism which may be stimulated due to high energy demand under such stress conditions.

1.1.3. Disruptive Effects of EPNs on the Carbohydrate Content in Insects:

In insects, carbohydrates represent an important energy source and perform a crucial role in the structure and function of tissues during development and metamorphosis, as well as for the maturation of reproductive organs and embryonic development (*cf.* Chippendale, 1978). In insects, also, the soluble carbohydrates are accumulated during larval stage and utilized in metamorphosis (Pant and Kumar, 1979). Also, they are stored in the fat body as glycogen, which is converted into trehalose before releasing into the haemolymph for utilization (Gilbert and Chino, 1974). It is important to mention that carbohydrates can be disturbed by xenobiotics (Kaufmann and Brown, 2008).

Total carbohydrates in haemolymph of 4th instar larvae of *Cx pipiens* was reduced after infection with the EPN *R. culicivora*x (Schmidt and Platzer, 1979). Infection with the EPN *Mermis nigrescens* resulted in the decrease of trehalose level and reduced carbohydrate metabolism in the fat body of its host, *S. gregaria* (Gordon *et al.*, 1971). The total carbohydrate content of *S. littoralis* larvae was significantly decreased by post-infection with EPNs *S. riobrave* and *H. bacteriophora* (Ahmed *et al.*, 2014). In Egypt, Shaurub *et al.* (2020) determined decreasing carbohydrate content in the 4th instar larvae of *S. littoralis* after infection with LD₅₀ of *S. riobrave* and *H. bacteriophora* for 24 h. Results obtained by Ghoneim *et al.* (2022) in Egypt were in accordance with these reported results of decreased carbohydrates in larvae of some insects after infection with certain EPNs, since carbohydrate content was prevalently reduced in haemolymph of the EPN-infected larvae of *A. ipsilon*. Also, the reducing potency of EPNs considerably increased with the time interval of exposure (41.38 and 31.03% carbohydrate reductions by *S. carpocapsae* and *H. bacteriophora*, respectively, at 48 h).

Schmidt and Platzer (1979) recorded a remarkable decline of glucose and trehalose levels in the haemolymph of 4th instar of the mosquito *Cx. pipiens*, after infection with EPN *R. culicivora*x. In contrast, *H. bacteriophora* AS1, *H. bacteriophora* HP88, *S. carpocapsae* ALL, and *S. riobrave* ML29 significantly enhanced the total glucose content in *C. capitata* 3rd instar larvae (Shaurub *et al.*, 2015).

Reduction of carbohydrate content in haemolymph of the insect after infection with EPNs may result from the nematode's nutritional demands for glucose and accelerated glycogenolysis and/or impaired glycogenesis. In other words, decrease of carbohydrates indicated that more sugar may be metabolized to meet the energy demands of EPNs and the host leading to consumption of sugar and carbohydrate contents (Sharma *et al.*, 2011; Yazdani *et al.*, 2014; Shaurub *et al.*, 2020). The interaction between EPNs and insect larvae appeared to be primarily nutritional. Growth of the EPNs proceeds while the nutritional requirements of the host larvae deteriorate resulting in a state of physiological starvation (Ahmed *et al.*, 2014). On the other hand, the EPNs may interfere with the hormonal regulation of carbohydrate metabolism in larvae (Gade, 2004; Sugumaran, 2010) or exhibited some effects on the carboxylase activity (Mukherjee and Sharma, 1996).

1.2. Disrupted Food Consumption and Utilization in Insects by EPNs:

In insects, the physiological events that are linked to food consumption and utilization appear to be regulated by neural, endocrine and secretagogue mechanisms (Chapman, 1985). Hormones produced by the brain neurosecretory cells, corpora cardiaca and corpora allata also regulate the digestive enzyme production (Prabhu and Sreekumar, 1994). For instance, in last instar larvae of the lawn armyworm *Spodoptera mauritia* (Lepidoptera: Noctuidae), maximal feeding activity is attained at high Juvenile hormone (JH) titer but when JH titer

declines and the subsequent release of ecdysteroids, the feeding activity decreases (Balamani and Nair, 1992; Mona, 2001).

On the other hand, feeding and reproduction in insects are closely related to nutritional factors, the qualitative and quantitative aspects of which have impact on the growth, development and fecundity (Slansky and Scriber, 1985). Therefore, an understanding of the nutritional indices in relation to the efficacies of ingestion, digestion, assimilation and food conversion by the growing larvae would be physiologically useful (Scriber and Slansky, 1981). More than three decades later, Lazarevi and Tomani (2015) reported the importance of understanding the digestive physiology of an insect for both fundamental studies of the mechanisms and patterns of dietary specialization and applied studies searching for efficient strategies of the pest control.

In this context, the most important parameters of food consumption and utilization are Consumption index (CI), Approximate digestibility (AD), Efficiency of conversion of ingested food to body substance (ECI), Efficiency of conversion of digested food to body substance (ECD), Growth rate (GR) (Waldbauer 1968) and Assimilation rate (AR) (Scriber and Slansky, 1981). Relative metabolic rate (RMR) was calculated according to Slansky (1985). These parameters may help to determine the nutritional efficiencies which can affect growth in insects (Johnson and Mundel, 1987; Hinks *et al.*, 1991).

Only, a little attention has been paid to the food consumption and utilization efficiencies in entomopathogen-infected insects (Tefera and Pringle, 2003; Hussain *et al.*, 2015). To our knowledge, scarce information exists in the current literature concerning the disruptive effects of EPNs on the food consumption and utilization of insect pests. Infection of the newly moulted 4th instar larvae of *S. littoralis* with *H. bacteriophora* resulted in a decrease of CI of 2- and 3-day-old larvae (Shaurubet *al.*, 2020). To understand this reduction of food consumption, IJs of EPNs penetrate the haemocoel of the insect host, they rapidly multiply, causing lethal septicaemia in the host (Wouts, 1984). Also, heterorhabditid EPNs can rupture the host cuticle (Poinar, 1990).

In addition, AD in the entire 6th instar of *S. littoralis* increased after infection of *S. littoralis* 4th instar larvae with *S. riobrave*. In contrast, infection with *H. bacteriophora* led to a decrease of AD of 2-day-old 4th instar larvae (Shaurubet *al.*, 2020). As reported by Barnby and Klocke (1987), the increasing AD may be due to a higher retention of the food bolus in the gut for a longer period and thereby longer exposure to digestive enzymes. This will allow for more digestion and absorption of nutrients for normal biomass production. Also, the considerable increase of AD enables the infected and nutrient-deficient larvae to meet their energy and developmental requirements (Shaurubet *al.*, 2020) (See Table 2).

Table 2. Mean ($\pm$ SE) approximate digestibility (AD; %) of *Spodoptera littoralis* infected as fourth instars with the LD₅₀ of *Steinernema riobrave*, *Heterorhabditis bacteriophora*, or *Beauveria bassiana*.

Instar	Age of instar (days)	Control	<i>S. riobrave</i>	<i>H. bacteriophora</i>	<i>B. bassiana</i>
Fourth	2 ¹	94.12 $\pm$ 5.80	92.70 $\pm$ 0.68	78.12 $\pm$ 2.30*	91.67 $\pm$ 2.10
	3	88.79 $\pm$ 6.60	89.30 $\pm$ 0.80	87.16 $\pm$ 4.80	91.78 $\pm$ 3.40
Fifth	1	89.00 $\pm$ 4.20	84.34 $\pm$ 3.40	88.69 $\pm$ 4.68	93.73 $\pm$ 7.80
	2	82.22 $\pm$ 3.40	88.20 $\pm$ 3.60	81.66 $\pm$ 6.70	86.28 $\pm$ 4.30
Sixth	1	75.88 $\pm$ 3.70	83.17 $\pm$ 2.00*	75.86 $\pm$ 2.00	85.16 $\pm$ 5.40*
	2	67.36 $\pm$ 2.00	77.66 $\pm$ 5.40*	68.87 $\pm$ 3.00	76.74 $\pm$ 3.40*
	3	68.62 $\pm$ 6.20	72.77 $\pm$ 3.70*	68.00 $\pm$ 6.20	80.91 $\pm$ 3.60*

¹Newly moulted 4th instar larvae were first infected for 24 hr before conducting the experiments. *Significant difference between treatment and the respective control (Student's *t*-test: $P < 0.05$) (After Shaurubet *et al.*, 2020).

With regard to the ECI, infection of the *S. littoralis* 4th instar larvae with *S. riobrave* resulted in increasing ECI of 2-day-old 4th instar larvae and 1-day-old 5th instar larvae but decreased ECI in 3-day-old 4th instar larvae. On the other hand, infection of larvae with *H. bacteriophora* led to increased ECI of 2-day-old 4th instar larvae and 1-day-old 6th instar larvae but decreased ECI of 2- and 3-day-old 6th instar larvae (Shaurubet *et al.*, 2020). In respect of ECD, its value increased in 1-day-old 5th instar larvae but was decreased in 1-day-old 6th instar larvae after infection of 4th instar larvae with *S. riobrave*. After infection with *H. bacteriophora*, ECD value of 2-day-old 4th instar larvae and 1-day-old 6th instar larvae increased but decreased during the remaining 6th instar duration (Shaurubet *et al.*, 2020). To explicate the considerable reduction of these food utilization efficiencies of *S. littoralis* late 6th and early 5th instars after infection with *H. bacteriophora*, it can be suggested that most of the ingested or digested food was metabolized for energy production, and it is likely that the major portion of this energy was utilized for defense against the invading pathogens (Shaurubet *et al.*, 2020).

However, these food utilization efficiencies are reflected on the larval growth of *S. littoralis*. After infection with *S. riobrave*, GR of 3-day-old 4th instar larvae was decreased but increased for 5th larval instar. After infection with *H. bacteriophora*, GR of 2-day-old 4th instar larvae and 1-day-old 6th instar larvae increased but decreased of 3-day-old 4th instar larvae and 2- and 3-day-old 6th instar larvae (Shaurubet *et al.*, 2020).

2. Drastic Impacts of EPNs on Developmental and Reproductive Physiology Of The Insect Pests:

Death of the insect host is usually reached within 24-72 h post-infection with EPNs (Shapiro-Ilan and Brown, 2013; Shaurubet *et al.*, 2014; Lacey *et al.*, 2015; Leonaret *et al.*, 2022; Kaliaskaret *et al.*, 2022). However, several authors (Peschiutta *et al.*, 2014; Cardoso *et al.*, 2015; Ansari and Hussain, 2020; Shah *et al.*, 2021) reported the success of some infected individuals to survive toxic effects of both EPN and its symbiotic bacteria with immune defences. These infected individuals can transform into the next developmental stage but are usually unhealthy because certain biological and physiological processes have been detrimentally affected. In this regard, there has been increasing evidence that some EPNs adversely affect certain metabolic processes in the insect host (Shaurubet *et al.*, 2020; Ghoneim *et al.*, 2022) and some important enzymatic activities (Shaurubet *et al.*, 2015; Ibrahim *et al.*, 2015; Shairra *et al.*, 2016; Vidhya *et al.*, 2016).

Ghoneim *et al.* (2023b) observed a few EPN-infected 4th instar larvae of *A. ipsilon* pupating at the lower two concentrations of *S. carpocapsae* and *H. bacteriophora*. In this study, also, a similar result was recorded after infection of 5th instar larvae with *S. carpocapsae*. In addition, some studies, on various agricultural insect pests, recorded the capability of some EPN-infected pupae to metamorphose into adults indicating some resistance of the pupal stage to EPNs (Malan *et al.*, 2011; Odendaal *et al.*, 2016; du Preez *et al.*, 2021a, b; Steyn *et al.*, 2021). In a study of Filgueiras and Willett (2021), EPNs can affect the developmental durations and changes in the risk of death of the non-susceptible pupal stage of the onion fly *Delia antiqua* (Diptera: Anthomyiidae), indicating an insect resistance to infection during the pupal stage. Inactivation of EPN during the pupal stage of an insect and re-establishment of its activity during the adult stage indicate the ability of this EPN to overcome the insect's immune system (Vallet-Geley *et al.*, 2008).

The available literature contains little information concerning the effects of EPNs on the biological, physiological and adult performance parameters of mosquitoes (Diptera) until 2022. Early, Welch and Bronskill (1962) found the pupation of some EPNs-infected larvae of the yellow fever mosquito *Aedes aegypti* (Culicidae) with *S. carpocapsae* 1 or 2 weeks longer than the normal congeners (*i.e.*, prolongation of larval period), while others failed to

pupate and then died after 3 or 4 weeks. Also, Obiamiwe and Mac Donald (1973) recorded a delay of development of the *Anopheles* (Culicidae) larvae after infection with the nematode *Octomyomermismuspratti*. According to Liu *et al.* (2020b), EPN *Steinernema abbasi* isolate could cause a delay in the next molt and pupation of the Asian tiger mosquito *Aedes albopictus* (Culicidae). Also, Elbrense *et al.* (2022) recorded significantly prolonged pupal duration of *Cx. pipiens* as an effect of infection with LC₅₀ of EPN *R. iyengari* pre-parasites. For more detail, see review of Ghoneim and Bakr (2024).

Prolongation of the pupal duration, or delay of the pupal ecdysis, of mosquito larvae infected with mermithid nematodes owing to a disturbance in their endocrine regulation or due to the insertion of some neurosecretory compounds by the mermithid nematode *Romanomer misiyengari* (Petersen and Willis, 1970). Also, this result may be related to the insufficient nutritional reserves, which are essential for building adult structures during the pupal stage, because EPNs absorb nutrients in the host's body through their body surface (Schmidt and Platzer, 1980; Gordon, 1981). In this regard, the haemolymph protein content of the mosquito *Cx. pipiens* larvae was depleted to one-sixth of the level in normal larvae after infection with the mermithid nematode *Romanomermis culicivorax* (Schmidt and Platzer, 1978). Also, the scarcity of nutritional reserves in mosquitoes may be due to the degeneration of the mid-gut epithelium leading to starvation, as recorded for the mosquito *Aedes aegypti* larvae after infection with *R. culicivorax* (Bailey and Gordon, 1973).

Infection of pupae (the soil dwelling stage) of Tobacco thrips *Frankliniella fusca* (Thysanoptera: Thripidae) with *H. bacteriophora* (F11-1), *S. feltiae*, *S. riobrave* or *S. rarum* resulted in reduction of the adult emergence (Gulzar *et al.*, 2021). After treatment of pupae of European grapevine moth *Lobesia botrana* (Lepidoptera: Tortricidae) with high IJ concentrations of four EPN species (*S. feltiae*, *S. carpocapsae*, *S. riojaense* and *H. bacteriophora*), adult emergence of the insect was significantly blocked (Vicente-Díez *et al.*, 2021). In Egypt, Elbrense *et al.* (2022) treated *Cx. pipiens* 4th instar larvae with LC₅₀ of *R. iyengari* pre-parasites and observed partial blockage of adult emergence but no effect on the adult longevity.

Gordon (1981) suggested that EPN-infected adults suffer from being sterilized or biologically castrated. According to Elbrense *et al.* (2022), the treatment of *Cx. pipiens* 4th larval instar with LC₅₀ of *R. iyengari* pre-parasites resulted in a considerable reduction of the female fecundity. This reduction of fecundity may result from non-selective use of mosquito energy reserves under stress of EPN infection (Baudoin, 1975). On the contrary, no effect was recorded on the egg-hatching rate of *Cx. pipiens* after infection of 4th instar larvae with LC₅₀ of *R. iyengari* pre-parasites (Elbrense *et al.* (2022).

3. Enzymatic Disturbance in Insect Pests Infected with EPNs:

Insect defenses and immune reactions in response to EPN infection have been studied only in a few EPN species-insect species combinations (Lewis and Clarke, 2012; Shapiro-Ilan *et al.*, 2018). Some authors (Brown *et al.*, 2006; Ffrench-Constant *et al.*, 2007; Toubarroet *et al.*, 2009) reported that EPNs in the families of Steinernematidae and Heterorhabditidae are lethal endoparasites of insects because they can secrete active substances, including toxins, proteases, and so on, contributing to the lethal effect on infected host insects. The lethal activity of EPNs is often closely related to the activity disturbance of some important enzymes in the host insects (Grewal *et al.*, 2005; Ahmed *et al.*, 2014; Ibrahim *et al.*, 2018, 2019). Many EPNs secrete chemicals that facilitate penetration and migration through host tissues, feeding, and avoidance of host immune responses. These chemicals include digestive enzymes and toxins (Lee and Atkinson, 1976). Beside these secretions, the EPN-symbiotic bacteria also secrete certain toxins for suppressing the immune responses of the host insect leading to the host death (Koppenhöfer, 2007; Richards and Blair, 2010). For instance, *Xenorhabdus nematophila*, the symbiont bacterium of *S. carpocapsae*, secretes several enzymes, including

hemolysins, lipases, and proteases, which are thought to contribute to the virulence or nutrient acquisition for the bacterium and its nematode host *in vivo* (Van Damme *et al.*, 2016; Askitosari *et al.*, 2021; El Aalaoui *et al.*, 2022). Disturbances in the enzymatic activity during the progress of pathogen play an important role for understanding the interaction between the pathogen and host as a part of a survival strategy (Ibrahim *et al.*, 2015).

3.1. Disturbed Activities of the Major Digestive Enzymes in Insects:

In insects, digestive enzymes are primarily generated and secreted by the midgut epithelium of the alimentary duct (Chapman, 1998). Two digestive enzymes, amylase and protease, were also found in the salivary gland complex (Li *et al.*, 2017; Holtof *et al.*, 2019). The digestive enzymes lipase, amylase, and protease, which metabolize sugars, lipids, cellulose, and proteins in the mid-gut of an insect, are extremely important for the energy production and the food metabolism in insects (Gökkuş *et al.*, 2016; Bonelli *et al.*, 2020). It is important to provide an insight into the disruptive effects of EPNs on activities of the main digestive enzymes of various categories.

3.1.1. Activity Disturbance of Carbohydrate Hydrolyzing Enzymes:

In insect physiology, it is well-known that the metabolism of carbohydrates is regulated mainly by carbohydrate hydrolyzing enzymes. Glycosidases catalyse cleavage of internal bonds in polysaccharides and hydrolyse oligosaccharides as well as disaccharides (Zibae *et al.*, 2008b). In the present review, we shed some light only on three digestive enzymes, amylase, trehalase and invertase.

Disturbed Amylase Activity by EPNs:

Amylases are necessary enzymes to hydrolyze carbohydrates present in the insect larvae. They catalyze the hydrolysis of α -D-(1,4)-glucan linkages in glycogen and other related carbohydrates (Franco *et al.*, 2000). Amylases are secreted by salivary glands and mid-gut of larvae (Ribeiro *et al.*, 2000). Depending on the available literature, amylase activity increased or decreased, depending on the insect species, the EPN species, applied concentration, the EPN-symbiotic bacteria and the time of infection. For example, Zółtowska (2004) reported an increase of the enzyme activity in 7th instar larvae of the greater wax moth *Galleria mellonella* (Lepidoptera: Pyralidae) at 48 h post-infection with 20 IJ/insect of the EPN *Heterorhabditis zealandica*. Two years later, Zółtowska and Łopieńska (2006) recorded a significant reduction in the enzyme activity after infection of last instar larvae of *G. mellonella* with EPNs *Steinernema affinis* and *S. feltiae*, at different exposure periods except a significant increase at 12, 18 and 24 h post-infection with *S. feltiae*. Also, amylase activity was decreased in *S. littoralis* larvae after infection with *H. bacteriophora* but increased after infection with *S. riobrave* or *S. feltiae* (Ahmed *et al.*, 2014). After infection of the 4th larval instar larvae of *A. ipsilon* with *H. zealandica* or *S. abbasi*, amylase activity was reduced (Ibrahim *et al.*, 2015).

Some years later, Shaurub *et al.* (2020) incubated the newly moulted 4th instar larvae of *S. littoralis* with LD₅₀ of EPNs *S. riobrave* and *H. bacteriophora* for 24 h and determined increasing amylase activity. Also, four EPNs, *H. bacteriophora*, *S. carpocapsae*, *Steinernema scapterisci*, *S. glaseri* were assessed against the last instar larvae of *G. mellonella* by Khater *et al.* (2020) in Egypt. Their results demonstrated that infection with the concentration 20 IJs of *H. bacteriophora* or *S. carpocapsae* resulted in a significant reduction of the amylase activity, but both *S. scapterisci* and *S. glaseri* caused a slight increase, 6 h post-infection. At concentration 50 IJs, a significant elevation in the enzyme activity was recorded at 6 h post-infection. Meanwhile, *H. bacteriophora* caused a significant reduction in enzyme activity at 12 and 48 h post-infection. After infection of the larvae of fruit flies *Bactrocera zonata* and *C. capitata* with the EPN *S. carpocapsae*, Elhadidy *et al.* (2021) recorded a significant decrease but a slight increase of the enzyme activity in these two fly species, respectively. In a recent study, Fathy and Abd El-Rahman (2023) infected the 5th

instar nymphs and adults of the African migratory locust *Locusta migratoria migratorioides* (Orthoptera: Acrididae) with the LC₅₀ of *Steinernema* spp. and *H. bacteriophora* and estimated reduced amylase activity. Due to the toxic effects of the EPNs, *Steinernema* spp., the *H. bacteriophora*, and their associated bacteria, considerably reduced amylase activity was determined (Muhammad *et al.*, 2022).

Disturbed Trehalase Activity by EPNs:

Because trehalase plays an important role in energy supply, it is one of the most important carbohydrases in insects, hydrolyzing trehalose (the main sugar in insect haemolymph) into two glucose molecules (Shukla *et al.*, 2015). As reported in the available literature, results of several studies revealed the inhibition and reduction of trehalase activity in several insects as a result to the infection with different EPNs. For instance, Dmitryjuk *et al.* (2001) recorded a slight decrease of enzyme activity in the 3rd instar larvae of *G. mellonella* after infection with the infective juvenile (IJs) of *S. affinis*. Also, Zóltowska and Łopieńska (2006) determined a significant decrease of the enzyme activity in 7th instar larvae of *G. mellonella* at 6 and 12 h post-infection with *S. affinis* or *S. feltiae*. After infection of the 4th larval instar larvae of *A. ipsilon* with *H. zealandica* and *S. abbasi*, trehalase activity was reduced (Ibrahim *et al.*, 2015).

In addition, infection of the last instar larvae of *G. mellonella* with concentration 100 IJs of *H. bacteriophora* resulted in an inhibition of trehalase activity at 6, 24 and 48 h post-infection (Khater *et al.*, 2020). In Egypt, Elhadidy *et al.* (2021) infected the larvae of the fruit flies *B. zonata* and *C. capitata* with *S. carpocapsae* and recorded remarkably decreased enzyme activity in *B. zonata* but a slight reduction in *C. capitata*. In Egypt, also, Fathy and Abd El-Rahman (2023) infected the 5th instar nymphs and adults of *L. migratoria migratorioides* with the LC₅₀ of *Steinernema* spp. and *H. bacteriophora* and estimated decreasing trehalase activity.

Energy demands are stepped up in the insect host in initial stage of EPN infection, when the physiology of the host is changed to combat the disease as a natural response. Therefore, the decrease of enzyme activity in the EPN-infected larvae can be attributed to decreased metabolic capabilities of these larvae. This is, also, due to decreased hydrolysis of trehalase to release glucose molecules under drastic stress conditions and high energy requirement (Hasegawa and Yamashita, 1970; Ibrahim *et al.*, 2015). In addition, this reduction may be due to the toxic effects of the EPNs and/or their symbiotic bacteria (Muhammad *et al.*, 2022).

Trehalase activity in some insects increased after infection with certain EPNs. For example, Zóltowska (2004) reported an increase of the enzyme activity in 7th instar larvae of *G. mellonella* at 18 and 24 h post-infection with 20 IJ/insect of *H. zealandica*. Also, the enzyme activity increased in the *S. littoralis* larvae after infection with *H. bacteriophora*, *S. riobrave* or *S. feltiae*. The highest increase was determined in case of infection with *S. riobrave* (Ahmed *et al.*, 2014). Depending on a study conducted by Shaurubet *et al.* (2020) in Egypt, the newly moulted 4th instar larvae of *S. littoralis* were infected with LD₅₀ of *S. riobrave* or *H. bacteriophora* for 24 hr. As a response to the EPN infection, trehalase activity was promoted. However, considerable activity increases of carbohydrate-hydrolyzing enzymes, particularly trehalase, may be expected to increase the availability of glucose, both for the defense mechanisms of the insect host and for development of the invading pathogen (Shaurubet *et al.*, 2020).

Disturbed Invertase Activity by EPNs:

From the physiological point of view, invertase cleave sucrose into the monosaccharides, glucose, and fructose. This enzyme is secreted in the gut of insect larvae (Heil *et al.*, 2005). Some authors reported a reduction of invertase activity in some insects by certain EPN species, such as Ibrahim *et al.* (2015) who recorded decreasing activity after

infection of the 4th larval instar larvae of *A. ipsilon* with the EPNs *H. zealandica* or *S. abbasi*. Also, the enzyme activity was remarkably inhibited in the larvae of *B. zonata* and *C. capitata* after infection with *S. carpocapsae* (Elhadidy *et al.*, 2021). Fathy and Abd El-Rahman (2023) infected the 5th instar nymphs and adults of *L. migratoriamigratorioides* with the LC₅₀ of *Steinernema* spp. and *H. bacteriophora* and estimated decreasing activity of invertase. The inhibition of invertase activity in these insects may be due to the EPNs themselves or/ and their associated symbiotic bacteria (Muhammad *et al.*, 2022).

In contrast, invertase activity increased in *S. littoralis* larvae after infection with *H. bacteriophora*, *S. riobrave* or *S. feltiae*. The highest increase in activity was observed in the case of infection with *S. riobrave* (Ahmed *et al.*, 2014). According to Shaurubet *et al.* (2020), incubation of the newly moulted 4th instar larvae of *S. littoralis* with the LD₅₀ of EPNs *S. riobrave* and *H. bacteriophora* for 24 h resulted in increasing invertase activity. Furthermore, increase and decrease of invertase activity had been reported in the same insect after infection with the same EPN species, depending on the time interval post-infection. For instance, Khater *et al.* (2020) assessed the efficiency of four EPNs, *H. bacteriophora*, *S. carpocapsae*, *S. scapterisci*, *S. glaseri* against the last instar larvae of *G. mellonella*. Depending on their results, the infection with concentration 20 IJs of all EPNs resulted in a significant decrease of the invertase activity at 6 and 12 h post-infection, but a significant increase at 24 h.

3.1.2. Activity Disturbance of Protein Hydrolyzing Enzymes:

Peptidases (peptide hydrolases) include proteinases (endopeptidases) and (exopeptidases) (Terra and Ferreira, 2005). In insects, proteinases, also known as proteases or proteolytic enzymes, are involved in the digestive processes, proenzyme activation, liberation of physiologically active peptides, complement activation and inflammation processes amongst others (Macedo and Freire, 2011). These enzymes catalyze the hydrolysis of peptide bonds in proteins and polypeptides (Zibae, 2012). Serine and cysteine proteinases are the major proteinase classes in the digestive systems of phytophagous insects (Haq *et al.*, 2004).

As previously described, IJs of many EPN species secrete chemicals to facilitate their penetration through the host tissues. These chemicals include digestive enzymes and toxins. Therefore, proteases are suggested as essential enzymes for the pathogenicity of the many EPNs. Haemolymph of larvae of many insects contains inhibitor (s) that inhibits proteases of both the invading IJs and their symbiotic bacteria. The inhibitor is produced during the second period of infection when the larval defense system has already been overcome and infection is established (Kucera and Mracek, 1989). For instance, when IJs of EPN *S. glaseri* were treated with protease inhibitors and injected into *G. mellonella* gut, the mortality% of larvae was reduced, and the nematode penetration of the larval gut was reduced (Abu Hatab *et al.*, 1995). Digestive enzyme inhibitors have been reported to be important in the control of insect pests (Franco *et al.*, 2000; Mehrabadi *et al.*, 2012).

Besides digestion, trypsin-like enzymes contribute in a wide range of other physiological processes in insects, such as molting (Wei *et al.*, 2007), tissue remodeling (Liu *et al.*, 2009), innate immunity (Kanost *et al.*, 2004), diapause (Chen *et al.*, 2005), fertilization (Friedlander *et al.*, 2001), and activation of enzyme precursors of trypsin, chymotrypsin (Parde *et al.*, 2012), chitinase (Royer *et al.*, 2002), and phenoloxidase (Yu and Kanost, 2004). In addition, trypsin can affect biology of the pathogenic organisms (Nakazawa *et al.*, 2004; Darpel *et al.*, 2011) and may have a role in preventing infection by parasites (Jochim *et al.*, 2008; Sant'Anna *et al.*, 2009).

The current literature contains very few studies investigating the inhibition of protease activity in some insects after infection with certain EPN species (Wang *et al.*, 2012; Ibrahim, *et al.*, 2018). The protease activity in haemolymph of the infected 4th instar larvae of *A. ipsilon* decreased after 16 h of infection with *H. zealandica* or *S. abbasi* (Ibrahim *et al.*, 2015).

The determined decrease in protease activity could be due to the symbiotic bacteria beginning to digest the proteins of the insect body 16 h post-infection (Ibrahim *et al.*, 2015). Fathy and Abd El-Rahman (2023) infected the 5th instars and adults of *L. migratoriamigratorioides* with LC₅₀ of *Steinernema* species and *H. bacteriophora* and estimated decreasing activity of protease.

3.1.3. Activity Disturbance of Lipid Hydrolyzing Enzymes:

Lipases catalyze the hydrolysis of fatty acid ester bonds (Grillo *et al.*, 2007). They are able to hydrolyze a variety of esters in organic solvent systems (Zibaeet *et al.*, 2008a). Also, hydrolysis of triacylglycerol into its free fatty acids and glycerol backbone is one of the many reactions that lipase is involved in (Yao *et al.*, 2021).

To the best of our knowledge, scarce results were available in the current literature about the impacts of EPNs on the lipid hydrolyzing enzymes in insects. The only available study was conducted in Egypt by Fathy and Abd El-Rahman (2023). They infected the 5th instar nymphs and adults of *L. migratoriamigratorioides* with the LC₅₀ of *Steinernema* spp. and *H. bacteriophora* and estimated increasing lipase activity. This result can be understood in view of the activity of EPN-symbiotic bacteria *X. nematophila* or *P. luminescens* which secrete various enzymes, such as hemolysis, lipases, and proteases, that contribute to pathogenicity or nutrient acquisition for the bacterium and its nematode host, lipase activity was dramatically boosted (Richards and Blair, 2010).

3.1.4. Activity Disturbance of Some Digestion-Related Enzymes:

As reported by several authors (Merzendorfer and Zimoch, 2003; Subbanna *et al.*, 2018; Henriques *et al.*, 2020), chitin is mostly produced by fungi, arthropods, and nematodes and makes up a significant portion of the insect cuticle (a complicated layer in the body wall). It supports the cuticles in the body wall and trachea, the peritrophic matrices lining the gut epithelium, and insect development and morphogenesis in insects. Chitinase protein secreted by the EPN-symbiotic bacteria, *Xenorhabdus* and *Photorhabdus* spp., plays a key role in the virulence of these bacteria by both degrading cuticle of the insect host and accelerating the binding process of toxins to the target sites (Liu *et al.*, 2020a; Mahmood *et al.*, 2020). Results of Gümüşsoy *et al.* (2022) on the codling moth *Cydia pomonella* (Lepidoptera: Tortricidae) suggested that toxic metabolites, including chitinase protein, are present in the cell-free supernatant of symbiotic bacteria and are responsible for mortality. Fathy and Abd El-Rahman (2023) infected the 5th instar nymphs and adults of *L. migratoriamigratorioides* with LC₅₀ of EPN *Steinernema* spp. and *H. bacteriophora* and estimated increasing activity of chitinase. Due to the presence of EPN-symbiotic bacteria, *X. nematophila* or *P. luminescens*, which create chitinase for their growth rate, the amount of chitinase was considerably increased (Chen *et al.*, 1996).

3.2. Disturbed Activities of the Major Detoxifying Enzymes in Insects Infected with EPNs:

In insects, detoxification enzymes have many functions for repairing physiological processes, detoxifying pathogenic products, and metabolizing the biologically active compounds (Visetson and Milne, 2001; Zhu-Salzman and Zeng, 2015). Also, they play a crucial role for decomposing toxic substances into non-toxic constituents, decreasing the toxicity of xenobiotics in order to maintain the normal physiological and biochemical criteria in the body (Mukanganyama *et al.*, 2003; Li and Liu, 2007) as well as they act as protectants against different oxidative stresses (Zibaeet *et al.*, 2011).

In insects, detoxification can be achieved by different families of enzymes including esterases (ESTs), phosphatases, transaminases, glutathione S-transferase (GSTs), glutathione peroxidase (GPx), glutathione reductase (GR), superoxide dismutase (SODs), catalases (CATs), monooxygenases (MOs), acetylcholinesterase (AChE), multi-function oxidases (MFOs), peroxidases (PODs), as well as tyrosinase (TYR) and carboxylesterase (CarE) (Li

et al., 2007; Zibaeet *et al.*, 2011). However, Li *et al.* (2022) reported SODs, PODs, and CATs as antioxidant enzymes and CarE, GST and AchE as detoxifying enzymes, since insects are known to use antioxidants and detoxifying enzymes in their immune systems as defense reactions against pathogen threats (Dubovskiyet *et al.*, 2012). On the other hand, antioxidant enzymes possess the function of scavenging excessive reactive oxygen species (ROS), degrading hydrogen peroxide (H₂O₂), and preventing free-radical-associated damage (Dubovskiyet *et al.*, 2008; Zhou *et al.*, 2019). In general, insects activate several defense mechanisms, viz., Phenoloxidase, hemocytes, detoxification and antioxidant enzymes (Lalitha *et al.*, 2018). Therefore, fluctuation in activities of these enzymes may have an impact on the insect adaptability to the surrounding environment (Serebrov *et al.*, 2001, 2006; Li *et al.*, 2016).

3.2.1. Disturbed Activities of Phosphatases in Insects:

As clearly reported by some authors (Srinivas *et al.*, 2004; Zheng *et al.*, 2007), phosphatases have been included in the category of detoxifying enzymes. As a lysosomal enzyme, acid phosphatase (ACP) may have a role in autophagy and cell turnover as well as defense (Xia *et al.*, 2000). ACP and alkaline phosphatase (ALP) are hydrolytic enzymes, and they are responsible for removing phosphate groups from many types of molecules, including nucleotides, proteins, and alkaloids in alkaline and acidic conditions, respectively in a process called 'dephosphorylation' (Zibaeet *et al.*, 2011). These two phosphatases not only hydrolyze phosphate groups from the ingested toxic molecules and naturally occurring ones, but also their activities may refer to the efficiency of digestion and transportation within insect body (Nation, 2008; Zibaeet *et al.*, 2011). Appearance of ALP in the plasma membrane, its activity is affected due to cell membrane damage, so ALP is also considered as a biomarker for cellular stress (Lalitha *et al.*, 2018).

Disturbed Activity of ACP in Insects:

According to the current literature, there are many reported results of increasing ACP activity in larvae of various insects after infection with different EPN species, such as larvae of *C. capitata* after infection with *S. riobrave* and *H. bacteriophora* (Soliman, 2002); larvae of *S. littoralis* after infection with *S. riobrave* and *H. bacteriophora* (Ahmed *et al.*, 2014); *A. ipsilon* 6th instar larvae at 6 hr post- infection with *S. glaseri* or *H. bacteriophora* (Hassan *et al.*, 2016); the 5th nymphs of *S. gregaria* after infection with 1000 IJs of *H. bacteriophora* (Gaber *et al.*, 2018); the 3rd instar larvae of *S. littoralis* after infection with *H. bacteriophora* or *S. carpocapsae* (Gomaa *et al.*, 2020); and the 4th instar larvae of *S. littoralis* after infection with LD₅₀ values of *S. riobrave* or *H. bacteriophora* for 24 h (Shaurubet *et al.*, 2020). Ghoneimet *et al.* (2023a) determined significantly increased ACP activity in haemolymph of 6 h-old last instar larvae of *A. ipsilon* after infection of the penultimate instar larvae with *S. carpocapsae* or *H. bacteriophora*. Moreover, *H. bacteriophora* exhibited a higher enhancing potency on the enzyme activity than *S. carpocapsae*, at this time (See Table 3). In addition, increased ACP activity had been reported in haemolymph of some insects after infection with some entomopathogens (Xia *et al.*, 2001; Mirhaghpourastet *et al.*, 2013; Vidhya *et al.*, 2016).

The increasing ACP activity in larvae of these insects after infection with certain EPNs may indicate a physiological adaptability to combat the oxidative stressor may be related to an inhibition of lipid peroxidation process and physiological response mechanism against the EPN toxic secretions. Also, the increasing activity of ACP may be attributed to its role in the transportation of lipids, such as monoacylglycerols, through low density lipoproteins to supply the energy demands of the treated larvae (Khorshidi *et al.*, 2019). In addition, the increase of ACP activity in larva may be due to the phagocytosis of certain hemocytes to the invading EPNs, since phagocytosis is known to stimulate the production of lysosomal enzymes, and ACP acts as a lysosome marker (Xia *et al.*, 2000; Hassan *et al.*, 2016). In other words, the increase of ACP activity in haemolymph may be an attempt by the

infected larvae to defend against the invading pathogens, where lysozymes are a component of humoral immunity (Mohamed *et al.*, 2016; Satyavathi *et al.*, 2018; Dorrah *et al.*, 2019).

Table 3. Acid phosphatase activity in haemolymph (U/L) of last (6th) instar larvae of *A. ipsilon* as influenced by infection of the newly moulted 5th instar larvae with LC₅₀ values of the tested Entomopathogenic nematodes.

Nematode species		Time interval		
		6 hr	24 hr	48 hr
<i>S. carpocapsae</i>	mean±SD	98.56±2.17 b	54.45±1.02 c	41.26±1.31 c
	Change (%)	+44.62	-30.38	-49.57
<i>H. bacteriophora</i>	mean±SD	115.18±1.93 c	70.72±1.21 b	57.29±1.10 b
	Change (%)	+69.01	-9.59	-29.98
Control	mean±SD	68.15±2.10	78.22±1.95	81.82±2.53

B, c: see footnote of table 1.(After Ghoneim *et al.*, 2023).

In contrast, ACP activity was remarkably decreased in haemolymph of 24 and 48 hr-old last instar larvae of *A. ipsilon*, after infection of the penultimate instar larvae with *S. carpocapsae* or *H. bacteriophora* (Ghoneim *et al.*, 2023a). To some extent, similar results of decreased ACP in some insect larvae after infection with some EPN species had been reported, such as the late 3rd instar larvae of *C. capitata* after infection with *H. bacteriophora* AS1, *H. bacteriophora* HP88, *S. carpocapsae* ALL, and *S. riobrave* ML29 (Shaurubet *et al.*, 2015) and 5th nymphs of *S. gregaria* after infection with 2000 IJs (high concentration) of *H. bacteriophora* (Gaber *et al.*, 2018). For some detail, see studies of Żółtowska *et al.* (2006), Wu *et al.* (2013) and Ahmed *et al.* (2014).

However, the decrease of ACP activity can be due to the inability of the cell to undergo enzymatically controlled reactions under EPN infection (Soliman, 2002). In addition, decreasing ACP activity may be due to the reduced phosphorus liberation for energy metabolism and decreased rate of metabolism, as well as decreased rate of transport of metabolites (Senthil Nathan *et al.*, 2005). Also, the decline of ACP level in larvae, as a response to infection with the present EPNs, may be due to strong inhibition of ecdysone which is followed by subsequent decrease in number of lysosomes and in turn declined level of ACP (Hassan, 2002).

Disturbed Activity of ALP in Insects:

Based on the available literature, there are many reported results of increasing ALP activity in several insects as response to the infection with certain EPNs. For instance, Soliman (2002) recorded an increase of ALP activity in the last instar larvae of *C. capitata* after infection with *S. riobrave* or *H. bacteriophora*. In Egypt, Ahmed *et al.* (2014) reported an increasing activity of ALP in *S. littoralis* larvae as response to the infection with *S. riobrave* or *H. bacteriophora*. Also, ALP activity increased in the 3rd instar larvae of *S. littoralis* after infection with two EPN isolates (*H. bacteriophora* and *S. carpocapsae*) and the entomopathogenic fungus *Beauveria bassiana*, separately or in combination (Gomaa *et al.*, 2020). In addition, increasing activity of ALP was recorded in 4th instar larvae of *S. littoralis* after incubation with the LD₅₀ values of *S. riobrave* and *H. bacteriophora* for 24 h (Shaurubet *et al.*, 2020). Recently, Ghoneim *et al.* (2023a) obtained results consistent with those reported results, since ALP activity increased in the haemolymph of *A. ipsilon* at 6 hr of last instar larvae after infection of penultimate instar larvae with *S. carpocapsae* or *H. bacteriophora*.

On the contrary, ALP activity in larvae of some insects was reduced after infection with certain EPNs, such as a slight decrease of ALP activity in *S. littoralis* larvae after infection with *S. feltiae* (Ahmed *et al.*, 2014); significantly declined ALP level in the late 3rd instar larvae of *C. capitata* after infection with four EPNs *H. bacteriophora* AS1, *H. bacteriophora*

HP88, *S. carpocapsae* ALL and *S. riobrave* ML29 (Shaurubet *et al.*, 2015); and decreasing activity of ALP in the 5th instar nymphs of *S. gregaria* at 1000 and 2000 IJs of *H. bacteriophora* (Gaber *et al.*, 2018). Also, significantly declined ALP activity was recorded in some insects infected with other EPNs (Żółtowska *et al.*, 2006; Wu *et al.*, 2013). In accordance with these reported results, Ghoneim *et al.* (2023a) found a considerable reduction of ALP activity in haemolymph of 24 h- and 48 h-old last instar larvae of *A. ipsilon* after infection of penultimate instar larvae with *S. carpocapsae* or *H. bacteriophora*. Depending on the reduction of ALP activity at 48 h, *S. carpocapsae* exhibited stronger reducing potency than *H. bacteriophora*.

3.2.2. Disturbed Activities of Transaminases in Insects:

Glutamic oxaloacetic transaminase (GOT, or Alanine aminotransferase, ALT) and Glutamic pyruvic transaminase (GPT, or aspartate aminotransferase, AST) are the key enzymes of transamination within the intermediary metabolism of insects to make amino acids available for essential biochemical requirements (Plant and Morris, 1972). Some authors (İçen *et al.*, 2005; Etebariet *et al.*, 2005) reported that these transaminases constitute a strategic link between carbohydrate and lipid metabolism. Activities of these enzymes are known to be disturbed under various physiological and pathological conditions. In other words, disturbance of these enzymes in insects denotes biochemical impairment and lesions of tissues and cellular function because they are involved in the detoxification processes and metabolism (Enan and Berberian, 1986).

There are many reported results of decreased activities of transaminases in some insects after infection with certain EPNs. For instance, GOT and GPT activities were decreased in *C. capitata* last instar larvae after infection with *S. riobrave* and *Heterorhabditis* sp. (Soliman, 2002). Activities of both enzymes were significantly decreased in the late 3rd instar larvae of *C. capitata* after infection with *H. bacteriophora* AS1 and *S. carpocapsae* ALL (Shaurubet *et al.*, 2015). The activities of these enzymes were remarkably decreased in the 4th instar larvae of *S. littoralis* at 48 h post-infection with *H. bacteriophora* and *S. riobrave* (Ahmed *et al.*, 2014). Also, their activities had been decreased in the 3rd instar larvae of *S. littoralis* after infection with EPNs *H. bacteriophora* and *S. carpocapsae* and the fungus *B. bassiana*, separately or in combination (Gomaa *et al.*, 2020). The newly moulted 4th instar larvae of *S. littoralis* were infected with LD₅₀ values of *S. riobrave* and *H. bacteriophora* for 24 h, leading to reduction in activities of these transaminases (Shaurubet *et al.*, 2020). In a recent study of Ghoneim *et al.* (2023a), the disturbance of transaminase activities depended on the time interval post-infection, since GOT activity significantly increased in the haemolymph of *A. ipsilon* last instar larvae at 6 h post-infection with *S. carpocapsae* or *H. bacteriophora*. At 24 h, GOT activity considerably increased by *H. bacteriophora* while decreased by *S. carpocapsae*. At 48 h time interval, GOT activity was remarkably decreased by both EPNs. With regard to GPT, its activity was predominantly declined after infection with *S. carpocapsae* or *H. bacteriophora*, at all-time intervals, with few exceptions.

To explain the reduction of GOT and GPT activities in the insect larvae after infection with some EPNs, some suggestions can be accepted. The decreasing activities of these enzymes may be attributed to the significant decline in the free amino acids content (Kaur *et al.*, 1985), the quantum of which directly influences the activity of transaminases at the time of protein synthesis (Soliman, 2002). Also, it may be due to a disruption of the link between carbohydrates and protein metabolism (Azmi *et al.*, 1998). Reduction in transaminase activities has been indicated to lead to the required reduction in the host protein synthesis, since these are generalized enzymes involved in dephosphorylation and energy transfer (Shaurubet *et al.*, 2020). Also, reduction of GPT activity may be due to the interference of the EPNs with the hormonal regulation of protein synthesis and neurosecretory hormones involved in the regulation of transaminase levels (Abulyazid *et al.*, 2005). In addition, this

reduction of GPT activity may be due to the effects of EPNs on the synthesis or functional levels of this enzyme directly or indirectly by altering the cytomorphology of the cells (Nath, 2000).

3.2.3. Disturbed Activity of Acetylcholinesterase in Insects:

Acetylcholinesterase (AChE, EC 3.1.1.7) is a key enzyme for catalyzing the hydrolysis of acetylcholine (ACh), a neurotransmitter, in the nervous system (Wang *et al.*, 2004; Zibae, 2011). AChE is primarily responsible for the termination of cholinergic neurotransmission at synapses in both insects and humans (Carlier *et al.*, 2008). Inhibition of AChE results in an excessive accumulation of ACh in the synapses which in turn leaves the ACh receptors permanently open, leading to hyperactivity and consequently paralysis and death (Soreq and Seidman, 2001; Fournier, 2005). AChE activity is one of the main resistance mechanisms in various insect species against the chemical insecticides, like organophosphorus compounds, since it degrades (through its hydrolytic activity) the neurotransmitter ACh, producing choline and an acetate group (Kozakiet *al.*, 2001; Li and Han, 2002; Yu, 2006).

Many studies have focused on the activity of AChE of insects for investigating insecticide toxicology, but few studies reported the effects of EPNs on AChE activity in insect hosts (Shairraet *al.*, 2016b). In this context, the AChE activity in some insects elevated after infection with certain EPNs, as reported by Soreq and Seidman (2001). Also, some investigators (Wu *et al.*, 2013; Han-dong *et al.*, 2013) recorded significantly increased AChE activity in *G. mellonella* larvae after infecting with EPN *H. beicherriana*.

In Egypt, Ibrahim *et al.* (2015) determined a significant increase of AChE activity in the 4th larval instar larvae of *A. ipsilon* during the first 8 hs of *H. zealandica* infection. In Egypt, also, Shairraet *al.* (2016b) reported that the activity of AChE was significantly enhanced in a dose-dependent manner in *A. ipsilon* larvae after infection with EPN *S. carpocapsae* or *S. scapterisci*. In China, Li *et al.* (2022) investigated the impact of EPNs and entomopathogenic bacteria, separately and in combination, on the AChE activity in the dark black chafer *Holotrichia parallela* (Coleoptera: Scarabaeidae). Based on their results, the enzyme activity was almost enhanced in larvae during the early period (24 h) after infection with *H. beicherriana*, *Bacillus thuringiensis* (Bt), or EPN+Bt combination. Moreover, the AChE activity after EPN+Bt treatment was higher than EPN or Bt infection alone.

Inhibition of the AChE activity was determined in 4th larval instar larvae of *A. ipsilon* after infection with *S. abbasi* (Ibrahim *et al.*, 2015). The enzyme activity in insects depends on the EPN concentration, since it was enhanced in the mealworm beetle *Tenebrio molitor* (Coleoptera: Tenebrionidae) larvae after infection with lower concentrations of EPN *H. beicherriana*, but was inhibited at higher concentrations (Li *et al.*, 2016). Enzyme activity was time post-infection dependent, since Elhadidy *et al.* (2021) recorded a significant increase of activity in the larvae of both fruit flies *B. zonata* and *C. capitata* 24 h post-infection with *S. carpocapsae* that decreased later on.

Increased activity of AChE may be the overreactive stress response to the EPNs infection. This overreaction often leads to death of the host insects (Ibrahim *et al.*, 2015; Shairraet *al.*, 2016b). Significant increase of AChE was determined in insect larvae 24 h post-infection with EPNs, which correspond to early infection and development of these nematodes and the mass production of the symbiotic bacteria, respectively (Wu *et al.*, 2013). According to Li *et al.* (2016), the increase in AChE activity may be correlated with secretions produced by the EPN/symbiotic bacteria complex and may be correlated with the virulence of the EPNs against host insects, although the mechanism of destruction is unknown!

3.2.4. Disturbed Activity of Carboxylesterase in Insects:

Carboxylesterase (CarE) is known as the primary metabolic and hydrolyzing enzyme that hydrolyzes various compounds, including thioester, aliesterase, sulfate, and amides, thus reducing the toxicity of exogenous xenobiotics (Oakeshott *et al.*, 2005; Montella *et al.*, 2012;

Hatfield *et al.*, 2016) or degrading the toxic substances (Zhang *et al.*, 2011). In other words, CarE is a crucial detoxifying enzyme playing an important role in insecticide resistance and has been associated with resistance to several insecticide classes in many insects (Ranson *et al.*, 2002). Due to its role in insecticide metabolism, determination of CarE activity is used as a biochemical indicator of insecticide resistance in many insects (de Carvalho *et al.*, 2006).

In addition to its role in insecticide resistance, the current literature contains some studies investigating the disturbance of CarE activity in several insect pests by the infection with EPNs. For example, activity of CarE was significantly enhanced in a dose-dependent manner in *G. mellonella* larvae after infection with EPN *H. beicherriana* (Wu *et al.*, 2013). The CarE activity increased in 4th larval instar larvae of *A. ipsilon* with the increase of time exposure to *S. abbasi* (Ibrahim *et al.*, 2015). Based on a study conducted by Li *et al.* (2022), activity of CarE in *H. parallela* larvae was almost enhanced during the early period (24 h) post-infection with EPN *H. beicherriana*, *B. thuringiensis* (*Bt*), or EPN+*Bt* combination. Moreover, the enzyme activity of EPN+*Bt* treatment was higher than EPN or *Bt* infection alone. Also, the CarE activity in insects depends on the EPN concentration, since it was enhanced in *T. molitor* larvae after infection with lower concentrations of EPN *H. beicherriana*, but was inhibited at higher concentrations (Li *et al.*, 2016).

According to Wu *et al.* (2013), the significantly enhanced activity of CarE may be the overreactive stress response to the EPNs infection and the overreaction can lead to the death of host insects because many lipids would be disintegrated, which would damage organs and tissues. It seems that the host insects overreacted to the EPNs infection by enhancing the CarE activity first, indicating that many lipids would be disintegrated, which would damage organs and tissues, and lead to the death of insect host. Then, the EPNs-infection response would consume the CarE resulting in decreased CarE activity. Ibrahim *et al.* (2015) explained the increase of CarE activity in larvae as inability to overcome the toxic protein released by the *S. abbasi*-symbiotic bacteria *Xenorhabdus* in the larval haemocoel. For more interpretation, Li *et al.* (2016) proposed that the insecticidal active substances, secreted by the symbiotic bacteria of EPNs during reproduction, can act as xenobiotic toxic substances and induce defensive enzymes, including CarE. Later, the EPNs-infection response can consume the CarE, resulting in decreased CarE activity. This is in accordance with Muñoz *et al.* (2006) who reported that a heavy parasitic burden with a long-established infestation would result in a generalized host weakness, inducing the immune depression and insecticide resistance in invertebrates. On the other hand, the decreases in CarE activity in the larvae infected with *H. zealandica* indicated that the insect immunity system is distributed so it cannot resist the toxic substances released by the symbiotic bacteria *Photorhabdus* in the larvae haemocoel (Ibrahim *et al.*, 2015).

3.2.5. Disturbed Activities of Esterases in Insects:

Esterases (ESTs) constitute a large class of enzymes, most of which are important in metabolism because they can break ester bond through hydrolysis (Sivakumarm and Maya, 1991; Hassan and Mohamed, 2008). Also, some ESTs play a critical role in the detoxification of synthetic chemicals (Shen and Dowd, 1991). Non-specific ESTs perform important functions in the degradation of toxins of different origins (Ibrahim *et al.*, 2019). Some of these enzymes show the strongest reaction to environmental stimulation (Hemming and Lindroth, 2000). The disintegration inactive poisonous particles with GSTs and ESTs through infections have an opener function in saving insects from pathogens (Dubovskiy *et al.*, 2012).

As clearly shown by the available literature, few studies concerned with the impacts of EPNs on the activities of ESTs in insects. For instance, Shairra and Awad (2011) evaluated the effect of EPN *H. bacteriophora* infection on the ESTs activities in nymphs of *S. gregaria*. Depending on their results, α -esterase activity significantly increased by the increasing time of infection, but β -esterase activity severely declined with increasing time post-infection.

Gaber *et al.* (2018) investigated the disruptive effect of the same EPN on the same enzymes in 5th nymphs of the same locust. According to their results, α -esterase activity significantly increased, in a dose-dependent course, but activity of β -esterase increased at the low concentration and dramatically decreased at high concentration. The 5th instar nymphs of another locust, *L. migratoria*, were infected with EPN *S. carpocapsae* by Abd-El Wahed and Elhadidy (2018). As a result, α -esterase activity significantly decreased at 48 h post-infection but not 24 h, while β -esterase activity significantly increased at 24 h post-infection.

Larvae of *G. mellonella* were infected with LC₅₀ concentrations of EPN (*H. zealandica*) and fungus (*B. bassiana*), both separately and in a combination, by Ibrahim *et al.* (2019). They recorded a significant increase of Non-specific EST activity in the case of both combined application and *H. zealandica* infection alone. Later on, last instar larvae of the same insect were infected with EPNs, *H. bacteriophora*, *S. carpocapsae*, *S. scapterisci* and *S. glaseri* by Khater *et al.* (2020). Depending on their results, infection with the concentration 50 IJ of *S. glaseri* led to an initial increase of α -esterase activity which was then significantly decreased later. Infection with *S. carpocapsae* led to a significant decrease of β -esterase activity 12 h post-infection. On the contrary, β -esterase activity increased at the highest concentrations, 24 and 48 h post-infection. Unfortunately, no scientific interpretation was provided for these reported results.

3.2.6. Disturbed Activity of Glutathione S-transferase in Insects:

Glutathione S-transferases (GSTs) constitute a large class of multifunctional intracellular enzymes and are widely distributed in prokaryote and eukaryote, such as animals, insects, plants, and microorganisms (Francis *et al.*, 2005). GSTs are known as biotransformation enzymes in all organisms and play a central role in the intracellular material transportation and hormone synthesis (Hayes *et al.*, 2005). Also, GST is one of the most important detoxification enzymes present in insects and represents the first of all enzymes associated with insecticide degradation. Its activity has been found to increase in insects resistant to insecticides (Clark, 1990; Papadopoulos *et al.*, 2000). For some detail, GST is involved in the detoxification of both endogenous and exogenous xenobiotics, natural and synthetic compounds in insects, *via* glutathione conjugation, dehydrochlorination and glutathione peroxidase activity (Francis *et al.*, 2005) to become more water soluble and less toxic (Hyslet *et al.*, 2007; Oruc, 2011; Erdem and Büyükgüzel, 2015). Thus, xenobiotics have increased solubility and are excreted from the insect body by the production of mercapturic acid derivatives (Enayati *et al.*, 2005; Li *et al.*, 2007; Ramsey *et al.*, 2010).

In addition, GST participates in the cellular antioxidant defenses against oxidative stress, since it contributes to metabolite removal and protection of tissues from damage by free radicals *via* the detoxification of lipid peroxides caused by oxidative stress (Leaver and George, 1998; Papadopoulos *et al.*, 2000). It is important for maintaining the homeostasis of oxidation and antioxidant in insects by removing redundant free radicals (Sheehan *et al.*, 2001; Wu *et al.*, 2013) beside its role as good biomarker to detect exposures to metals, organic pollutants (Yang *et al.*, 2002) and pesticides (Taysse *et al.*, 1998). The disintegration inactive poisonous particles with GSTs and ESTs through infections have an opener function in save insects from pathogens (Dubovskiy *et al.*, 2012).

According to some results reported in the current literature, GST activity increased or decreased in insects after infection with certain EPN species depending on some factors. After infection of *T. molitor* larvae with different concentrations of EPN *H. beicherriana*, Li *et al.* (2016) determined remarkable enhancement of the GST activity at lower EPN concentrations, but inhibition at higher concentrations. This finding was consistent with a previous study which reported an induction of GST in the *G. mellonella* larvae after infection by *H. beicherriana* (Wu *et al.*, 2013). Also, larvae of *G. mellonella* had been infected with the LC₅₀ of EPN (*H. zealandica*) and entomopathogenic fungus (*B. bassiana*), separately and

in a combination. GST activity significantly increased in 36 hr, then decreased to 48 hr post-combined infection, in all treatments (Ibrahim *et al.*, 2019). Based on the results of Li *et al.* (2022), activity of GST in larvae of *H. parallela* was almost enhanced during the early period (24 hr) after EPN *H. beicherriana*, Entomopathogenic bacterium *B. thuringiensis* (*Bt*), or EPN+*Bt* combination exposure. Moreover, the enzyme activity of EPN+*Bt* treatment was higher than EPN or *Bt* infection alone. As recorded by Shairra *et al.* (2016b), the activity of GST was significantly enhanced in a dose-dependent manner in *A. ipsilon* larvae after infection with *S. carpocapsae* or *S. scapterisci*. Also, an increased GST activity was observed in a time dependent manner in *S. litura* larvae after infection with *H. indica* (Lalitha *et al.*, 2018).

The significantly enhanced activity of GST at early time interval of EPN infection may be the overreactive stress response to the EPNs infection and the overreaction could lead to the death of host insects (Shairra *et al.*, 2016b). However, the decreased GST activity 16 h post-infection indicates that EPNs infection may consume GST through detoxification reactions of glutathione-dependent enzymes (Wu *et al.*, 2013). In other words, the increase in GST activity in larvae may be due to the detoxification reactions and adaptive responses to neutralize EPN-induced oxidative stress (Li *et al.*, 2016).

3.3. EPN Infection Interferes with Activities of the Major Humoral Immunity-Related Enzymes in Insects:

3.3.1. Humeral Immune Defences in Insects- A Synopsis:

Insects have several defense reactions against the invading pathogens, including the morphological and immunological defenses (Moret and Siva-Jothy, 2003; Kunc *et al.*, 2017). The innate immunity-defense strategies can be classified into two major classes, cellular immune reactions and humoral immunereactions (Berger and Jurcova, 2012). Cellular immune response is provided by certain types of the freely circulating hemocytes, whose population increases in response to the microbial pathogenicity, since hemocytes are responsible for the formation of cell aggregates; nodulation, phagocytosis and encapsulation (Chapman, 1998; Hoffmann, 2003; Irving *et al.*, 2005). In this regard, the pathogen infection was reported to activate a host of immune activities, most prominent among them being phagocytosis and encapsulation by hemocytes (Hoffmann, 1996). For some details of cellular immune defences in insects, see Lavine and Strand (2002), Kanost *et al.* (2004), Kanost *et al.* (2004), Ribeiro and Brehelin (2006), Strand (2008) and Ghoneim *et al.* (2021).

Humoral innate immune responses in insects involve many antioxidant enzymes and antibacterial proteins produced in the fat body and hemocytes of insects (Wang and Zhang, 2008). In insects, some of the major antioxidant defense enzymes, involved in scavenging of free radicals, are peroxidases, Superoxide dismutases, glutathione peroxidase, catalases, reduced glutathione and glutathione reductase (Mates, 2000; Cytrynska *et al.*, 2007; Lalitha *et al.*, 2018). In insects, also, there are some secondary metabolites, produced by the EPN-symbiotic bacteria, active against insects, and the pathogenicity of these toxins to insects occurs *via* the suppression of these immune responses in the insect host (Ullah *et al.*, 2014). These toxic metabolites result in generation of reactive oxygen species (ROS). These free radicals are highly reactive and result in harmful effects on cells and tissues in the host organism. Living organisms must, therefore, remove or scavenge ROS before cell damage occurs (Feig *et al.*, 1994). Also, certain components of the insect immune system produce ROS as a tool to limit microbial growth (Lalitha *et al.*, 2018).

It is important to point out that the low levels of ROS are not harmful to living cells and may even perform useful functions, whereas high levels of ROS are dangerous through reactions with many intracellular targets, including proteins, lipids, and DNA. In other words, the lethal process of an Entomopathogenic organism is related to considerably elevated levels of ROS (Wang *et al.*, 2001; Wu and Liu, 2012), as well as activity disturbance of some

proteolytic enzymes in the insect host (Grewal *et al.*, 2005). However, excess production of ROS and other free radical intermediates produced in response to environmental stress, such as pathogen infections, can cause significant damage to cells due to oxidative stress (Rahman and Macnee, 2000) because ROS-induced damage result in the cell death or sublethal injury, such as mutations, chromosomal aberrations, or carcinogenesis (Wang *et al.*, 2001).

EPNs and their symbiotic bacteria collaborate to suppress the immune response of the insect host for achieving a successful infection (Dowds and Peters, 2002). For instance, EPN *Steinernema carpocapsae* and its symbiotic bacterium *Xenorhabdus nematophila* can inhibit the antibacterial peptide immune reaction of insects (Binda-Rossetti *et al.*, 2016). In a study, Toubarro *et al.* (2013) found *S. carpocapsae* displaying destructive approaches for host immunity through proteolytic secretion which inhibits host immunological defenses. The EPN-symbiotic bacteria, *Photorhabdus* spp. and *Xenorhabdus* spp. displayed similar lifestyles but have different molecular defensive mechanisms (Goodrich-Blair and Clarke, 2007). The symbiotic bacteria *Xenorhabdus* spp. inhibits the host's immune system by producing a variety of toxins and carrying type III effector molecules that may interfere with the actin cytoskeleton and prevent phagocytosis (Dillman *et al.*, 2012). *Photorhabdus* spp. use lipopolysaccharide (LPS) modification to resist the action of the host-derived antimicrobial peptide (AMPs) (Eleftherianos *et al.*, 2006), while *Xenorhabdus* spp. prevents induction of insect AMP expression altogether (Istkharet *et al.*, 2019). In insects, the pathogenic impacts of symbiotic bacteria and the anti-bacterial resistance mechanisms have been well described, however, EPN-associated defenses are nowadays primarily the focus of research (Sikandar *et al.*, 2021).

3.3.2. Effects of EPNs on the Activity of Peroxidase:

From the function point of view, peroxidase (POD) protects the living cells from oxidative damage induced by xenobiotics and pathogenic infection in insects (Zhao *et al.*, 2001). Also, Lalitha *et al.* (2018) reported that POD activity was significantly low in *S. litura* larvae for 3, 6, 9 and 12 h post-infection with *H. indica*, *i.e.*, POD activity did not change in response to the EPN infection. On the other hand, the POD activity in *G. mellonella* larvae infected with *H. beicherriana* gradually increased from 0 to 24 h of treatment, followed by a sharp decrease at 40 h (Wu and Liu, 2012).

Similar result was obtained by Li *et al.* (2016) for the *T. molitor* larvae after infection with different concentrations of EPN *H. beicherriana*, since POD activity quickly increased at higher concentrations after 24 h but decreased later on. Recently, Li *et al.* (2022) found that the activity of POD in larvae of the beetle *H. parallela* was almost enhanced during the early period (24 h) after EPN *H. beicherriana*, *B. thuringiensis* (*Bt*), or EPN+*Bt* combination exposure. However, the enzyme activity was suppressed after infection with the EPN-*Bt* combination for 4 days. This rising and then decline of POD activity may be associated with the persistent infection of EPN, the rapid propagation of its symbiotic bacteria, and the increase of *Bt* intake, which destroyed the cellular structure of the mid-gut and fat body of larvae and hindered the normal physiological metabolism (Vachon *et al.*, 2012; Wu *et al.*, 2015). In contrast, no fluctuation of POD activity was detected by Abd-El Wahed and Elhadidy (2018) but permanent decrease, since infection of *L. migratoria* 5th instar nymphs with EPN *S. carpocapsae* resulted in a remarkable decrease of POD activity at 24 h post-infection, but a slight decrease of activity at 48 h post-infection.

Apart from EPNs, some authors (Müller *et al.*, 2007; Wu *et al.*, 2011) reported that the increase in the activity of POD was found in connection with the resistance of insects to pesticides. Also, exposure to chemical pesticides usually results in remarkable increase in POD activity in the cowpea seed beetle *Callosobrochus maculatus* (Coleoptera: Chrysomelidae) (Kolawole and Kolawole, 2014). In addition, Elbanna *et al.* (2012) reported

that the POD activity in *S. gregaria* nymphs was altered between rising and declining by entomopathogenic fungus *M. anisopliae* infection.

3.3.3. Effects of EPNs on the Activity of Catalase:

After exposure of an insect to pesticides, catalase (CAT) activity promotes removal of hydrogen peroxide (H_2O_2) and prevention of oxidative damage. Its activity in the tissue surrounding the primary infection site is reported to have a close relation to programmed cell death (Schenk *et al.*, 2000). The current literature contains different fluctuations of CAT activity in insects under EPN infection. For instance, Li *et al.* (2016) infected the *T. molitor* larvae with different concentrations of EPNs *H. beicherriana* and observed a quick increase of CAT activity at higher concentrations after 24 h but decreased later. This result was consistent with the result of Wu and Liu (2012), since the activity of CAT in *G. mellonella* larvae infected with *H. beicherriana* gradually increased from 0 to 24 h of treatment, followed by a sharp decrease at 40 hr. In a study of Lalitha *et al.* (2018), a high catalase activity was observed at 6 h exposure of EPN *H. indica* on *S. litura* larvae. Recently, Li *et al.* (2022) found the activity of CAT in larvae of the beetle *H. parallela* almost enhanced during the early period (24 h) of EPN *H. beicherriana*, *B. thuringiensis* (Bt), or EPN+Bt combination exposure. Then, the enzyme activity was suppressed after infection with the EPN-Bt combination for 4 days. These authors tried to explain their result by the rapid propagation of EPN-symbiotic bacteria, and the increase of Bt intake, which destroyed the cellular structure of the mid-gut and fat body of larvae and hindered the normal physiological metabolism (Vachon *et al.*, 2012; Wu *et al.*, 2015). In addition to EPNs, the CAT activity was decreased after exposure of *B. thuringiensis* alone on the *G. mellonella* larvae (Dubovskiy *et al.*, 2008).

3.3.4. Effects of EPNs on the Activity of Superoxide Dismutases:

Superoxide dismutase (SOD) is an important enzyme among the antioxidant defense enzymes of insects which are responsible for the scavenging of free radicals. It scavenges superoxide anions and detoxifies them by converting them to H_2O_2 and oxygen. The H_2O_2 is then transformed to the water and oxygen by other enzymes, such as POD, CAT and GPx (Nordberg and Arner, 2001) and the imbalance between oxidative stress and antioxidant responses leads to disease and even death of insects (Felton and Summers, 1995).

A few studies shed some light on the disturbance of SOD activity in insects under stress of the EPN infection. After infection of *T. molitor* larvae with different concentrations of *H. beicherriana*, Li *et al.* (2016) recorded a quick increase of SOD activity at higher concentrations after 24 h but the enzyme activity was decreased later on. A few years before this study, Wu and Liu (2012) reported that the SOD activity in *G. mellonella* larvae infected with *H. beicherriana* gradually increased from 0 to 24 h post-treatment, followed by a sharp decrease at 40 h. In the same trend, Li *et al.* (2022) reported that the SOD activity in larvae of *H. parallela* increased during the first 24 h post-infection with *H. beicherriana*, *B. thuringiensis* (Bt), or EPN+Bt combination. Then, the enzyme activity decreased after infection with the EPN-Bt combination for 4 days.

3.3.5. Effects of EPNs on the Activity of Tyrosinase:

Pathogen infection normally activates multiple systemic responses in insect hosts, including phagocytosis and encapsulation by hemocytes (Xia *et al.*, 2000; Hassan *et al.*, 2016), accompanied with melanization reactions (Hoffmann *et al.*, 1996; Söderhäll and Cerenius, 1998). Insects defend themselves against attacking parasites and pathogens via the melanin biosynthetic pathway as a necessary mechanism of defense reaction (Sugumaran, 2002) because they do not have antibodies (Hoffmann *et al.*, 1996). Tyrosinase (TYR) is often considered as an essential component of invertebrate's immune system. TYR has been known as phenoloxidase for three physiologically important processes (cuticle sclerotization, defense reaction, and wound healing). Through a series of reactions, the pathogen or any foreign organism is encapsulated and often melanized (Sugumaran *et al.*, 2000). Including

TYR, melanization is based on the pro-phenoloxidase (pro-PO) cascade, which is a common and generalized response to immune defense (Hoffmann *et al.*, 1996). The first step in generating catechol products is the hydroxylation of tyrosine (a monophenol) to produce 3,4-dihydroxyphenylalanine (DOPA, an o-diphenol). DOPA and its derivatives, such as dopamine, are oxidized to their corresponding quinones, which undergo subsequent reactions leading to melanization (Gorman *et al.*, 2007).

To investigate the effect of EPN infection on TYR activity in insects, Balasubramanian *et al.* (2010) conducted a study in which they found that a trypsin-like serine protease extracted from secretions of the EPN *S. carpocapsae* could inhibit the TYR activity in *G. mellonella* larvae. According to these authors, their result indicates that the changes of TYR activity in the EPN-infected larvae are context dependent. Moreover, activity of TYR was significantly enhanced in a dose-dependent course in *G. mellonella* larvae early after infection with *H. beicherriana*. On the other hand, the enzyme activity decreased 16 h post-infection (Wu *et al.*, 2013). After infection of the *T. molitor* larvae with different concentrations of *H. beicherriana*, Li *et al.* (2016) found that the TYR activity increased slowly at lower concentrations for 16 h, followed by a slight decrease, and then increasing from 32 to 40 h.

Wu *et al.* (2013) explained the increasing TYR activity in *G. mellonella* larvae infection with *H. beicherriana* by an overreactive stress response to EPNs infection which could lead to death of the insect host, while the decreased TYR activity 16 h post-infection indicated that EPN infection may consume superfluous TYR. For some details, Li *et al.* (2016) suggested that the activity of TYR first increased shortly after the EPN infection to the insect host and then considerably increased in bulk during the mass production of the EPN-symbiotic bacteria released by EPN. These reactions not only prevent the multiplication of the invading IJs of the EPN but also prevent the damage caused by it (Sugumaran, 2002). In addition, when immune defense is expressed in the infected insect at TYR high levels, it is often traded off against other fitness parameters (Moret, 2006) because immune effector systems are costly to maintain and use while beneficial against parasitic attacks (Sugumaran, 2000).

In general, the increasing activities of antioxidant enzymes in *G. mellonella* larvae after infection with EPNs suggested that EPNs' infection increases the level of oxidative stress and antioxidative responses in larvae, and the oxidative damage contributes to the cell death (Wu and Liu, 2012). Some years later, Li *et al.* (2016) suggested that the host antioxidative response and detoxification reactions played a central role in the defensive reaction to EPNs, and that this stress which was reflected by the higher-level enzymes activity contributed to the death of hosts.

3.3.6. Effects of EPNs on the Activity of Immune Response-Mediated Phenoloxidase in Insects:

In invertebrate immunology a complex array of host defenses has been reported including phagocytosis and melanization (i.e., synthesis and deposition of melanin around the pathogen) (Bulet *et al.*, 2004; Nappi and Christensen, 2005). The innate immune response in insects involves both humoral and cellular components. Humoral responses comprise the antimicrobial melanizing enzyme, phenoloxidase (PO) (Meister and Lagueux, 2003; Shelby and Popham, 2006; Zhao *et al.*, 2007). The melanization is achieved as a consequent to the activation of the inactive proPO to PO (Cerenius and Soderhall, 2004; Shelby and Popham, 2006; Cytrynska *et al.*, 2007). This enzymatic activation requires protease cascades triggered by the detection of specific microbial patterns (Cerenius *et al.*, 2008; Eliáš *et al.*, 2020). Shortly, melanization response in insects, as a humoral immune defence, involves the deposition of melanin to invading pathogens. The melanization reaction needs PO which

catalyzes the oxidation of mono- and diphenols to orthoquinones (Eleftherianos and Revenis, 2011; Lu *et al.*, 2014).

It may be important to shed some light on the enzyme PO in some detail. It is a copper containing enzyme (Lu and Jang, 2007). In insect haemolymph, PO is responsible for melanotic immune responses that occur immediately against invading pathogenic microbes into insects (Castillo *et al.*, 2011). For some detail, see the recent review of Mahanta *et al.* (2023). However, PO participates in encapsulation, clotting of haemolymph, wound healing (Cerenius *et al.*, 2008) and stimulates phagocytosis. Also, PO is involved in other physiologically important processes, such as sclerotization of the cuticle, an essential step for the survival of all insects (Sugumaran *et al.*, 2000). PO is produced in response to infection by activated hemocytes of the insects (Lavine and Strand, 2002; Kanost and Gorman, 2008; Castillo *et al.*, 2011; González-Santoyo and Córdoba-Aguilar, 2012). Therefore, PO activity is a good indicator of protective response against invading entomopathogens and parasitoids in insects (Freitake *et al.*, 2007). Furthermore, PO has main role for melanogenesis in the immune system of insects for converting the phenols to short lived chemically reactive quinones, which subsequently polymerize to form melanin coat around the invading pathogen (Cerenius and Soderhall, 2004; Eleftherianos and Revenis, 2011; Castillo *et al.*, 2011) and generating toxic radicals (Shelby and Popham, 2006).

For some details, the enzyme PO is the active form of zymogenic proPO. Although proPO can be activated in the insect plasma by proteolytic cleavage at a specific site near its amino terminus by serine proteases, the zymogen is localized generally in the oenocytoids, plasmatocytes, spherulocytes and granulocytes (Lavine and Strand, 2002; Ling *et al.*, 2005; Neuwirth, 2005). In the defense process, peptidoglycan recognition protein binds to its respective elicitors and results in the production of inactive proPO (Eleftherianos and Revenis, 2011). Activation of proPO to produce PO occurs immediately against invading microbes in the haemolymph of insects in response to the immune challenge (Marmaras *et al.*, 1996; Gillespie *et al.*, 1997), such as the melanin formation around the invading pathogens (Christensen *et al.*, 2005; Xue *et al.*, 2006) that participate in sequestering and/or killing the pathogens (Nappi and Vass, 2001).

As reported in the current literature, a few studies investigated the infectious effects of EPNs on the PO activity in insect hosts (Walter *et al.*, 2008; Kramarz *et al.*, 2016). Injection of *H. bacteriophora* into the 3rd instar larvae of the flesh fly *Parasarcophagaturcoufi* (Diptera: Sarcophagidae) significantly suppressed the haemolymph PO activity even in the presence of the activators laminarin, alpha-chymotrypsin and methanol. This suppression was dose-dependent and reached its maximum at 30 h post-injection (Ayyad *et al.*, 2001). Shairra *et al.* (2016a) infected the *P. gossypiella* larvae with the EPN *S. riobrave* or *H. bacteriophora* and found decreasing PO activity followed by increased activity, indicating the important role of PO in defeating the immune system in larvae of *P. gossypiella*. At the same year, results of Shairra *et al.* (2016b) revealed that PO activity was significantly enhanced in a dose-dependent manner in haemolymph of *A. ipsilon* larvae infected with EPN *S. carpocapsae* or *S. scapterisci*. The same authors interpreted their result by the overreactive stress response of larvae to the EPNs infection. This overreaction often leads to death of the insect host.

Ebrahimi *et al.* (2014) recorded an increase of PO activity in the haemolymph of *S. carpocapsae*-injected larvae of Colorado potato beetle *Leptinotarsa decemlineata* (Coleoptera: Chrysomelidae). They found this increase in an EPN dose-dependent manner, and the increasing EPN concentration led to increased PO activity which was coincident with the appearance of EPN-symbiotic bacteria in the haemolymph of these larvae. Abd-El Wahed and Elhadidy (2018) reported that the infection of *L. migratoria* 5th instar nymphs with EPN *S. carpocapsae* resulted in increasing PO activity, 24 & 48 h post-infection. Depending on

the results of Ebrahimi *et al.* (2018), PO activity assay revealed higher levels of the enzyme activity in EPN-injected *H. armigera* larvae, which suggested the *H. armigera* PO activation by *S. feltiae* infection. Releasing the symbiotic bacteria and increasing the bacterial cells in haemolymph of *H. armigera* during the experimental time interval were reasonable explanation for increasing trend of PO activity in nematode-injected insects. Also, similar results were obtained by Yang *et al.* (2012) in larvae of the same insect.

On the contrary, Brivio *et al.* (2002) reported that the infection of *G. mellonella* larvae with *S. feltiae* led to the inhibition of PO activity. They showed that IJs of *S. feltiae* induced the speedy suppression of PO tend to avoid host humoral encapsulation. Also, infection of the 4th larval instar larvae of *A. ipsilon* with EPNs *H. zealandica* and *S. abbasi* led to decreasing PO activity in the infected larvae (Ibrahim *et al.*, 2015). Reduction of the enzyme activity was remarkable in *H. zealandica*-infected larvae may be due to the highly toxic compounds released by the *Photorhabdus* symbiotic bacteria which induce PO inactivation in the haemolymph of *A. ipsilon* larvae (Ibrahim *et al.*, 2015). In Egypt, also, Ibrahim *et al.* (2019) investigated the effects LC₅₀ concentrations of EPN (*H. zealandica*) and entomopathogenic fungus (*B. bassiana*), separately and in a combination, on immune and antioxidant enzymes in larvae of *G. mellonella*. Depending on their results, the combination of *H. zealandica*+*B. bassiana* or *H. zealandica* alone exhibited a significant suppressive effect on the PO activity over time.

To investigate the correlation between proPO and PO, Brivio *et al.* (2002) reported that the *S. carpocapsae*-secreted chymotrypsin proteases can inhibit proPO leading to reduced PO activation and encapsulation in the *G. mellonella* larvae, while *S. feltiae* induces proPO inactivation in larvae by down-regulating the proPO pathway. Similar result was obtained by Balasubramanian *et al.* (2009), since EPN *S. abbasi* induces poPO inactivation in *A. ipsilon* larvae. Depending on the results of Lalitha *et al.* (2018), a significant increase in PO levels were observed in haemolymph of *S. litura* larvae suggesting that *H. indica* cell wall components are recognized by ProPO and results in PO activation.

Summary Points:

- * The main body metabolites (proteins, lipids and carbohydrates) in insect larvae were reported to be disrupted by the infection with EPNs. The interaction between nematode and insect larvae appeared to be primarily nutritional, since growth of the EPN proceeds while the nutritional requirements of the host larvae deteriorate, i.e., the host become in a state of physiological starvation. The protein content in haemolymph of larvae is usually reduced by the EPN infection suggesting the main cause of the host quick death is the suppression of immune response of the infected larvae.
- * Many results revealed the reduction of lipid content, regardless of the EPN species. This may be due to the shift in energy metabolism towards lipid catabolism as a result of physiological stress caused by EPNs. Also, the carbohydrate content in EPN-infected insects was reported to decrease due to the accelerated glycogenolysis and/or impaired glycogenesis.
- * Not only the main body metabolites but also the food consumption and metabolic efficiencies of insects have been disrupted by EPN infection. The considerable reduction of the food utilization efficiencies of the insect larvae after infection with EPN suggested that most of ingested or digested food was metabolized for energy production, and it is likely that the major portion of this energy was utilized for defense against the invading pathogens.
- * Some EPNs adversely affect the developmental physiology, such as the developmental durations and some adult performance parameters, as well as the major reproductive parameters, fecundity and egg fertility.
- * In respect of the major digestive enzymes in insects, activities of different carbohydrate hydrolyzing enzymes, such as amylase, trehalase and invertase, were disturbed by EPN infections.

- * Concerning protein hydrolyzing enzymes, very few studies revealed the inhibition of protease activity in some insects after infection with certain EPN species which could be due to the symbiotic bacteria start to digest the proteins of the insect body 16 hr post-infection.
- * In connection with lipid hydrolyzing enzymes, increasing lipase activity was observed in insects after EPN infection. This reported result can be understood in view of the activity of EPN- symbiotic bacteria which secrete various enzymes, such as hemolysis, lipases, and proteases, that contribute to pathogenicity or nutrient acquisition for the bacterium and its nematode host, lipase activity was dramatically boosted.
- * The detoxification enzymes in insects have many functions for repairing the physiological processes, detoxifying pathogenic products, and metabolizing the biologically active compounds. In insects, also, the detoxification can be achieved by different families of enzymes. Some authors reported superoxide dismutase, peroxidases and catalases as antioxidant enzymes and carboxylesterase, glutathione S-transferase and acetylcholinesterase as detoxifying enzymes, since insects are known to use antioxidants and detoxifying enzymes in their immune systems as defense reactions against pathogen threats.
- * With regard to disturbed activities of the detoxifying enzymes, phosphatases, in insects by the EPN infection, increasing activity of acid phosphatase (ACP) in the EPN-infected larvae may indicate a physiological adaptability to combat the oxidative stress or may be due to the phagocytosis of certain hemocytes to the invading EPNs. In contrast, decrease of ACP activity can be due to the inability of the cell to undergo enzymatically controlled reactions under nematode infection, or may be due to the reduced phosphorus liberation for energy metabolism and decreased rate of metabolism, as well as decreased rate of transport of metabolites. In addition, there are many reported results of increasing alkaline phosphatase (ALP) activity in several insects as response to the infection with certain EPNs. On the contrary, ALP activity in larvae of some insects was reduced after infection with certain EPNs.
- * Dealing with the disturbed activities of transaminases in insect larvae by EPNs, decreasing activities of these enzymes may be attributed to the significant decline in the free amino acids content, the quantum of which directly influences the activity of transaminases at the time of protein synthesis. Also, reduction of transaminase activity may be due to the interference of the EPNs with the hormonal regulation of protein synthesis and neurosecretory hormones involved in the regulation of transaminase levels.
- * Few studies investigated the effects of EPNs on the activity of acetylcholinesterase (AChE) in insect larvae. Many studies recorded the increasing enzyme activity, but few studies recorded decreasing activity of this enzyme. Increased AChE activity in insects may be the overreactive stress response to the EPN infection. This overreaction often leads to death of the insect. Also, the increase in AChE activity may be correlated with secretions produced by the nematode-bacterium complex.
- * In addition to the role of Carboxylesterase (CarE) in insecticide resistance, some studies reported the enhancement of its activity early after infection with EPNs and it was inhibited later on. It seems that the insects overreacted to the EPNs infection by enhancing the CarE activity first, indicating that many lipids would be disintegrated, which would damage organs and tissues, and lead to the death of host insects. Then, the EPNs-infection response would consume the CarE resulting in the decreased CarE activity.
- * Esterases (ESTs) constitute a large class of enzymes. Most enzymes of this group are important in metabolism because they can break ester bond through hydrolysis. Also, some ESTs play a critical role in the detoxification of synthetic chemicals. Only few studies focused on the impacts of EPNs on the activities of ESTs in insects.
- * Glutathione S-transferase (GST) is involved in the detoxification of both endogenous and exogenous xenobiotics, natural and synthetic compounds in insects. In addition, GST

participates in the cellular antioxidant defenses against oxidative stress. GST activity increased or decreased in the insect haemolymph after infection with certain EPN species depending on some factors. The enhanced activity of GST at early time interval of EPN infection may be the overreactive stress response to the EPNs infection while the decreased activity 16 h post-infection indicates that EPNs infection may consume GST through detoxification reactions of glutathione-dependent enzymes.

* Some research works have been conducted aiming at the understanding of physiological and biochemical mechanisms involved in the insect infection. Insects have several defense reactions against invading pathogens. The innate immunity-defense strategies can be classified into two major classes: cellular and humoral. Humoral innate immune responses in insects involve many antioxidant enzymes and antibacterial proteins produced in the fat body and hemocytes of insects. Some of the major antioxidant defense enzymes are peroxidases, Superoxide dismutases, glutathione peroxidase, catalases, reduced glutathione and glutathione reductase.

* Peroxidase (POD) protects cells from oxidative damage induced by xenobiotics and pathogenic infection in insects. Some studies reported rising (or enhancement) during the early period (24 hr) and then declination (or inhibition) of POD activity in insects after infection with EPNs. Other studies reported continuous decrease of the enzyme level.

* The activity of Catalase (CAT) in the tissue surrounding the primary infection site is reported to have a close relation to programmed cell death. Some studies reported increasing CAT activity (enhanced) in insects at early time interval (6-24 hr) post-infection with EPNs, but decreased (inhibited) activity some days later.

* Superoxide dismutase (SOD) is an important enzyme among the antioxidant defense enzyme EPNs butts which are engaged in the scavenging of free radicals. A number of studies shed some light on the altered activity of SOD in insects under stress of EPN infection.

* Tyrosinase (TYR) is often considered as an essential component of invertebrate's immune system. TYR has been known as phenoloxidase for three physiologically important processes (sclerotization, defense reaction, and wound healing). The early increasing TYR activity in the EPN-infected larvae may be due to an overreactive stress response to EPNs ending in the insect death, while decreased TYR activity 16 hr post-infection indicate that EPNs infection may consume superfluous TYR.

* Among the necessary components of the humoral innate immunity in insects is Phenoloxidase (PO) via catalyzing the biosynthesis of quinones and other reactive intermediates to eliminate the invading pathogens and parasites. Also, PO is involved in other physiologically important processes, such as sclerotization of the cuticle, an essential step for the survival of all insects. Therefore, PO activity is a measure of protective response against invading microbes in insects.

Conclusions and Prospectives:

As clearly shown in the present review, different EPNs exert various suppressive actions on the innate immune defenses of insects, while the infected insects try to overcome these stresses by several humoral immune defenses, reflecting on the disruption of different physiological and biochemical processes, including the intermediary metabolism, detoxifying enzymes and antioxidant enzymes. The disruptive effects of the EPNs on insects have suggested the potential of these entomopathogens for pest control. However, some points of research need more investigation in future, such as the link between humoral immunity defenses and cellular immunity defenses to understand the defense integration in insects against EPNs.

Declarations

Ethical Approval: Not applicable.

Competing Interests: The authors declare that they have no competing interests.

Authors' Contributions: All authors prepared and wrote this article equally. They read and approved the final manuscript.

Funding: No funding was received.

Availability of Data and Materials: All datasets analyzed and described during the present study are available.

Acknowledgment: The authors would like to express their appreciation and sincere thanks to Dr. El-Sayed H. Shaurub, Professor of Entomology, Faculty of Science, Cairo University, Egypt, and Dr. James L. Nation, Professor Emeritus, University of Florida, Gainesville, USA, for the critical revision of draft of the present review.

REFERENCES

- Abd-El Wahed, S.M.N. and Elhadidy, N.M. (2018): Immunity Changes in *Locusta migratoria* Linnaeus (Orthoptera: Acrididae) Infected by Entomopathogenic Nematode *Steinernema carpocapsae* (Rhabditida: Steinernematidae). *Journal of Plant Protection and Pathology (Egypt)*, 9(12): 877-881. DOI:10.21608/JPPP.2018.44100
- Abd-Elgawad, M.M.M. (2020): Can rational sampling maximise isolation and fix distribution measure of entomopathogenic nematodes? *Nematology*, 22(8): 907–916. <https://doi.org/10.1163/15685411-00003350>
- Abdel-Razek, A.; Kamel, K.E. and Salama H.S. (2004): Biochemical effects of the nematode-bacteria complex on the red palm weevil, *Rhynchophorus ferrugineus* (Olivier) (Coleoptera: Curculionidae). *Archives of Phytopathology and Plant Protection*, 37: 205–214. <https://doi.org/10.1080/0323540042000218754>
- Abu Hatab, M.; Selvan, S. and Gaugler, R. (1995): Role of proteases in penetration of insect gut by the entomopathogenic nematode, *Steinernema glaseri* (Nematoda: Steinernematidae). *Journal of Invertebrate Pathology*, 66: 125-130.
- Abulyazid, I.; Mahmoud, S.M.; Elshafei, A.M. and Taha, R.H. (2005): Physiological changes of irradiated and diseased mulberry silkworm. *Bombyx mori*. *Egyptian Journal of Agricultural Research*, 83(4): 1431-1445.
- Adams, B.J. and Nguyen, K.B. (2002): Taxonomy and systematic. In: "Entomopathogenic Nematology"(Gauglar, R., ed.). CABI, New York, NY, pp: 1-34.
- Ahmed, N.F.; Maklad, A.M.H.; Yassin, S.A. and Abolmaaty, S.M. (2014): Biochemical effects of *Steinernema feltiae*, *Steinernema riobrave* and *Heterorhabditis bacteriophora* on *Spodoptera littoralis* larvae. *Egyptian Academic Journal of Biological Sciences (C Physiology and Molecular Biology)*, 6(1):23–34. <https://doi.org/10.21608/eajbsc.2014.16044>
- Ali, M.; Allouf, N. and Ahmad, M. (2022): First report of entomopathogenic nematode *Steinernema affine* (Nematoda: Steinernematidae) in Syria and its virulence against *Galleria mellonella* L. (Lepidoptera: Pyralidae). *Egyptian Journal of Biological Pest Control*, 32: 101. <https://doi.org/10.1186/s41938-022-00602-x>
- Amizadeh, M.; Hejazi, M.J.; Niknam, G. and Askari-Saryazdi, G. (2019): Interaction between the entomopathogenic nematode, *Steinernema feltiae* and selected chemical insecticides for management of the tomato leaf miner, *Tuta absoluta*. *Biocontrol*, 64: 709–721. Doi: 10.1007/s10526-019-09973-x
- Amutha, V.; Vengateswari, G. and Shivakumar, M.S. (2021): Entomopathogenicity of nematode *Panagrolaimus* spp. (Rhabditida: Panagrolaimidae) against lepidopteran pest *Spodoptera litura*. *International Journal of Pest Management*, 67(4): 320-327. DOI: 10.1080/09670874.2020.1776415

- Ansari, M.A. and Hussain, M.A. (2020): Indiscriminate feeding of commercial entomopathogenic nematodes caused death of *Aedes aegypti* larvae: potential for vector control of chikungunya, dengue and yellow fever. *Biocontrol Science and Technology*, 30(8): 840-854. <https://doi.org/10.1080/09583157.2020.1776215>
- Arthurs, S.; Heinz, K.M. and Prasifka, J.R. (2004): An analysis of using entomopathogenic nematodes against above-ground pests. *Bulletin of Entomological Research*, 94(4): 297-306.
- Askary, T.H. and Abd-Elgawad, M.M.M. (2021): Opportunities and challenges of entomopathogenic nematodes as biocontrol agents in their tripartite interactions. *Egyptian Journal of Biological Pest Control*, 31:42, 10pp. <https://doi.org/10.1186/s41938-021-00391-9>
- Askitosari, Th.D.; Pantjajani, T.; Nathania, S.; Wahyudi, A.F. and Sugianto, N.Ch. (2021): Identification of entomopathogenic nematode-associated bacteria originating from Mojokerto. *Indonesian Journal of Biotechnology and Biodiversity*, 5(1): 31-38. Doi: <https://doi.org/10.47007/ijobb.v5i1.76>
- Ayyad, T.H.; Dorrah, M.A.; Shaurub, E.H. and El-Saadawy, H.A. (2001): Effect of the entomopathogenic nematode, *Heterorhabditisbacteriophora* HP88 and azadirachtin on the immune defense response and prophenoloxidase of *Parasarcophagasurcoufi* larvae (Diptera: Sarcophagidae). *Journal of the Egyptian Society of Parasitology*, 31:295–325.
- Azmi, M.A.; Naqvi, S.N.H.; Khan, M.F.; Akhtar, K. and Khan, F.Y. (1998): Comparative toxicological studies of RBa (Neem extract) and Coopex (permethrin+bioallethrin) against *Sitophilus oryzae* with reference to their effects on oxygen consumption and GOT, GPT Activity. *Journal of Zoology*, 22: 307–310.
- Bailey, C.H. and Gordon, R. (1973): Histopathology of *Aedes aegypti* (Diptera: Culicidae) larvae parasitized by *Reesimermisnielsenii* (Nematoda: Mermithidae). *Journal of Invertebrate Pathology*, 22(3): 435–441. [https://doi.org/10.1016/0022-2011\(73\)90174-2](https://doi.org/10.1016/0022-2011(73)90174-2)
- Balamani, E. and Nair, V.S.K. (1992): Inhibitory effects of a juvenile hormone analogue on prothoracic gland activity in the penultimate and last instar larval development of *Spodoptera mauritia*Boisd. (Lepidoptera: Noctuidae). *Zoologischer Anzeiger*, 228: 182.
- Balasubramanian, N.; Hao, Y.J.; Toubarro, D.; Nascimento, G. and Simões, N. (2009): Purification, biochemical and molecular analysis of a chymotrypsin protease with prophenoloxidase suppression activity from the entomopathogenic nematode *Steinernema carpocapsae*. *International Journal of Parasitology*, 39: 975-984.
- Balasubramanian, N.; Toubarro, D. and Simões, N. (2010): Biochemical study and *in vitro* insect immune suppression by a trypsin-like secreted protease from the nematode *Steinernema carpocapsae*. *Parasite Immunology*, 32:165-175.
- Barnby, M.A. and Klocke, J.A. (1987): Effects of azadirachtin on the nutrition and development of the tobacco budworm *Heliothis virescens* (Fabr.) (Noctuidae). *Journal of Insect Physiology*, 33: 69–75. [https://doi.org/10.1016/0022-1910\(87\)90076-X](https://doi.org/10.1016/0022-1910(87)90076-X)
- Baudoin, M. (1975): Host castration as a parasitic strategy. *Evolution*, 29(2): 335–352. <https://doi.org/10.1111/j.1558-5646.1975.tb00213.x>
- Beck B.E.; Brusselman D.; Nuyttens M.; Moens S.; Pollet F. and Spanoghe P. (2013): Improving foliar applications of entomopathogenic nematodes by selecting adjuvants and spray nozzles. *Biocontrol Science and Technology*, 23(5): 507-520. <https://doi.org/10.1080/09583157.2013.777692>

- Berger, J. and Jurčová, M. (2012): Phagocytosis of insect haemocytes as a new alternative model. *Journal of Applied Biomedicine*, 10: 35-40. <https://doi.org/10.2478/v10136-012-0003-1>
- Bernstein, C. and M. Jervis, (2008): Food searching in parasitoids: the dilemma of choosing between 'intermediate' or future fitness gains. "Behavioural Ecology of Parasitoids". Blackwell, U.K. pp: 129-171.
- Bhat, A.H.; Chaubey, A.K. and Askary, T.H. (2020): Global distribution of entomopathogenic nematodes, *Steinernema* and *Heterorhabditis*. *Egyptian Journal of Biological Pest Control*, 30:31. <https://doi.org/10.1186/s41938-020-0212-y>
- Binda-Rossetti, S.; Mastore, M.; Protasoni, M. and Brivio, M.F. (2016): Effects of an entomopathogen nematode on the immune response of the insect pest red palm weevil: Focus on the host antimicrobial response. *Journal of Invertebrate Pathology*, 133: 110-119. Doi: 10.1016/j.jip.2015.11.001.
- Bonelli, M.; Bruno, D.; Brilli, M.; Gianfranceschi, N.; Tian, L.; Tettamanti, G.; Caccia, S. and Casartelli, M. (2020): Black soldier fly larvae adapt to different food substrates through morphological and functional responses of the midgut. *International Journal of Molecular Science*, 21(14):4955. <https://doi.org/10.3390/ijms21144955>
- Boz, A. and Gülel, A. (2012): The effects of temperature and time after parasitization on total amount of protein, lipid and carbohydrate in hemolymph of host larvae, *Ephestiakuethniella* Zeller (Lepidoptera: Pyralidae). *Turkish Entomology Journal*, 36(2): 239-247.
- Brivio, M.F.; Pagani, M. and Restelli, S. (2002): Immune suppression of *Galleria mellonella* (Insecta, Lepidoptera) humoral defences induced by *Steinernema feltiae* (Nematoda Rhabditida): involvement of the parasite cuticle. *Experimental Parasitology*, 101:149–156. Doi:10.1016/s0014-4894(02)00111-x.
- Brown, S.E.; Cao, A.T.; Dobson, P.; Hines, E.R.; Akhurst, R.J. and East, P.D. (2006): Txp40, a ubiquitous insecticidal toxin protein from *Xenorhabdus* and *Photorhabdus* bacteria. *Applied and Environmental Microbiology*, 72:1653-1662. Doi:10.1128/AEM.72.2.1653-1662.2006.
- Brusselman, E.; Beck, B.; Pollet, S.; Temmerman, F.; Spanoghe, P.; Moens, M. and Nuyttens, D. (2012): Effect of the spray application technique on the deposition of entomopathogenic nematodes in vegetables. *Pest Management Science*, 68: 444-453. Doi:10.1002/ps.2290.
- Bulet, P.; Stocklin, R. and Menin, L. (2004): Anti-microbial peptides: from invertebrates to vertebrates. *Immunological Reviews*, 198: 169–184. Doi: 10.1111/j.0105-2896.2004.0124.x.
- Çağlayan, A.; Ata0124. x.nd Kepenekci, İ. (2021): Efficacy of some native Entomopathogenic nematodes against the alfalfa weevil, *Hypera postica* (Gyllenhal) (Coleoptera: Curculionidae), and the lucerne beetle, *Goniocetena fornicata* (Brüggemann) (Coleoptera: Chrysomelidae), adults under laboratory conditions. *Egyptian Journal of Biological Pest Control*, 31:89, 7pp. <https://doi.org/10.1186/s41938-021-00436-z>
- Canavoso, L.E.; Jouni, Z.E.; Karnas, K.J.; Pennington, J.E. and Wells, M.A. (2001): Fat metabolism in insects. *Annual Review of Nutrition*, 21: 23-46. Doi: 10.1146/annurev.nutr.21.1.23
- Capinera, J.L. (2001): Handbook of vegetable pests. *Academic Press*, New York, 729 pp.
- Cardoso, D.O.; Gomes, V.M.; Dolinski, C. and Souza, R.M. (2015): Potential of entomopathogenic nematodes as biocontrol agents of immature stages of *Aedes aegypti*. *Nematoda*, 2(1): e092015. <http://dx.doi.org/10.4322/nematoda.09015>

- Carlier, P.R.; Andersonb, T.D.; Wonga, D.M.; Hsua, J.H.; Maa, M.; Wongc, E.A.; Choudhuryc, R.; Lamd, P.C.H.; Totrovd, M.M. and Bloomquist, J.R. (2008): Towards a species-selective acetylcholinesterase inhibitor to control the mosquito vector of malaria, *Anopheles gambiae*. *Chemico-Biological Interactions*, 175(1-3):368-375. Doi: 10.1016/j.cbi.2008.04.037
- Castillo, J.C.; Reynolds, S.E. and Eleftherianos, I. (2011): Insect immune responses to nematode parasites. *Trends in Parasitology*, 27(12): 537–547. [https://doi.org/ 10.1016/j.pt.2011.09.001](https://doi.org/10.1016/j.pt.2011.09.001)
- Cerenius L and Soderhll K (2004): The prophenoloxidase-activating system in invertebrates. *Immunological Reviews*, 198:116–126. Doi:10.1111/j.0105-2896.2004.00116.x
- Cerenius, L.; Lee, B.L. and Soderhall, K (2008): ProPOsystem: pros and cons for its role in invertebrate Immunity. *Trends in Immunology*, 29: 263–271. Doi: 10.1016/j.it.2008.02.009.
- Chapman R.F. (1985): Structure of digestive system. In "Comprehensive insect physiology, biochemistry and pharmacology" (Kerkut, G.A. and L.I. Gilbert, eds.). *Pergamon Press*, Oxford), 4: 165.
- Chapman, R.F. (1998): The insects: structure and function. 4th ed. *Cambridge: Cambridge University Press*, United Kingdom, pp:116-118.
- Chaston, J.M.; Murfin, K.E.; Heath-Heckman, E.A. and Goodrich-Blair, H. (2013): Previously unrecognized stages of species-specific colonization in the mutualism between *Xenorhabdus* bacteria and *Steinernema* nematodes. *Cell Microbiology*, 15: 1545–1559. Doi: 10.1111/cmi.12134.
- Chen, G.; Zhang, Y.; Li, J.; Dunphy, G.B.; Punja, Z.K. and Webster, J.M. (1996): Chitinase activity of *Xenorhabdus* and *Photorhabdus* species, bacterial associates of entomopathogenic nematodes. *Journal of Invertebrate Pathology*, 68(2):101–108. <https://doi.org/10.1006/JIPA.1996.0066>
- Chen, B.; Kayukawa, T.; Jiang, H.; Monteiro, A.; Hoshizaki, S. and Ishikawa, Y. (2005): Da trypsin, a novel clip-domain serine protease gene up-regulated during winter and summer diapauses of the onion maggot, *Delia antiqua*. *Gene*, 347: 115–123. Doi: 10.1016/j.gene.2004.12.026.
- Chippendale, G.M. (1978): The functions of carbohydrates in insect life processes. In: "Biochemistry of Insects"(Rockstein, M., ed.). Academic Press, New York, pp: 2-54.
- Chowanski, S.; Lubawy, J.; Spochacz, M.; Ewelina, P.; Grzegorz, S.; Rosinski, G. and Slocinska, M. (2015): Cold induced change in lipid, protein and carbohydrate levels in the tropical insect *Gromphadorhinacoquereliana*. *Comparative Biochemistry and Physiology*, Part A, 183: 57-63. <https://doi.10.1016/j.cbpa.2015.01.007>
- Christensen, B.M.; Li, J.; Chen, C.-C. and Nappi, A.J. (2005): Melanization immune responses in mosquito vectors. *Trends in Parasitology*, 21: 192-199. Doi: 10.1016/j.pt.2005.02.007.
- Clark, A.G. (1990): The glutathione S-transferases and resistance to insecticides. In: "Glutathione S-transferases and drug resistance" (Hayes, J.D.; Pickett, C.B. and Mantle, T.J., eds). Taylor and Francis, London, pp: 369–378.
- Cytrynska, M.; Mak, P.; Zdybicka-Barabas, A.; Suder, P. and Jakubowicz, T. (2007): Purification and characterization of eight peptides from *Galleria mellonella* immune hemolymph. *Peptides*, 28: 533-546. Doi: 10.1016/j.peptides.2006.11.010.
- Dapporto, L.; Lambardi, D. and Turillazzi, S. (2008): Not only cuticular lipids: first evidence of differences between foundresses and their daughters in polar substances in the paper wasp *Polistes dominulus*. *Journal of Insect Physiology*, 54: 89-95. <https://doi.org/10.1016/j.jinsphys.2007.08.005>

- Darpel, K.E.; Langner, K.F.A.; Nimtz, M.; Anthony, S.J.; Brownlie, J.; Takamatsu, H.-H.; Mellor, Ph.S. and Mertens, P.P.C. (2011): Saliva proteins of vector culicoides modify structure and infectivity of bluetongue virus particles. *PLoS ONE*, 6: e17545. Doi: 10.1371/journal.pone.0017545
- de Carvalho, R.A.; Torres, T.T. and de Azeredo-Espin, A.M. (2006): A survey of mutations in the *Cochliomyia hominivorax* (Diptera: Calliphoridae) esterase E3 gene associated with organophosphate resistance and the molecular identification of mutant alleles. *Veterinary Parasitology*, 140(3-4): 344-351. Doi: 10.1016/j.vetpar. 2006.04.010.
- Demok, S.; Endersby-Harshman, N.; Vinit, R.; Timinao, L.; Robinson, L.J.; Susapu, M.; Makita, L.; Laman, M.; Hoffmann, A. and Karl, S. (2019) Insecticide resistance status of *Aedes aegypti* and *Aedes albopictus* mosquitoes in Papua New Guinea. *Parasite Vectors*, 12(1): 333. Doi:10.1186/s13071-019-3585-6
- Derbalah, A.S.; Khidr, A.A.; Moustafa, H.Z. and Taman, A. (2014): Laboratory evaluation of some non-conventional pest control agents against the pink bollworm *Pectinophora gossypiella* (Saunders). *Egyptian Journal of Biological Pest Control*, 24(2): 363-368.
- Dillman, A.R.; Chaston, J.M.; Adams, B.J.; Ciche, T.A.; Goodrich-Blair, H.; Stock, S.P. and Sternberg, P.W. (2012): An entomopathogenic nematode by any other name. *PLoS Pathogens*, 8(3): e1002527. Doi: 10.1371/journal.ppat.1002527
- Dmitryjuk, M.; Lopieńska, E. and Zołtowska, K. (2001): The concentration of trehalose and activity of trehalase from *Galleria mellonella* larvae infected by *Steinernema affinis*, Bovien 1937 (Nematoda: Rhabditida: Steinernematidae). *Wiadomości parazytologiczne*, 47(3): 305-309.
- Dorrah, M.A.; Mohamed, A.A. and Shaurub, E.H. (2019): Immunosuppressive effects of the limonoid azadirachtin, insights on a nongenotoxic stress botanical, in flesh flies. *Pesticide Biochemistry and Physiology*, 153: 55–66. Doi: 10.1016/j.pestbp. 2018. 11.004.
- Dowds, B.C.A. and Peters, A. (2002): Virulence mechanisms. In: "Entomopathogenic Nematology"(Gaugler, R., ed). CABI International, Wallingford, pp: 79–98.
- Du Preez, F.; Malan, A.P. and Addison, P. (2021): Potential of *in vivo*-and *in vitro* cultured entomopathogenic nematodes to infect *Lobesia vanillana* (Lepidoptera: Tortricidae) under laboratory conditions. *PLoS ONE*, 16(8): e0242645.https:// doi.org/10. 1371/journal.pone.0242645
- Du Preez, G.; Daneel, M.; De Goede, R.; Du Toit, M.J.; Ferris, H.; Fourie, H.; Geisen, S.; Kakouli-Duarte, T.; Korthals, G.; Sanchez-Moreno, S. and Schmidt, J.H. (2021): Data: nematode based indices in soil ecology - application, utility and future directions. *Open AgrarRespository*. https://doi.org/10.5073/20211217-170559.
- Dubovskiy, I.M.; Martemyanow, V.V.; Vorontsova, Y.L.; Rantala, M.J.; Gryzanova, E.V. and Glupov, V.V. (2008): Effect of bacterial infection on antioxidant activity and lipid peroxidation in the midgut of *Galleria mellonella* L. larvae (Lepidoptera, Pyralidae). *Comparative Biochemistry and Physiology*, C, 148:1–5. Doi: 10.1016/j.cbpc.2008.02.003
- Dubovskiy, I.M.; Slyamova, N.D.; Kryukov, V.Y.; Yaroslavl'tseva, O.N.; Levchenko, M.V.; Belgibaeva, A.B.; Adilkhankyzy, A. and Glupov, V.V. (2012): The activity of nonspecific esterases and glutathione-S- transferase in *Locusta migratoria* larvae infected with the fungus *Metarhizium anisopliae* (Ascomycota, Hypocreales). *Entomological Review*, 92: 27-31. https:// doi.org/10.1134/s0013873812010022
- Ebrahimi, L.; Niknam, G.; Dunphy, G.B., and Toorchi, M. (2014): Side effects of immune response of Colorado potato beetle, *Leptinotarsa decemlineata* against the

- entomopathogenic nematode, *Steinernema carpocapsae* infection. *Invertebrate Survival Journal*, 11: 132-142.
- Ebrahimi, L.; Shiri, M. and Dunphy, G.B. (2018): Effect of entomopathogenic nematode, *Steinernema feltiae*, on survival and plasma phenoloxidase activity of *Helicoverpa armigera* (Hb) (Lepidoptera: Noctuidae) in laboratory conditions. *Egyptian Journal of Biological Pest Control*, 28:12, 4pp. Doi: 10.1186/s41938-017-0016-x
- El Aalaoui, M.; Mokrini, F.; Dababat, A.A.; Lahlali, R. and Sbaghi, M. (2022): Moroccan entomopathogenic nematodes as potential biocontrol agents against *Dactylopius opuntiae* (Hemiptera: Dactylopiidae). *Scientific Reports*, 12: 1–17. <https://doi.org/10.1038/s41598-022-11709-4>
- Elbanna, Sh.M.; Elhadidy, N.M.; Semida, F.M. and Abdel-Rasool, T. (2012): Physiological and biochemical effect of entomopathogenic fungus *Metarhizium anisopliae* on the 5th instar of *Schistocerca gregaria* (Orthoptera: Acrididae). *Journal of Scientific Research in Environmental Science and Toxicology*, 1(1): 007-018.
- Elbrense, H.; Shamseldean, M.; Meshrif, W.S.; Seif, A. (2022): The parasitic impact of *Romanomermis yengari* (Nematoda: Mermithidae) on the survival and biology of *Culex pipiens* (Diptera: Culicidae). *African Entomology*, 30, 6pp <http://dx.doi.org/10.17159/2254-8854/2022/a11687>
- El-Bishry, M.H. (1989): Studies the utilization of the entomogenous nematode in controlling some pest in Egypt. Ph.D. Thesis, Faculty of Agriculture, Cairo University, Egypt, 81pp.
- Eleftherianos, I. and Revenis, C. (2011): Role and importance of phenoloxidase in insect hemostasis. *Journal of Innate Immunity*, 3: 28–33. Doi: 10.1159/000321931
- Eleftherianos, I.; Millichap, P.J. and Reynolds, S.E. (2006): RNAi suppression of recognition protein mediated immune responses in the tobacco hornworm *Manduca sexta* causes increased susceptibility to the insect pathogen *Photobacterium*. *Developmental and Comparative Immunology*, 30: 1099-1107. Doi: 10.1016/j.dci.2006.02.008.
- Elhadidy, N.M.; Badr, F.A. and Azzazy, A.M. (2021): Potential and bio-Chemical effects of *Steinernema carpocapsae* (Rhabditida: Steinernematidae), an entomopathogenic nematode, against *Bactrocera zonata* and *Ceratitidis capitata* (Diptera: Tephritidae). *Journal of Plant Protection and Pathology*, Mansoura University (Egypt), 12 (10): 725-732.
- Eliáš, S.; Hurychová, J.; Toubarro, D.; Frias, J.; Kunc, M.; Dobeš, P.; Simões, N. and Hyršl, P. (2020): Bioactive excreted/secreted products of entomopathogenic nematode *Heterorhabditis bacteriophora* inhibit the phenoloxidase activity during the infection. *Insects*, 11, 353. Doi: 10.3390/insects11060353
- Enan, E.E. and Berberian, I.G. (1986): Interaction of pesticide exposure level with some biochemical enzymes among field workers. *Journal of the Egyptian Society of Parasitology*, 3: 76-90.
- Enayati, A.A.; Ranson, H. and Hemingway, J. (2005): Insect glutathione transferases and insecticide resistance. *Insect Molecular Biology*, 14: 3-8. <https://doi.org/10.1111/j.1365-2583.2004.00529.x>
- Erdem, M. and Büyükgüzel, E. (2015): The effects of xanthotoxin on the biology and biochemistry of *Galleria mellonella* L. (Lepidoptera: Pyralidae). *Archives of Insect Biochemistry and Physiology*, 89: 193-203. Doi:10.1002/arch.21236
- Etebari, K.; Mirhoseini, S.Z. and Matindoost, L. (2005): A study on interspecific biodiversity of eight groups of silkworms (*Bombyx mori*) by biochemical markers. *Insect Science*, 12(2): 87–94. Doi: 10.1111/j.1744-7917.2005.00010.x

- Etebari, K.; Bizhannia, A.; Sorati, R. and Matindoost, L. (2007): Biochemical changes in haemolymph of silkworm larvae due to pyriproxyfen residue. *Pesticide Biochemistry and Physiology*, 88: 14–19. <https://doi.org/10.1016/j.pestbp.2006.08.005>
- Fagan, W.F.; Siemann, E.; Mitter, C.; Denno, R.F.; Huberty, A.F.; Woods, H.A. and Elser, J.J. (2002): Nitrogen in insects: implications for trophic complexity and species diversification. *American Naturalist*, 160: 784–802. Doi: 10.1086/343879.
- Fathy, Z. and Abd El-Rahman, R.M. (2023): Effect of entomopathogenic nematodes *Steinernema* species (steinernematidae: rhabditida) and *Heterorhabditis bacteriophora* (heterorhabditidae: rhabditida) on the digestive enzymes and midgut histology of the African migratory locust *Locusta migratoria migratorioides* (Acrididae: Orthoptera). *International Journal of Tropical Insect Science*, <https://doi.org/10.1007/s42690-023-00979-8>
- Feig, D.I.; Sowers, L.C. and Loeb, L.A. (1994): Reverse chemical mutagenesis: Identification of the mutagenic lesions resulting from reactive oxygen species-mediated damage to DNA. *Proceedings of the National Academy of Sciences (PNAS)*, USA. 91: 6609–6613.
- Felton, G.W. and Summers, C.B. (1995) Antioxidant systems in insects. *Archives of Insect Biochemistry*, 29:187–197. Doi:10.1002/arch. 940290208
- Ferreira, T. and Malan, A.P. (2014): *Xenorhabdus* and *Photorhabdus*, bacterial symbionts of the entomopathogenic nematodes *Steinernema* and *Heterorhabditis* and their *in vitro* liquid mass culture: a review. *African Entomology*, 22: 1-14. <https://hdl.handle.net/10520/EJC150979>
- Ffrench-Constant, R.H.; Dowling, A. and Waterfield, N.R. (2007): Insecticidal toxins from *Photorhabdus* bacteria and their potential use in agriculture. *Toxicon*, 49: 436-451. Doi: 10.1016/j.toxicon. 2006.11. 019.
- Filgueiras, C.C. and Willett, D.S. (2021): Non-lethal effects of entomopathogenic nematode infection. *Scientific Reports*, 11: 17090. <https://doi.org/10.1038/s41598-021-96270-2>
- Fournier, D. (2005): Mutations of acetylcholinesterase which confer insecticide resistance in insect populations. *Chemico-Biological Interactions*, 157: 257-261. Doi: 10.1016/j.cbi.2005.10.040.
- Francis, F.; Vanhaelen, N. and Haubruge, E. (2005): Glutathione S-transferases in the adaptation to plant secondary metabolites in the *Myzus persicae* aphid. *Archives of Insect Biochemistry and Physiology*, 58: 166–174. Doi: 10.1002/arch.20049
- Franco, O.L.; Rigden, D.J.; Melo, F.R.; Bloch, Jr C.; Silva, C.P. and Grossi-de-Sa, M.F. (2000): Activity of wheat α -amylase inhibitors towards bruchid α -amylases and structural explanation of observed specificities. *European Journal of Biochemistry*, 267: 2166–2173. Doi: 10.1046/j.1432-1327.2000.01199. x.
- Freitak, D.; Wheat, C.W.; Heckel, D.G. and Vogel, H. (2007): Immune system responses and fitness costs associated with consumption of bacteria in larvae of *Trichoplusia*. *BMC Biology*, 5:1–13. <https://doi.org/10.1186/1741-7007-5-56>
- Friedlander, M.; Jeshtadi, A.; Stuart, E. and Reynolds, S.E. (2001): The structural mechanism of trypsin-induced intrinsic mobility in *Manduca sexta* spermatozoa *in vitro*. *Journal of Insect Physiology*, 47: 245–255. Doi: 10.1016/s0022-1910(00)00109-8.
- Gaber, M.A.M.; Shamseldean, M.S.M.; Ibrahim, N.M. and Rabia, M.M. (2018): Impact and biochemical changes of Egyptian and imported entomopathogenic nematodes on the desert locust *Schistocerca gregaria* (Forsk., 1775) (Orthoptera: Acrididae). *Middle East Journal of Agriculture Research*, 7(4): 1528-1544.
- Gade, G. (2004): Regulation of intermediary metabolism and water balance of insects by neuropeptides. *Annual Review of Entomology*, 49: 93-113. Doi: 10.1146/ annurev. ento.49.061802.123354.

- Ghally, S.E.; Serag El-Din, O.S. and Amin, M.A. (1988): Effects of the parasitic nematodes on total proteins and total lipids of *Ceratitis capitata* Wied. (Diptera, Tephritidae). *Journal of the Egyptian Society of Parasitology*, 18(2): 619–627.
- Ghoneim, K. (2024): Suppressive strategies of entomopathogenic nematodes and their symbiotic bacteria against the hemocyte-mediated immune defenses of insects: a brief review. *African Research Journal of Biosciences*, 1(2): 41-61. <https://doi.org/10.62587/AFRJBS.1.2.2024.41-61>
- Ghoneim, K. and Bakr, R.F.A. (2024): Entomopathogenic Nematodes and their symbiotic bacteria as bioagents to combat the mosquito vectors of human diseases in the world: A comprehensive review. *Egyptian Academic Journal of Biological Sciences (E. Medical Entomology & Parasitology)*, 16(1): 41–126. Doi: 10.21608/EAJBSE.2024.340637
- Ghoneim, K. and Hamadah, Kh. (2024): Compatibility of Entomopathogenic nematodes with agrochemicals and biocontrol potential of their combinations against insect pests: an updated review. *Egyptian Academic Journal of Biological Sciences (A. Entomology)*, 17(2):107-171. Doi:10.21608/EAJBSA.2024.365899
- Ghoneim, K. and Hassan, H.A. (2024): Virulence of entomopathogenic nematodes and their symbiotic bacteria against insect pests, with special reference to *Agrotis ipsilon* (Lepidoptera: Noctuidae): a comprehensive review. *Egyptian Academic Journal of Biological Sciences (F. Toxicology & Pest Control)*, 16(1):63-122. Doi: 10.21608/EAJBSF.2024.347351
- Ghoneim, K.; Bakr, R.F.A. and Hamadah, Kh. (2021): Disturbing effects of botanicals on the haemogram and immune parameters of insects: recent progress of the search for effective biopesticides. *Egyptian Academic Journal of Biological Sciences (A. Entomology)*, 14(1): 147-193. Doi: 10.21608/EAJBSA.2021.157363
- Ghoneim, K.; Tanani, M.; Hassan, H.A. and Bakr, N.A. (2022): Comparative efficiency of the entomopathogenic nematodes, *Steinernema carpocapsae* and *Heterorhabditis bacteriophora*, against the main body metabolites of *Agrotis ipsilon* (Lepidoptera: Noctuidae). *Egyptian Academic Journal of Biological Sciences (C. Physiology & Molecular Biology)*, 14(2): 57-72. Doi: 10.21608/EAJBSC.2022.260471
- Ghoneim, K.; Hassan, H.A.; Tanani, M. and Bakr, N.A. (2023a): Enzymatic disturbance in larvae of the black cut worm, *Agrotis ipsilon* (Lepidoptera: Noctuidae), by infection with the entomopathogenic nematodes, *Steinernema carpocapsae* and *Heterorhabditis bacteriophora*. *Egyptian Academic Journal of Biological Sciences (C. Physiology & Molecular Biology)*, 15(1):121-142. Doi: 10.21608/EAJBSC.2023.285633
- Ghoneim, K.; Tanani, M.; Hassan, H.A. and Bakr, N.A. (2023b): Pathogenicity of the Entomopathogenic Nematodes, *Steinernema carpocapsae* and *Heterorhabditis bacteriophora*, against the Black Cutworm *Agrotis ipsilon* (Hufnagel) (Lepidoptera: Noctuidae). *European Journal of Applied Sciences*, 11(2): 526-562. Doi:10.14738/aivp.112.14356.
- Gilbert, L.I. and Chino, H. (1974): Transport of lipids in insects. *Journal of Lipid Research*, 15(5): 439-456.
- Gill, H.K. and Garg, H. (2014): Pesticides: Environmental impacts and management strategies. *Pesticides - Toxic Aspects*. <https://doi.org/10.5772/57399>.
- Gillespie, J.P.; Kanost, M.R. and Trenczek, T. (1997): Biological mediators of insect immunity. *Annual Review of Entomology*, 42: 611-643. Doi: 10.1146/annurev.ento.42.1.611.

- Glare, T.R.; Gwynn, R.L. and Moran-Diez, M.E. (2016): Development of biopesticides and future opportunities. In: "Microbial-Based Biopesticides: Methods and Protocols"(Glare, T.R. and Moran-Diez, M.E., eds.). New York: Springer Science+Business Media BV: 211-221.
- Gökkuş, A.; Kahriman, F.; Alatürk, F. and Ali, B. (2016): Variation of nutritional values in leaves and stalks of different maize genotypes having high protein and high oil during vegetation. *Agricultural Science Procedia*, 10: 18–25. <https://doi.org/10.1016/J.AASPRO.2016.09.004>
- Gomaa, Sh.I.; Halawa, S.M.; Shalaby, F.F. and Abdel-Hafez, H.F. (2020): Toxicological and biochemical parameters of microbial preparations on the cotton leafworm *Spodoptera littoralis* (Lepidoptera: Noctuidae). *Egyptian Journal of Plant Protection Research Institute*, 3(1): 204-214.
- González-Santoyo, I. and Córdoba-Aguilar, A. (2012): Phenoloxidase: a key component of the insect immune system. *Entomologia Experimentalis et Applicata*, 142 (1): 1-16. Doi: 10.1111/j.1570-7458.2011.01187.x
- Goodrich-Blair, H. and Clarke, D.J. (2007): Mutualism and pathogenesis in *Xenorhabdus* and *Photorhabdus*: two roads to the same destination. *Molecular Microbiology*, 64: 260–268. Doi: 10.1111/j.1365-2958.2007.05671. x.
- Gordon, R. (1981): Mermithid nematodes: physiological relationships with their insect hosts. *Journal of Nematology*, 13(3): 266–274.
- Gordon, R.; Webster, J.M. and Mead, D.E. (1971): Some effects of the nematode *Mermis nigrescens* upon carbohydrate metabolism in the fat body of its host, the desert locust *Schistocerca gregaria*. *Canadian Journal of Zoology*, 49: 431–434. Doi: 10.1139/z71-066.
- Gordon, R.; Webster, J.M. and Hislop, T.J. (1973): Mermithid parasitism, protein turnover vitellogenesis in the desert locust, *Schistocerca gregaria* Forskal. *Comparative Biochemistry and Physiology B*, 46: 575–593. [https://doi.org/10.1016/0305-0491\(73\)90098-9](https://doi.org/10.1016/0305-0491(73)90098-9)
- Gorman, M.J.; An, Ch. and Kanost, M.R. (2007): Characterization of tyrosine hydroxylase from *Manduca sexta*. *Insect Biochemistry and Molecular Biology*, 37(12): 1327–1337. Doi: 10.1016/j.ibmb.2007.08.006
- Gozel, U. and Gozel, C. (2016): Entomopathogenic Nematodes in Pest Management. In: "Integrated Pest Management (IPM): Environmentally Sound Pest Management"(Gill, H.K. and Goyal, G. eds.). Doi: 10.5772/63894
- Grewal, P.S.; Koppenhöfer, A.M. and Choo, H.Y. (2005): Lawn, turfgrass, and pasture applications. In: "Nematodes as Biocontrol Agents"(Grewal, P.S; Ehlers, R.U. and Shapiro-Ilan, D.I., eds.) Wallingford: CABI Publishing. Cambridge, MA USA, pp: 115-146.
- Grillo, L.A.; Majerowicz, D. and Gondim, K.C. (2007): Lipid metabolism in *Rhodnius prolixus* (Hemiptera: Reduviidae): Role of a midgut triacylglycerol-lipase. *Insect Biochemistry and Molecular Biology*, 37:579–588. Doi: 10.1016/j.ibmb.2007.03.002.
- Gulcu, B.; Cimen, H, Raja, R.K. and Hazir, S. (2017): Entomopathogenic nematodes and their mutualistic bacteria: their ecology and application as microbial control agents. *Biopesticides International*, 3(2): 79-112.
- Gulzar, S.; Usman, M.; Wakil, W.; Wu, Sh.; Oliveira-Hofman, C. R.; Srinivasan, Toews, M. and Shapiro-Ilan, D. (2021): Virulence of entomopathogenic nematodes to pupae of *Frankliniella fusca* (Thysanoptera: Thripidae). *Journal of Economic Entomology*, 114(5): 2018- 2023. <https://doi.org/10.1093/jee/toab132>

- Gümüşsoy, A.; Yüksel, E.; Özer, G.; Imren, M.; Canhilal, R.; Amer, M. and Dababat, A.A. (2022): Identification and biocontrol potential of entomopathogenic nematodes and their endosymbiotic bacteria in apple orchards against the codling moth, *Cydia pomonella* (L.) (Lepidoptera: Tortricidae). *Insects*, 13: 1085. <https://doi.org/10.3390/insects13121085>
- Gunstone, T.; Cornelisse, T.; Klein, K.; Dubey, A. and Donley, N. (2021): Pesticides and soil invertebrates: A hazard assessment. *Frontiers of Environmental Science*, 122. <https://doi.org/10.3389/FENVS.2021.643847>/ BIBTEX
- Hahn, D.A. and Denlinger, D.L. (2007): Meeting the energetic demands of insect diapause: nutrient storage and utilization. *Journal of Insect Physiology*, 53: 760-773. Doi: 10.1016/j.jinsphys.2007.03.018.
- Han-dong, W.; Liu, Q.Z.; Li, X.; and Zhang, H. (2013): Laboratory of Entomology and Nematology, China Agricultural University, Beijing 100193, P.R. China, 8(25): 3245-3250.
- Haq, H.S.; Shaikh, M.A. and Khan, R.H. (2004): Protein proteinase inhibitor genes in combat against insects, pests and pathogens: natural and engineered phytoprotection. *Archives of Biochemistry and Biophysics*, 431: 145-159. Doi: 10.1016/j.abb.2004.07.022.
- Hasegawa, K. and Yamashita, O. (1970): Mode d'action de l'hormone de diapause dans le métabolisme glucidique de ver à soie *Bombyx mori* L. *Annals of Endocrinology*, 31: 631-636.
- Hassan, H.A. (2002): Biological and biochemical studies on the effect of some botanical extracts on cotton leafworm *Spodoptera littoralis* (Boisd.) (Lepidoptera: Noctuidae). M.Sc. Thesis, Faculty of Science, Ain Shams University, Egypt, 173pp.
- Hassan, H.A. and Mohamed, S.A. (2008): Multiple forms of esterase in the larvae of *Pectinophora gossypiella* treated with three volatile oils. *Egyptian Academic Journal of Biological Sciences (A: Entomology)*, 1(2): 145-156. Doi: 10.21608/EAJBSA.2008.15746
- Hassan, H.A.; Shairra, S.A. and Ibrahim, S.S. (2016): Virulence of entomopathogenic nematodes *Steinernema glaseri* and *Heterorhabditis bacteriophora* Poinar (HP88 strain) against the black cutworm, *Agrotis ipsilon*. *Egyptian Academic Journal of Biological Sciences (A: Entomology)*, 9(1): 33-48. Doi: 10.21608/EAJBSA.2016.12853
- Hatfield, M.J.; Umans, R.A.; Hyatt, J.L.; Edwards, C.C.; Wierdl, M.; Tsurkan, L.; Taylor, M.R. and Potter, PM (2016): Carboxylesterases: General detoxifying enzymes. *Chemico-Biological Interactions*, 259: 327-331. <https://doi.org/10.1016/j.cbi.2016.02.011>
- Hayes, J.D.; Flanagan, J.U. and Jowsey, I.R. (2005): Glutathione transferases. *Annual Review of Pharmacology and Toxicology*, 45:51-88. <https://doi.org/10.1146/annurev.pharmtox.45.120403.095857>
- Heil, M.; Buchler, R. and Boland, W. (2005): Quantification of invertase activity in ants under field conditions. *Journal of Chemical Ecology*, 31(2): 431-437. Doi: 10.1007/s10886-005-1352-y.
- Hemming, J.D. and Lindroth, R.L. (2000): Effects of phenolic glycosides and protein on gypsy moth (Lepidoptera: Lymantriidae) and forest tent caterpillar (Lepidoptera: Lasiocampidae) performance and detoxication activities. *Environmental Entomology*, 29(6):1108–1115. <https://doi.org/10.1603/0046-225X-29.6.1108>
- Henriques, B.S.; Garcia, E.S.; Azambuja, P. and Genta, F.A. (2020): Determination of Chitin Content in Insects: An Alternate Method Based on Calcofluor Staining. *Frontiers in Physiology, Section Invertebrate Physiology*, 11. <https://doi.org/10.3389/fphys.2020.00117>

- Hietakangas, V. and Cohen, S.M. (2009): Regulation of tissue growth through nutrient sensing. *Annual Review of Genetics*, 43: 389-410. <http://dx.doi.org/10.1146/annurev-genet-102108-134815>
- Hinks, C.f; Cheeseman, M.T.; Erlandson, M.A.; Olfert, O. and Westcott, N.D. (1991): The effects of Kochia, wheat and oats on digestive proteinases and the protein economy of adult grasshopper *Melanoplus sanguinipes*. *Journal of insect Physiology*, 37: 417-430. [https://doi.org/10.1016/0022-1910\(91\)90051-Z](https://doi.org/10.1016/0022-1910(91)90051-Z)
- Hoffmann, J.A. (1996): Innate immunity higher insects. *Current Opinion in Immunology*, 7:410. Doi: 10.1016/s0952-7915(96)80098-7.
- Hultmark, D. (2003): *Drosophila* immunity: Paths and patterns. *Current Opinion in Immunology*, 15:12–19. Doi: 10.1016/s0952-7915(02)00005-5.
- Holtof, M.; Lenaerts, C.; Cullen, D. and Vanden Broeck, J. (2019): Extra-cellular nutrient digestion and absorption in the insect gut. *Cell Tissue Research*, 377(3): 397–414. <https://doi.org/10.1007/S00441-019-03031-9>.
- Hussain, A.; Rizwan-ul-Haq, M.; Al-Ayedh, H.; Ahmed, S.; Al-Jabr, A.M. (2015): Effect of *Beauveria bassiana* infection on the feeding performance and antioxidant defence of red palm weevil, *Rhynchophorus ferrugineus*. *BioControl*, 60: 849–859. <https://doi.org/10.1007/s10526-015-9682-3>
- Hyršl, P.; Buyukguzel, E. and Buyukguzel, K. (2007): The effects of boric acid-induced oxidative stress on antioxidant enzymes and survivorship in *Galleria mellonella*. *Archives of Insect Biochemistry and Physiology*, 66:23-31. Doi: 10.1002/arch.20194.
- Ibrahim, S.A.M.; Taha, M.A.; Salem, H.H.A. and Farghaly, D.S. (2015): Changes in enzyme activities in *Agrotisipsilon* (Lepidoptera, Noctuidae) as a response to entomopathogenic nematode infection. *International Journal of Advanced Research*, 3(5): 111-118.
- Ibrahim, E.; Dobeš, P.; Kunc, M.; Hyršl, P. and Kodrík, D. (2018): Adipokinetic hormone and adenosine interfere with nematobacterial infection and locomotion in *Drosophila melanogaster*. *Journal of Insect Physiology*, 107:167–174. <https://doi.org/10.1016/j.jinsphys.2018.04.002>.
- Ibrahim, S.A.M, Salem H.H.A. and Taha M.A. (2019): Dual application of entomopathogenic nematodes and fungi on immune and antioxidant enzymes of the greater wax moth, *Galleria mellonella* L. *Egyptian Journal of Biological Pest Control*, 29: 20. <https://doi.org/10.1186/s41938-019-0125-9>
- Içen, E.; Armutçu, F.; Büyükgüz el, K. and Gürel, A. (2005): Biochemical stress indicators of greater wax moth exposure to organophosphorus insecticides. *Journal of Economic Entomology*, 98(2): 358-66. Doi: 10.1603/0022-0493-98.2.358
- Irving, P.; Ubeda, J.; Doucet, D.; Troxler, L.; Lagueux, M.; Zachary, D.; Hofmann, J.; Hetru, C. and Meister, M. (2005): New insights into *Drosophila* larval haemocyte functions through genome wide analysis. *Cell Microbiology*, 7:335–350. Doi: 10.1111/j.1462-5822.2004.00462.x.
- Ismail, S.M. (2018): Joint action of certain insecticides by sublethal dose effect on the cotton leafworm *Spodoptera littoralis* (Lepidoptera: Noctuidae) larvae. *Egyptian Journal of Plant Protection Research*, 1: 43-50.
- Ismail, S.M. (2021): Field persistence of certain new insecticides and their efficacy against black cutworm, *Agrotisipsilon* (Hufnagel). *Bulletin of the National Research Center*, 45:17, 7pp. <https://doi.org/10.1186/s42269-020-00481-y>
- Istkhar, and Chaubey, A.K. (2019): Changes in protein profile and encapsulation avoiding responses of entomopathogenic nematode in the American bollworm *Helicoverpa armigera* (Hübner) (Lepidoptera: Noctuidae). *Egyptian Journal of Biological Pest Control*, 29: 69, 8pp. <https://doi.org/10.1186/s41938-019-0173-1>

- Istkhar, R.; Chaubey, A.K. and Garg, A.P. (2019): Entomopathogenic nematodes in the biological control of insect pests with reference to insect Immunity. In: (Varma, A.; Tripathi, S. and Prasad, R., eds.) "Plant Biotic Interactions". Springer, Switzerland AG.
- Jagodič, A.; Trdan, S. and Laznik, Ž. (2019): Entomopathogenic nematodes: can we use the current knowledge on belowground multitrophic interactions in future plant protection programmes?- Review. *Plant Protection Science*, 55(4): 243- 254. Doi: 10.17221/24/2019-PPS
- James, M.; Malan, A.P. and Addison, P. (2018): Surveying, and screening South African entomopathogenic nematodes for the control of the Mediterranean fruit fly, *Ceratitidis capitata* (Wiedemann). *Crop Protection*, 105: 41-48. <https://doi.org/10.1016/j.cropro.2017.11.008>
- Jeschke, P.; Nauen, R.; Schindler, M. and Elbert, A. (2011): Overview of the status and global strategy for neonicotinoids. *Journal of Agricultural and Food Chemistry*, 59(7): 2897–2908. <https://doi.org/10.1021/jf101303g>
- Jochim, R.J.; Teixeira, C.R.; Laughinghouse, A.; Mu, J.; Oliveira, F. Gomes, R.B.; Elnaiem, D. and Valenzuela, J.G. (2008): The midgut transcriptome of *Lutzomyia longipalpis*: comparative analysis of cDNA libraries from sugar-fed, blood-fed, post-digested and *Leishmania infantumchagasi*-infected sandflies. *BMC Genomics*, 9: 15. Doi: 10.1186/1471-2164-9-15.
- Johnson, D. and Mundel, H. (1987): Grasshopper feeding rats, preferences and growth on sawflower. *Annals of Applied Biology*, 11(1): 43-52. Doi: 10.1111/j.1744-7348.1987.tb01431.x
- Kaliaskar, D.; Shibaeva, A.; Zhappar, N.; Shaikhutdinov, V.; Asherbekova, L.; Bekbulatov, S. and Kalyaskarova, A. (2022): The efficiency of aboriginal entomopathogenic nematodes from semi-arid zone against Tenebrionidae larvae with comparison to commercial bio-insecticides. *AGRIVITA Journal of Agricultural Science*, 44(3): 526-536. <http://doi.org/10.17503/agrivita.v44i3.3760>
- Kamruzzaman, A.S.M.; Mikani, A.; Mohamed, A.A.; Elgendy, A.M. and Takeda, M. (2020): Crosstalk among indoleamines, neuropeptides and JH/20E in regulation of reproduction in the American cockroach *Periplaneta americana*. *Insects*, 11: 155. Doi:10.3390/insects11030155.
- Kanost, M. and Gorman, M.J. (2008): Phenoloxidases in insect immunity. In: "Insect Immunology"(Beckage, N., ed.), pp: 69-96. Academic Press, San Diego, CA, USA. Doi: 10.1016/B978-012373976-6.50006-9
- Kanost, M.R.; Jiang, H. and Yu, X.Q. (2004): Innate immune responses of a lepidopteran insect, *Manduca sexta*. *Immunology Review*, 198: 97-105. <https://doi.org/10.1111/j.0105-2896.2004.0121.x>
- Kaufmann, C. and Brown, M.R. (2008): Regulation of carbohydrate metabolism and flight performance by a hypertrehalosaemic hormone in the mosquito *Anopheles gambiae*. *Journal of Insect Physiology*, 54: 367-377. Doi: 10.1016/j.jinsphys.2007.10.007
- Kaur, S.P.; Sidhu, D.S.; Dhillon, S.S. and Kumar, N.K. (1985): Transaminases during development and aging of the bruchid *Zabrotes subfasciatus* (Boh.) (Coleoptera: Bruchidae). *Insect Science and its Application*, 6(5): 585-590.
- Kaya, H.K. and Gaugler, R. (1993): Entomopathogenic nematodes. *Annual Review of Entomology*, 38: 181–206. <https://doi.org/10.1146/annurev.en.38.010193.001145>
- Khan, R.R.; Ali, R.A.; Ali, A.; Arshad, M.; Majeed, Sh.; Ahmed, S.; Khan, S.A. and Arshad, M. (2018): Compatibility of entomopathogenic nematodes (Nematoda: Rhabditida) and the biocide, spinosad for mitigation of the armyworm, *Spodoptera litura* (F.)

- (Lepidoptera: Noctuidae). *Egyptian Journal of Biological Pest Control*, 28: 1–6. <https://doi.org/10.1186/s41938-018-0063-y>
- Khater, K.S.; El-lakwah, S.F.; Abd-Elmonem, H.M.; Ahmed, F.A. and Shoukry, I.F. (2020): Biochemical Virulence of Some Entomopathogenic Nematodes on *Galleria mellonella* larvae (Lepidoptera: Galleridae). *Egyptian Academic Journal of Biological Sciences (C. Physiology & Molecular Biology)*, 12(2): 97-109. Doi: 10.21608/EAJBSC.2020.125099
- Khorshidi, M.; Abad, R.F.P.; Saber, M. and Zibae, A. (2019): Effects of hexaflumuron, lufenuron and chlorfluazuron on certain biological and physiological parameters of *Helicoverpa armigera* (Hübner) (Lepidoptera: Noctuidae). *Biocatalysis and Agricultural Biotechnology*, 21, 101270: 8pp. <https://doi.org/10.1016/j.bcab.2019.101270>
- Kim, K.; Kim, Y. and Kim, Y. (2002): Biochemical evidence of the inhibitory effect of diflubenzuron on the metamorphosis of the silkworm, *Bombyx mori*. *Journal of Asia-Pacific Entomology*, 5: 175-180. [https://doi.org/10.1016/S1226-8615\(08\)60149-1](https://doi.org/10.1016/S1226-8615(08)60149-1)
- Kolawole, A.O. and Kolawole, A.N. (2014): Insecticides and bio-insecticides modulate the glutathione-related antioxidant defense system of cowpea storage bruchid (*Callosobruchus maculatus*). *International Journal of Insect Science*, 6: 79–88. Doi: 10.4137/IJIS.S18029.eCollection 2014.
- Koppenhöfer, H.S. (2007): Bacterial symbionts of *Steinernema* and *Heterorhabditis*. In: "Entomopathogenic nematodes: systematics, phylogeny and bacterial symbionts"(Nguyen K, Hunt D.M., eds.). Brill, Leiden. https://doi.org/10.1163/ej.97890_04152_939.i-816.43
- Koppenhöfer, A.M.; Shapiro-Ilan, D.I. and Hiltbold, I. (2020): Entomopathogenic nematodes in sustainable food production. *Frontiers in Sustainable Food Systems*, 4:125. Doi: 10.3389/fsufs.2020.00125
- Kozaki, T.; Shono, T.; Tomita, T.; and Kono, Y. (2001): Fenitroxon insensitive acetylcholinesterases of the housefly, *Musca domestica* associated with point mutations. *Insect Biochemistry and Molecular Biology*, 31: 991-997. Doi: 10.1016/s0965-1748(01)00047-9.
- Kramarz, P.; Malek, D.; Gawel, M.; Drobniak, S.M. and Homa, J. (2016): Reproductive status of *Tribolium castaneum* (Coleoptera: Tenebrionidae) affects its response to infection by *Steinernema feltiae* (Rhabditida: Steinernematidae). *European Journal of Entomology*, 113:309–314. Doi: 10.14411/eje.2016.039
- Krishnayya, P.V. and Grewal, P.S. (2002): Effect of neem and selected fungicides on viability and virulence of the entomopathogenic nematode *Steinernema feltiae*. *Biocontrol Science and Technology*, 12: 259-266. <https://doi.org/10.1080/09583150210388>
- Kucera, M. and Mracek, Z. (1989): Proteolytic enzymes of the invasive larvae of entomopathogenic Steinernematidae nematodes. *Acta Entomologica Bohemoslovaca*, 86:193-202.
- Kumar, P.; Ganguly, S. and Somvanshi, V.S. (2015): Identification of virulent entomopathogenic nematode isolates from a countrywide survey in India. *International Journal of Pest Management*, 61(2): 135-143. <https://doi.org/10.1080/09670874.2015.1023869>
- Kumar, R.; Pandey, S. and Singh, R. (2022): Evaluation of the entomopathogenic nematode, *Steinernema asiaticum* against the diamondback moth, *Plutellaxylostella* (Linnaeus) (Lepidoptera: Plutellidae) under screen house and field conditions. *Egyptian Journal of Biological Pest Control*, 32:90 <https://doi.org/10.1186/s41938-022-00589-5>

- Kunbhar, S.; Rajput, L.B.; Ahmed, G.A.; Akber, C.G. and Sahito, J.G.M. (2018): Impact of botanical pesticides against sucking insect pests and their insect predators in brinjal crop. *Journal of Entomology and Zoology Studies*, 6: 83–87.
- Kunc, M.; Badrul, A.; Pavel, H. and Ulrich, T. (2017): Monitoring the effect of pathogenic nematodes on locomotion of *Drosophila* larvae. *Fly*, 3:1–10. Doi: 10. 1080/19336934.2017.1297350
- Lacey, L.A. and Georgis, R. (2012): Entomopathogenic nematodes for control of insect pests above and below ground with comments on commercial production. *Journal of Nematology*, 44(2): 218–225.
- Lacey, L.A.; Grzywacz, D.; Shapiro-Ilan, D.I.; Frutos, R.; Brownbridge, M. and Goettel, M.S. (2015): Insect pathogens as biological control agents: back to the future. *Journal of Invertebrate Pathology*, 132: 1-41. <https://doi.org/10.1016/j.jip.2015.07.009>
- Lalitha, K.; Karthi, S.; Vengateswari, G.; Karthikraja, R.; Perumal, P. and Shivakumar, M.S. (2018): Effect of entomopathogenic nematode of *Heterorhabditis indica* infection on immune and antioxidant system in lepidopteran pest *Spodoptera litura* (Lepidoptera: Noctuidae). *Journal of Parasitic Diseases*, 42(2): 204–211. <https://doi.org/10.1007/s12639-018-0983-1>
- Lavine, M.D. and Strand, M.R. (2002): Insect haemocytes and their role in immunity. *Insect Biochemistry and Molecular Biology*, 32: 1295-1309. [https://doi.org/10.1016/S0965-1748\(02\)00092-9](https://doi.org/10.1016/S0965-1748(02)00092-9)
- Lazarevi, J. and Tomani, M.J. (2015): Dietary and phylogenetic correlates of digestive trypsin activity in insect pests. *Entomologia Experimentalis et Applicata*, 157:123–151. <https://doi.org/10.1111/eea.12349>
- Laznik, Ž. and Trdan S. (2011) Entomopathogenic nematodes (Nematoda: Rhabditida) in Slovenia: from tabula rasa to implementation into crop production systems. 2011. In: "Insecticides - advances in integrated pest management"(Perveen, F., ed). InTech, Rijeka, pp: 627–656
- Laznik, Z. and Trdan, S. (2012): Entomopathogenic nematodes (Nematoda: Rhabditida) in Slovenia: from tabularasa to implementation into crop production systems. In: "Insecticides Advances in Integrated Pest Management" (Perveen F, ed.), pp: 627-656. Doi: 10.5772/29540
- Laznik, Z.; Vidrih, M. and Trdan, S. (2012): The effect of different entomopathogens on white grubs (Coleoptera: Scarabaeidae) in an organic hay producing grassland. *Archives of Biological Sciences*, 64(4):1235-1246. Doi: 10.2298/ABS1204235L
- Leaver, M.J. and George, S.G. (1998): A piscine glutathione S-transferase which efficiently conjugates the end-products of lipid peroxidation. *Marine Environmental Research*, 46: 71-74. [http://dx.doi.org/10.1016/S0141-1136\(97\)00071-8](http://dx.doi.org/10.1016/S0141-1136(97)00071-8)
- Lee, D.L. and Atkinson, H.J. (1976): Physiology of nematodes, 2nd ed. Columbia University Press, New York.
- Leonar, A.L.; Nimkingrat, P.; Aryal, S.; Martinez, J.G.; Bhat, A.H. and Sumaya, N.H. (2022): Natural association of the entomopathogenic nematode *Heterorhabditis indica* (Rhabditida: Heterorhabditidae) from the Philippines with the non-symbiotic bacterium *Ochrobactrum anthropi* (Proteobacteria: Brucellaceae). *Egyptian Journal of Biological Pest Control*, 32:83 <https://doi.org/10.1186/s41938-022-00576-w>
- Lewis, E.E. and Clarke, D.J. (2012): Nematode parasites and entomopathogens. In: "Insect Pathology" (Vega FE and Kaya HK., eds.). 2nd ed. Elsevier, Netherlands. pp: 395-424 Doi: 10.1016/B978-0-12-384984-7.00011-7
- Li, F.L. and Han, Z.J. (2002): Purification and characterization of acetylcholinesterase from cotton aphid (*Aphis gossypii* Glover). *Archives of Insect Biochemistry and Physiology*, 51(1): 37-45. Doi: 10.1002/arch.10048

- Li, X.Z. and Liu, Y.H. (2007): Diet influences the detoxification enzyme activity of *Bactrocera tau* (Walker) (Diptera: Tephritidae). *Acta Entomologica Sinica*, 50(10): 989-995. <http://www.insect.org.cn/EN/Y2007/V50/I10/989>
- Li, X.Y.; Cowles, R.S.; Cowles, E.A.; Gaugler, R. and Cox-Foster, D.L. (2007): Relationship between the successful infection by entomopathogenic nematodes and the host immune response. *International Journal for Parasitology*, 37: 365–374. Doi: 10.1016/j.ijpara.2006.08.009.
- Li, X.; Q. Liu, E.E. Lewis, and E. Tarasco, (2016): Activity changes of antioxidant and detoxifying enzymes in *Tenebrio molitor* (Coleoptera: Tenebrionidae) larvae infected by the entomopathogenic nematode *Heterorhabditis beicherriana* (Rhabditida: Heterorhabditidae). *Parasitology Research*, 115(12):4485-4494. Doi: 10.1007/s00436-016-5235-7..
- Li, W.; Zhao, X.; Yuan, W. and Wu, K. (2017): Activities of digestive enzymes in the omnivorous pest *Apolygus lucorum* (Hemiptera: Miridae). *Journal of Economic Entomology*, 110(1):101–110. <https://doi.org/10.1093/JEE/TOW263>.
- Li, E.; H. Wu, Zh. Wang, K. L. Li, Sh. Zhang, Y. Cao, and J. Yin, (2022): Implication of antioxidant and detoxifying enzymes in the resistance of *Holotrichia parallela* larvae to EPN-*Bt* infection. *Research Square*, 19pp. Doi: <https://doi.org/10.21203/rs.3.rs-1541715/v1>
- Liao, M.; Xiao, J.J.; Zhou, L.J.; Yao, X.; Tang, F.; Hua, R.M.; Wu, X.W. and Cao, H.Q. (2017): Chemical composition, insecticidal and biochemical effects of *Melaleuca alternifolia* essential oil on the *Helicoverpa armigera*. *Journal of Applied Entomology*, 141: 721–728. <https://doi.org/10.1111/jen.12397>
- Ling, E.; Shirai, K.; Kanekatsu, R. and Kiguchi, K. (2005): Hemocyte differentiation in the hematopoietic organs of the silkworm, *Bombyx mori*: prohemocytes have the function of phagocytosis. *Cell Tissue Research*, 320: 535-543. <https://doi.org/10.1007/s00441-004-1038-8>
- Liu, Y.; Sui, Y.P.; Wang, J.X. and Zhao, X.F. (2009): Characterization of the trypsin-like protease (HA-TLP2) constitutively expressed in the integument of the cotton bollworm, *Helicoverpa armigera*. *Archives of Insect Biochemistry and Physiology*, 72: 74–87. Doi: 10.1002/arch.20324.
- Liu, J.; Bai, H.; Song, P.; Nangong, Z.; Dong, Z.; Li, Z. and Wang, Q. (2020a): Insecticidal activity of chitinases from *Xenorhabdus nematophila* HB310 and its relationship with the toxin complex. *Toxins*, 14: 646. Doi: 10.3390/toxins14090646
- Liu, W.T.; Chen, T.L.; Hou, R.F.; Chen, C.C. and Tu, W.C. (2020b): The invasion and encapsulation of the entomopathogenic nematode, *Steinernema abbasi*, in *Aedes albopictus* (Diptera: Culicidae) larvae. *Insects*, 11(12): 832. <https://doi.org/10.3390/insects11120832>
- Lu, Z. and Jiang, H. (2007): Regulation of phenoloxidase activity by high-and low- molecular weight inhibitors from the larval haemolymph of *Manduca sexta*. *Insect Biochemistry and Molecular Biology*, 37:478-485. <https://doi.org/10.1016/j.ibmb.2007.02.004>
- Lu, A.; Zhang, Q.; Zhang, J. Yang, B.; Wu, K.; Xie, W.; Luan, Y.-X. and Ling, E. (2014): Insect prophenoloxidase: the view beyond immunity. *Frontiers in Physiology*, 5: 252. <https://doi.org/10.3389/fphys.2014.00252>
- Macedo, M.L.R. and Freire, M.G.M. (2011): Insect digestive enzymes as a target for pest control. *Invertebrate Survival Journal*, 8: 190-198.
- Mahanta, D.K.; Bhoi, T.K.; Komal, J.; Samal, I.; Nikhil, R.M.; Paschapur, A.U.; Singh, G.; Kumar, P.V.D.; Desai, H.R.; Ahmad, M.A.; Singh, P.P.; Majhi, P.K.; Mukherjee, U.; Singh, P.; Saini, V.; Srinivasa, N. and Yele, Y. (2023): Insect-pathogen crosstalk and the cellular-molecular mechanisms of insect immunity: uncovering the underlying

- signaling pathways and immune regulatory function of non-coding RNAs. *Frontiers in Immunology*, 14: 1169152. Doi: 10.3389/fimmu.2023.1169152
- Mahmood, S.; Kumar, M.; Kumari, P.; Mahapatro, G.K.; Banerjee, N. and Sarin, N.B. (2020): Novel insecticidal chitinase from the insect pathogen *Xenorhabdus nematophila*. *International Journal of Biological Macromolecules*, 159: 394–401. Doi: 10.1016/j.ijbiomac. 2020.05.078.
- Mahmoud M.F. (2014): Efficacy of entomopathogenic nematodes to certain insect pests infesting oilseed rape in the laboratory and greenhouse. *Egyptian Journal of Biological Pest Control*, 24(2): 387–391.
- Mahmoud, M.F.; Mahfouz, H.M. and Mohamed, K.M. (2016): Compatibility of entomopathogenic nematodes with neonicotinoids and azadirachtin insecticides for controlling the black cutworm, *Agrotis ipsilon* (Hufnagel) in Canola Plants. *International Journal of Research in Environmental Science*, 2(1): 11–18. Doi: dx.doi.org/10. 20431/2454-9444.0201002
- Malan, A.P.; Knoetze, R. and Moore, S.D. (2011): Isolation and identification of entomopathogenic nematodes from citrus orchards in South Africa and their biocontrol potential against false codling moth. *Journal of Invertebrate Pathology*, 108:115–25. <https://doi.org/10.1016/j.jip.2011.07.006>
- Marmaras, V.J.; Charalambidis, N.D. and Zervas, C.G. (1996): Immune response in insects: the role of phenoloxidase in defense reactions in relation to melanization and sclerotization. *Archives of Insect Biochemistry and Physiology*, 31: 119–133. Doi: 10.1002/(SICI)1520-6327(1996)31:2<119::AID-ARCH1>3.0.CO;2-V
- Mates, J.M. (2000): Effects of antioxidant enzymes in the molecular control of reactive oxygen species toxicology. *Toxicology*, 153(1): 83–104. Doi: 10.1016/s0300-483x(00)00306-1.
- Mehrabadi, M.; Bandani, A.R.; Mehrabadi, R. and Alizadeh, H. (2012): Inhibitory activity of proteinaceous α -amylase inhibitors from triticale seeds against *Eurygaster integriceps* salivary α -amylases: Interaction of the inhibitors and the insect. *Pesticide Biochemistry and Physiology*, 102: 220–228. <https://doi.org/10.1016/j.pestbp.2012.01.008>
- Meister, M. and Lagueux, M. (2003): *Drosophila* blood cells. *Cellular Microbiology*, 5: 573–580. Doi: 10.1046/j.1462-5822.2003.00302.x
- Merzendorfer, H. and Zimoch, L. (2003): Chitin metabolism in insects: structure, function and regulation of chitin synthases and chitinases. *Journal of Experimental Biology*, 206(24): 4393–4412. <https://doi.org/10.1242/JEB.00709>
- Meslin, C.; Bozzolan, F.; Braman, V.; Chardonnet, S.; Pionneau, C.; François, M.-Ch.; Severac, D.; Gadenne, Ch.; Anton, S.; Maibèche, M.; Jacquin-Joly, E. and Siaussat, D. (2021): Sublethal exposure effects of the neonicotinoid clothianidin strongly modify the brain transcriptome and proteome in the male moth *Agrotis ipsilon*. *Insects*, 12(2):152. Doi: 10.3390/insects12020152.
- Mirhaghparast, S.K.; Zibae, A. and Hajizadeh, J. (2013): Effects of *Beauveria bassiana* and *Metarhizium anisopliae* on cellular immunity and intermediary metabolism of *Spodoptera littoralis* Boisduval (Lepidoptera: Noctuidae). *Invertebrate Survival Journal*, 10: 110–119.
- Mohamed, A.A.; Zhang, L.; Dorrah, M.A.; Elmogy, M. and Yousef, H.A. (2016): Molecular characterization of a c-type lysozyme from the desert locust, *Schistocerca gregaria* (Orthoptera: Acrididae). *Developmental and Comparative Immunology*, 61: 60–69. Doi: 10.1016/j.dci.2016.03.018.

- Mona, P.M. (2001): On the developmental profile of hormones in *Spodoptera mauritia* Boisd. (Lepidoptera: Noctuidae), Ph.D Thesis, University of Calicut, Calicut, India, 137 pp.
- Montella, I.R.; Schama, R. and Valle, D. (2012): The classification of esterases: an important gene family involved in insecticide resistance—a review. *Memórias do Instituto Oswaldo Cruz*, 107: 437–449. <https://doi.org/10.1590/S0074-02762012000400001>
- Moret, Y. (2006): ‘Trans-generational immune priming’: specific enhancement of the antimicrobial immune response in the mealworm beetle, *Tenebrio molitor*. *Proceedings of the Biological Sciences*, 273:1399–1405. Doi: 10.1098/rspb.2006.3465.
- Moret, Y. and Siva-Jothy, M.T. (2003): Adaptive innate immunity? Responsive-mode prophylaxis in the mealworm beetle, *Tenebrio molitor*. *Proceedings of the Royal Society of London. Series B*, 270: 2475–2480. Doi:10.1098/rspb.2003.2511
- Mostafalou, S. and Abdollahi, M. (2012): Concerns of environmental persistence of pesticides and human chronic diseases. *Clinical and Experimental Pharmacology*, 01(S5). <https://doi.org/10.4172/2161-1459.s5-e002>.
- Muhammad, J.; Fathy, Z. and Moussa, S. (2022): Entomopathogenic bacteria as natural enemy against the African migratory locust, *Locusta migratoria migratorioides* (Reiche & Fairmaire, 1849) (Orthoptera: Acrididae). *Egyptian Journal of Biological Pest Control*, 32:92. <https://doi.org/10.1186/s41938-022-00592-w>
- Mukanganyama, S.; Figueroa, Ch.C.; Hasler, J.A. and Niemeyer, H.M. (2003): Effects of DIMBOA on detoxification enzymes of the aphid. *Journal of Insect Physiology*, 49(3): 223–229. Doi:10.1016/S0022-1910(02)00269-X
- Mukherjee, S.N. and Sharma, R.N. (1996): Azadirachtin induced changes in feeding, dietary utilization and midgut carboxylesterase activity of the final instar larvae of *Spodoptera litura* (Fabricius) (Lepidoptera: Noctuidae). *Journal of Environmental Science and Health*, B31: 1307–1319.
- Müller, P.; Donnelly, M.J. and Ranson, H. (2007): Transcription profiling of a recently colonised pyrethroid resistant *Anopheles gambiae* strain from Ghana. *BMC Genomics*, 8:36–47. <https://doi.org/10.1186/1471-2164-8-36>
- Muñoz, P.; Meseguer, J. and Esteban, M.A. (2006): Phenoloxidase activity in three commercial bivalve species: Changes due to natural infestation with *Perkinsus atlanticus*. *Fish and Shellfish Immunology*, 20:12–19. Doi: 10.1016/j.fsi.2005.02.002
- Nakazawa, H.; Tsuneishi, E.; Ponnuvel, K.M.; Furukawa, S.; Asaoka, A. Tanaka, H.; Ishibashi, J. and Yamakawa, M. (2004): Antiviral activity of a serine protease from the digestive juice of *Bombyx mori* larvae against nucleopolyhedrovirus. *Virology*, 321: 154–162. Doi: 10.1016/j.virol.2003.12.011.
- Nappi, A.J. and Christensen, B.M. (2005): Melanogenesis and associated cytotoxic reactions: applications to insect innate immunity. *Insect Biochemistry and Molecular Biology*, 35: 443–459. Doi: 10.1016/j.ibmb.2005.01.014.
- Nappi, A.J. and Vass, E. (2001): Cytotoxic reactions associated with insect immunity. *Advances in Experimental Medicine and Biology*, 484: 329–348. Doi: 10.1007/978-1-4615-1291-2_33.
- Nath, B.S. (2000): Changes in carbohydrate metabolism in hemolymph and fat body of the silkworm, *Bombyx mori* L., exposed to organophosphorus insecticides. *Pesticide Biochemistry and Physiology*, 68(3): 127–137. <https://doi.org/10.1006/pest.2000.2509>
- Nation, J.L. (2008): *Insect Physiology and Biochemistry*, 2nd ed. CRC Press, London, 560pp. <https://doi.org/10.1201/9781420061789>

- Nawaz, F.; Khan, N.; Shah, J.A.; Khan, A.; Liaqat, A.; Ullah, S.; Khalil, A. U.; Jan, T.; Ullah, S.; Ali, M. and Ali, M. (2017): Yield and yield components of chickpea as affected by various levels of FYM and rhizobium inoculation. *Pure and Applied Biology*, 6(1): 346-351. <http://dx.doi.org/10.19045/bspab.2017.60033>
- Neuwirth, M. (2005): The structure of the haemocytes of *Galleria mellonella*. *Journal of Nematology*, 139: 105-123.
- Nickle, W.R. and Welch, H.E. (1984): Nematode parasites of Lepidoptera. In: "Plant and insect nematodes" (pp. 655-696). New York and Basel: Marcel Decker Inco.
- Nordberg, J. and Arner, E.S.J. (2001): Reactive oxygen species, antioxidants, and the mammalian thioredoxin system. *Free Radical Biology and Medicine*, 31(11):1287–1312. Doi:10.1016/s0891-5849(01)00724-9.
- Oakeshott, J.G.; Claudianos, C.; Campbell, P.M.; Newcomb, R.D. and Russell, R.J. (2005): Biochemical genetics and genomics of insect esterases. In: "Comprehensive Molecular Insect Science" (Eds. Gilbert, L.I., Latrou, K. & Gill, S.S.), Vol. Elsevier BV, Oxford, UK, pp: 309-381.
- Obiamiwe, B.A. and MacDonald, W.W. (1973): A new parasite of mosquitoes, *Reesimermismuspratti* sp. nov. (Nematoda: Mermithidae), with notes on its life cycle. *Annals of Tropical Medical Parasitology*, 67:439–444. <https://doi.org/10.1080/00034983.1973.11686912>
- Odendaal, D.; Addison, M.F. and Malan, A.P. (2016): Control of diapausing codling moth, *Cydia pomonella* (Lepidoptera: Tortricidae), in wooden fruit bins using entomopathogenic nematodes (Heterorhabditidae and Steinernematidae). *Biocontrol Science and Technology*, 26(11): 1504-1515. Doi: <https://doi.org/10.4001/003.023.0224>
- Oruc, E. (2011): Effects of diazinon on antioxidant defense system and lipid peroxidation in the liver of *Cyprinus carpio* (L.). *Environmental Toxicology*, 26: 571-578. Doi: 10.1002/tox.20573.
- Pant, R. and Kumar, S. (1979): Metabolic fate of carbohydrates and lipids during moulting cycle of *Philosamiaricini* (Lepidoptera: Saturniidae). *Insect Biochemistry*, 9: 577-582. [https://doi.org/10.1016/0020-1790\(79\)90095-7](https://doi.org/10.1016/0020-1790(79)90095-7)
- Papadopoulos, A.I.; Boukouvala, E. and Kakaliouras, G. (2000): Effect of organophosphate and pyrethroid insecticides on the expression of GSTs from *Tenebrio molitor* pupae. *Pesticide Biochemistry and Physiology*, 68: 26–33. <https://doi.org/10.1006/pest.2000.2489>
- Parde, V.D.; Sharma, H.C. and Kachole, M.S. (2012): Inhibition of *Helicoverpa armigera* gut pro-proteinase activation in response to synthetic protease inhibitors. *Entomologia Experimentalis et Applicata*, 142: 104–113. <https://doi.org/10.1111/j.1570-7458.2011.01209.x>
- Peçen, A. and Kepenekci, İ. (2022): Efficacy of entomopathogenic nematode isolates from Turkey against wheat stink bug, *Aelia rostrata* Boheman (Hemiptera: Pentatomidae) adults under laboratory conditions. *Egyptian Journal of Biological Pest Control*, 32:91 <https://doi.org/10.1186/s41938-022-00590-y>
- Peschiutta, M.; Cagnolo, S.R. and Almirón, W.R. (2014): Susceptibility of larvae of *Aedes aegypti* (Linnaeus) (Diptera: Culicidae) to entomopathogenic nematode *Heterorhabditis bacteriophora* (Poinar) (Rhabditida: Heterorhabditidae). *Revista de la Sociedad Entomológica Argentina*, 73 (3-4): 99-108.
- Petersen, J.J. and Willis, O.R. (1970): Some factors affecting parasitism by mermithid nematodes in southern house mosquito larvae. *Journal of Economic Entomology*, 63(1): 175–178. <https://doi.org/10.1093/jee/63.1.175>.

- Plant, R. and Morris, I.D. (1972): A comparative study on the variation of aminotransferase activity and its total free amino acids in the fat body, haemolymph, intestine and haemolymph protein content in *Philosamia ricini* during larval-pupal development. *Indian Journal of Biochemistry and Biophysics*, 9: 199-202.
- Poinar, G.O. (1990): Taxonomy and biology of Steinernematidae and Heterorhabditidae. In: "Entomopathogenic Nematodes in Biological Control" (R Gaugler & H K Kaya, eds.), pp. 23–61. CRC Press, Boca Raton, FL, USA
- Prabhu, V.K.K. and Sreekumar S. (1994): Endocrine regulation of feeding and digestion in insects. In "Perspectives in entomological research" (Agarwal, O.P., ed.). Scientific Publishers, Jodhpur, 117.
- Pugazhvendan, S.R. and Soundararajan, M. (2009): Effects of Penfluronon total hemocyte count of *Chrysocoris purpures*. *Middle East Journal of Science Research*, 4: 338-340.
- Rahman, I. and Macnee, W. (2000): Regulation of redox glutathione levels and gene transcription in lung inflammation: therapeutic approaches. *Free Radical Biology and Medicine*, 28:1405–1420. Doi:10.1016/S0891-5849(00)00215-X
- Ramsey, J.S.; Rider, D.S.T.K.; Walsh, M.; De Vos, K.H.J.; Gordon, L.; Ponnala, S.L.; Macmil, B.A.; Roe. and Jander, G. (2010): Comparative analysis of detoxification enzymes in *Acyrtosiphon pisum* and *Myzus persicae*. *Insect Molecular Biology*, 19: 155–164. Doi: 10.1111/j.1365-2583.2009.00973. x.
- Ranson, H.; Claudianos, C.; Ortelli, F.; Abgrall, C.; Hemingway, J.; Sharakhova, M.V.; Unger, M.F.; Collins, F.H. and Feyereisen, R. (2002): Evolution of multigene families associated with insecticide resistance. *Science*, 298(5591): 179-181. Doi:10.1126/science. 1076781.
- Ribeiro, C. and Brehelin, M. (2006): Insect haemocytes: what type of cell is that? *Journal of Insect Physiology*, 52: 417- 429. [https://doi.org/ 10.1016/j.jinsphys.2006.01.005](https://doi.org/10.1016/j.jinsphys.2006.01.005)
- Ribeiro, J.M.; Rowton, E.D. and Charlab, R. (2000): Salivary amylase activity of the phlebotomine sand fly, *Lutzomyia longipalpis*. *Insect Biochemistry and Molecular Biology*, 30(4): 271-277. Doi: 10.1016/s0965-1748(99)00119-8
- Richards, G.R. and Blair, H.G. (2010): Examination of *Xenorhabdus nematophila* lipases in pathogenic and mutualistic host interactions reveals a role of xlpA in nematode progeny production. *Applied and Environmental Microbiology*, 76: 221–229. Doi:10.1128/AEM. 01715-09
- Rodriguez-Ortega, M.J.; Grosvik, B.E.; Rodriguez-Ariza, A.; Goksoyr, A. and Lopez-Barea, J. (2003): Changes in protein expression profiles in bivalve molluscs (*Chamaelea gallina*) exposed to four model environmental pollutants. *Proteomics*, 3: 1535-1543. Doi: 10.1002/pmic.200300491.
- Royer, V.; Fraichard, S. and Bouhin, H. (2002): A novel putative insect chitinase with multiple catalytic domains: hormonal regulation during metamorphosis. *Biochemical Journal*, 366: 921–928. Doi: 10.1042/BJ20011764.
- Sant'Anna, M.R.V.; Diaz-Albiter, H.; Mubarak, M.; Dillon, R.J. and Bates, P.A. (2009): Inhibition of trypsin expression in *Lutzomyia longipalpis* using RNAi enhances the survival of *Leishmania*. *Parasites & Vectors*, 2: 62. <https://doi.org/10.1186/1756-3305-2-62>
- Satyavathi, V.V.; Mohamed, A.A.; Kumari, S.; Mamatha, D.M. and Duvic, B. (2018): The IMD pathway regulates lysozyme-like proteins (LLPs) in the silkworm *Antheraea mylitta*. *Journal of Invertebrate Pathology*, 154: 102–108. Doi: 10.1016/j.jip.2018.04.006
- Schenk, P.M.; Kazan, K.; Wilson, I.; Anderson, J.P.; Richmond, T.; Somerville, S.C. and Manners, J.M. (2000): Coordinated plant defense responses in *Arabidopsis* revealed

- by cDNA microarray analysis. *Proceedings of the National Academy of Sciences of the United States of America*, 97: 11655–11660. <https://doi.org/10.1073/pnas.97.21.11655>
- Schmidt, S.P. and Platzer, E.G. (1979): Hemolymph composition of mosquito larvae infected with a mermithid nematode. *Journal of Nematology*, 10: 299.
- Schmidt, S.P. and Platzer, E.G. (1980): Changes in body tissues and hemolymph composition of *Culex pipiens* in response to infection by *Romanomermisculicivora*. *Journal of Invertebrate Pathology*, 36, 240-254. [https://doi.org/10.1016/0022-2011\(80\)90030-0](https://doi.org/10.1016/0022-2011(80)90030-0)
- Scriber, J.M. and Slansky, F.Jr. (1981): The nutritional ecology of immature insects. *Annual Review of Entomology*, 26:183-211. <https://doi.org/10.1146/annurev.en.26.010181.001151>
- Senthil Nathan, S.; Kalaivani, K. and Chung, P.G. (2005): The effects of azadirachtin and nucleopolyhedrovirus on midgut enzymatic profile of *Spodoptera litura* Fab. (Lepidoptera: Noctuidae). *Pesticide Biochemistry and Physiology*, 83: 46–57. <https://doi.org/10.1016/j.pestbp.2005.03.009>
- Serebrov, V.V.; Alekseev, A.A. and Glupov, V.V. (2001): Changes in the activity and pattern of hemolymph esterases in the larvae of wax moth *Galleria mellonella* L. (Lepidoptera, Pyralidae) during mycosis. *Biology Bulletin of the Russian Academy of Sciences*, (5):588-92. (Article in Russian).
- Serebrov, V.V.; Gerber, O.N.; Malyarchuk, A.A.; Martemyanov, V.V.; Alekseev, A.A. and Glupov, V.V. (2006): Effect of entomopathogenic fungi on detoxification enzyme activity in greater wax moth *Galleria mellonella* L. (Lepidoptera, Pyralidae) and role of detoxification enzymes in development of insect resistance to entomopathogenic fungi. *Izvestiya Akademii Nauk, Seriya Bio-logicheskaya*, 6: 712–718. *Biology Bulletin*, 33(6): 581–586. Doi: 10.1134/S1062359006060082
- Shah, F.A.; Abdorrahman, M.M.; Berry, C.; Touray, M.; Hazir, S. and Butt, T.M. (2021): Indiscriminate ingestion of entomopathogenic nematodes and their symbiotic bacteria by *Aedes aegypti* larvae: a novel strategy to control the vector of Chikungunya, dengue and yellow fever. *Turkish Journal of Zoology*, 45: 372-383. Doi:10.3906/zoo-2107-2
- Shahzad, M.; Qu, Y.; Zafar, A.; Ur Rehman, S. and Islam, T. (2020): Exploring the influence of knowledge management process on corporate sustainable performance through green innovation. *Journal of Knowledge Management*, 24(9): 2079-2106. Doi: 10.1108/JKM-11-2019-0624
- Shairra, S.A. and Awad, H.H. (2011): Effect of the Entomopathogenic Nematode *Heterorhabditis bacteriophora* (Hp 88) and the Garlic Extract *Allium sativum* on the Immune Challenge of the Desert Locust, *Schistocerca gregaria* (Forsk.). *Egyptian Journal of Biological Pest Control*, 21(1), p.11.
- Shairra, S.A.; El-Sharkawy, M.A.A.; Hassan, K.A. and Ahmed, D.A. (2016): The Efficacy of entomopathogenic nematodes on the pink bollworm, *Pectinophora gossypiella*. *Egyptian Academic Journal of Biological Sciences (F. Toxicology & Pest control)*, 8(2): 103 – 113. Doi: 10.21608/EAJBSF.2016.17123
- Shapiro-Ilan, D.I. and Brown, I. (2013): Earthworms as phoretic hosts for *Steinernema carpocapsae* and *Beauveria bassiana*: Implications for enhanced biological control. *Biological Control*, 66(1), 41–48. <https://doi.org/10.1016/j.biocontrol.2013.03.005>
- Shapiro-Ilan, D.I.; Hiltbold, I. and Lewis, E.E. (2018): Ecology of invertebrate pathogens: nematodes, pp. 415–440. In: "Ecology of invertebrate diseases"(Hajek, A.E. and Shapiro-Ilan, D.I. eds.). John Wiley and Sons, Hoboken, N.J.
- Sharma, P.; Mohan, L.; Kumar, K.D. and Srivastava, C.N. (2011): Status of carbohydrate, protein and lipid profile in the mosquito larvae treated with certain phytoextracts.

- Asian Pacific Journal of Tropic Medicine*, 4(4): 301-304. Doi: 10.1016/S1995-7645(11)60090-4.
- Shaurub, E.H. (2023): Review of entomopathogenic fungi and nematodes as biological control agents of tephritid fruit flies: current status and a future vision. *Entomologia Experimentalis et Applicata*, 171:17–34. Doi: 10.1111/eea.13244
- Shaurub, E.H.; Abd El-Meguid, A. and Abd El-Aziz, N.M. (2014): Quantitative and ultrastructural changes in the haemocytes of *Spodoptera littoralis* (Boisd.) treated individually or in combination with *Spodoptera littoralis* multicapsid nucleopolyhedrovirus (SpliMNPV) and azadirachtin. *Micron*, 65: 62–68. Doi:10.1016/j.micron.2014.04.010
- Shaurub, E.H.; Soliman, N.A.; Hashem, A.G. and Abdel-Rahman, A.M. (2015): Infectivity of four entomopathogenic nematodes in relation to environmental factors and their effects on the biochemistry of the Medfly *Ceratitis capitata* (Wied.) (Diptera: Tephritidae). *Neotropical Entomology*, 44: 610–618. <https://doi.org/10.1007/s13744-015-0332-3>.
- Shaurub, E.H.; Zohdy, N.Z.; Abdel-Aal, A.E. and Emara, S.A. (2018): Effect of chlorfluazuron and flufenoxuron on development and reproductive performance of the black cutworm, *Agrotis ipsilon* (Hufnagel) (Lepidoptera: Noctuidae). *Invertebrate Reproduction & Development*, 62(1): 27–34. <https://doi.org/10.1080/07924259.2017.1384407>
- Shaurub, E.H.; Reyad, N.F. and Mohamed, A.A. (2020): Pathogen-mediated modulation of host metabolism and trophic interactions in *Spodoptera littoralis* larvae. *Entomologia Experimentalis et Applicata*, 168: 956–966. Doi: 10.1111/eea.12998
- Sheehan, D.; Meade, G.; Foley, V.M. and Dowd, C.A. (2001) Structure, function and evolution of glutathione transferases: implications for classification of non-mammalian members of an ancient enzyme superfamily. *Biochemistry Journal*, 360:1–16. Doi:10.1042/0264-6021:3600001
- Shelby, K.S. and Popham, H.J.R. (2006): Plasma phenoloxidase of the larval tobacco budworm, *Heliothis virescens*, is virucidal. *Journal of Insect Science*, 6: 13: 1-12. https://doi.org/10.1673/2006_06_13.1
- Shen, S.K., and Dowd, P.F. (1991): 1-Naphthyle acetate esterase activity from cultures of the symbiont yeast of the cigarette beetle (Coleoptera: Anobiidae). *Journal of Economic Entomology*, 84(2): 402-407. <https://doi.org/10.1093/jee/84.2.402>.
- Shukla, E.; Thorat, L.J.; Nath, B.B. and Gaikwad, S.M. (2015): Insect trehalase: physiological significance and potential applications. *Glycobiology*, 25: 357–367. Doi: 10.1093/glycob/cwu125.
- Sikandar, A.; Yuan, R.H.; Lian, X.L.; Zhen, M.A.; Zhao, P.; Li, F.; Liu, X.P. and Wang, Y.Y. (2021): Entomopathogenic nematodes as bioinsecticides - A review. *Applied Ecology and Environmental Research*, 19(3):2459-2476. Doi: http://dx.doi.org/10.15666/aeer/1903_24592476
- Sivakumarm, S. and Maya, Z.B. (1991): Electrophoretic characterization of esterase in the greenbug, *Schizaphis germanium*. *Journal of the Kansas Entomological Society*, 64: 357-362.
- Slansky Jr F. and Scriber, J.M. (1985): Food consumption and utilisation. In: "Comprehensive insect physiology, biochemistry and pharmacology" (Kerkut, G.A. and Gilbert, L.I., eds). vol. 4. pp. 87-163. Pergamon, Oxford.
- Söderhäll, K. and Cerenius, L. (1998): Role of the prophenoloxidase-activating system in invertebrate immunity. *Current Opinion in Immunology*, 10:23-28. Doi: 10.1016/S0952-7915(98)80026-5.

- Soliman, N.A. (2002): Pathological and biochemical effects of some entomopathogenic nematodes on Mediterranean fruit fly, *Ceratitis capitata* (Wiedemann) (Diptera: Tephritidae). PhD. Thesis, Cairo University, Giza, Egypt, 114pp.
- Soreq, H. and Seidman, S. (2001): Acetylcholinesterase-new roles for an old actor. *Nature Reviews Neurosciences*, 2: 294-302. <https://doi.org/10.1038/35067589>
- Srinivas, R.; Udikeri, S.S.; Jayalakshmi, S.K. and Sreeramulu, K. (2004): Identification of factors responsible for insecticide resistance in *Helicoverpa armigera*. *Comparative Biochemistry and Physiology, C Toxicological Pharmacology*, 137: 261-269. Doi: 10.1242/dev.01005.
- Steyn, V.M.; Malan, A.P. and Addison, P. (2021): Efficacy of entomopathogens against *Thaumotibialeucotreta* under laboratory conditions. *Entomologia Experimentalis et Applicata*, 169: 449–461. <https://doi.org/10.1111/eea.13044>
- Strand, M.R. (2008): The insect cellular immune response. *Insect Science*, 15: 1-14. <https://doi.org/10.1111/j.1744-7917.2008.00183.x>
- Suarez, R.K.; Darveau, C.A. and Welch, K.C. (2005): Energy metabolism in orchid bee flight muscles: carbohydrate fuels all. *Journal of Experimental Biology*, 208: 3573–3579. Doi: 10.1242/jeb.01775.
- Subbanna, A.R.N.S.; Rajasekhara, H.; Stanley, J.; Mishra, K.K. and Pattanayak, A. (2018): Pesticidal prospectives of chitinolytic bacteria in agricultural pest management. *Soil Biology & Biochemistry*, 116:52–66. <https://doi.org/10.1016/J.SOILBIO.2017.09.019>.
- Sugumaran, M. (2000): Insect Melanogenesis. *Archives of Biochemistry and Biophysics*, 378(2): 393-403.
- Sugumaran, M. (2002): Comparative biochemistry of eumelanogenesis and the protective roles of phenoloxidase and melanin in insects. *Pigment Cell Research*, 15:2–9. Doi:10.1034/j.1600-0749.2002.00056.x
- Sugumaran, M. (2010): Chemistry of cuticular sclerotization. *Advances of Insect Physiology*, 39: 151-209. <https://doi.org/10.1016/B978-0-12-381387-9.00005-1>
- Sugumaran, M.; Nellaiappan, K. and Valivittan, K. (2000): A new mechanism for the control of phenoloxidase activity: inhibition and complex formation with quinoneisomerase. *Archives of Biochemistry and Biophysics*, 379: 252-260. <https://doi.org/10.1006/abbi.2000.1884>
- Sujatha, P.C.K. and Jeyasankar, A. (2018): Entomopathogenic nematode as biocontrol agent-recent trends- a review. *International Journal of Entomology and Nematology Research*, 2(1): 10-24. Doi: 10.22192/ijarbs
- Sunanda, B.S.; Jeyakumar, P. and Jacob, V.V. (2014): Bioefficacy of different formulations of entomopathogenic nematode *Steinernema carpocapsae* against diamondback moth (*Plutellaxylostella*) infesting cabbage (*Brassica oleracea* var. *capitata*). *Journal of Biopesticides*, 7: 210–215.
- Takeda, M. (2008): Current research of pest insects of vegetables in last decade. *Annual Report of the Kansai Plant Protection*, 50: 39-44.
- Taşkın, A.D. and Aksoylar, M.Y. (2011): Total lipid and total fatty acid percentages of pre-adult stages and adults of *Itopectis melanocephala* (Gravenhorst 1829) (Hymenoptera: Ichneumonidae). *Turkish Entomology Journal*, 35(4): 641-649.
- Taysse, L.; Chambras, C.; Marionnet, D.; Bosgiraud, C. and Deschaux, P. (1998): Basal level and induction of cytochrome P450, EROD, UDPGT, and GST activities in carp (*Cyprinus carpio*) immune organs (spleen and head kidney). *Bulletin of Environmental Contamination and Toxicology*, 60: 300–305. Doi: 10.1007/s001289900625.

- Tefera, T. and Pringle, K.L. (2003): Food consumption by *Chilo partellus* (Lepidoptera: Pyralidae) larvae infected with *Beauveria bassiana* and *Metarhizium anisopliae* and effects of feeding natural versus artificial diets on mortality and mycosis. *Journal of Invertebrate Pathology*, 84: 220–225. Doi: 10.1016/j.jip.2003.11.001.
- Terra, W.R. and Ferreira, C. (2005): Biochemistry of digestion. In: "Comprehensive molecular insect science"(Gilbert, L.I.; Iatrou, K. and Gill, S.S, ed.s), vol. 3. San Diego, California, USA: Elsevier. Pp: 171–224.
- Tiryaki, D. and Temur, C. (2010): The fate of pesticide in the environment. *Journal of Biological and Environmental Sciences*, 4(10): 29-32.
- Toledo, A.V.; Remeslenicov, A.M. and Lopez-Lastra, C.C. (2010): Histopathology cause by the entomopathogenic fungi, *Bauveria bassiana* and *Metarhizium anisopliae*, in the adult planthopper, *Peregrinus maidis*, a maize virus vector. *Journal of Insect Science*, 10: 1-10. <https://doi.org/10.1673/031.010.3501>
- Toubarro, D.; Lucena-Robles, M.; Nascimento, G.; Costa, G.; Montiel, R.; Coelho, A.V. and Simões, N. (2009): An apoptosis-inducing serine protease secreted by the entomopathogenic nematode *Steinernema carpocapsae*. *International Journal of Parasitology*, 39:1319-1330. Doi: 10.1016/j.ijpara.2009.04.013.
- Toubarro, D.; Martinez, Avila, M.; Montiel, R. and Simoes, N. (2013): A pathogenic nematode targets recognition proteins to avoid insect defenses. *PLoS ONE*, 8: e75691. Doi:10.1371/journal.pone.0075691
- Ullah, I.; Khan, A.L.; Ali, L.; Khan, A.R.; Waqaset, M.; Lee, I. and Shin, J. (2014): An insecticidal compound produced by an insect-pathogenic bacterium suppresses host defenses through phenoloxidase. *Molecules*, 19: 20913–20928. Doi:10.3390/molecules191220913.
- Vachon, V.; Laprade, R. and Schwartz, J. (2012): Current models of mode of action of *Bacillus thuringiensis* insecticidal crystal proteins: A critical review. *Journal of Invertebrate Pathology*, 111: 1-12. Doi: 10.1016/j.jip.2012.05.001.
- Vallet-Gely, I.; Lemaitre, B. and Boccard, F. (2008): Bacterial strategies to overcome insect defences. *Nature Reviews Microbiology*, 6: 302–313. Doi: 10.1038/nrmicro1870.
- Van Damme, V.M.; Beck, B.K.E.G.; Berckmoes, E.; Moerkens, R.; Wittemans, L.; De Vis, R.; Nyuttens, D.; Casteels, H.F.; Maes, M.; Tirry, L. and De Clerq, P. (2016): Efficacy of entomopathogenic nematodes against larvae of *Tuta absoluta* in the laboratory. *Pest Management Science*, 72: 1702-1709. Doi: 10.1002/ps.4195
- Vashisth, S.; Chandel, Y.S. and Sharma, P.K. (2013): Entomopathogenic nematodes - a review. *Agricultural Reviews*, 34(3): 163-175. Doi: 10.5958/j.0976-0741.34.3.001
- Vattikonda, S.R. and Sangam, S.R. (2017): Effect of forskolin on the growth and differentiation of the ovary of *Papilio demoleus* L. (Lepidoptera: Papilionidae). *International Research Journal of Environmental Science*, 6: 13-17.
- Veres, P.R.; Neuman, A.; Bertram, T.H. and Ryerson, T.B. (2020): Global airborne sampling reveals a previously unobserved dimethyl sulfide oxidation mechanism in the marine atmosphere. *Proceedings of the National Academy of Sciences (PNAS), USA*, 117(9): 4505-4510. <https://doi.org/10.1073/pnas.1919344117>
- Vicente-Díez, I.; Blanco-Pérez, R.; Chelkha, M.; Puellas, M.; Pou, A.; Campos-Herrera, R. (2021): Exploring the use of entomopathogenic nematodes and the natural products derived from their symbiotic bacteria to control the grapevine moth, *Lobesia botrana* (Lepidoptera: Tortricidae). *Insects*, 12, 1033. 14 pp. <https://doi.org/10.3390/insects12111033>
- Vidhya, D.; Rajiv, P. and Padmanabhan, N (2016): Impact of entomopathogenic fungal infection on the detoxifying enzyme in cotton leafworm *Spodoptera litura* (Fabricius).

- International Journal of Pharmaceutical and Biological Sciences*, 7: 943–948. Doi: 10.22376/ijpbs.2016.7.4.b943-948
- Visetson, S. and Milne, M. (2001): Effects of root extract from derris (*Derris elliptica* Benth) on mortality and detoxification enzyme level in diamond back moth larvae (*Plutella xylostella* Linn). *Agriculture and Natural Resources*, 35: 157–163.
- Viteri, D.M.; Linares, A.M. and Flores, M. (2018): Use of the entomopathogenic nematode *Steinernema carpocapsae* in combination with low-toxicity insecticides to control fall armyworm (Lepidoptera: Noctuidae) larvae. *Florida Entomologist*, 101: 327–329. <https://doi.org/10.1653/024.101.0228>
- von Brando, T. (1973): "Biochemistry of parasites" 2nd ed. Academic Press, New York.
- Waldbauer, G.P. (1968): The consumption and utilization of food by insects. *Advances in Insect Physiology*, 5: 229–288. [https://doi.org/10.1016/S0065-2806\(08\)60230-1](https://doi.org/10.1016/S0065-2806(08)60230-1)
- Walkowiak, K.; Spochacz, M. and Rosinski, G. (2015): Peptidomimetics- A new class of bioinsecticides. *Postępy Biologii Komórki*, 42(2): 235–254.
- Walter, N.T.; Dunphy, G.B. and Mandato, C.A. (2008): *Steinernema carpocapsae* DD136: metabolites limit the non-self-adhesion responses of hemocytes of two lepidopteran larvae, *Galleria mellonella* (F. Pyralidae) and *Malacosoma disstria* (F. Lasiocampidae). *Experimental Parasitology*, 120: 161–174. Doi: 10.1016/j.exppara.2008.07.001.
- Wang, W. and Zhang, X. (2008): Comparison of antiviral efficiency of immune responses in shrimp. *Fish and Shellfish Immunology*, 25:522–527. Doi: 10.1016/j.fsi.2008.07.016.
- Wang, Y.; Oberley, L.W. and Murhammer, D.W. (2001): Antioxidant defense systems of two lepidopteran insect cell lines. *Free Radical Biology and Medicine*, 30: 1254–1262. Doi:10.1016/S0891-5849(01)00520-2
- Wang, J.J.; Cheng, W.X.; Ding, W. and Zhao, Z.M. (2004): The effect of insecticide dichlorvos on esterase activity extracted from the psocids, *Liposcelis bostrychophila* and *L. entomophila*. *Journal of Insect Science*, 4(23): 1–5. Doi: 10.1093/jis/4.1.23
- Wang, Q.Y.; Nangong, Z.Y.; Yang, J.; Song, P. Wang, Y.; Cui, L. and Cui, L. (2012): Toxic activity of a protein complex purified from *Xenorhabdus nematophila* HB310 to *Plutella xylostella* larvae. *Insect Science*, 19(3): 329–336. <https://doi.org/10.1111/j.1744-7917.2011.01472.x>.
- Wee, K.E.; Yonan, C.R. and Chang, F.N. (2000): A new broad spectrum protease inhibitor from the entomopathogenic bacterium *Photobacterium luminescens*. *Microbiology*, 146: 3141–3147. Doi: 10.1099/00221287-146-12-3141.
- Wei, Z.; Yin, Y.; Zhang, B.; Wang, Z.; Peng, G.; Cao, Y. and Xia, Y. (2007): Cloning of a novel protease required for the molting of *Locusta migratoria manilensis*. *Development Growth and Differentiation*, 49: 611–621. Doi:10.1111/j.1440-169X.2007.00957.x
- Welch, H.E. and Bronskill, J.F. (1962): Parasitism of mosquito larvae by the nematode, DD136 (Nematoda: Neoaplectanidae). *Canadian Journal of Zoology*, 40: 1263–1268. <http://dx.doi.org/10.1139/z62-102>.
- Wilson, J.K.; Ruiz, L. and Davidowitz, G. (2019): Dietary protein and carbohydrates affect immune function and performance in a specialist herbivore insect (*Manduca sexta*). *Physiological and Biochemical Zoology*, 92: 58–70. Doi: 10.1086/701196.
- Wouts, W.M. (1984): Nematode parasites of lepidopterans. In: "Plant and Insect Nematodes" (Nickle, W.R., ed.), pp.: 655–696. Marcel Dekker, New York, NY, USA
- Wu, H. and Liu, Q. (2012): Antioxidative responses in *Galleria mellonella* larvae infected with the entomopathogenic nematode *Heterorhabditis* sp. *beicherriana*. *Biocontrol Science and Technology*, 22: 601–606. Doi:10.1080/09583157.2012.670803

- Wu, Q.J.; Zhang, Y.J.; Xu, B.Y. and Zhang, W.J. (2011): The defending enzymes in abamectin resistant *Plutellaxylostella*. *Chinese Journal of Applied Entomology*, 48(2): 291–295.
- Wu, H.; Liu, Q.; Li, X.; Wang, Y. and Zhang, H. (2013): Activities of four enzymes in *Galleria mellonella* larvae infected with entomopathogenic nematode *Heterorhabditisbeicherrianan* sp. *African Journal of Agricultural Research*, 8:3245–3250. Doi: 10.5897/AJAR11.743
- Wu, W.D.; Sun, H.Y.; Xi, J.H.; Yin, J.; Zhang, S.; Cao, Y.Z.; Li, K.B. and Xiao, C. (2015): Observations of the ultrastructure of the fat body and midgut tissues of two white grub species, *Holotrichia parallela* and *H. oblita* (Coleoptera: Melolonthidae), infected by entomopathogenic nematode *Heterorhabditisbacteriophora*. *Acta Entomologica Sinica*, 58:836–845. <https://doi.org/10.16380/j.kcxb>
- Xia, Y.; Dean, P.; Judge, A.J.; Gillespie, J.P.; Clarkson, J.M. and Charnley, A.K. (2000): Acid phosphatases in the haemolymph of the desert locust, *Schistocerca gregaria*, infected with the entomopathogenic fungus *Metarhizium anisopliae*. *Journal of Insect Physiology*, 46: 1249–1257. Doi: 10.1016/S0022-1910(00)00045-7.
- Xia, Y.; Clarkson, J.M. and Charnley, K.A. (2001): Acid phosphatases of *Metarhizium anisopliae* during infection of the tobacco hornworm *Manduca sexta*. *Archives of Microbiology*, 176: 427–434. Doi: 10.1007/s002030100342.
- Xue, C.B.; Luo, W.C.; Chen, Q.X.; Wang, Q. and Ke, L.N. (2006): Enzymatic properties of phenoloxidase from *Pieris rapae* (Lepidoptera) larvae. *Insect Science*, 13: 251–256. <https://doi.org/10.1111/j.1744-7917.2006.00091.x>
- Yağcı, M.; A. Özdem, F.D. Erdoğan, and E. Ayan, (2021a): Efficiency of entomopathogenic nematodes (Rhabditida: Heterorhabditidae and Steinernematidae) on the codling moth (*Cydia pomonella* L.) (Lepidoptera: Tortricidae) under controlled conditions. *Egyptian Journal of Biological Pest Control*, 31:75, 5pp. <https://doi.org/10.1186/s41938-021-00399-1>
- Yağcı, M.; Fırat, T.A.; Erdoğan, F.D. and Şahin, M. (2021b): Virulence of four entomopathogenic nematode against diferent stages of the Mediterranean fruit fy, *Ceratitis capitata* Wiedemann (Diptera: Tephritidae). *Egyptian Journal of Biological Pest Control*, 31:126, 5pp. <https://doi.org/10.1186/s41938-021-00472-9>
- Yang, Y.; Sharma, R.; Zimniak, P. and Awasthi, Y.C. (2002): Role of Alpha class glutathione S-transferases as antioxidant enzymes in rodent tissues. *Toxicological and Applied Pharmacology*, 182:105–115. Doi: 10.1006/taap.2002.9450.
- Yang, J.; Zeng, H.M.; Lin, H.F.; Yang, X.F.; Liu, Z.; Guo, L.H.; Yuan, J.J. and Qiu, D.W. (2012): An insecticidal protein from *Xenorhabdusbudapestensis* that results in prophenoloxidase activation in the wax moth, *Galleria mellonella*. *Journal of Invertebrate Pathology*, 110(1): 60–67. Doi: 10.1016/j.jip.2012. 02.006.
- Yao, W.; Liu, K.; Liu, H.; Jiang, Y.; Wang, R.; Wang, W. and Wang, T. (2021): A valuable product of microbial cell factories: Microbial lipase. *Frontiers in Microbiology*, 12:743377. https://doi.org/10.3389/FMICB.2021.743377/BIBTEX_448
- Yazdani, B.; Nikbakht, A.; Etemadi, N.A. (2014): Physiological effects of different combinations of humic and fulvic acid on gerbera. *Communications in Soil Science and Plant Analysis*, 45: 1357–1368. <http://dx.doi.org/10.1080/00103624.2013.875200>
- Yu, S.J. (2006): Insensitivity of acetylcholinesterase in a field strain of the fall armyworm, *Spodoptera frugiperda* (J.E. Smith). *Pesticide Biochemistry and Physiology*, 84(2): 135–142. <https://doi.org/10.1016/j.pestbp.2005.06.003>
- Yu, X.Q. and Kanost, M.R. (2004): Immulectin-2, a pattern recognition receptor that stimulates hemocyte encapsulation and melanization in the tobacco hornworm,

- Manduca sexta*. *Developmental and Comparative Immunology*, 28: 891–900. Doi: 10.1016/j.dci.2004.02.005.
- YuDong, W.; Chun, X.; Jiao, Y.; YaZhong, C. and KeBin, L. (2012): Evaluation of the impacts on infection ability of entomopathogens nematodes to Grub, *Holotrichia parallela* for three chemical pesticides. [Chinese]. *Journal of Biological Control*, 28(1): 67–73.
- Yüksel, E.; Imren, M.; Özdemir, E.; Bozbuğa, R. and Canhilal, R. (2022): Insecticidal effect of entomopathogenic nematodes and the cell-free supernatants from their symbiotic bacteria against different larval instars of *Agrotis segetum* (Denis & Schiffermüller) (Lepidoptera: Noctuidae). *Egyptian Journal of Biological Pest Control*, 32:54 <https://doi.org/10.1186/s41938-022-00555-1>
- Zhang, B.; Helen, H.S.; Wang, J. and Liu, H. (2011) Performance and enzyme activity of beet armyworm *Spodoptera exigua* (Hübner) (Lepidoptera: Noctuidae) under various nutritional conditions. *Agricultural Sciences in China*, 10:737–746. Doi:10.1016/S1671-2927(11)60057-6
- Zhao, X.; Smart, C.T.; Li, J. and Christensen, B.M. (2001): *Aedes aegypti* peroxidase gene characterization and developmental expression. *Insect Biochemistry and Molecular Biology*, 31: 481-90. Doi: 10.1016/s0965-1748(00)00155-7.
- Zhao, P.; Jiajing, L.; Yang, W. and Jiang, H. (2007): Broad-spectrum antimicrobial activity of the reactive compounds generated *in vitro* by *Manduca sexta* phenoloxidase. *Insect Biochemistry and Molecular Biology*, 37:952–959. Doi: 10.1016/j.ibmb.2007.05.001.
- Zheng, Y.Z.; Lan, W.S.; Qiao, C.L.; Mulchandani, A. and Chen, W. (2007): Decontamination of vegetables sprayed with organophosphate pesticides by organophosphorus hydrolase and carboxylesterase (BI). *Applied Biochemistry and Biotechnology*, 136(3): 233-242. Doi: 10.1007/s12010-007-9022-x.
- Zhou, G. and Miesfeld, R.L. (2009): Energy metabolism during diapause in *Culex pipiens* mosquitoes. *Journal of Insect Physiology*, 55: 40-46. Doi: 10.1016/j.jinsphys.2008.10.002.
- Zhou, C.; Yang, H.; Wang, Z.; Long, G.Y. and Jin, D.C. (2019): Protective and detoxifying enzyme activity and ABCG subfamily gene expression in *Sogatella furcifera* under insecticide stress. *Frontiers in Physiology*, 9:1890. <https://doi.org/10.3389/fphys.2018.01890>
- Zhu, Q.; He, Y.; Yao, J.; Liu, Y.; Tao, L. and Huang, Q. (2012): Effects of sublethal concentrations of the chitin synthesis inhibitor, hexaflumuron, on the development and hemolymph physiology of the cutworm, *Spodoptera litura*. *Journal of Insect Science*, 12(27): 1-13. Doi: 10.1673/031.012.2701.
- Zhu-Salzman, K. and Zeng, R.S. (2015): Insect response to plant defensive protease inhibitors. *Annual Review of Entomology*, 60:233-252. <https://doi.org/10.1146/annurev-ento-010814-020816>
- Zibae, A. (2011): Botanical insecticides and their effects on insect biochemistry and immunity. In: "Pesticides in the Modern World-Pests Control and Pesticides Exposure and Toxicity Assessment", (Stoytcheva, M., ed.). Chapter 4: 55-68., InTechOpen, Croatia.
- Zibae, A. (2012): Digestive enzymes of large cabbage white butterfly, *Pieris brassicae* L. (Lepidoptera: Pieridae) from developmental and site of activity perspectives. *Italian Journal of Zoology*, 79:1, 13-26, Doi: 10.1080/11250003.2011.607190
- Zibae, A.; Bandani, A.R. and Ramzi, S. (2008): Lipase and invertase activities in midgut and salivary glands of *Chilo suppressalis* (Walker) (Lepidoptera, Pyralidae), rice striped stem borer. *Invertebrate Survival Journal*, 5:180–189.

- Zibae, A.; Bandani, A.R. and Ramzi, S. (2008b): Characterization of α -amylase in the midgut and the salivary glands of rice striped stem borer, *Chilo suppressalis* Walker (Lepidoptera: Pyralidae). *Journal of Asia Pacific Entomology*, 11: 201-205. <https://doi.org/10.1016/j.aspen.2008.09.003>
- Zibae, A.; Zibae, I. and Sendi, J.J. (2011): A juvenile hormone analog, pyriproxyfen, affects some biochemical components in the hemolymph and fat bodies of *Eurygaster integriceps* Puton (Hemiptera: Scutelleridae). *Pesticide Biochemistry and Physiology*, 100(3): 289-298. Doi: 10.1016/j.pestbp.2011.05.002
- Żółtowska, K. (2004): Content of saccharides and activity of alpha-glycosidases in *Galleria mellonella* larvae infected with entomopathogenic nematodes *Heterorhabditis zealandica*. *Wiadomości parazytologiczne*, 50(3):495-501.
- Żółtowska, K. and Łopieńska E. (2006): Content of glycogen and trehalose and activity of alpha-amylase and trehalase in *Galleria mellonella* larvae infected with entomopathogenic nematodes *Steinernema affinis* and *S. feltiae*. *Wiadomości parazytologiczne*, 52(2):103-7.
- Żółtowska, K.; Grochla, P. and Łopieńska-Biernat, E. (2006): Activity of superoxide dismutase in *Galleria mellonella* larvae infected with entomopathogenic nematodes *Steinernema affinis* and *S. feltiae*. *Wiad Parazitology*, 4:283–286.