

RESEARCH ARTICLE

BALANCE TRAINING WITH CORONARY ARTERY BYPASS GRAFT SURGERY PATIENTS

MINI REVIEW

Aliaa Abd El Maged Ahmed¹, Heba Ali Abd ElGhafar Ali ¹, Hosam Fathy Ali², Hady Atef

Labib Mohamed¹

¹Department of Physical Therapy for Cardiovascular /Respiratory Disorder and geriatrics,
Faculty of Physical Therapy, Cairo University, Egypt

²Cardiothoracic Surgery Department Faculty of Medicine Cairo University

Corresponding Author:

Aliaa Abd El Maged Ahmed, Faculty of Physical Therapy, Cairo University, Address: Egypt,
Cairo. Telephone: 00201552279700.Email: aliaamaged99@gmail.com

ABSTRACT

The surgical operation known as a "cabbage" or Coronary Artery Bypass Graft (CABG) is performed to treat coronary artery disease (CAD). In order to bypass a blocked or partially obstructed artery in the heart, a healthy blood vessel is transplanted during this surgery. By increasing blood flow to the heart muscle, the new channel lessens symptoms like breathlessness, chest pain, disturbed balance and inability to walk. However, there are numerous side effects from surgery, including postural instability and risk of fall. which can eventually lead to decrease quality of life and functional capacity and increase length of hospital stay. It has been demonstrated that cardiac rehabilitation and balance exercise interventions may change length of hospital stay in patients after CABG. Regular exercise has been reported to improve these patients' balance, functional capacity, length of hospital stays, and quality of life. So, balance exercises and cardiac rehabilitation exercises are used to improve postural balance and functional capacity in post CABG patients.

Keywords: Balance exercise, cardiac rehabilitation, CABG, CAD.

INTRODUCTION

Atherosclerosis, a process when plaques or atheroma obstruct coronary arteries, causes cardiovascular illnesses. [1]

The goal of coronary bypass grafting surgery (CABG) is to supply an appropriate coronary artery (≥ 1.5 mm in diameter) with increased blood flow to ischemic heart disease-affected areas. [2]

The preferred course of treatment for multivessel coronary artery disease (CAD), particularly in individuals with diabetes or ischemic heart failure, is coronary artery bypass graft (CABG) surgery [3].

Over half of the 9 million deaths a year are attributable to atherosclerotic cardiovascular disease, which is in turn caused by environmental exposures. [4]

claimed Using the support and coordinated action of muscle groups, postural control is the act of maintaining, obtaining, or restoring a state of equilibrium throughout any posture or activity [5].

Impaired balance can lead to a dread of falling, which in turn lowers quality of life and makes it harder to carry out everyday tasks. [6]

Balance exercise improves balance, cognitive functions, as well as quality of life, balance and coordination exercise may result in greater improvement in the quality of life. [7].

Both physical performance and balance, which are critical components in preventing falls, can benefit from balance training. [8]

Enhanced functional mobility, less discomfort, better cognitive function, increased cardiovascular endurance, and overall functional independence were the goals of physical therapy treatment. [9]

For individuals with coronary artery disease, cardiac rehabilitation is an effective non-pharmacological strategy that can enhance their general health and cardiopulmonary fitness. Following a coronary bypass graft, patients can restore their voluntary physiological, psychological, vocational, and educational status with the use of planned cardiac rehabilitation. When compared to regular medical care, cardiac rehabilitation plays a significant role in reducing mortality rates by 26% [10].

Healthcare professionals should be aware of the significance of cardiac rehabilitation and should use it more regularly [11].

DISCUSSION

The balance training enhances walking ability (step length, gait speed, cadence, and double-limb support period) as well as increase balance berg scale (BBS) ($p = 0.001$). [12]

the sit-to-stand (STS) test has been widely employed across a wide range of chronic conditions, and its properties have been investigated. [13].

The SF-36 questionnaire is also a commonly used generic quality of life instrument and includes 36 questions over 8 domains. the physical and mental domains and transforming these into summary t scores ranging from 0 to 100 with higher scores denoting higher quality of life [14].

early rehabilitation beginning within 3 days of CABG prescribed by physicians or therapists was significantly associated with improved ADLs at discharge after the CABG [15]

post-CABG balance deficits is physical deconditioning, which can last for several weeks. Furthermore, cardiac surgery can worsen pulmonary function, and respiratory

muscle dysfunction and postoperative pulmonary complications may, in turn, influence functional capacity and balance. Additionally, postsurgical disability has been shown to be associated with cardiac disease. Therefore, this disability may also contribute to poor balance control. A diminished ability to maintain balance is associated with an increased risk of falling, bone fracture, joint dislocation, concussion, and even death [16]. So early balance training can affect the functional capacity, quality of life and length of hospital stay.

CONCLUSION it was concluded that early rehabilitation after CABG surgery via balance exercises could have beneficial effect on improving postural balance, patient's functional capacity which in turn reflect to better quality of life. Balance exercises should be viewed as a safe, non-invasive therapeutic option which could be implemented as a routine practice along with traditional cardiac rehabilitation physiotherapy in CABG patients.

REFERENCES

- 1- Basil, K. U. M., Orélien Sylvain MTOPI BOPDA, C. H. U. Franklin, Ferdinand TINGOM, and Samuel Honoré MANDENGUE. "The prevalence and associated risk factors of work-related cardiovascular disorders among healthcare workers in five reference hospitals in the city of Douala, Cameroon." *GSC Advanced Research and Reviews* 14, no. 1 (2023): 036-046.
- 2- Belyayev, L., Stock, E. M., Hattler, B., Bakaeen, F. G., Kinlay, S., Quin, J. A., ... & Zenati, M. A. (2024). Complete Coronary Revascularization and Outcomes in Patients Who Underwent Coronary Artery Bypass Grafting: Insights from The REGROUP Trial. *The American Journal of Cardiology*.
- 3- Pölzl, L., M. Thielmann, S. Cymorek, F. Nägele, J. Hirsch, M. Graber, N. Sappeler et al. "Impact of myocardial injury after coronary artery bypass grafting on long-term prognosis." *European Heart Journal* 43, no. 25 (2022): 2407-2417.
- 4- Deo, Salil V., Yakov Elgudin, Issam Motairek, Frederick Ho, Robert D. Brook, Jason Su, Stephen Fremes et al. "Air pollution and adverse cardiovascular events after coronary artery bypass grafting: a 10-year nationwide study." *JACC: Advances* 3, no. 2 (2024): 100781.
- 5- Patel, Jagruti K., Vandana G. Patel, Neha R. Desai, Neha R. Patel, Mohammed S. Mukkhi, and Palak K. Ribadiya. "Effect of 4 Weeks Balance Training Program in Healthy Young

- Adults: A Randomized Clinical Trial Study." *Indian Journal of Physiotherapy & Occupational Therapy* Print-(ISSN 0973-5666) and Electronic-(ISSN 0973-5674) 14, no. 2 (2020): 74-79.
- 6- Cha, Eunsil, and Jung Hyun Park. "Spinopelvic alignment as a risk factor for poor balance function in low back pain patients." *Global Spine Journal* 13, no. 8 (2023): 2193-2200.
- 7- Dunskey, A. (2019). The effect of balance and coordination exercises on quality of life in older adults: a mini-review. *Frontiers in aging neuroscience*, 11, 318.
- 8- Phu, Steven, Sara Vogrin, Ahmed Al Saedi, and Gustavo Duque. "Balance training using virtual reality improves balance and physical performance in older adults at high risk of falls." *Clinical interventions in aging* (2019): 1567-1577.
- 9- Babchanik, Svyatoslav. "Physical rehabilitation of a patient status post coronary artery bypass graft surgery with intensive care unit delirium in an acute care setting." PhD diss., California State University, Sacramento, 2020.
- 10- Persis Selvamani, S. "Effect of Supervised Moderate Intensity Exercise with Forced Expiratory Technique in Improving Endurance Capacity and Quality of Life in Subjects with Coronary Artery Bypass Graft." PhD diss., KG College of Physiotherapy, Coimbatore, 2019.
- 11- Osailan, A., & Abdelbasset, W. K. (2020). Exercise-based cardiac rehabilitation for postcoronary artery bypass grafting and its effect on hemodynamic responses and functional capacity evaluated using the Incremental Shuttle Walking Test: A retrospective pilot analysis. *Journal of the Saudi Heart Association*, 32(1), 25.
- 12- Noh, Hyeon-Jeong, Soon-Hyun Lee, and Dae-Hyouk Bang. "Three-dimensional balance training using visual feedback on balance and walking ability in subacute stroke patients: A single-blinded randomized controlled pilot trial." *Journal of Stroke and Cerebrovascular Diseases* 28, no. 4 (2019): 994-1000.
- 13- O'Grady, H. K., Edbrooke, L., Farley, C., Berney, S., Denehy, L., Puthucherry, Z., & Kho, M. E.

- (2022). The sit-to-stand test as a patient-centered functional outcome for critical care research: a pooled analysis of five international rehabilitation studies. *Critical Care*, 26(1), 175.
- 14- Butterly, Elaine W., Peter Hanlon, Anoop SV Shah, Laurie J. Hannigan, Emma McIntosh, Jim Lewsey, Sarah H. Wild et al. "Comorbidity and health-related quality of life in people with a chronic medical condition in randomised clinical trials: An individual participant data meta-analysis." *PLoS medicine* 20, no. 1 (2023): e1004154.
- 15- Ohbe, H., Nakamura, K., Uda, K., Matsui, H., & Yasunaga, H. (2021). Effect of early rehabilitation on physical function in patients undergoing coronary artery bypass grafting: A nationwide inpatient database study. *Journal of clinical medicine*, 10(4), 618.
- 16- Abolahrari-Shirazi, Sara, Farahnaz Emami, Zahra Rojhani-Shirazi, Mansour Janati, and Mersa Razeghi. "The Effects of Coronary Artery Bypass Grafting on Postural Balance." *Journal of Rehabilitation Sciences & Research* 9, no. 1 (2022): 42-46.
- 1)