


EGYPTIAN ACADEMIC JOURNAL OF **BIOLOGICAL SCIENCES** BOTANY



ISSN 2090-3812

www.eajbs.com

Vol. 16 No.2 (2025)



Effects of Sources of Phosphorus and Watering Regimes on the Growth of African Star Apple (*Chrysophyllum albidum* G. Don) Seedlings

Olusegun D. Adelani

Forestry and Environmental Technology, Federal College of Forestry Ibadan, P.M.B 5087, Dugbe, Ibadan, Nigeria

*E. Mail: adelani.olusegun@yahoo.com

ARTICLE INFO

Article History

Received: 1/8/2025

Accepted: 11/10/2025

Available: 15/10/2025

Keywords:

Phosphorus,
Moisture
application
interval,
Plantlets, Soil
amendment,
Enrichers.

ABSTRACT

There is an insufficiency of examined and corroborated findings on the outcome of phosphorus-based enrichers on the improvement of *Chrysophyllum albidum*. To advance the time-consuming development of *Chrysophyllum albidum*, studied was accomplished. A 2 x 3 split-plot experimental arrangement following three imitations was approved to establish the influences of fountains of phosphorus and watering periods on the advancement of *Chrysophyllum albidum* plantlets. Two origins of nutrient specifically; poultry manure (25g) and single superphosphate (10g) accomplished the main plot investigation. The daily watering (200ml), 3-day interval (200ml) and 5 days interval (200ml) established sub plot investigation. The investigation encompassed two bases of phosphorus and three moisture application intervals reciprocated thrice. Twelve (12) plantlets denoted a copy. Two hundred and sixteen (216) plantlets were explored. A month-old *C. albidum* plantlets were transferred into pots with enhancers and exposed to 200ml at different day's meantime. The assembled data were subjected to two-way Analysis of Variance (ANOVA) at a 5% significance level. The outcomes disclosed that bases of phosphorus and watering stretches tremendously ($p < 0.05$) improved the growth of *C. albidum* plantlets. A distinguished height (41.41cm), girth (4.03cm), leaf number (16.18), collar diameter (2.02cm), chlorophyll a (0.6092 mg/g), total chlorophyll a+b (0.3753mg/g) and phosphorus uptake (2.02%) were obtained from plantlets developed in the soil modified with single super phosphate and watered (200ml) at 3 days interval relative to control experiment. Improvement of soil accompanying single super phosphate and application of 3 days watering intervening time enriches the growth of *C. albidum* plantlets.

INTRODUCTION

The tropical forests are sources of useful timber and non-timber species essential to the well-being of the local people (Areo, 2018; Iroko *et al.*, 2020). Nadrowski *et al.* (2010) stated that forests perform critical roles in maintaining and providing ecosystem services and functions. Africa is privileged to have forests and forest resources of ecological and socio-economic potentials. Unfortunately, the reality of the situation is that these resources are being milked in an unsustainable manner and consequently vanish at appalling rates, mostly due to human activities (Adeyoju, 2001). Lamb *et al.* (2005) stated that the great direct and indirect benefits of forest assets are threatened owing to combined influences of clear felling, forest fragmentation and degradation.

The yearly clear cutting in Nigeria is 3.5%, which is nearly 350,000–400,000 hectares per year (FAO, 2005; Lapido, 2010). In a report by IITA (2011), Nigeria was established as the country accompanying the worst deforestation rate in the world, which can be accredited to her escalating population and associated leading poverty rate. Rotowa *et al.* (2020) disclosed that the high rate of deforestation and a fall in afforestation of the species has led to yearly decline in formation and successful management of forest. Thus, result to incessant erosion and decrease of natural gene pool of significant forest tree species that are critical for the survival of the current and forthcoming generation (Iroko *et al.*, 2020) as African star apple in Nigeria (Adelani *et al.*, 2023).

African star apple or White star apple, *Chrysophyllum albidum* synonymns of *Gambeya albida* (Adelani *et al.*, 2022) is a climax tree species of tropical rainforest that is a member of the family Sapotaceae (Olaoluwa *et al.*, 2012; Wole, 2013) that produces accurately half of the order accompanying 800 species (Ehiagbonare *et al.*, 2008). It is named “Osan Agbalumo,” “Udara” or “Udala” and “Agwaluma or Agwaluba” in Yoruba, Igbo and Hausa accents individually (Rahaman, 2012; Wole, 2013; Adelani *et al.*, 2018). It is in the company of forest tree species incorporated in the long-established agroforestry systems (Ureigho and Ekeke, 2010; Laurent *et al.*, 2012), that furnishes NTFPs of enormous household significance to rural and urban dwellers in West Africa accompanying excellent export prospects (Nwoboshi, 2000). The *C. albidum* has been recognized as having considerable economic (Onyekwelu *et al.*, 2011), nutritional and medicinal (Onyekwelu and Stimm, 2011) and industrial (Olaoluwa *et al.*, 2012) values.

Comprehensive human activities exhaust the populace of unregenerated *C. albidum* (Onyekwelu *et al.*, 2011) during the permanent transformation of forest regions to other uses (Adelani *et al.*, 2020). Adelani *et al.* (2014) mentioned that despite the significance of *C. albidum*, it has been considerably ignored specifically regarding its regeneration. This requires intentional measures to save the genetic erosion of this significant species. It is appropriately comprehended that the disappearance of diversification possibly reduced in a maintainable approach, by multiplying the quantity of varieties adopted in productive forestry through breeding of native timber varieties (Hoque *et al.*, 2004a) and fruit tree types as *C. albidum*.

Adelani *et al.* (2023) published that the time-consuming growth of *C. albidum* and low soil fertility that connect to clear felling as well as degeneration are difficulties to its reproduction and planned management. Fertilizer applications are fundamental to address these challenges. Upgraded soil fertility through the administration of fertilizers is an indispensable determinant in plant growth (Brady and Weil, 2004). Fertilizer employment is widely established in nurseries to boost plant vigor and productivity (Shen *et al.*, 2010). Moreover, fertilizer enrichment can upgrade plant growth (King *et al.*, 2006) by either intensifying soil resources or by embellishing the ability of seedlings to accumulate materials (Lincoln *et al.*, 2007) and by improving soil pH (Jose *et al.*, 2003).

In the last few decades, the implementation of enrichers in forest nurseries has enticed accelerating awareness all over the globe on the account of escalated request for fiber, wood (Kauter *et al.*, 2003), CO₂ offsets as well as foods. Excellent growth of the plants through application of fertilizer assists to surmount food insecurity of teeming populace growth. Sarfraz, *et al.* (2023) established that the estimated universal population of 9.9 billion by 2050 warrants an over 70% surge in food creation. Consequently, the effective employment of fertilizers to embellish cultivation of *Chrysophyllum albidum* to meet population demand of its abundant potentiality is extremely crucial. The types of fertilizer to use are also vital.

Farmers employ synthetic fertilizers independently for adequating or providing nutrients that are low in supply (Jafer and Gebresilasie, 2018). Purbajanti *et al.* (2019) announced that farmers use synthetic fertilizers due to the fact that they are more reasonable,

economical, accessible, users friendly and swift in reciprocation; while natural enricher accomplishes gradual reaction on plant yield, even though they are capable of sustaining soil attributes. It is well established that administration of synthetic plant foods strengthened the seedling and substantial growth in biomass of many tree categories (Bungard *et al.*, 2000; Ong *et al.*, 2004; Hoque *et al.*, 2004b; Uddin *et al.*, 2007); while the natural soil amendment advances the material and synthetic attributes of the soils (Ojelabi *et al.*, 2018); enhancing soil water holding capacity and operations of soil organism (Mbah and Mbagwu, 2003); promising long time remaining effects of nutrient (Makinde and Ayoola, 2008; Ogunade *et al.*, 2020) and as well as entices low cost (Bayu *et al.*, 2006).

Assessment of the effect of natural and synthetic origins of phosphorus on plants is crucial because phosphorus is restricted in availability in tropical soil. Thompson *et al.* (2019) stated that tropical forest productivity is limited by soil phosphorus (P) accessibility. Richardson and Simpson (2011) expressed that its obtainability for plants is primarily controlled by mineral weathering, organic matter decomposition, microbial activity, and interactions accompanying soil particles. In highly weathered tropical soils, P is frequently bound to iron and aluminium oxides, making it less approachable to forests (Dalia *et al.*, 2024). The inefficiency and unavailability of phosphorus can be traced to their properties accompanying only 15–25% being taken up by plants; while the remaining amount is leached, causing soil deterioration and water pollution directly (Conley *et al.*, 2009; Johnston *et al.*, 2014). Paz-Ares *et al.* (2022) substantiated that phosphorus is detected in abundance in the lithosphere, but the form used by plants, inorganic orthophosphate (Pi), is indissoluble and distributes gradually in soils. This leads to P inadequacy in farming land, other ecological communities (Fahad *et al.*, 2023) and marginal land pushed to forestry practices in tropics.

Tropical forestlands are always identified by small soil phosphorus (P) obtainability, indicating that P limits plant achievement (Laura *et al.*, 2023). Nutritional studies entailing phosphorus accessibility principally disclosed it limits plant advancement (Epstein and Bloom, 2004; Campo and Vazquez-Yanes, 2004; Sardans *et al.*, 2004; Villar-Salvador *et al.*, 2004; Condit *et al.*, 2013). Various bioassay investigations on seedling growth have corroborated that phosphorus is the chief limiting nutrient in tropical forest soils (Hinsinger, 2001; Lawrence, 2001; Sardans *et al.*, 2004; Villar-salvador *et al.*, 2004; Dean *et al.*, 2009; Wan Juliana *et al.*, 2009; Nilus *et al.*, 2011; Thompson *et al.*, 2019).

Phosphorus (P) is a vital macronutrient that engages in a pivotal role in the growth and development of plants (Fahad *et al.*, 2023). However, Fahad *et al.* (2023) asserted that the insufficient accessibility of phosphorus in soil presents a meaningful difficulty for crop yield. Khan *et al.* (2023) affirmed that phosphorus is pivotal for the growth of plants. Nevertheless, the difficult attainability of P in soil forms the foremost problem to challenge for crop output, especially after plants suffer from drought. A significant section of soils universal shorts of phosphorus (P), leading to a serious restriction of crop yields and give rise to a monumental threat to universal food guarantee (Heuer *et al.*, 2017). Forest crops also need rudimental soil nutrients to furnish food security. The productivity and stability of forest ecosystems are fundamentally governed by nutrient obtainability particularly phosphorus (P) (Shi *et al.*, 2016; Wang *et al.*, 2022). Dalia *et al.* (2024) emphasized that, while P is a limiting nutrient in forest ecosystems, its fertilization upgrades biomass productivity, thereby donating to higher carbon sequestration in international forests.

Low P availability in forest soils is necessitating plants to acquire extra P from diverse origins for the plant community to function (Olander and Vitousek, 2004; Condit *et al.*, 2013). Lambers *et al.* (2018) avowed that when P is incorporated into forest soils, plants often demonstrate various responses, such as modifications in morphology, phosphatase enzyme production, and embellished interactions accompanying mycorrhizal fungi. Reciprocations of plants to phosphorus nutrient availability in the soil result in

vigorous growth. P is vital for ATP synthesis, nucleic acid formation, and membrane function straight forwardly modifying plant growth, photosynthesis (Ewuketu and Wenxing, 2005), carbon assimilation, and ribulose-1,5-biphosphate regeneration in the Calvin cycle (Vance *et al.*, 2003; Vitousek *et al.*, 2010). Phosphorus is vital nourishment for plant growth and productivity (Dalia *et al.*, 2024).

Salami *et al.* (2020) expressed that application of phosphorus fertilizers is more important practice in nursery formation. The utilization of P-fertilizer remarkably magnified the seedling enhancement of chosen agroforestry tree species in nursery (Mohammad *et al.*, 2007). Siddhuraju (2007) opined that throughout the primary stages of maturation, trees are very well dependent on soil nutrients provide. Negligence to handle nursery soil adequately by way of not furnishing it with the essential nutrient can lead to exhaustion of site attribute and a decline of seedling growth (Hoque *et al.*, 2004b). Soil nutrient (P) is significant for a healthy seedling growth in the nursery. Not only to supply *C. albidum* seedlings with essential nutrient (P) through fertilizer application is vital, but also to expose it to suitable watering interval that embellishes its maturation for standard planting stock (Adelani *et al.*, 2023).

Inadequacy of soil moisture can result in nutritional shortages, even within the soil provided with fertilizer (da Silva *et al.*, 2011). Homyak *et al.*, (2017) opined that water shortage has an effect on the movement and disappearance of both nitrogen and phosphorus nutrients. Water is indispensable for germination, dividing cells and expansion, metabolic activities, and other functions (da Silva *et al.*, 2013). Adelani (2019) established that of all the determinants influencing nutrient uptake, growth and development of the plant, water is most crucial. Regardless of the quality and quantity of nutrients existing in the soil, only water helps the nutrient to dissolve and form ions for the assimilation of the root for plant growth and development. The significance of water cannot be over overstated as it facilitates in biochemical, physiological and hormonal process in the plants (Adelani, 2019).

Growth and biomass productivity is straightway commensurate to furnish and use water in plants (Cao, 2000; Olajuyigbe *et al.*, 2013). Water is an influential determinant in the growth, development and productivity of plants (Gbadamosi, 2014; Ogidan *et al.*, 2018). Yisau *et al.* (2020) observed that productivity of vigorous seedlings within good time frame is of essentials in silviculture as well as forestry in general. Standstill in seedling growth or subsequent mortality as a result of lack of water converts into financial forfeiture to a nursery operator (Mhango *et al.*, 2008). Aderounmu *et al.* (2017) cited that water has major impacts on growth of plants, whereas watering needs of different species vary. Even with variations in species watering preference, fertilizer and water complement each other for plant growth and productivity.

Adequate fertilizer utilization and watering interval-gap are paramount determinants and outstanding components that regulate production and growth of quality young plants (Adelani *et al.*, 2023). Dalia *et al.* (2024) noted that both nutrients and water are two of the most critical components influencing tree growth, and they collaborate with each other. The production of healthy seedlings necessitates the proper recognition of specific limiting factors (Fox, 2000). There is a scarcity of assessed findings on the moisture and nutrient specification (Okafor, 2003; Adelani *et al.*, 2014) for majority of tree species as *Chrysophyllum albidum*. Little work is obtainable on the effect of sources of phosphorus and moistening regimes on the growth of *C. albidum* seedlings. In this light, the fact finding was channelled on the effect of sources of phosphorus and moisture application regimes on the enhancement of *C. albidum* seedlings.

MATERIALS AND METHODS

Description of Experimental Site:

The observational trial location was at the forest nursery of the Federal University of Agriculture Abeokuta. It is sited along Alabata Road, North-East of Abeokuta. It is positioned within latitudes 7° N and 7° 55 ' N and longitudes 3° 20 'E and 3° 37 ' E. The Federal University of Agriculture Abeokuta is established within the rain forest zone of South Western Nigeria (Amujoyegbe *et al.*, 2008). It is next to Ogun-Osun River Basin Development Authority (OORBDA), along Osiele-Abeokuta Road, off Abeokuta-Ibadan Road. It is in the North Eastern end of Abeokuta and lies almost on latitude 7° 30 ' N and longitude 3° 54 ' E. It lies within the humid lowland rain forest region accompanying two distinctive seasons. The wet season extends from March to October; while the dry season extends from November to February (Aiboni, 2001). The rainfall has a characteristic bimodal distribution accompanying peaks in July and September and breaks in August. Generally, the rainfall could be heavy and erosive sometimes accompanied by lightning and thunderstorm at the beginning and end of rainy season.

Seedling Growth:

Chrysophyllum albidum seedlings were raised from seeds for the study. Established seedlings of *C. albidum* of regular measurement at 2-4 leaf periods after two weeks of watering were introduced into 0.75 litre polypot that accommodated sand washed accompanying acid.

Effects of Sources of Phosphorus and Watering Regimes on the Growth of *C. albidum* Seedlings:

The effect of origins of phosphorus and watering intervals on the growth of *Chrysophyllum albidum* plantlets was studied. A 2x3 split-plot experimental representation along with three duplications was selected to evaluate the influences of beginnings of phosphorus and watering interval gaps on the growth of *C. albidum* plantlets. Two sources of nutrient particularly; poultry manure (25g) and single superphosphate (10g) formed the main plot investigation. The daily watering (200ml), 3 days interval (200ml) and 5 days interval (200ml) made up sub plot investigation. The investigation incorporated two inceptions of phosphorus and three watering breaks repeated thrice. Twelve (12) plantlets expressed a clone. Two hundred and sixteen seedlings were engaged. Seedling under each base of phosphorus was split into 3 groups and exposed to different regimes of moisture application. A month-old *C. albidum* plantlets were moved into pots with boosters and exposed to 200ml at varying day's interval.

Growth constants were taken fortnightly for 12 months. Growth variants studied include seedling height accompanying the use of meter rule., girth accompanying the use of veneer caliper., collar diameter was calculated by meter rule., the number of leaves were enumerated manually, and Leaf area was attained by linear measurement of leaf length and leaf width as presented by Clifton-Brown and Lewandowski (2000).

$$LA = 0.74 \times L \times W \quad (1)$$

Where LA, Leaf area= Product of linear dimension of the length and width at the broadest part of the leaf.

The average of the growth criterion for duration of examination was used for categorization. Total fresh weight, total dry weight, relative turgidity, net assimilation rate, absolute growth rate, relative growth rate, chlorophyll a, b and a+b and phosphorus uptake were investigated during and after 48 weeks.

Formula for AGR, RGR

$$\text{Absolute Growth Rate} = \frac{W_2 - W_1}{T_2 - T_1} \text{ gwk}^{-1} \quad (2)$$

W_2 and W_1 are plant weight at corresponding time T_1 and T_2 .

RGR=LAR*NAR

$$\text{Relative Growth Rate} = \frac{\log_e W_2 - \log_e W_1}{T_2 - T_1} g g^{-1} w k^{-1} \quad (3)$$

T1=Initial time (Weeks)

T2= Final or second time (Weeks)

W₂= Total dry weight at T₂

W₁=Total dry weight at T₁

$$\text{NAR} = \frac{w_2 - w_1}{t_2 - t_1} \times \frac{\log_e A_2 - \log_e A_1}{A_2 - A_1} g c m^{-2} w k^{-1} \quad (4)$$

Where W_2 and W_1 are plant dry weights at times t_1 and t_2 , $\log_e A_2$ and $\log_e A_1$ are the natural logs of leaf areas A_1 and A_2 at times t_1 and t_2 .

Relative turgidity, which is synonymous to relative water content, was detected by interval between fresh weight and dry weight divided by turgid weight subtracting dry weight multiply by 100. Turgid weight was found out by weight of the plant before and after soaking in water for 24hours. Plantlet of high vigour was detached into three parts particularly: leaf, stem and root. Each part was weighed (fresh weight, FW), then left soaked in distilled water for twenty-four (24hrs) (under normal room light and temperature) (Turgid Weight) and after hydration, the samples were then dehydrated in an oven at 70°C for 48 hours and weighed (DW) (after being cool down in desiccators).

The RWC/RT is determined as follows:

$$\text{Relative turgidity} = \frac{FW - DW}{TW - DW} \times 100 \quad (5)$$

FW= Fresh weight

DW= Dry weight

TW= Turgid weight

Evaluations of Chlorophyll were established by straight forward calculations of the absorbance at different wavelengths, using Model 6405 uv/vis Spectrophotometer, serial number 1364. The concentrations were computed by accumulating 20.2 A₆₄₅, 8.02 A₆₆₃ and divided by length of light path in cell (usually 1cm), fresh weight in grams and is 1000. The outcome was multiplied by the volume of chlorophyll solution in mL. A₆₄₅ and A₆₆₃ is the absorbance at 645 and 663 nm.

$$\text{Chlorophyll a (mg g}^{-1}\text{)} = \frac{12.7(A_{663}) - 2.69(A_{645}) \times VC / LLP \times FW \times 1000}{12.7(A_{663}) - 2.69(A_{645}) \times VC} \quad (6)$$

$$\text{Chlorophyll b (mg g}^{-1}\text{)} = \frac{22.9(A_{645}) - 4.86(A_{663}) \times VC / LLP \times FW \times 1000}{22.9(A_{645}) - 4.86(A_{663}) \times VC} \quad (7)$$

$$\text{Total Chlorophyll a+b (mg g}^{-1}\text{)} = \frac{20.2 A_{645} + 8.02 A_{663} \times VC}{LLP \times FW \times 1000} \quad (8)$$

Where:

A= absorbance at the given wavelength

C.C= Concentration of chlorophylls

VC = Volume of chlorophyll in mL

LLP = Length of light path usually 1cm

FW = Fresh Weight in grams

Tissue analyses were assessed for the whole plant of *C. albidum* seedlings to calculate the nutrient uptake. Membrane analysis was also carried out for the specimen of entire plant of *C. albidum* plantlets prior to establishing in the beginning and after establishing at end of the investigation respectively to ascertain nutrient uptake. Phosphorus content was detected by Bray-1 method.

Nutrient Uptake was evaluated by Method of

$$\text{Sharma et al. (2012)} = \frac{\%N \times \%P \times \%K \times \text{Dry matter kg ha}^{-1}}{100} \quad (9)$$

Effects of Sources of Phosphorus and Watering Regimes on the Growth of African Star Apple

Actual nutrient uptake was ascertained by variations in nutrient uptake at the beginning and the end of the experiment

Data Analysis

Data were amassed and put through analysis of variance (ANOVA) using SAS (2003). An analyzing of outstanding means was achieved using Fishers' Least Difference LSD at 5% level of significance.

RESULTS

Prominent height (29.83cm), leaf area (145.3cm²), total fresh weight (8.01g), total dry weight (3.65g), relative turgidity (58.58%) and higher absolute growth rate (0.397gwk⁻¹) were gotten from seedlings sown in the soil modified following poultry manure. Notable girth (3.11cm), leaf number (14.60), collar diameter (1.56cm) and relative growth rate (0.465gg⁻¹wk⁻¹) were traced from seedlings established in the soil improved accompanying single superphosphate. The lower values of height (29.45cm), leaf area (107.1cm²); total fresh weight (5.90g), total dry weight (2.85g), relative turgidity (56.73%) and absolute growth rate (0.366gwk⁻¹) were reported from seedlings put up in the soil fertilized with single superphosphate fertilizer. The smaller value of 1.04cm (girth), 12.10 (leaf number), 0.52cm (collar girth) and 0.405gg⁻¹wk⁻¹ (relative growth rate) were gotten from seedlings established in poultry manure.

Remarkable height (39.97cm), girth (2.48cm), collar diameter (1.24cm) and total dry weight (3.32g) were reported from seedlings implanted in the soil watered at 3days interval. Outstanding leaf number (14.53) and relative growth rate (0.494gg⁻¹wk⁻¹) were produced from seedlings watered daily. Meaningful leaf area (156.7cm²), total dry weight (3.46g) and relative turgidity (58.61%) were derived from seedlings watered at 5 days interval. A prominent height (39.97cm), girth (2.48cm), collar diameter (1.24cm) and total dry weight (3.32g) were written from seedlings watered (200ml) at 3 day's interval.

The least value of height (18.41cm), leaf number (11.67), relative growth rate (0.403gg⁻¹wk⁻¹) were put down from seedlings watered at 5days interval. The least value of 1.69cm (girth), 0.85cm (collar diameter), 6.51g (total fresh weight), 2.56g (total dry weight), 56.76% (relative turgidity) and 0.336gwk⁻¹ (absolute growth rate) were documented in seedlings watered daily (Table 1).

Table 1: Effects of Sources of Phosphorus and Watering Regimes on the Growth of *C. albidum* Seedlings

S.O.P	H(cm)	G(cm)	LN	LA(cm ²)	CD(cm)	TFW(g)	TDW(g)	RT(%)	NARgcm ⁻² wk ⁻¹	AGR(gwk ⁻¹)	RGR(gg ⁻¹ wk ⁻¹)
POM	29.83 ^a	1.04 ^b	12.10 ^b	145.1 ^a	0.52 ^b	8.01 ^a	3.65 ^a	58.58 ^a	0.003 ^a	0.397 ^a	0.405 ^b
SSP	29.45 ^a	3.11 ^a	14.60 ^a	107.1 ^b	1.56 ^a	5.90 ^b	2.85 ^a	56.73 ^a	0.003 ^a	0.366 ^a	0.465 ^a
SE±	0.52	0.07	0.70	4.85	0.05	0.54	0.38	4.93	0.001	0.07	0.02
WR											
1	30.55 ^b	1.69 ^c	14.53 ^a	126.69 ^b	0.85 ^c	6.51 ^a	2.56 ^b	56.76 ^a	0.003 ^a	0.336 ^a	0.494 ^a
3	39.97 ^a	2.48 ^a	13.86 ^a	95.07 ^c	1.24 ^a	6.96 ^a	3.32 ^a	57.59 ^a	0.003 ^a	0.336 ^a	0.408 ^b
5	18.41 ^c	2.06 ^a	11.67 ^b	156.7 ^a	1.03 ^b	7.39 ^a	3.46 ^a	58.61 ^a	0.002 ^a	0.449 ^a	0.403 ^b
SE±	0.64	0.11	0.86	5.94	0.06	0.35	0.27	6.04	0.001	0.09	0.03

Means on the same column having different superscript are significantly different (p<0.05).

S.O.P= Source of phosphorus, WR= Watering regime, H= Height, G=Girth, LN= Leaf Number, LA= Leaf area, CD=Collar diameter, TFW=Total Fresh Weight, TDW=Total Dry Weight, R.T= Relative Turgidity, NAR= Net Assimilation Rate, Absolute Growth Rate = AGR, RGR= Relative Growth Rate.

Significant height (41.41cm), girth (4.03cm), leaf number (16.18) and collar diameter (2.02cm) were entered from plantlets sown in the soil corrected accompanying single super phosphate and watered at 3 days interval. Superior leaf area (210.76cm²) was acquired from seedlings established in the soil enriched with poultry manure and watered at 5days interval. Excellent absolute growth rate (0.584gwk⁻¹) and relative growth rate (0.540gg⁻¹wk⁻¹) were

obtained from seedlings placed in the soil upgraded with single super phosphate and watered at 5 and 1 days interval respectively.

The lowest values of height (15.07cm), girth (0.71cm), leaf number (10.50), collar diameter (0.36cm), net assimilation rate ($0.002\text{gcm}^{-2}\text{wk}^{-1}$) and relative growth rate ($0.337\text{gg}^{-1}\text{wk}^{-1}$) were noted for seedlings cultivated in poultry manure and watered at 5 days interval. The minimum value of 85.34cm^2 (leaf area) was traced from seedlings put in soil helped accompanying poultry manure and watered at 3 days interval. The meanest values of 5.13g (total fresh weight) and 2.15g (total dry weight) were produced from seedlings transplanted in the soil boosted accompanying single superphosphate and watered daily. The minimal value of 48.92% (relative turgidity); $0.002\text{gcm}^{-2}\text{wk}^{-1}$ (net assimilation rate) and 0.214gwk^{-1} (absolute growth rate) were recounted from seedlings introduced into the soil amended with single superphosphate and watered at 3 days interval (Table 2).

Table 2: Interactive Effect of Sources of Phosphorus and Watering Regimes on the Growth of *C. albidum* Seedlings

S.O.P	WR	H(cm)	G(cm)	LN	LA(cm^2)	CD(cm)	TFW(g)	TDW(g)	RT (%)	NAR ($\text{gcm}^{-2}\text{wk}^{-1}$)	AGR(gwk^{-1})	RGR($\text{gg}^{-1}\text{wk}^{-1}$)
P.O.M	1	35.89 ^b	1.48 ^d	14.26 ^{ab}	139.37 ^b	0.74 ^d	7.90 ^{ab}	2.98 ^{ab}	56.82 ^a	0.003 ^b	0.418 ^{ab}	0.448 ^{ab}
	3	38.52 ^{ab}	0.94 ^a	11.54 ^b	85.34 ^d	0.47 ^e	8.18 ^a	3.60 ^{ab}	66.26 ^a	0.005 ^a	0.459 ^{ab}	0.431 ^{ab}
	5	15.07 ^e	0.71 ^a	10.50 ^c	210.76 ^a	0.36 ^a	7.95 ^{ab}	4.38 ^a	52.67 ^a	0.002 ^b	0.314 ^{ab}	0.337 ^b
SSP	1	25.20 ^c	1.90 ^c	14.80 ^{ab}	114.01 ^c	0.95 ^c	5.13 ^b	2.15 ^b	56.71 ^a	0.003 ^b	0.300 ^{ab}	0.540 ^a
	3	41.41 ^a	4.03 ^a	16.18 ^a	104.79 ^{cd}	2.02 ^a	5.75 ^b	3.85 ^{ab}	48.92 ^a	0.002 ^b	0.214 ^b	0.386 ^b
	5	21.75 ^d	3.41 ^b	12.8 ^b	102.71 ^b	1.71 ^b	6.83 ^{ab}	2.55 ^{ab}	64.56 ^a	0.003 ^b	0.584 ^a	0.469 ^{ab}
SE±		1.20	0.15	0.86	8.40	0.06	0.94	0.82	8.55	0.001	0.12	0.05

Means on the same column having different superscript are significantly different ($p < 0.05$)

S.O.P= Source of phosphorus, WR= Watering regime, H= Height, G=Girth, LN= Leaf Number, LA= Leaf area, CD=Collar diameter, TFW=Total Fresh Weight, TDW=Total Dry Weight, R.T= Relative Turgidity, NAR= Net Assimilation Rate, Absolute Growth Rate = AGR, RGR= Relative Growth Rate.

Significant values of 0.6092 mg/g (chlorophyll a), 0.3753mg/g (total chlorophyll a+b) and 2.02% (phosphorus uptake) were written from stems and leaves of seedlings initiated in the soil developed with single superphosphate and watered (200ml) at 3 days interval. Substantial values of 0.2386mg/g (chlorophyll b) and 2.02% (phosphorus uptake) were entered from leaves of plantlets instituted in the soil controlled with poultry manure and watered at 5days interval.

Table 3: Interactive Effect of Sources of Phosphorus and Watering Regimes on the Chlorophyll and Phosphorus Uptake

S.O.P	WR	P.P	Cl (a)mg/g	Cl (b)mg/g	Cl (a+b) (mg/g)	P Uptake (%)
POM	1	Leaf	0.2147 ^{bc}	0.1468 ^c	0.3616 ^{ab}	1.18 ^{de}
	1	Stem	0.0948 ^d	0.0769 ^d	0.1717 ^b	0.16 ^{de}
	1	Root	0.0283 ^{ef}	0.021 ^{ef}	0.0493 ^d	0.62 ^{cd}
POM	3	Leaf	0.1707 ^c	0.1653 ^b	0.3334 ^b	1.47 ^b
	3	Stem	0.0849 ^d	0.0808 ^d	0.1657 ^{cd}	0.21 ^{de}
	3	Root	0.274 ^b	0.0216 ^{ef}	0.049 ^d	0.53 ^d
POM	5	Leaf	0.1249 ^{cd}	0.2386 ^a	0.3635 ^{ab}	2.02 ^a
	5	Stem	0.0877 ^d	0.0942 ^d	0.1819 ^{bc}	0.20 ^{de}
	5	Root	0.0245 ^f	0.0241 ^e	0.0486 ^d	0.74 ^{cd}
SSP	1	Leaf	0.2355 ^b	0.1288 ^c	0.3642 ^{ab}	0.78 ^{cd}
	1	Stem	0.101 ^d	0.0742 ^d	0.1752 ^c	0.09 ^e
	1	Root	0.0314 ^e	0.0183 ^f	0.0497 ^d	0.32 ^{de}
SSP	3	Leaf	0.2486 ^b	0.1268 ^c	0.3753 ^a	2.02 ^a
	3	Stem	0.6092 ^a	0.0751 ^d	0.1771 ^c	0.13 ^e
	3	Root	0.0301 ^e	0.0183 ^f	0.0484 ^d	0.76 ^{cd}
SSP	5	Leaf	0.1565 ^c	0.1847 ^b	0.3411 ^b	0.96 ^c
	5	Stem	0.0908 ^d	0.0861 ^d	0.1768 ^c	0.18 ^e
	5	Root	0.0302 ^e	0.0221 ^{ef}	0.0523 ^d	0.32 ^b
SE±			0.02	0.01	0.01	0.16

Means on the same column having different superscript are significantly different ($p < 0.05$).

S.O.P= Source of phosphorus, WR= Watering regime, P.P= Plant part, Cl(a) = Chlorophyll a, Cl(b)= Chlorophyll b, Cl(a+b)= Total chlorophyll, P uptake= Phosphorus uptake.

The least value of 0.0245mg/g (chlorophyll a) and 0.0484 (total chlorophyll a+b) were noted down from roots of plantlets originating in poultry manure amended soil and watered at 5days interval and root of those developed in soil modified accompanying single superphosphate and water at 3 days interval respectively. The least value of 0.0183mg/g (Chlorophyll b) was established from root of seedlings set up in soil reinforced with single superphosphate and watered at day 1 and 3 intervals. The least value of 0.09% (P uptake) was reported from stem of seedlings produced in the soil advanced accompanying single superphosphate and watered daily (Table 3).

DISCUSSION

Significant growth constants recorded from *C. albidum* seedlings cultivated in poultry manure revealed its better source of phosphorus. A source of phosphorus motivates the growth of plants. Several investigators have recounted the superior performance of origins of phosphorus in improving the growth of trees (Rickard, 2000; Abedin *et al.*, 2002; Khatik *et al.*, 2004; Liu *et al.*, 2005; Broschat, 2006; Ogunwole *et al.*, 2006; Suthiphasilp, 2009; Balal *et al.*, 2011; Pereira *et al.*, 2014; Larwanou *et al.*, 2014; Anamayi *et al.*, 2016). Dalia *et al.* (2024) discovered that sources of phosphorus (H_3PO_4) meaningfully improved the growth of *Swietenia mahagoni*. Some of agroforestry trees increased appreciably with the application of P fertilizer (Triple Super Phosphate TSP) (Mohammad *et al.*, 2008). Administration of triple superphosphate (77kg P ha⁻¹) on the topsoil of Chittagong University Campus is mandatory to boost the diameter and total dry matter production of *Michelia champaca* seedlings (Hoque *et al.*, 2004b).

Source of phosphorus, organic manure furnished *C. albidum* seedlings with better phosphorus. Phosphorus is a rudimentary element grouped as a macronutrient because of the approximately large amounts of phosphorus mandatory for plants. The greatest role of phosphorus inside a plant is energy store and transport in adenosine di and tri phosphate as well as engages in a principal responsibility in the establishment of DNA and proteins (Abod and Siddiqui, 2002; Taiz and Zeiger, 2002; Epstein and Bloom, 2004; Havlin *et al.*, 2005; Nathan, 2009; Busman *et al.*, 2009; CFF, 2011; Anderson, 2015; Malhotra *et al.*, 2018; Sezer and Seda, 2019; Isidra-Arellano *et al.*, 2021; Dalia *et al.*, 2024). Phosphorus is a paramount component of nucleic acids, phospholipids, high-energy phosphate bond complexes, and many coenzymes (Wyngaard *et al.*, 2016; Fahad *et al.*, 2023; Dalia *et al.*, 2024). Yao *et al.* (2012) observed that phosphorus is inevitable for glucose and nitrogen metabolism, and the mutual conversion of protein and carbohydrate metabolism. It is a supreme constituent of ATP, the chemical that accords energy to the plant for nutrition translocation, nutrient uptake, and respiration (Dalia *et al.*, 2024).

Owing to its role in proteins and nucleotides, adequate phosphorus is connected to improve root growth, strong stem development and appropriate development of reproductive parts (Williamson *et al.*, 2001; Zapata and Zaharah, 2002 ; Haggai *et al.*, 2003, Lopez-Bucio *et al.*, 2003; Epstein and Bloom, 2004; Anghinoni and Bissani, 2004; Jin *et al.*, 2005; Havlin *et al.*, 2005; Hill *et al.*, 2006; Pandey *et al.*, 2006; Alabi, 2006; Huda *et al.*, 2007; Mohammad *et al.*, 2007; Xu *et al.*, 2012; Wu *et al.*, 2015; Waraich *et al.*, 2015; Muhammad *et al.*, 2017; Wen *et al.*, 2017; Lai *et al.*, 2018; Sezer and Seda, 2019 ; Thompson *et al.*, 2019; Dalia *et al.*, 2024). Moreover, various investigators (Ramos *et al.*, 2000; Nicoloso *et al.*, 2001; Schumacher *et al.*, 2004; Gomes *et al.*, 2004; Melo *et al.*, 2005) relate the P effect on the primary growth of forest species. Ramos *et al.* (2000) pronounced a general positive effect of phosphorus from nitrogen, phosphorus and potassium employment in the improvement of *Bauhinia forficata* seedlings. Gomes *et al.* (2004) noticed that *Anadenantheta colubrine* offered favourable reaction to the phosphorus

administration. Nicoloso *et al.* (2001) noted that from 60 to 120 days subsequently, *Apuleia leiocarpa* seedlings emergence, phosphorus fertilization escalated sequentially to seedling height in conditions of nitrogen and potassium inadequacy. Bhuiyan *et al.* (2000) has reported the speed up of growth factors on the administration of P-fertilizer, specifically on *Casuarina* spp. The responsibilities of phosphorus in plants are very basic and indispensable. It assists a plant to convert other nutrients into useful building blocks with which to grow (Heather, 2021). Graciano *et al.* (2006) registered that P addition is accelerating soil N absorption in *Eucalyptus grandis* seedlings.

Phosphorus is vital to flowering, fruiting and to transfer of hereditary (Vance *et al.*, 2003; Epstein and Bloom, 2004; Nathan, 2009; Haque and Khan, 2012; Anderson, 2015; Sezer and Seda, 2019). It also intensifies disease resistance in plants (Anderson, 2015; Dalia *et al.*, 2024). It alleviates the toxic effects of arsenite in the soils (Soriano and Fereres, 2003; Quaghebeur and Rengel 2003; Liu *et al.*, 2005; Tao *et al.*, 2006; William *et al.*, 2007; Pigna *et al.*, 2010). Fahad *et al.* (2023) established that phosphorus comes up as a key player in bolstering plant resilience against unfavourable atmospheric situations. Phosphorus increased reduced soil pH by 0.06 (P+) and 0.14(P++) units equated accompanying the control ($P < 0.01$) (Laura *et al.*, 2023). Fahad *et al.* (2023) observed that phosphorus is imperative for the synthesis and activation of enzymes concern with the nutrient obtaining and transfer and its dearth give rise to nutrient instability and poisonousness signs.

For tree cultivated in soil without phosphorus addition, there was a reduction in biomass production equated to the plants that obtained phosphorus fertilization on Padula, Vertricella and La Bolte soils of Italy respectively (Pigna *et al.*, 2010). Tariq *et al.* (2023) noticed that lacking of P affects the uptake of other nutrients like nitrogen (N), potassium (K) and calcium (Ca) by plants, changing their architecture contributing to things such as diminished stem elongation and the production of shorter and thicker stems. P insufficiency also influences biomass allocation, with a drop in above-ground biomass and a rise in root biomass (Fahad *et al.*, 2023). Overall, P inadequacy not only affects nutrient acquisition and transport, but also stimulates variations in plant morphology, biomass allocation, leaf morphology and photosynthesis, underscoring its significance in diverse physiological processes and overall plant growth and development (Fahad *et al.*, 2023). Jose (2003) also announced that high deprivation of phosphorus and mineral uptake may lead to poor cell differentiation and multiplication of plant. Nevertheless, some species, such as *Arabidopsis thaliana* establishes larger roots in P-deficient conditions (Ma *et al.*, 2001).

Outstanding growth parameters produced from seedlings planted in poultry manure relative to single superphosphate could also be traced to potentials of natural manure in embellishing the growth of seedlings. Organic manures are fundamental in the growth and development of plant because they serve as soil modification (Abou El-Magd *et al.*, 2005; Ojelabi *et al.*, 2018; Adedokun *et al.*, 2020c), binding nutrients vulnerable to leaching (Abou El –Magd *et al.*, 2006), balance of soil pH fluctuation (Eifediyi and Remison, 2010) and reduction of soil pH (Laura *et al.*, 2023). Poultry manure is natural by characteristics. Poultry manure comprises nutrients and essential elements (nitrogen, phosphorus, potassium, calcium, magnesium, sulphur, copper, zinc) that can boost plant productivity and strengthen the physical and chemical attributes of the soil (Telkamp, 2015; Usman *et al.*, 2022). Alabi (2006) declared that poultry manure accommodates enormous contents of all the mineral nutrients essential to the plant. Poultry manure utilization upgrades soil retention and uptake of plant nourishments. This intensifies plant health by increasing water and nutritious accessibility as well as overpowering harmful levels of plant parasitic nematodes, fungi and bacteria (Agbede *et al.*, 2008). Salami *et al.* (2020) announced that P in poultry manure is swiftly hydrolyzed and chemically accelerated or assimilated for plant growth.

The excellent performance of poultry manure in improving the growth of tree seedlings have been reported by Olajire-Ajayi *et al.* (2018) (*Mansonia altissima*) and

Adedokun *et al.* (2020a) (*Tamarindus indica*). Contrary to this finding, Adedokun *et al.* (2020b) recommended 1g of urea fertilizer for upbringing *Blighia sapida* seedlings. Iroko *et al.* (2020) approved 0.09g/pot (141kg/ha) urea plant food employment for successful nurturing of *Pterocarpus erinaceus* seedlings. Salami (2015) discovered that application of 15g urea to soil embellished the healthy growth of *Enterolobium cyclocarpum* seedlings.

Significant growth criterion written from seedlings moistened at 3 days revealed that modest and adequate supply of water is basic. Moderate supply of water that prevents the excesses and insufficiencies varies from one species to another. Quantity of water is adequate when it can successfully improve the growth of tree seedlings better than others. Various researchers such as Isah *et al.* (2012) (*Acacia senegalensis*), Olajuyigbe *et al.* (2013) (*Diospyros mespiliformis*) and Usman *et al.* (2013) (*Moringa oleifera*) published the efficacy of adequate provision of water in upgrading the growth of tree seedlings. Mukhtar (2012) disclosed that plant water level has a powerful impact on plant growth and biomass productivity through its influence on leaf and root expansion. This also authenticates the announcements of earlier investigators who have published highest growth variables in seedlings subjected to adequate watering (Akinyele, 2007; Dauda *et al.*, 2009; Gbadamosi, 2014; Oboho and Igharo, 2017; Ogidan *et al.*, 2018). This consequently implies that growth and biomass output is straight-forwardly proportional to the present and utilize of water (Sale, 2015; Mukhtar, 2016) and it also stresses the importance of instituting maximum water specifications for tree seedlings to enhance growth (Mukhtar *et al.*, 2016).

Only the moderate and adequate supply of water advances the plant growth. Lesser growth variables written from seedlings treated with 5days watering intervals compared to that of 3 days is an indication of low availability of water to seedlings. Inferior water availability damages seedling and sapling growth and survival (Cao, 2000; Engelbrecht and Kursar, 2003). Insufficient water supply is dangerous to the growth of plants. A decrease in water consumption negatively influences plant production (Kang *et al.*, 2024). Scarcity can have a negative effect not only on the morphological features of plant, but also on the physiological, biochemical, and molecular features (Fathi and Tari, 2016). The menace of inadequacy of water accounts for morphological and physiological alterations in higher plants (Ghorbani *et al.*, 2019). On the other hand, reduced growth constant recorded from seedlings watered daily relative to that of 3 days interval revealed the effect of unrestrained supply of water on plant growth. Superabundance supply of water is harmful. In an unrestricted water supply condition, available oxygen in the soil is usually low, thus restricting nutrient assimilation for species not acclimatized to this condition. This is in conformity with the discoveries of Predick *et al.* (2009) who established that growth of flood-uncompromising varieties is often depressed in overabundance of water. Similar observation has been reported by Gbadamosi (2014) who stated that extravagant supply of water had adverse influence on the growth and biomass production of the *Picralima nitida* seedlings. Gonzales *et al.* (2009) reported the unfavorable consequence of surplus watering on the enhancement of *Chenopodium quinoa*.

A significant morphological and physiological growth specifications entered from seedlings embedded in the soil amended with single superphosphate and watered (200ml) at 3 days interval revealed the appropriate collaboration of sources of phosphorus and watering regimes. Appropriate association of fountains of phosphorus and watering stretches enriches the enlargement of tree seedlings. The excellent performance of seedlings implanted in the soil modified following single superphosphate in the presence of adequate watering regime of three days interval could also be attributed to ability of 3 days watering regime to dissolve the sources of phosphorus correctly for plant uptake for growth to take place. This is in consonance with the report of John (2013) who disclosed that when the water is sufficiently supplied to sources of phosphorus; it is effective in supplying the nutrients to the plant for growth. Similar pronouncement has been registered by Adelani (2019) who noticed that

daily watering considerably ($P < 0.05$) embellishes plantlet growth and nutritional uptake (P) of *Citrus tangelo* planted in the combination of sand and pulverized *Jacaranda mimosifolia* leaf litters as source of nitrogen, phosphorus and potassium.

CONCLUSION

Some of our indigenous priority trees species of ample potentials are slow to meet the population request of their benefits. Inadequate water and nutrient supply is also a threat to propagation as well as biodiversity preservation of our native fruit tree types. Investigation on the effects of fountains of phosphorus and watering seasons exposed that the improvement of soil with single super phosphate and application of 3 days watering periods enriches the increase of *C. albidum* seedlings. The essential roles of adequate combinations of fertilizer and watering application to seedlings of agroforestry species as *C. albidum* for afforestation, reforestation, agroforestry systems as well as for biodiversity conservation programmes cannot be over emphasized.

Declarations:

Ethical Approval: This study did not involve any live animals. It was based solely on experimental laboratory and analysis of plant extracts

Author's Contributions: Main Author.

Funding: This research did not receive external funding.

Availability of Data and Materials: All data generated or analyzed during this study are included in this article.

Acknowledgements: Not applicable.

REFERENCES

- Abedin, M. J., Cresser, M.S., Meharg, A. A., Feldmann, J., Cotter-Howells, J. (2002). Arsenic accumulation and metabolism in rice (*Oryza sativa* L). *Environmental Science and Technology*, 36, 962-968. doi: 10.1021/es0101678
- Abod, S.A., Siddiqui, M.T. (2002). Growth response of teak (*Tectona grandis* L.f) seedlings to nitrogen, phosphorus and potassium fertilizers. *Pertanika Journal of Tropical Agricultural Science*, 25(2), 107-113.
- Abou, El-Magd, M.A., Hoda, M., Mohammed, A., Fawzy, Z.F. (2005). Relationship growth and yield of broccoli with increasing N, P or K ratio in a mixture of NPK fertilizers. *Annals of Agricultural Science*, Moshtohor, 43, 791-805.
- Abou, El-Magd., El-Bassiony, M.A., Fawzy, Z.F. (2006). Effect of organic manure with and without chemical fertilizers on growth, yield and quantity of some varieties of Broccoli plant. *Journal of Applied Sciences Research*, 2(10), 791-798.
- Adedokun, S.A., Adelusi, F.T., Agbeja, A.O., Odewale, M.O., Eniola, O. (2020b). Influence of urea application on the growth of *Blighia sapida* K.D. Koenig seedlings. *Forestry Development in Nigeria: Fifty years of Interventions and Advocacy*. In: Ogunsanwo, O.Y., Adewole, N.A., Oni, P.I and Idumah, F (Eds). *Proceedings of the 42nd Annual Conference of the Forestry Association of Nigeria held in Ibadan. 23rd -28th November, 2020*. pp 20-25.
- Adedokun, S.A., Adelusi, F.T., Agbeja, A.O., Odewale, M.O., Eniola, O. (2020c). Comparative study of leaf compost of *Gmelina arborea* (Roxb), *Terminalia catappa* (Linn) and poultry manure on the early growth of *Albizia lebbek* (L.Benth) seedlings. *Forestry Development in Nigeria: Fifty years of Interventions and Advocacy*. In: Ogunsanwo, O.Y., Adewole, N.A., Oni, P.I and Idumah, F (Eds). *Proceedings of the 42nd Annual Conference of the Forestry Association of Nigeria held in Ibadan. 23rd -28th November, 2020*. Pp273-279.

Effects of Sources of Phosphorus and Watering Regimes on the Growth of African Star Apple

- Adedokun, S.A., Adelusi, F.T., Odewale, M.O., Eniola, O., Onifade, A.O. (2020a). Influence of potting mixture of poultry manure and horse dung on the growth of *Tamarindus indica*. *Forestry Development in Nigeria: Fifty years of Interventions and Advocacy*. In: Ogunsanwo, O.Y., Adewole, N.A., Oni, P.I and Idumah, F (Eds). *Proceedings of the 42nd Annual Conference of the Forestry Association of Nigeria held in Ibadan*. 23rd -28th November, 2020. Pp 13-19.
- Adelani, D.O. (2019). Effect of watering regimes on the growth and nutrient uptake of *Citrus tangelo* J. W. seedlings grown in a mixture of sand and pulverized *Jacaranda mimosifolia* D. don leaves. *Journal of Research in Forestry, Wildlife and Environment* 11(3), 172-179 <http://www.ajol.info/index.php/jrfwe> (Nigeria)
- Adelani, D.O. Aduradola, A.M., Suleiman, R.T., Oyelowo, O.J. (2023). Effects of sources of nitrogen and watering regimes on the growth of African star apple (*Chrysophyllum albidum* G. Don) seedlings. *Ethiopian Journal of Environmental Studies and Management*, 16(4), 454-473. doi: <https://ejesm.org/doi/v16i4.5>
- Adelani, D.O., Amos, O., Oyebamiji, N.A. (2020). Effect of selected nitrogen fixing trees leaf litters on the growth of *Chrysophyllum albidum* G. Don: A step to enhance the growth of an endangered species. *Biodiversity Conservation and Current Environmental Challenges*. In: Babalola, F.D., Olaleru, F., Arabomen O and Ringim, A.S. *Proceedings of 7th Biodiversity Conservation Conference, Nigeria Chapter of Society for Conservation Biology (NSCB) and Society for Conservation Biology (SCB)*, 27-29th October, 2020.pp229-234.
- Adelani, D.O., Ogunsanwo, J.A., Awobona, T.A. (2018). Effect of seed weights on the germination and early seedling growth of African star apple (*Chrysophyllum albidum* G. Don). *Nigerian Journal of Forestry*, 48(1), 33-38.
- Adelani, D.O., Suleiman, R.A., Akesode, H.A., Akande, M.T. (2014). Effect of sources and rates of organic fertilizer on the growth of *Chrysophyllum albidum* seedlings. *Organic Agriculture Research: A Catalyst for Sustainable National Agricultural Transformation Agenda*. In: Olabiyi, T.I and Bolarinwa, I.F (Eds). *Proceedings of the 10th National Conference on Organic Agriculture* held at Ladoke Akintola University of Technology, P.M.B 4000, Ogbomoso, Oyo State, Nigeria from 17th - 20th November, 2014. Printed by A- Treasurer Digital Press LTD, Nigeria. Pp 65-73.
- Adelani, D.O., Suleiman, R.T., Oni, B.O., Ogunkalu, A.O., Ariyo, O.C., Mohammed, R., Ogunsanwo, J.A., Ariyo, M.O. (2022). Effect of leaf litters of selected nitrogen-fixing albizia species on the growth of African star apple (*Gambeya albida* G.Don) Aubrev& Pellegr.) seedlings. *Forestry and the Challenges of Insecurity, Climate Change and Covid-19 Pandemic in Nigeria*. In: Oluwadare, A.O., Akinyemi, O.F., Idumah, O., Akintunde-Alo, D.A and Lawal, A. (Eds). *Forestry and the Challenges of Insecurity, Climate Change and Covid -19 Pandemic in Nigeria. Proceedings of 8th Biennial National Conference of the Forests and Forest Products Society*, held at Forestry Research Institute of Nigeria, Ibadan, Nigeria between 14th-20th August, 2022. Pp 225-232.
- Aderounmu, A.F., Adenuga, D.A., Ogidan, O.A., Alonge, O.O. (2017). Effect of different watering regimes on the early growth of *Terminalia superba* ENGL and DIELS. In: Adekunle, V.A.J., Ogunsanwo, O.Y and Akinwale, A.O (Eds). *Harnessing the Uniqueness of Forest for Sustainable Development in a Diversifying Economy. Proceedings of the 39th Annual Conference of the Forestry Association of Nigeria*. Pp183-189.
- Adeyoju, S.K. (2001). Forestry for National Development: A critique of the Nigeria Situation. In: L. Popoola, Abu, J.E and Oni, P.I (Eds). *Proceedings of the 27th*

- Annual Conference of the Forestry Association of Nigeria (FAN)*, Abuja, FCT, Nigeria. Pp55-68.
- Agbede, T.M; Ojeniyi, S.O., Adeyemo, A.J.(2008). Effect of poultry manure on soil physical and chemical properties, growth and grain yield of sorghum in South West, Nigeria. *American-Eurasian Journal of Sustainable Agriculture*, 2, 72-77.
- Aiboni, V.U. (2001). Characteristics and classification of representative location in soil of a topographical University of Agriculture, Abeokuta. *Asset Series A*, 1(1), 51- 61.
- Akinyele, A.O. (2007). Silvicultural requirement of seedlings of *Buchholzia coriacea* Engler. Unpublished Ph.D Thesis. *Dept of Forest Resources Management, University of Ibadan*. Pp 20-34.
- Alabi, D. A. (2006). Effects of fertilizer phosphorus and poultry droppings treatments on growth and nutrient components of pepper (*Capsicum annuum* L). *African Journal of Biotechnology*, 5(8), 671-677. 18 April, 2006 Available online at <http://www.academicjournals.org/AJB>
- Amujoyegbe, B.J., Bamire, A.S., Elemo, K.O. (2008). Agronomic analysis of fertilizer effect on maize/ cowpea intercrop in Ile-Ife and Abeokuta, South-Western Nigeria. *Asset Series, A*, 8(1), 62-72
- Anamayi, S. E., Oladele, O. N., Suleiman, R. A., Oloyede E. O., Yahaya, U. (2016). Effects of cow dung and N.P.K fertilizer at different levels on the growth performance and nutrient composition of *Moringa oleifera*. *Annals of Experimental Biology*, 2016, 4 (1), 35-39.
- Anderson, P. (2015). N.P.K and Fe, what are they good for? *www.Limbwalker Tree Service*, 1(1): 1-4. Accessed on 1/16/2015.
- Anghinodni, I., Bissani, C.A. (2004). Fosforo e . adubos fosfatados. In: Bissani, C.A (Ed). *Fertilidade dos solos e manejo de adubacao de culturas*. V.I. Porto Alegre: *Genesis*, pp 117-137.
- Areo, O.S. (2018). Awareness of forestry as a career in secondary schools in Oyo State, Nigeria (A case study of Ido L.G.A of Oyo State). Nigeria. *Journal of Life Science*, 8(2), 190-198.
- Balal, R.M., Gimeno, V., Shahid, M.A., lidon, V., Garcia,A. L; AAbbas T. Garcia-Sanchez, F., Ghazanfer, U. (2011). Effects of phosphorus fertilization on growth, leaf mineral concentration and xylem-phloem nutrient mobility in two root stock of prunus (*Prunus persica x Prunus amygdalus*) and (*Prunus insititia*) in the Mediterranean area. *Australian Journal of Crop Science*, 5(12), 1542-1549.
- Bayu, W., Rethman, N.F.G., Hammers, P.S., Alemu, G. (2006). Effects of farm yard manure and inorganic fertilizers on sorghum growth, yield and nitrogen use in a semiarid area of Ethiopia. *Journal of Plant Nutrition*, 29(2), 391-407. doi:10. 1080/01904160500320962
- Bhuiyan, M.Z.A., Hossain, M.K., Osman, K.T. (2000). Effect of inorganic fertilizers on the initial growth performance of *Casuarina equisetifolia* seedlings in the nursery. *Indian Journal of Forestry*, 23(3), 296-300.
- Brady, N.C., Weil, R.R. (2004). *The Nature and Properties of Soils*. Prentice Hall, Upper Saddle River, New Jersey, 638-666.
- Broschat, T.K. (2006). Effects of phosphorus and phosphoric acids on growth and phosphorus concentrations in container-growth tropical ornamental plants. *Horticultural Technology*, 16(1), 105-108. doi:10.21273/horttech.16.1.0105
- Bungard, R.A., Press, M.C., Scholes, J.D. (2000). The influence of nitrogen on rain forest Dipterocarp seedlings exposed to large increase irradiance. *Plant Cell and Environment*, 23(11), 1183-1194. <https://doi.org/10.1046/j.1365-3040.2000.00642.x>

- Busman, L., Lamb J., Randall, G., Rehm, G., Schmitt, M. (2009). The nature of phosphorus in soil. *Nutrient Management. Soil Phosphorus in the Agriculture, Regents of Minnesota, Minnesota* Fo-06795-Go. Pp 7.
- California Fertilizer Foundation (CFF) (2011). Plant nutrients-phosphorus. www.calfertilizer.org. pp4. Accessed on 30/09/14.
- Campo, J., Vazquez-Yanes, C. (2004). Effects of nutrient limitation on above ground carbon dynamics during tropical dry forest regeneration in Yucatan, Mexico. *Ecosystems*, 7, 311–319. <https://doi.org/10.1007/s10021-003-0249-2>
- Cao, K. F. (2000). Water relations and gas exchange of tropical saplings during a prolonged drought in a Bornean Health Forest, with reference to root architecture. *Journal of Tropical Ecology*, 16, 101-116. doi: <https://doi.org/10.1017/S0266467400001292>
- Clifton-Brown, J.C., Lewandowski, I. (2000). Water use efficiency and biomass partitioning of three different *Miscanthus* genotypes with limited and unlimited water supply. *Annals of Botany*, 86(1), 191- 200. <https://doi.org/10.1006/anbo.2000.1183>
- Condit, R., Engelbrecht, B. M. J., Pino, D., Pérez, R., Turner, B. L. (2013). “Species distributions in response to individual soil nutrients and seasonal drought across a community of tropical trees.” *Proceedings of the National Academy of Sciences of the United States of America*, 110, 5064–5068.
- Conley, D.J., Paerl, H.W., Howarth, R.W., Boesch, D.F., Seitzinger, S.P., Havens, K.E., Lancelot, C., Likens, G.E. (2009). Ecology-controlling eutrophication: nitrogen and phosphorus. *Science*, 323, 1014–1015. doi: 10.1126/science.1167755
- da Silva, E. C., Nogueira, R. J. M. C., da Silva, M. A., de Albuquerque, M. B. (2011). Drought stress and plant nutrition. *Plant stress*, 5(1), 32-41.
- da Silva, E.C., de Albuquerque, M.B., Azevedo Neto, A.D., da Silva Junior, C.D. (2013). Drought and its consequences to plants-from individual to ecosystem. In book: *Responses of Organisms to Water Stress* (pp.17-47), Edition: First Edition, Chapter 2: Drought and Its Consequences to Plants – From Individual to Ecosystem, In Tech Publisher, Şener Akınc (Eds). <https://doi.org/10.5772/53833> Tech.
- Dalia, A.A., Ahmed, S.E., Ahmed, A.E., Mohamed, M., Mohamed, E. (2024). The effect of fertilizing with different N and P sources on the growth of *Swietenia mahagoni* (L.) Jacq seedlings under water stress. *Delta Journal of Science*, 49(2), 16-38.
- Dauda, T.O., Asiribo, O.E., Akinbode, S.O., Saka, J.O., Salahu, B.F. (2009). An assessment of the roles of irrigation farming in the Millenium Development Goals. *African Journal of Agricultural Research*, 4(5), 445-450. <https://doi.org/10.5897/AJAR.9000037>
- Dean, F. M., Travis, W. I., Friday, J.B., Paul, G.S (2009). Effects of fertilization on phosphorus pools in the volcanic soil of a managed tropical forest. *Forest Ecology and Management*, 258(10), 2199-2206. doi:10.4067/S0718-95162010000200004
- Ehiagbonare, J.E., Onyibe, H.I., Okoegwale, E. (2008). Studies on the isolation of normal and abnormal seedlings of *Chrysophyllum albidum*: a step towards sustainable management of the Taxon in the 21st Century. *Scientific Research and Essay*, 3(12), 567-570. <http://www.academicjournals.org/SRE>
- Eifediyi, E.K., Remison, S.U. (2010). Growth and yield of cucumber (*Cucumis sativum* L) as influenced by farmyard manure and inorganic fertilizer. *Journal of Plant Breeding and Crop Science*, 2(7), 216-220. <http://www.academicjournals.org/jpbcs>
- Engelbrecht, B.M.J., Kursar, T.A. (2003). Comparative drought-resistance of seedlings of 28 species of co-occurring tropical woody plants. *Oecologia*, 136(3), 383-393. doi: 10.1007/s00442-003-1290-8
- Epstein, E ., Bloom, A.J. (2004). *Mineral nutrition of plants: Principles and perspectives* (Second Edition). Sunderland, M A: Sinauer Associates, Inc.; 2004. 402p.

- Ewuketu, L., Wenxing, L. (2005). Phosphorus limitation constrains global forest productivity directly and indirectly via forest community structural attributes: meta-analysis. *Global Ecology and Biogeography*, 34, 1-15, 34:e70048 <https://doi.org/10.1111/geb.70048>,
- Fahad, K., Abu, B. S., Sergey S., Meixue, Z., Chenchen, Z. (2023). Phosphorus plays key roles in regulating plants' physiological responses to abiotic stresses. *Plants* 12(15), 2861; <https://doi.org/10.3390/plants12152861>, Pp 29.
- FAO. (2005). *Annual Rate of Deforestation*. Retrieved from [http://green.wikia.com/wiki/Deforestation in Nigeria](http://green.wikia.com/wiki/Deforestation_in_Nigeria). Pp.5. Accessed 17/11/ 2005.
- Fathi, A., Tari, D. B. (2016). Effect of drought stress and its mechanism in plants. *International Journal of Life Sciences*, 10 (1), 1-6. <https://api.semanticscholar.org/CorpusID:49233092>
- Fox, T. R. (2000). Sustained productivity in intensively managed forest plantations. *Forest Ecology and Management*, 138(1-3), 187–202. [https://doi.org/10.1016/S0378-1127\(00\)00396-0](https://doi.org/10.1016/S0378-1127(00)00396-0)
- Gbadamosi, A. E. (2014). Effect of watering regimes and water quantity on the early seedling growth of *Picralima nitida* (Stapf). *Sustainable Agriculture Research*, 3 (2), 35-42. doi:10.5539/sar.v3n2p35
- Ghorbani, P., Eshghi, S., Ershadi, A., Shekafandeh, A., Razzaghi, F. (2019). The possible role of foliar application of manganese sulfate on mitigating adverse effects of water stress in grapevine. *Communications in Soil Science and Plant Analysis*, 50(13), 1550-1562. <https://doi.org/10.1080/00103624.2019.1626873>
- Gomes, K.C.O., Paiva, H.N., Neves, J.C.L., Barros, N.F., Silva, S.R. (2004). Influencia da Saturacao por bases e do fosforo no crescimento de mudas de angicobranco. *Revista Arvore, Viosa*, 28(6), 785-792.
- Gonzales, J. A., Gallardo, M., Hilal, M., Rosa, M., & Prade, F. E. (2009). Physiological responses of quinoa (*Chenopodium quinoa* Willd.) to drought and waterlogging stresses: dry matter partitioning. *Botanical Studies*, 50, 35-42.
- Graciano, C., Goya, F., Frangi, L., Guamet, J. (2006). Fertilization with phosphorus increases soil nitrogen absorption in young plants of *Eucalyptus grandis*. *Forest Ecology Management*. 236 (2-3), 202-210. <https://doi.org/10.1016/j.foreco.2006.09.005>
- Haggai, P.T., Singh, L., Oseni, T.O.(2003). Effect of nitrogen and phosphorus application on the growth characters of sesame (*Sesamum indicum* L) in Northern Guinea Savanna of Nigeria. Samaru. *Journal of Agricultural Research*, 19, 79-90.
- Haque, M.A., Khan, M.K. (2012). Effect of phosphatic biofertilizer with inorganic and organic sources of phosphorus on growth and yield of lentil. *Journal of Environmental Sciences and Natural Resources*, 5(2), 225-230.
- Havlin, J. L., Tisdale, S.L., Beaton, J.D., Nelson, W.L.(2005). Soil Fertility and Fertilizers: *An Introduction to Nutrient Management*. Pearson Education, Inc, Upper Saddle River, New Jersey 07458. Pp. 244-254.
- Heather, R. (2021). The Importance of Phosphorus In Plant Growth. <https://www.gardeningknowhow.com/garden-how-to/soil-fertilizers/phosphorus-plant-growth.htm>. Accessed 16/05/25. pp 4.
- Heuer, S., Gaxiola, R., Schilling, R., Herrera-Estrella, L., López-Arredondo, D., Wissuwa, M., Delhaize, E., Rouached, H. (2017). Improving phosphorus use efficiency: a complex trait with emerging opportunities. *Plant Journal*, 90 (5), 868–885. <https://doi.org/10.1111/tpj.13423>
- Hill, J.O., Simpson, R.J., Moore, A.D., Chapman, D.F. (2006). Morphology and response of roots of pasture species to phosphorus and nitrogen nutrition. *Plant and Soil*, 286(1–2), 7–19. <https://doi.org/10.1007/s11104-006-0014-3>

- Hinsinger, P. (2001). Bioavailability of soil inorganic P in the rhizosphere as affected by root-induced chemical changes: a review. *Plant and Soil*, 237(2), 173–195. <https://doi.org/10.1023/A:1013351617532>
- Homyak, M., Allison, D., Huxman, E., Goulden, L., Treseder, K. (2017). Effects of drought manipulation on soil nitrogen cycling: A meta-analysis. *Journal of Geophysical Research: Biogeosciences*, 122 (12), 3260–3272. <https://doi.org/10.1002/2017JG004146>
- Hoque, R. A.T. M., Hossain, M.K., Mohiuddin, M., Hoque, M.M. (2004a). Effect of Inorganic fertilizers on the initial growth performance of *Anthocephalus chinensis* (Lam.) Rich. Ex. Walp. seedlings in the nursery. *Journal of Applied Sciences*, 4(3), 477–485. <https://doi.org/10.3923/jas.2004.477.485>
- Hoque, A.T.M.R., Hossain, M. K., Mohiuddin, M., Hoque, M.M. (2004b). Effect of inorganic fertilizers on initial growth performance of *Michelia champaca* Linn. seedling in nursery. *Journal of Biological Sciences*, 4(4), 489–497.
- Huda, S.M.S., Sujauddin, M., Shafinat, S., Uddin, M.S. (2007). Effects of phosphorus and potassium addition on growth and nodulation of *Dalbergia sissoo* in the nursery. *Journal of Forestry Research*, 18(4), 279–282. <https://doi.org/10.1007/s11676-007-0056-2>
- IITA (2011). Deforestation: Nigeria ranked worst in the World. *A publication of International Institute of Tropical Agriculture*. Retrieved from <http://www.thisdaylive.com/articles/deforestation-nigeria-ranked-worst-in-the-world/103321/5.pp2>. Accessed 21/11/2021.
- Iroko, O.A., Bolanle-Ojo, O.T., Agboola, R.O., Wahab, W.T. (2020). Effect of nitrogen fertilization on the early growth and development of *Pterocarpus erinaceus* Poir (Fabaceae) seedlings. *Forestry Development in Nigeria: Fifty years of Interventions and Advocacy*. In: Ogunsanwo, O.Y., Adewole, N.A., Oni, P.I and Idumah, F (Eds). *Proceedings of the 42nd Annual Conference of the Forestry Association of Nigeria* held in Ibadan 23rd–28th November, 2020. Pp599–606.
- Isah, A.D., Bello, A.G., Maishanu, H.M., Abdullahi, S. (2012). Effect of watering regime on the early growth of *Acacia senegal* (Linn) wild provenance. *International Journal of Plant, Animal and Environmental Sciences*, 1, 2231–4490. [url=\(https://api.semanticscholar.org/CorpusID:130275732\)](https://api.semanticscholar.org/CorpusID:130275732)
- Isidra-Arellano, M.C., Delaux, P.M., Valds-Lpez, O. (2021). The phosphate starvation response system: its role in the regulation of plant-microbe interactions. *Plant and Cell Physiology*, 62(3), 392–400. doi: 10.1093/pcp/pcab016.
- Jafer, D., Gebresilasie, H. (2018). Inorganic fertilizer requirement of coffee (*Coffea arabica* L) – Review. *Journal of Biology, Agriculture and Health care*, 8(21): 1–11.
- Jin, J., Wang, G. H., Liu, X., Pan, X., Herbert, S. J. (2005). Phosphorus application affects the soybean root response to water deficit at the initial flowering and full pod stages. *Soil Science and Plant Nutrition*, 51(7), 953–960. <https://doi.org/10.1111/j.1747-0765.2005.tb00133.x>
- John, W. (2013). How plants get water and nutrient, pp1–2. <http://www.dummies.com/how-to/content/how-plants-get-water-and-nutrient>; access 6/11/2013.
- Johnston, A.E., Poulton, P.R., Fixen, P.E., Curtin, D. (2014). Phosphorus. its efficient use in agriculture. *Advances in Agronomy*, 123, 177–228. <https://doi.org/10.1016/B978-0-12-420225-2.00005-4>
- Jose, L. M. (2003). Nitrogen and phosphorus reabsorption in tree of Neo-tropical rain forest. *Journal of Tropical Ecology*, 19(4), 465–468. doi:10.1017/S026646740300350X
- Jose, S., Merritt, S., Ramsey, C. L. (2003). Growth, nutrition, photosynthesis and transpiration responses of long leaf pine seedlings to light, water and nitrogen.

- Forest Ecology and Management*. 180(1–3), 335–344. [https://doi.org/10.1016/S0378-1127\(02\)00583-2](https://doi.org/10.1016/S0378-1127(02)00583-2)
- Kang, S. M., Shaffique, S., Peter, O., Injamum-Ul-Hoque, M. D., Lee, I. J. (2024). Physiological and molecular adaptation response of soybean seedlings under osmotic stress. *Pakistan Journal of Botany*, 56 (3), 1-8.
- Kauter, D., Lewandowski, I., Claupein, W. (2003). Quantity and quality of harvestable biomass from *Populus* short rotation coppice for solid fuel use: a review of the physiological basis and management influences. *Biomass and Bioenergy*, 24(6), 411–427. [https://doi.org/10.1016/S0961-9534\(02\)00177-0](https://doi.org/10.1016/S0961-9534(02)00177-0)
- Khan, F., Siddique, A. B., Shabala, S., Zhou, M., Zhao, C. (2023). Phosphorus plays key roles in regulating plants' physiological responses to abiotic stresses. *Plants*, 12 (15), 2861. <https://doi.org/10.3390/plants12152861>
- Khatik, S.K., Deshmukh, K.K., Sharma, G.D., Dwivedi, B.S. (2004). Effect of farm-yard manure , rock phosphate and biofertilizer on utilization of phosphorus by gram (*Cicer arietinum*) and its transformation in soil. *Journal of Nuclear Agriculture and Biology*, 33(1), 39-48.
- King, N., Seiler, J., Fox, T.R., Johnsen, K. (2006). The short-term effects of fertilization on loblolly pine photosynthesis and biomass In: Conner K.F, editor. *Proceedings of the 13th Biennial Southern Silvicultural Research Conference*. Ashville, NC: U.S. Department of Agriculture, Forest Service, Southern Research Station; 2006. p 113–116.
- Lai, J., Di, T., Suhui, M., Xuli, Z., Longchao, X., Jianxiao, Z., Xin, J., Chengyang, Z., Haihua, S., Zhang, Z., Yide, L., Biao, Z., Jingyun, F. (2018). The response of tree growth to nitrogen and phosphorus additions in a tropical montane rainforest. *Science of the Total Environment*, 618, 1064-1070.
- Lamb, D., Erskine, P.D., Parrotta, J. A. (2005). Restoration of degraded tropical forest landscapes. *Science*, 310, 16-28. doi:10.1126/science.1111773.
- Lambers, H., Albornoz, F., Kotula, L., Laliberté, E., Ranathunge, K., Teste, F. P., Zemunik, G. (2018). How belowground interactions contribute to the coexistence of mycorrhizal and non-mycorrhizal species in severely phosphorus-impooverished hyper diverse ecosystems. *Plant and Soil*, 424(1-2), 11-33. <https://doi.org/10.1007/s11104-017-3427-2>
- Lapido, D. (2010). The State of Nigeria forest in conserving endangered species in Nigeria. *Nigerian Field Society 80th Anniversary*, IITA, Ibadan p. 4.
- Larwanou, M., Adamou, M. M., Abasse, T. (2014). Effects of fertilization and watering regimes on early growth and leaf biomass production for two food tree species in the Sahel: *Moringa oleifera* Lam. and *Adansonia digitata* L. *Journal of Agricultural Science and Applications* 3(4), 82-88. doi:10.14511/jasa.2014.030401
- Laura, T., Damaris, P., Daniel, P., German, V. G., Fiona, M. S., Jessica, G., Jennifer, S. P.(2023). Phosphorus limitation of early growth differs between nitrogen-fixing and non-fixing dry tropical forest tree species. *New Phytologist*, 237(3), 766-779. <https://doi.org/10.1111/nph.18612>
- Laurent, G.H., Toussaint, O.L., Francis, G.H.G., Lisette, E.S.A., Brice, S. (2012). Ethnobotanical study of the African star apple (*Chrysophyllum albidum* G. Don) in the Southern Benin (West Africa). *Journal of Ethnobiology and Ethnomedicine*, 8, 40-47.
- Lawrence, D. (2001). Nitrogen and phosphorus enhance growth and consumption of four secondary forest tree species in Bomeo. *Journal of Tropical Ecology*, 17(6), 859-869. doi: <https://doi.org/10.1017/S0266467401001638>
- Lincoln, M. C., Will, R. E., Morris, L. A., Carter, E. A., Markewitz, D., Britt, J.R., Cazell, B., Ford, V. (2007). Soil change and loblolly pine (*Pinus taeda*) seedling

- growth following site preparation tillage in the Upper Coastal Plain of the southeastern United States. *Forest Ecology and Management*, 242 (2-3), 558–568. doi:10.1016/j.foreco.2007.01.069
- Liu, X., Zhang, S., Shan, X., Zhu, Y.G. (2005). Toxicity of arsenate on germination, seedling growth and amylolytic activity of wheat. *Chemosphere*, 61, 293-301. doi: 10.1016/j.chemosphere.2005.01.088.
- Lopez, G., Ahmadi, S.H., Amelung, W., Athmann, M., Ewert, F., Gaiser, T., Gocke, M.I., Kautz, T., Postma, J., Rachmilevitch, S., Schaaf, G., Schnepf, A., Stoschus, A., Watt, M., Yu, P., Seidel, S.J. (2023). Nutrient deficiency effects on root architecture and root-to-shoot ratio in arable crops. *Frontiers Plant Science*, 13, 1-18, 13, 1067498. doi: 10.3389/fpls.2022.1067498.
- López-Bucio, J., Cruz-Ramírez, A., Herrera-Estella, L. (2003). The role of nutrient availability in regulating root architecture. *Current Opinion in Plant Biology*, 6(3), 280–287. doi: 10.1016/s1369-5266(03)00035-9.
- Ma, Z., Bielenberg, D.G., Brown, K.M., Lynch, J.P. (2001). Regulation of root hair density by phosphorus availability in *Arabidopsis thaliana*. *Plant, Cell and Environment*, 24(4), 459–467. <https://doi.org/10.1046/j.1365-3040.2001.00695.x>
- Makinde, E.A., Ayoola, A.A. (2008). Residual influence of early season crop fertilization and cropping system on growth and yield of cassava. *American Journal of Agricultural and Biological Sciences*, 3(4), 712-715. doi: 10.3844/ajabssp.2008.712.715, <http://www.scipub.org/fulltext/ajab/ajab34712-715.pdf>.
- Malhotra, H., Vandana., Sharma, S., Pandey, R. (2018). Phosphorus nutrition: plant growth in response to deficiency and excess. In *Plant Nutrients and Abiotic Stress Tolerance*; Springer: Berlin/Heidelberg, Germany, 2018; pp. 171–190. doi:10.1007/978-981-10-9044-8_7
- Mbah, C.N., Mbagwu, J.C.C. (2003). Studies on decomposition, mineralogical rate and biochemical oxygen demand of organic wastes. *International Journal of Agriculture and Biological Services*, 2, 51-54.
- Melo, A.S., Gois, M.P.P., Brito, M.E.B., Viegas, P.R.A., Araujo, F.P., Melo, D.L.M.F., Mendonca, M.C. (2005). Desenvolvimento de porta-enxertos de umbuzeiro em resposta a adubacao com nitrogenio e fosforo. *Cicenc. Rural, Santa Maria*, 35(2), 324-331. <http://www.alice.cnptia.embrapa.br/alice/handle/doc/132904>
- Mhango, J., Akinnifesi, F.K, Mng'omba, S.A., Sileshi, G. (2008). Effect of growing medium on early growth and survival of *Uapaca kirkiana* Müell Arg. seedlings in Malawi. *African Journal of Biotechnology*, 7(13), 2197-2202.
- Mohammad, B. U., Mohammed, A. S. A. K., Sharif, A. M., Mohammed, K. H. (2008). Effects of inorganic fertilizers on biological nitrogen fixation and seedling growth of some agroforestry trees in Bangladesh. *Journal of Forestry Research*, 19(4), 303–306. doi 10.1007/s11676 -008-0054-z
- Mohammad, B.U., Sharif, A. M., Mohammed, A. S. A. K., Mohammed, K. H. (2007). Effects of phosphorous fertilizer on seedlings growth and nodulation capabilities of some popular agroforestry tree species of Bangladesh. *Journal of Forestry Research*, 18 (4), 283- 286.
- Muhammad, R., Peng, Z., Hai-long, S., Salahuddin .(2017). Influence of nitrogen and phosphorous on the growth and root morphology of *Acer mono*. *PLoS One*, 12(2), e0171321 doi: 10.1371/journal.pone.0171321, pp 24
- Mukhtar, R. B.(2012). Effect of some silvicultural treatments on early growth of *Diospyros mespiliformis*. *M.Sc Dissertation of the Department of Forest Resources Management, University of Ibadan, Nigeria*. Pp48.

- Mukhtar, R.B. (2016). Effect of drought stress on early growth of *Adansonia digitata* (L) in semi-arid region of Nigeria. *Journal of Research in Forestry, Wildlife and Environment*, 8(4), 109-115. <http://www.ajol.info/index.php/jrfwe>
- Mukhtar, R.B., Mansur, M.A., Abdullahi, S., Bunza, M.S. (2016). The growth of *Balanites aegyptiaca* (L) seedlings under varied watering intervals in the nursery. *Journal of Tropical Agriculture, Food, Environment and Extension*, 15(3), 30-33. <https://dx.doi.org/10.4314/as.v15i3>.
- Nadrowski, K., Wirth, C., Sherer-Lorenzen, M. (2010). Is forest diversity driving ecosystem and functions and services?. *Current Opinions in Environmental Sustainability*, 2(1-2), 75-79. <https://doi.org/10.1016/j.cosust.2010.02.003>
- Nathan, M.V.(2009). *Soils, Plant Nutrition and Nutrient Management, Master Gardener Core Manual*. Pp 20
- Nicoloso, F.T., Fogaca, M.A. de F., Zanchetti, F. (2001). Nutricao mineral de mudas de grapia (*Apuleia leiocarpa*) em Argissolo vermelho Distrofico arenico: (1) efeito da adubacao NPK no Crescimento. *Cienc. Rural, Santa Maria*, 31(6), 991-998.
- Nilus, R., May Cock, C.R., Majalap-lee, N., Burslem, D.F.R.P. (2011). Nutrient limitation of tree seedling growth in three soil types found in Sabah. *Journal of Tropical Forest Science*, 23(2), 133-142. <http://www.jstor.org/stable/23616913>
- Nwoboshi, L.C. (2000). *The Nutrient Factor in Sustainable Forestry*. Ibadan University Press, Nigeria, 303pp
- Oboho, E.G., Igharo, B. (2017). Effect of pre-germination treatments on germination and watering regimes on the early growth of *Pycnanthus angolensis* (Welw) Warb. *Journal of Agriculture and Veterinary Science*, 10(3), 62-68. doi:10.9790/2380-1003026268
- Ogidan, O. A., Olajire-Ajayi, B. L., Adenuga, D. A. (2018). Assessment of watering regimes on seedling growth performance of *Kigelia africana* (Lam) Benth. *Biodiversity Conservation and National Development Potentials and Challenges*. In: Gabriel, S.U., Folaranmi, D.B and Edem, A.E (eds). *Proceedings of 6th Nigeria Chapter of Society for Conservation Biology (NSCB) Biodiversity Conference*. pp 341-345.
- Ogunade, J.O., Gbadebo, J.O., Ajekigbe, J.M., Banjo, T.A., Bolanle-Ojo, O.T. (2020). Early growth performance of *Parkia biglobosa* (Jacq) Benth on plantation as influenced by fertilizer application. *Forestry Development in Nigeria: Fifty years of Interventions and Advocacy*. In: Ogunsanwo, O.Y., Adewole, N.A., Oni, P.I and Idumah, F (Eds). *The proceedings of the 42nd Annual Conference of the Forestry Association of Nigeria* held in Ibadan, 23-28th November, 2020. Pp555-563.
- Ogunwole, J.O., Lawal, A.B., Olarewaju, J.D., Audu, K., Adekpe, D.I., Ugbabe, O.O., Yaro, D.T., Oyinlola, E.Y. (2006). Integrated soil waterlogging and nutrient management for late season crop production systems in Nigeria. *Journal of Agronomy*, 5, 314-320. doi: 10.3923/ja.2006.314.320
- Ojelabi, O.K., Musa, F.B., Akinleye, F.O., Oyewumi, R.V. (2018). Effects of different organic manure on the growth of *Pterocarpus erinaceus* Poir. *Emerging issues in Forest Management: Experiences and Lessons for Nigeria*. In: Akinwole, A.O., Adekunle, V.A.J and Ogunsanwo, O.Y (Eds). *The proceedings of the 40th Annual Conference of the Forestry Association of Nigeria* held in Lagos, Lagos State, 12-16th March, 2018. Pp 614-624.
- Okafor, J.C.(2003). Promising Trees of Agroforestry in Southern Nigeria. Forestry Development and Investigation Branch, Forestry Commission, Enugu, Nigeria. 13pp.
- Olajiire-Ajayi, B.L., Okeleke, S.O., Omoyeni, K.O. (2018). Effect of organic manure on the growth and fibre characteristics of *Mansonia altissima* A.Chev seedlings. *Emerging issues in Forest Management: Experiences and Lessons for Nigeria*. In: Akinwole,

- A.O., Adekunle, V.A.J and Ogunsanwo, O.Y (Eds). *The proceedings of the 40th Annual Conference of the Forestry Association of Nigeria* held in Lagos, Lagos State , 12-16th March, 2018. Pp.605-613.
- Olajuyigbe, S.O., Jimoh, S.O., Adegeye, A.O., Mukhtar, R.B. (2013). Drought stress on early growth of *Diospyros mespiliformis* Hochst ex A. Rich in Jega, Northern Nigeria. *Nigerian Journal of Ecology*, 12(1), 71-76.
- Olander, L. P., Vitousek, P. M. (2004). "Biological and geochemical sinks for phosphorus in soil from a wet tropical forest." *Ecosystems*, 7, 404–419. <https://doi.org/10.1007/s10021-004-0264-y>
- Olaoluwa, T.A., Muhammad, N.O., Oladiji, A.T. (2012). Biochemical assessment of the mineral and some antinutritional constituents of *Aspergillus niger* fermented *Chrysophyllum albidum* seed meal. *African Journal of Food Science*, 6(1), 20-28. doi:10.5897/ajfs11.144
- Ong, K., Lim, M.T., Labit, S. (2004). Growth and photosynthetic response of *Shorea platyclados* seedlings to nitrogen application. In: H. Aminah., S. Ani, H.C. Sim and B. Krishnapillay, Eds. *Proceedings of the seventh round table conference on Dipterocarps, Asia Pacific Association of Forestry Research Institutions*, Kuala Lumpur, 2004, pp. 89-100.
- Onyekwelu, J.C., Stimm, B. (2011). *Chrysophyllum albidum*. In: Roloff, A; Weisgerber, H; Lang, U; Stimm, B. *Enzyklopadie Holzgewachse*, Willey der VCH, Weinheim, 59. Erg.Lfg.10/11, 12pp.
- Onyekwelu, J.C., Stimm, B., Mosandi, R., Olusola, J.A. (2011). Domestication of *Chrysophyllum albidum* from rainforest and derived savannah ecosystem – Phenotype variation and selection of elite trees. *Conference on International Research on Food Security, Natural Resource Management and Rural Development*. Tropentage, pp. 7.
- Pandey, S.T., Singh, P., Pandey, P. (2006). Site specific nutrient management for *Withania somnifera* at subtropical belt of Uttaranchal. *International Journal of Agricultural Sciences*, 2, 626–628.
- Paz-Ares, J., Puga, M.I., Rojas-Triana, M., Martinez-Hevia, I., Diaz, S., Poza-Carrión, C., Miñambres, M., Leyva, A. (2022). Plant adaptation to low phosphorus availability: core signaling, crosstalks, and applied implications. *Molecular Plant*, 15(1), 104–124. doi: 10.1016/j.molp.2021.12.005.
- Pereira, B.F.F., Tucci, C.A.F., Santos, J.Z.L., Silva, T.A.F. (2014). Phosphorus sources: Effect on the tropical cedar nutrition and growth. *Journal of Tropical Forest Science*, 26(4), 513-521.
- Pigna, M., Cozzolino, V., Giandonato Caporale, A., Mora, M.L., Di Meo, V., Jara, A.A., Violante, A. (2010). Effects of phosphorus fertilization on arsenic uptake by wheat grown in polluted soils. *Journal of Soil Science and Plant Nutrition*, 10(4), 428-442.
- Predick, K.I., Gergel, S.E., Turner, M.G. (2009). Effect of flood regime on tree growth in the flood plain and surrounding uplands of Wisconsin River. *River Research and Applications* 25, 283-296. <http://dx.doi.org/10.1002/rra.1156>.
- Purbajanti, E.D., Slamet, W., Fuskah, E., Rosyida, R.(2019). Effects of organic and inorganic fertilizers on growth, activity of nitrate reductase and chlorophyll contents of peanuts (*Arachis hypogaea* L). *IOP Conf. Series: Earth and Environmental Science*, 250 (2019) 012048. *IOP Publishing*. pp1-8. doi 10.1088/1755-1315/250/1/012048
- Quaghebeur, M., Rengel, Z. (2003). The distribution of arsenate and arsenite in shoots and roots of *Holcus lanatus* as influenced by arsenic tolerance and arsenate and phosphate supply. *Plant Physiology*, 132(3), 1600-1609. doi: 10.1104/ pp.103.021741.

- Rahaman, O. (2012). A review of medicinal value of *Chrysophyllum albidum* (African star apple). *African Traditional Medicine*, 1(1), 1-3.
- Ramos, M.R.C., Pinto, J.E.P.B., Furtini Neto, A.E., Davido, A.C.(2000). Influencia da Aplicac ao de nitrogenio fosforo e potassio no crescimento e composicao mineral de mudas de patade-vaca (*Bauhinia forficata*). *Revista Brasileira de Plantas Medicinai*s: Botucatu, 3(1), 79-86.
- Richardson, A. E., Simpson. R. J. (2011). "Soil microorganisms mediating phosphorus availability." *Plant Physiology*, 156(3), 989–996. doi: 10.1104/pp.111.175448
- Rickard, D.A. (2000). A review of phosphorus acid and its salts as fertilizer materials. *Journal of Plant Nutrition*, 23(2), 161-180. <https://doi.org/10.1080/01904160009382006>
- Rotowa, O. J., Ibeh, K.G., Anozie, E. L. (2020). Management of *Khaya senegalensis* (Desr.) A. Juss Intercropped with Maize and Groundnut under Agroforestry Scheme. *International Journal of Applied Research and Technology*, 9(2), 8 – 14.
- Salami, K. D. (2015). Comparative effect of nitrogenous and phosphorous fertilizers on the growth of *Enterolobium cyclocarpum* Jacq seedlings. *Nigeria Tropical Biology Association (NTBA), Conservation in 21st Century Nigeria: Transcending Disciplinary Boundaries, 5th Annual Biodiversity Conference* held at Federal University of Technology, Akure (FUTA) from 19 – 20 May, 2015, pp. 265-268.
- Salami, K. D., Lawal, A. A., Oyebamiji, N A., Yisau, J .A. (2020). Effect of nitrogen, phosphorus fertilizers and mycorrhizal nodulation on the growth and yield of tree seedlings. *FUW Trends in Science and Technology Journal*, 5(1), 296-299. www.ftstjournal.com 296 e-ISSN: 24085162
- Sale, F.A.(2015). Evaluation of watering regime and different pot sizes on the growth of *Parkia biglobosa* seedlings under nursery condition. *European Scientific Journal*, 11(12), 313-325.
- Sardans, J., Roda, F., Penuelas, J. (2004). Phosphorus limitation and competitive capacities of *Pinus halepensis* and *Quercus ilex* subsp *rotundifolia* on different soils. *Plant Ecology*, 174, 305-317. <https://doi.org/10.1023/B:VEGE.0000049110.88127.a0>
- Sarfraz, S., Ali, F., Hameed, A., Ahmad, Z., Riaz, K. (2023). Sustainable agriculture through technological innovations. In: *Sustainable Agriculture in the Era of the OMICs Revolution*; Springer: Cham, Switzerland, pp. 223–239.
- SAS (2003). Statistical analysis system. SAS release 9. 1 for windows, SAS Institute Inc. Cary, NC, USA
- Schumacher, M.V., Ceconi, D.E., Santana, C.A. (2004). Influencia de diferentes doses de fosforo no crescimento de mudas de angicovermelho (*Parapiptadenia rigida* (Bentham) Brenan). *Revista Arvore vici*osa, 28(1), 149-155. <https://doi.org/10.1590/S0100-67622004000100019>
- Sezer, S., Seda B. A. (2019). The effect of different phosphorus doses on the growth of pepper seedling (*Capsicum annuum* L.). *Asian Journal of Advances in Agricultural Research*, 11(4), 1-4. doi: 10.9734/AJAAR/2019/v11i430067
- Sharma, N. K., Singh, R. J., Kumar, K. (2012). Dry matter accumulation and nutrient uptake by wheat (*Triticum aestivum* L) under poplar (*Populus deltoids*) based Agroforestry System. *International Scholarly Research Notices*, 12, 1-9.
- Shen, J. P., Zhang, L. M., Guo, J. F., Ray, J. L., He, J. Z. (2010). Impact of long-term fertilization practices on the abundance and composition of soil bacterial communities in Northeast China. *Applied Soil Ecology*, 46(1), 119–124.
- Shi, S., Peng, C., Wang, M., Zhu, Q., Yang, G., Yang, Y., Xi, T., Zhang, T. (2016). "A Global meta-analysis of changes in soil carbon, nitrogen, phosphorus and sulfur and stoichiometric shifts after forestation." *Plant and Soil*, 407, 323–340. <https://doi.org/10.1007/s11104-016-2889-y>

- Siddhuraju, P. (2007). Antioxidant activity of polyphenolic compounds extracted from defatted raw and dry heated *Tamarindus indica* seed coat. *LWT, Food Science and Technology*, 40(6), 982-990. <https://doi.org/10.1016/j.lwt.2006.07.010>
- Soriano, M.A., Fereres, E. (2003). Use of crops for insitu phytoremediation of polluted soils foiling a toxic flood from mine spill. *Plant Soils*, 256, 253-264. <https://doi.org/10.1023/A:1026155423727>
- Suthiphasilp, R. (2009). Effect of manure and chemical fertilizer on solubility of phosphate of phosphorus in soils. *Asian Journal of Food Agro-Industry*, S205-S210.
- Taiz, L., Zeiger, E. (2002). Plant physiology. 3rd edn. Sunderland: Sinauer Associates., 690 pp.
- Tao, Y., Zhang, S., Yuan, W. J., Shan, X.Q. (2006). Effect of oxalate and phosphate on the release of arsenic accumulation in wheat. *Chemosphere*, 65, 1281-1287. doi: 10.1016/j.chemosphere.2006.04.039
- Tariq, A., Zeng, F., Graciano, C., Ullah, A., Sadia, S., Ahmed, Z., Murtaza, G., Ismoilov, K., Zhang, Z. (2023). Regulation of metabolites by nutrients in plants. *Plant Ionomics*, 1–18. <https://doi.org/10.1002/9781119803041.ch1>
- Telkamp, M. (2015). The straight poop of using chicken manure as fertilizer. Retrieved 30th, 2017.
- Thompson, J. B., Slot, M., Dalling, J. W., Winter, K., Turner, B. L., Zalamea, P. (2019). Species-specific effects of phosphorus addition on tropical tree seedling response to elevated CO₂. *Functional Ecology*, 33 (10), 1871-1881. <https://doi.org/10.1111/1365-2435.13421>
- Uddin, M. B., Mukul, S.A., Khan, M.A.S.A., Hossain, M.K. (2007). Effects of phosphorous fertilizer on seedlings growth and nodulation capabilities of some popular agroforestry tree species of Bangladesh. *Journal of Forestry Research*, 18(4), 283-286. doi: 10.1007/s11676-007-0054-4
- Ureigho, V.N., Ekeke, B.A. (2010). Nutrient values of *Chrysophyllum albidum* Linn African star apple as a domestic income plantation species. *An International Multi-disciplinary Journal, Ethiopia*, 4(2), 50-56. doi:10.4314/afrev.v4i2.58288
- Usman, A., Wakawa, D.L., Adeogun, P.F. (2013). The influence of soil types and watering regimes on early growth of *Moringa oleifera* (LAM) in semi-arid region of Nigeria. In: Labode Popoola., Idumah, F.O., Ogunsanwo, O.Y and Azeez, I.O (Eds). *Forest Industry in a Dynamic Global Environment. Proceedings of the Annual Conference of the Forestry Association of Nigeria*. PP751-755.
- Usman, I.A., Afera, S.I.N., Awua, N.(2022). Effect of growing media on germination and early growth of *Erythrophleum suaveolens* (Perr). *Securing the Nigeria's Forest Estates for Sustainable Development*. In: Ogunsanwo, O.Y, Adewole, N.A, Oni, P.I and Azeez, I.O (Eds). *The proceedings of the 43rd Annual Conference of the Forestry Association of Nigeria held in Akure 14-18th March, 2022*. Pp 122-130.
- Vance, C. P., Uhde-Stone, C., Allan, D. L.(2003). "Phosphorus acquisition and use: critical adaptations by plants for securing a nonrenewable resource." *New Phytologist*, 157, 423–447. doi: <https://doi.org/10.1046/j.1469-8137.2003.00695.x>
- Villar-Salvador, P., Planelles, R., Enriquez, E., Penuelas, R.J.(2004). Nursery cultivation regimes, plant functional attributes and field performance relationship in the Mediterranean oak, *Quercus ilex* L. *Forest Ecology and Management*, 196 (2-3), 257-266
- Vitousek, P. M., Porder, S., Houlton, B. Z., Chadwick, O. A. (2010). "Terrestrial phosphorus limitation: mechanisms, implications, and nitrogen-phosphorus interactions." *Ecological Applications*, 20, 5–15. doi: 10.1890/08-0127.1. PMID: 20349827.

- Wan Juliana, W. A., Burslem, D.F.R.P., Swaine, M.D. (2009). Nutrient limitation of seedling growth on contrasting soils from Pasoh Forest Reserve, Peninsular Malaysia. *Journal of Tropical Forest Science*, 21, 316-327.
- Wang, R., Bicharanloo, B., Hou, E., Jiang, Y., Dijkstra, F. A. (2022). "Phosphorus supply increases nitrogen transformation rates and retention in soil: a global meta-analysis." *Earth's Future*, 10 (3), e2021EF002479. <https://doi.org/10.1029/2021EF002479>.
- Waraich, E. A., Ahmad, Z., Ahmad, R., Saifullah., Ashraf, M.Y. (2015). Foliar applied phosphorous enhanced growth, chlorophyll contents, gas exchange attributes and PUE in wheat (*Triticum aestivum* L.). *Journal of Plant Nutrition*, 38(12), 1929–1943.
url=(<https://api.semanticscholar.org/CorpusID:96604972>)
- Wen, Z., Li, H., Shen, J., Rengel, Z. (2017). Maize responds to low shoot P concentration by altering root morphology rather than increasing root exudation. *Plant Soil*, 416, 377–389. <https://doi.org/10.1007/s11104-017-3214-0>
- Williams, P.N., Villana, A., Deacon, C., Raab, A., Figuerola, J., Green, A.J., Feldmann, J.B., Meharg, A.A. (2007). Greatly enhanced arsenic shoot assimilation in rice leads to elevated grain levels compared to wheat and barley. *Environmental Science and Technology*, 41, 6854-6859. <http://dx.doi.org/10.1021/es070627i>
- Williamson, L., Ribrioux S., Fitter A., Leyser, O. (2001). Phosphate availability regulates root system architecture in Arabidopsis. *Plant Physiology*, 126(2), 875–882.<https://doi.org/10.1104/pp.126.2.875>
- Wole, O. (2013). Unlimited nutritional benefits of African star apple. *Natural Health*, 1(1), 1-4.
- Wu, C., Fan, Z., Wang, Z. (2015). Effect of phosphorus stress on chlorophyll biosynthesis, photosynthesis and biomass partitioning pattern of *Fraxinus mandchurica* seedlings. *Abstract of Unbound Medline*. 1pp.
- Wyngaard, N., Cabrera, L., Jarosch, A., Benemann, K. (2016). Phosphorus in the coarse soil fraction is related to soil organic phosphorus mineralization measured by isotopic dilution. *Soil Biology and Biochemistry*, 96, 107-118. <https://doi.org/10.1016/j.soilbio.2016.01.022>
- Xu, B. C., Niu, F.R., Duan, D.P., Xu, W.Z., Huang, J. (2012). Root morphological characteristics of *Lespedeza davurica* (L.) intercropped with *Bothriochloa ischaemum* (L.) Keng under water stress and P application conditions. *Pakistan Journal of Botany*, 44(6), 1857 -1864.
- Yao, Y., Wu, D., Gong, Z., Ma, C. (2012). Effect of phosphorus level on nitrogen accumulation and yield in soybean. *Journal of Nuclear Agricultural Sciences*, 26, 947–951.
- Yisau, J. A., Salami, K. D., Aduradola, A. M. (2020). Effects of types of (organic and inorganic) fertilizer and quantities on the early growth of *Albizia zygia* seedlings. *FUDMA Journal of Agriculture and Agricultural Technology*, 6(1), 147 – 152.
- Zapata, F., Zaharah, A. R. (2002). Phosphate availability from phosphate rock and sewage sludge as influenced by addition of water soluble phosphate fertilizers. *Nutrient Cycling in Agroecosystems*, 1 (63), 43–48. <https://doi.org/10.1023/A:1020518830129>