

Psychological distress and Coping Strategies among Women with Breast Cancer receiving Chemotherapy at Assiut university Hospital

*Nadia Ebraheem Sayied **Sahar Ali Abd Elmohsen

*Psychiatric Nursing Department,** Adult Nursing Department, Faculty of Nursing, Assiut University

ABSTRACT

The study aimed to assess depression, anxiety symptoms as psychological distress and coping strategies among women with breast cancer receiving chemotherapy, and to investigate the relationship between them. A descriptive research design was utilized. The study was carried out in the oncology unit at Assiut University Hospital and South Egypt Cancer Institute. The sample comprised of all women with breast cancer receiving chemotherapy attending the oncology unit and South Egypt Cancer Institute during three months period from January to March 2015. **Methodology:** comprised of personal data questionnaire, the Hospital Anxiety Depression Scale (HADS), developed by, Zigmond and Snaith (1983); contains seven items that assess anxiety and seven items that assess depression rated on a four-point Likart scale (0 to 3) and Brief COPE scale developed by, Carver, (1997). It consists of 28-item self report measure of both adaptive and maladaptive coping skills .Individuals respond to each item on a four-point Likart scale, indicating the frequency with which each strategy is used (1to4). **Results:** 62.2%, 32.4% of the studied group were had high level of symptoms of anxiety, and depression respectively. The highest mean scores of coping strategies were related to religion use of emotional support, active coping, use of instrumental support, planning, and positive reframing. **Conclusion:** The highest mean score of the studied group used Problem-focused Strategies and emotional focused strategies, as a method of coping with anxiety and depression symptoms. **Recommendations:** Designing and implementing psycho-educational programs about stressors related to patients with breast cancer.

Key words: psychological distress, Coping Strategies, Breast Cancer, Chemotherapy.

Castro, Schlebusch , 2006 and Ballenger , et al. 2001).

INTRODUCTION

Cancer diagnosis is usually apparent as a crisis for both patients and their families (Taleghani et al. 2006). The most common cancer affecting women is breast cancer. Women who have had cancer often experience significant psychological and physical stress, including the stress of the diagnosis, debilitating treatments and chronic pain. Distress can compromise compliance with treatment and negatively affect prognosis and survival rates, so the significance of patients' psychological status forms an essential element of oncological treatment (Carayol M. et al. 2013, Lo

For those diagnosed in the late stages, they too have to come to terms with their diagnosis, the fact that they will have to receive palliative as opposed to curative care and the fears and vagueness about end of life (Kearney, & Richardson, 2006, Lackey, et al. 2001). Breast cancer rates are increasing in developing countries, including Egypt, and are largely recognized to aging of the population (Jemal, et al. 2010, Freedman et al. 2006). Although breast cancer incidence rates in Egypt are substantially lower than the rates in the United States and other developed countries (Shulman, 2010). Breast

cancer is the most common cancer among women in Egypt (**Omar, et al. 2003**).

Women who have had a breast cancer often experience significant psychological and physical stress, as well as physical problems. They have to change their lifestyle following a long period of treatment, and this may well influence their quality of life. Their daily life is full of stress and worry regarding their family/sexual roles disrupting physical status, emotional and spiritual well-being and personal relationships for the patient and family. In addition to the feeling of worries about their future life in terms of their general functionality status. (**Mohaghghi, 2003**).

The majority cancer patients receiving chemotherapy experience psychological distress as a result of negative effects of chemotherapy agents, the uncertainty of post-treatment, and the occurrence of psychosocial problems (**Del Mastro et al. 2002 and Nor Zuraida 2010**). The same author added that, anxiety is widespread at the beginning of treatment, worrying of the potential side effects of the agents and fear of recurrence after completion of treatment. The incidence of depression varies between 8% and 36% depending on the site of cancer, diagnostic criteria.

Several studies have revealed that depression, anxiety, stress and poor quality of life are often psychological consequences of living with cancer, and cancer patients face the double challenge of learning to manage the physical as well as psychological effects of cancer. Furthermore, previous studies suggest that depression and poor quality of life have been associated with 5 year survival rates as well as increased mortality as a result of cancer (**Montazeri, 2008**).

Coping with cancer requires dealing with a number of specific problems such as pain and fatigue, anger, changes in

appearance, and relationships with partners and family. Successful, coping strategies may need to be different according to specific stressor. Previous studies of coping have not looked at particular stressors other than chemotherapy, yet these specifics may help target interventions. For instance, coping with fatigue and pain may need special strategies than coping with changes in appearance (**Janeen, et al. 2007**).

Proper coping and adjustment is essential in facing chronic diseases, especially during the treatment period such as chemotherapy, employed four main styles in coping with the treatment crisis - think positively/fighter, acceptance, fearfulness and hopelessness, and had "confrontive" element in coping style, were found to experience less psychological and physical symptoms, as compared to the patients who had "avoidant" element in the coping strategy used, (**John D and MacArthur, 1998, Shapiro, et al. 1997 Payne, 1990**).

Another investigator pointed out that behavioral escape avoidance and cognitive escape-avoidance as the most significant coping mechanisms which contribute to the psychological distress of the cancer patients receiving chemotherapy (**Zabalegui, 1999**). Therefore, the coping style with a fighting spirit has been observed to associate with a greater adherence to the chemotherapy regimen (**Ayres, et al. 1994**).

Significance of the study:-

Breast cancer rates are increasing in developing countries, including Egypt. Although breast cancer incidence rates in Egypt are substantially lower than the rates in the United States and other developed countries, breast cancer is the most common cancer among women in Egypt. Patients undergoing treatment for cancer face major physical and emotional challenges such as, stress, anxiety and depression that adversely

affect the quality of life and daily functioning of those patients. So that it seems important to investigate how women with breast cancer cope with their illness and related stressors. Data generated from this study may help in designing and implementing program for those patients to assist them in coping with disease and its consequence.

Aim of the study:-

This study aimed to assess depression, anxiety symptoms as psychological distress and coping strategies among women with breast cancer receiving chemotherapy.

Investigate the relationship between anxiety, depression and coping strategies used by those patients.

Research question :- Is there is a relation between depression, anxiety and coping strategies among women with breast cancer receiving chemotherapy?

Subjects and methods:-

Research Design: A descriptive research design was utilized in this study.

Setting:-

The study was conducted in the oncology unit at Assiut University Hospital and South Egypt Cancer Institute.

Subjects:-

All women with breast cancer receiving chemotherapy attending the oncology unit and South Egypt Cancer Institute during three months period from January to March 2015.

Methodology:-

- 1- **Personal data questionnaire** which comprised of name, age, sex, level of education and marital status.

- 2- **The Hospital Anxiety Depression Scale (HADS)** developed by, Zigmond and Snaith (1983); score was used to report anxiety symptoms and psychological depressive symptoms. The instrument was designed for medically ill patients and does not include physical symptoms. It has been validated in patients with cancer. The HADS contains seven items that assess anxiety and seven items that assess depression rated on a four-point Likart scale (0 to 3), with 3 indicating higher symptom frequencies. Score for each subscale anxiety and depression and range from 0-21 with scores categorized as follows: normal (0-7), mild (8-10), moderate (11-14), severe (15-21). Scores for the entire scale (emotional distress) range from 0-42, with higher scores indicating more distress. Cronbach's alpha reliability coefficient of the (HADS) was found to be 0.72.

- 3- **Brief COPE scale**, it was developed by, Carver, (1997). It consists of 28-item self report measure of both adaptive and maladaptive coping skills. As an abbreviated version of the COPE Inventory, it can be used to assess trait coping (the manner in which people cope with stress in everyday life) and state coping (the way people cope with a specific stressful situation). In total, 14 dimensions are covered by this scale. These are self distraction, active coping, denial, substance use, use of emotional support, use of instrumental support, behavioral disengagement, venting, positive reframing, planning, humor, acceptance, religion and self-blame. Every dimension has two items. The coping dimensions also can be divided into two major categories: Problem-focused Strategies (i.e. active coping, planning and using instrumental support) and Emotion focused Strategies (i.e. positive reframing, acceptance, religion, using emotional support and denial). Individuals respond to each item on a four-point Likart scale, indicating the

frequency with which each strategy is used (1) = I have not been doing this at all; 4 = I have been doing this a lot). With alpha coefficients of 0.79. The previous two questionnaires were translated into Arabic language. This translation was revised by five experts; three assistant professors in psychiatry and two assistant professors in neurology. A back translation into English language and compared with the original sentences to ensure content validity.

Methods of data collection:-

1- An official approval was obtained from the Dean of the faculty of Nursing Assiut University directed to the Head of oncology unit at Assiut University Hospital and Dean of South Egypt Cancer Institute in order to get permission to conduct the study. The purpose of the study was explained to the patients before starting data collection. Each session ranged from 30 to 45 minutes.

Ethical consideration:-

Ethical approval for the protocol of research was obtained from post graduate and research committee in the faculty of nursing-Assiut University. The researchers obtained informed consent for participation from all patients who reassured about the confidentiality of the obtained information to avoid misunderstanding. The patient's have freedom to withdrawal from the study at any time.

Pilot study:-

A pilot study was conducted at the beginning of the study. It included 10% of the total sample to investigate the feasibility of data collection tools and their clarity.

They were included in the study because no modification was done.

Statistical analysis:-

Statistical analysis was done by using SPSS version 20 Software Package. Data collected were coded and analyzed. The results were tabulated and statistically compared using categorical variables and were described by number and percent (N, %), where continuous variables described by mean and standard deviation (Mean, SD). t-test and ANOVA to compare between continuous variables. A two-tailed $p < 0.05$ was considered statistically significant.

Results:-

Table (1) shows, socio-demographic characteristics of breast cancer patients on chemotherapy. The mean age was (52.9 + 12) years, ranged from 18 to 80 years old, 66.7% of them were married, 25% were divorced, 52% from urban and the rest of them from rural area. According to education, 61.8% of the study group were illiterate and have primary education and 78.4% of them were housewives.

Relation between brief COPE scale and depression are presented in Table 2. It reveals that, patients on moderate depression had higher score 55.±119, Followed by severe depression 53.4±116. Also there were no statistical significant association between brief COPE scale, and depression.

Relation between brief COPE scale and anxiety are presented in Table 3. It illustrates that, patients on high anxiety had higher score (54±116). Also there were no statistical significant association between brief COPE scale and anxiety.

Figure (1) shows that, 62.2%, 32.4% of studied group were had high level of anxiety, and depression respectively. This result suggests that, women with breast cancer are vulnerable for anxiety and depression.

Table (4) reveals distribution of the mean scores and standard deviations of Brief COPE scale. It was found that, the highest mean scores were related to religion, use emotional support, active coping, use instrumental support, planning, and positive reframing. This results suggests that women with breast cancer used Problem-focused Strategies and Emotion focused Strategies as a methods of coping. The lowest scores were related to, substance use, behavioral disengagement, and self-blame.

Correlation between brief COPE scale, depression and anxiety are presented in Table 5. It indicates that, patients with anxiety

symptoms had significant higher scores on, denial, positive reframing, acceptance, use of instrumental support, self - distraction and self blame as their coping strategies with ($P<0.001$), ($p=0.002$), ($p=0.004$), ($p=0.012$), ($p=0.19$), respectively. While patients who were depressed had significantly higher scores on coping strategies such as planning, denial, positive reframing, acceptance, use of instrumental support, active coping, self-distraction with ($P<.001$) ($p=0.001$) ($p=0.002$) (0.002) ($p=0.12$) ($p=0.38$) respectively. This result suggests that, those patients used Problem-focused Strategies and emotion focused strategies to cope with their disease.

Table 1: Sociodemographic characteristics of the studied group (N.102)

	No.	%
Age (in years)		
Range	18 – 80	
Mean+SD	52.9 ± 12	
Age groups		
18 – 40	17	16.7
40 – 60	57	55.9
60 – 80	28	27.5
Marital status		
Married	68	66.7
Single	5	4.9
Divorced	4	3.9
Widow	25	24.5
Address		
Rural	49	48.0
Urban	53	52.0
Education		
Illiterate & primary	63	61.8
Secondary	26	25.5
University	13	12.7
Occupation		
House wife	80	78.4
Employee	22	21.6

Table 2: Relation between COPE and depression (N.102)

Total depression level	Mean ± SD	P. value
Normal	50±12.1	0.143
Moderate	55.7±11.9	
severe	53.4±11.6	
Total	52.9±12	

Ns: No statistically significant difference (p level >0.05)

**Psychological Distress and Coping Strategies among Women with Breast Cancer receiving
Chemotherapy at Assiut University Hospital**

Table 3: Relation between COPE and anxiety (N.102)

Total Anxiety level	Mean ± SD	P. value
Normal	51.4±11.2	0.421
Moderate	50.4±14	
severe	54±11.6	
Total	52.9±12	

Ns: No statistically significant difference (p level >0.05)




Figure (1)

Table 4: Mean scores and standard deviations of coping strategies among women with breast cancer receiving chemotherapy(N.102)

Items	Range	Mean±SD
Self-distraction	2 – 8	5.8 ± 1.7
Active coping	2 – 8	6.7 ± 1.7
Denial	2 – 8	5.5 ± 1.9
Substance use	2 – 8	2.7 ± 1.3
Use of emotional support	2 – 8	6.8 ± 1.4
Use of instrumental support	2 – 8	6.5 ± 1.7
Behavioral disengagement	2 – 8	4.2 ± 1.8
Venting	2 – 8	5.7 ± 1.6
Positive reframing	2 – 8	6 ± 1.7
Planning	3 – 8	6.3 ± 1.6
Humor	2 – 8	5 ± 1.6
Acceptance	2 – 8	5.7 ± 1.8
Religion	3 – 8	6.8 ± 1.3
Self-blame	2 – 8	4.5 ± 2.2
Total depression	3 – 19	9.5 ± 3.3
Total Anxiety	5 – 20	11.8 ± 3.5
Brief COPE score	60 – 93	78.1 ± 7.6

Table 5: Relation between coping strategies, depression and anxiety symptoms among women with breast cancer receiving chemotherapy(N.102)

COPE Items	Total depression		Total Anxiety	
	r	P. value	r	P. value
Self-distraction	-0.21	0.038*	-0.25	0.012*
Active coping	-0.25	0.012*	-0.09	0.371
Denial	0.35	0.000**	0.47	<.001**
Substance use	0.07	0.494	0.19	0.060
Use of emotional support	-0.13	0.209	-0.17	0.093
Use of instrumental support	-0.31	0.002**	-0.28	0.004**
Behavioral disengagement	0.09	0.388	0.05	0.607
Venting	0.17	0.080	0.25	0.012*
Positive reframing	-0.32	0.001**	-0.37	<.001**
Planning	-0.37	0.000**	-0.16	0.109
Humor	-0.05	0.631	-0.02	0.870
Acceptance	-0.30	0.002**	-0.30	0.002**
Religion	-0.15	0.122	-0.13	0.208
Self-blame	0.19	0.058	0.23	0.019*
Brief COPE score	-0.24	0.014*	-0.10	0.319

* Statistically significant correlation (p level <0.05)** Statistically significant correlation (p level <0.01)

Discussion:-

Breast cancer patients on chemotherapy experienced high level of depressive and anxiety symptoms. These could be normal responses to the threat of the illness, uncertainty, side effects of chemotherapy agents, loss of control or an underlying psychiatric disorder such as depression and anxiety disorders.

The present study showed that more than half of the studied sample were from urban area and aged from 40 to 60 years old; more than two thirds of them were married. These findings were partially supported by, **Kelly et al (2008)** who reported that, breast cancer incidence rates in, Gharbiah, Egypt, during their study period was 31.8 % among women aged 40–49 years, and 29.8 % among women aged 50–59 years. Around 6.6 % of all breast cancer cases were diagnosed in women less than 40 years of age.

Zawilla ,(2011) ,stated that, in Egypt, breast cancer, it is the most common cancer among women, representing 18.9% of total

cancer cases 35.1% in women and 2.2% in men among the Egypt National Cancer Institute’s (NCI) series of 10,556 patients during the year 2001, with an age-adjusted rate of 49.6 per 100,000 people.

In this respect, Cancer in **Norway (2009)**, found that, the most common cancer diagnosis in women aged from 25 to 69 years old and constitutes 34% of all cancers in women aged 25 to 49 years old. It affected 2,745 women in 2009, representing 22% of all female cancers, and mainly affects women over 50 years old, Norwegian Breast Cancer **Group (2011)**. Also, **National Board of Health and Welfare (2002)**, reported that the risk of getting breast cancer increases significantly with age, and approximately 4700 women over 55 years developed the disease during 2002.

In addition, **Parkin, et al. (2002) and Soliman, et al. (2004)**, reported that, developing countries show a higher incidence of breast cancer in urban than in rural areas, a pattern that has not been fully explained. Other studies have shown that, women living in higher socio-economic status (SES) or urban communities have an increased risk of

developing breast cancer that cannot be explained by their exposure to individual risk factors (**Robert, et al. 2004**). In our opinion breast cancer incidence is higher in urban than in rural areas due to environmental and dietary factors, also breast cancer more common in middle age.

The present study, revealed that, more than one third of the study sample had severe depression, followed by moderate depression. While more than two third of them had severe anxiety. These findings are partially supported by, **Saniah and Zainal (2010)**, who reported that, Prevalence of depression was 19.1% and prevalence for anxiety was 24.1%. The total prevalence of psychological distress (i.e. depression and/anxiety) as determined by HADS among patients was 29%. In addition, **Ando et al. (2009)**, **Montgomery & McCrone (2010)**, reported that anxiety and/or depression assessed prior to diagnosis were significant predictors of psychological distress after breast cancer diagnosis.

In this respect, **Hanan ,etal(2014)**, reported that, breast cancer patients have high degrees of depression and anxiety (25.9% & 25.6%) respectively.

Joulaee et al, (2012) indicated that women's life as a spouse is very important in the family, and all intimate relations with men are in this format. For this reason, when a woman feels that she is not sexually attractive to her spouse, this might be the end of her sexual life. The spouse's reaction to the physical deficits caused by breast cancer may negatively affect women's self-esteem and confidence. These psycho-emotional problems can subsequently increase the physical problems.

In this respect. **Montgomery (2010)**, **Blow et al.(2011)**, **Steffens et al(2011)** found that, majority of women with breast cancer has been identified as stressful,

characterized by increased perceived stress, uncertainty, anxiety, fear and mood disturbance.

In our opinion higher incidence of depression and anxiety may be attributed to, breasts are viewed mostly as the women's unique sexual appearance in Muslim cultures; and greater concerns about body image and sexuality; some subsequent problems in relation to their spouse. Or may be attributed to women concerns about children; concerns about having children when faced with a life-threatening illness; premature menopause leading to infertility. Also, this may be due to, changes in their appearance has been a main concern primarily for younger women.

The current study showed that, women with breast cancer have used a wide variety of coping strategies as; religion, use emotional support, active coping, use instrumental support, planning, and positive reframing. The lowest scores were on: substance use, behavioral disengagement, and self-blame. Similar findings reported by, **Saniah and, Zainal (2010)**, who reported that, the highest scores were on religion, acceptance, use emotional support and use instrumental support. The lowest score were on substance use, behavioral disengagement and self-blame.

In addition **Culver et al. (2002)**, in investigating coping and distress among women undergoing treatment of early stage breast cancer, reported four main coping responses: use of humor, religious coping, self-distraction and venting. The investigators compared the coping responses among women from three ethnic groups; non-Hispanic Whites, Hispanic and African Americans. Non-Hispanic Whites reported more use of humor, Hispanic women used more, venting and self-distraction while religious coping was the common strategy among African Americans.

In this respect, a study by, **Taleghani, et al. (2006)**, showed that the majority of strategies used by Iranian women to cope with breast cancer were being positive on religious faith. Similarly, **Overcash, (2004)**, found that believing in God was of paramount importance for these women, which could help them in not giving up. Other researchers also suggest that the religious beliefs can be a supportive resource for people with cancer or other incurable diseases (**Taleghani et al. 2008**), but in Muslims, this is more pronounced.

This may be attributed to a positive relationship and full reliance on God gave participants hope that God was in control and that the disease was from him. This beliefs in God helped them to accept the diagnosis and cope with the disease. Also emotional support plays a key role in how women who are receiving a chemotherapy adjust and living with disease.

The current study demonstrated that, patients with anxiety symptoms had significantly higher score on, denial, positive reframing, and acceptance, use of instrumental support, self- distraction and self-blame as their coping strategies. While Patients who were depressed had significantly higher score on coping strategies such as planning, denial, positive reframing, acceptance, use of instrumental support, active coping, self-distraction, respectively. This finding is partially supported by; **Saniah and, Zainal (2010)** who reported that; patients with anxiety symptoms used denial, behavioral disengagement and venting. This may be explained by, Anxiety was considerably related with realistic, family, emotional problems and spiritual or religious concerns whereas depression was related with practical problems such as transportation, financial and work problems.

Similarly, **Lazarus & Folkman (1985)** stated that; denial in the form of avoiding all thoughts about the possible devastating

effects of cancer may particularly benefit some patients at the time of diagnosis. This may be attributed to, depressed patients tend to internalize their problems and blame them. This can be part of the symptoms of depression. Behavioral disengagement is a dimension that reduces one's effort to deal with the stressor, even giving up the effort to accomplish objectives with which the stressor is interfering. Also, **Folkman and Moskowitz (2004)**, reported that, "problem-focused coping" is associated with greater personal control than "emotion-focused coping".

This may be attributed to, broad difference in women's responses to cancer, with many women adjusting somewhat well and others having more problems. The inconsistency in adjustment to breast cancer diagnosis and treatment has led researchers to examine how factors such as demographics, disease and treatment characteristics, social support, and personality (e.g., coping style) are associated with adjustment. Of these factors, coping strategies in particular have been shown to be highly associated with adjustment and quality of life (QOL) after breast cancer diagnosis.

This may be explained by; denial may not eliminate negative mood states but may help a woman with breast cancer distance herself from negative thoughts and feelings, thereby fostering feelings of hope for a positive health outcome.

Conclusion:-

The highest mean score of the studied group used Problem-focused Strategies and emotional focused strategies, as a method of coping with anxiety and depression symptoms.

Women with breast cancer are vulnerable for anxiety and depression.

Recommendations:-

Designing and implementing psycho-educational programs about stressor related to the patients with breast cancer.

- Liaison psychiatric nurse should be available to deal with problems of patients with breast cancer.

References:-

Ando, Iwamitsu, Kuranami, Okazaki, Wada, Yamamoto, Todoroki, Watanabe & Miyaoka. (2009): Psychological characteristics and subjective symptoms as determinants of psychological distress in patients prior to breast cancer diagnosis. *Support Care Cancer* 17(11), 1361-1370

Ayres, Hoon, Franzoni, Matheny, Cotanch and Ballenger, Davidson, Lecrubier, (2001): Consensus statement on depression, anxiety and oncology. *J Clin Psychiatry*;62(8):64-67.

Blow, Swiecicki, Haan, Osuch, Symonds, Smith, Walsh & Boivin (2011) The emotional journey of women experiencing a breast abnormality. *Qualitative Health Research* 21(10), 1316-1334.

Cancer in Norway (2009): Cancer incidence, mortality, survival and prevalence in Norway. Retrieved from <http://www.kreftregisteret.no/en/Cancer-prevention/Breast-Cancer-Screening-Programme> on 22 May 2012.

Carayol, Bernard, Boiche,(2013). Psychological effect of exercise in women with breast cancer receiving adjuvant therapy: what is the optimal dose needed? *Ann Oncol.*;24(2):291-300.

Carver,(1997). You want to measure coping but your protocol's too long. Consider the Brief COPE. *Int J Behav Med*; 4 (1):92-100

Culver, Arena, Antoni, and Carver,(2002). coping and distress among women under treatment of early stage breast cancer: Comparing African Americans, Hispanic and Non-Hispanic Whites. *Psycho-oncology*; 11 495-505.

Del Mastro, Costantini, Morasso (2002). Impact of two different dose intensity chemotherapy regimens on psychological distress in early breast cancer patients. *Eur J Cancer*; 38: 359-66.

Folkman S. & Moskowitz (2004) Coping: pitfalls and promise. *Annual Review of Psychology* 55, 745-774.

Freedman, Edwards, Ries, and Young,(2006) ,Cancer Incidence in Four Member Countries (Cyprus, Egypt, Israel, and Jordan) of the Middle East Cancer Consortium (MECC) Compared with US SEER, NIH Publication No. 06-5873, National Cancer Institute, Bethesda, Md, USA,.

Hanan E. H., Magda M. M., Azza M El.A.,(2014), Emotional Distress Associated with Gynecologic and Breast Cancer in Beni-Suef City , *International Journal of Science and Research (IJSR)*, 6(14)

Janeen , Manuel, PhD; Stephanie , Burwell, Sybil , Crawford, Renee , Lawrence, Deborah , Farmer, Anita Hege, Kimberly, Phillips,Nancy , Avis,(2007), Younger Women's Perceptions of Coping With Breast Cancer, *Cancer Nurs.* ;30(2):85-94.

Jemal, Center, DeSantis, and Ward,(2010). "Global patterns of cancer

- incidence and mortality rates and trends,” *Cancer Epidemiology Biomarkers and Prevention*, vol. 19, no. 8, pp. 1893–1907, View at Publisher · View at Google Scholar · View at Scopus.
- John and MacArthur** (1998) Research Network, on Socioeconomic Status and Health .
- Joulaee , Joolae , Kadivar, & Hajibae,** (2012) Living with breast cancer: Iranian women’s lived experiences. *International Nursing Review*.
- Kearney, and Richardson,** (2006). *Nursing Patients with Cancer: Principles and Practice*, Elsevier, London .
- Kelly, Hirko, Amr Soliman, Ahmed Hablas, Ibrahim , Seifeldin, Mohamed Ramadan, Mousumi Banerjee, Joe , Harford, Robert , Chamberlain, and Sofia , Merajver,**(2013). Trends in Breast Cancer Incidence Rates by Age and Stage at Diagnosis in Gharbiah, Egypt, over 10 Years (1999–2008), *Journal of Cancer Epidemiology* Volume (2013), Article ID 916394, 7 pages.
- Lackey, Gates, ad Brown,** (2001). African Americans Women’s Experiences with the initial Discovery, Diagnosis and Treatment of Breast Cancer; *Oncol Nurs Forum*.;
- Lazarus & Folkman** (1985) . Coping and adaptation. In: Gentry WD (ed) *Handbook of Behavioral Medicine*. Guilford, New York.
- Lo Castro and Schlebusch** (2006), The measurement of stress in breast cancer patients. *S. Afr. J Psychol.*;36:102-119.
- Mohaghghi,** (10 October 2003) Posted in Family Physician – News. National Board of Health and Welfare (2002) Cancer incidence in Sweden ,2002. Stockholm: Socialstyrelsen, The National Board of Health and Welfare, Centre for Epidemiology, Official Statistics of Sweden.
- Montazeri,** (2008) Health related quality of life in breast cancer patients: Abibliographic review of the literature from 1974 to 2007. *Journal of Experimental and Clinical Cancer Research*, 27(1), 32-36.
- Montgomery & McCrone** (2010) Psychological distress associated with the diagnostic phase for suspected breast cancer: systematic review. *Journal of Advanced Nursing* 66(11), 2372-2390.
- National Board of Health and Welfare** (2002) Cancer incidence in Sweden 2002. Stockholm: Socialstyrelsen, The National Board of Health and Welfare, Centre for Epidemiology, Official Statistics of Sweden.
- Nor Zuraida, Ng CG,**(2010). Psychological distress among cancer patients on chemotherapy. *JUMMEC*; 13(1): 13-19.
- Norwegian Breast Cancer Group** (2011) *Brystkreft. Diagnostikk og behandling. En Veiledning (Breast Cancer. Diagnosis and Treatment. A Guide)*. Norsk kirurgisk Forening. Norsk Onkologisk Forening. Den Norske kreftforening. Oslo. Retrieved from <http://www.nbcbg.no/nbcbg.blaaboka.html#Anchor-49575> on 22 May 2012.
- Omar, Khaled, Gaafar, Zekry, Eissa, and El-Khatib,**(2003), “Breast cancer in Egypt: a review of disease presentation and detection strategies,” *Eastern Mediterranean Health Journal*, 9, (3) pp. 448–463,. View at Scopus.
- Overcash,** (2004) Using narrative research to understand the quality of life of older women with breast cancer. *Oncology Nursing Forum*, 31, 1153–1159.

- Parkin, Whelan, Ferlay, Teppo, Thomas,** (2002). Cancer incidence in five continents, volume VIII. International Agency for Research on Cancer, Lyon.
- Payne** (1990). Coping with palliative chemotherapy. *J Advance Nurs*;15(6): 652-658 .
- Robert, Strombom, Trentham-Dietz,** (2004). Socioeconomic risk factors of breast cancer: distinguishing individual- and community-level effects. *Epidemiology*;15(4):442-450.
- Saniah and Zainal** (2010), , Anxiety, Depression and Coping Strategies in Breast Cancer Patients on Chemotherapy MJP-02-08-10, University of Malaya, 50603 Kuala Lumpur, Malaysia .
- Shapiro , Boggs, Rodrigue, Urry, Algina, Hellman and Ewen** (1997). Stage II breast Cancer: Differences between four patterns in side effects during adjuvant chemotherapy. *J Psychosom Res*; 43(2):143-157.
- Shulman ,Willett, Sievers, and Knaul,** (2010) “Breast cancer in developing countries: opportunities for improved survival,” *Journal of Oncology*, vol. 2010, Article ID 595167, 6 pages,. View at Publisher · View at Google Scholar · View at Scopus.
- Soliman, Vulimiri, Kleiner** ,(2004). High levels of oxidative DNA damage in lymphocyte DNA of premenopausal breast cancer patients from Egypt. *Int J Environ Health Res*;14(2):121-34.
- Steffens , Wright , Hester & Andrykowski** (2011) Clinical, demographic, and situational factors linked to distress associated with benign breast biopsy. *Journal of Psychosocial Oncology* 29(1), 35-50.
- Takayanagis** (1994). Influence of mood and adjustment to cancer on compliance with chemotherapy among breast cancer patients. *J Psychosom Res*; 38 (5):393-402.
- Taleghani, Yekta, Nasrabadi** (2006). Coping with breast cancer in newly diagnosed Iranian women. *J Adv Nurs*;54:265-272
- Taleghani, Yekta, Nasrabadi, & Käppeli,** (2008) Adjustment process in Iranian women with breast cancer. *Cancer Nursing*, 31 (3), 32–41.
- Zabalegui** (1999). Coping strategies and psychological distress in patients with advanced cancer. *Onco Nurs Forum*; 26:1511-1518.
- Zawilla N. Breast cancer in Egypt: A fact sheet. *The Health*. 2011; 2 (1): 8- 10.
- Zigmund, Snaith,** (1983). The Hospital Anxiety and Depression Scale. *Acta Psychiatrica Scand.*; 67:361–370.
-