

Revisional Laparoscopic Mini-Gastric Bypass for Weight Loss Failure after Restrictive Procedures

Hossam El-Din Hassan Hussein, Ahmed Elnabil Mortada, Haitham Mostafa Elmaleh,
Ahmed Adel Abbas, Amir Nagui Abdalla Iskandar

General Surgery Department, Faculty of Medicine, Ain Shams University

Corresponding Author: Amir Nagui Abdalla Iskandar, E-mail: amir_nagui@live.com

ABSTRACT

Background: bariatric surgery has proven its effectiveness in achieving and maintaining weight loss and improving obesity-related co-morbidities, quality of life, and survival. As demand for bariatric surgery increases, so too will the need for revisional surgeries. The revision rate following primary bariatric surgery is reported to be between 10% and 25%. To facilitate weight loss surgically, many different types of bariatric procedures have been developed and established.

Objectives: to assess the effectiveness of revisional laparoscopic mini-gastric bypass for management of inadequate weight loss after different restrictive procedures.

Patients and Methods: the present study is a prospective study that was conducted at Ain Shams University Hospitals in Egypt, between July 2016 and July 2018. It included Forty (40) patients who underwent a restrictive bariatric procedure 2 years ago or more with inadequate weight loss. Operative time, intraoperative complications, rate of conversion. Postoperative pain, consumption of analgesics, length of hospital stay, start of oral feeding.

Results: the patients series involved 40 patients, 12 of which were male patients (40%) whereas, 24 were female patients (60%). The patients' age at the time of revisional surgery ranged from 23 to 57 years old with a mean \pm SD of 38.50 ± 8.42 years. The preoperative BMI ranged from 31.25 to 52.62 kg/m² with a mean \pm SD of 41.59 ± 4.99 kg/m² with an excess weight ranging from 19.79 to 86 kg with a mean \pm SD of 46.51 ± 14.27 kg. The mean time interval between the initial restrictive surgery and the revisional LMGB was 41.63 ± 16.92 months. 18 patients (45%) had a VBG as their primary restrictive surgery, 10 patients (25%) had LAGB, 8 patients (20%) had LSG and 4 patients (10%) had LGCP. 28 patients (70%) underwent LMGB for IWL compared to 12 patients (30%) for WR.

Conclusion: the revisional Laparoscopic Mini-Gastric Bypass (r-LMGB) appears to be a feasible and safe option after failed restrictive bariatric surgery. However, additional studies with larger population and longer follow up period are required to evaluate longer-term success.

Keywords: Revisional Laparoscopic Mini-Gastric, Weight Loss Failure, Restrictive Procedures

INTRODUCTION

Bariatric surgery has proven its effectiveness in achieving and maintaining weight loss and improving obesity-related co-morbidities, quality of life, and survival. As demand for bariatric surgery increases, so too will the need for revisional surgeries. The revision rate following primary bariatric surgery is reported to be between 10% and 25%⁽¹⁾. To facilitate weight loss surgically, many different types of bariatric procedures have been developed and established. They fall into three main categories: A) Restrictive procedures that lead to fixed or adjustable physical reduction in the size of the upper gastrointestinal tract. B) Malabsorptive procedures that bypass a proportion of the intestine with less physical restriction of food intake. C) A combination of the restrictive and malabsorptive, which combines restriction of the upper food pathway with intestinal bypass⁽²⁾.

There are four major restrictive surgical procedures that were used over the last years after patient selection, assessment, and evaluation. These

procedures are the Vertical Banded Gastroplasty (VBG) which is no longer performed nowadays, the Laparoscopic Adjustable Gastric Banding (LAGB), the Laparoscopic Greater Curvature Gastric Plication (LGCP) and the Laparoscopic Sleeve Gastrectomy (LSG)⁽³⁾. The Vertical Banded Gastroplasty (VBG), involved vertically partitioning the stomach at the angle of His through a window created near to the lesser curvature at the base of the pouch. A silastic ring was then placed around this window to secure the narrow, tabularized stomach reservoir. However, another technique has been adapted by some surgeons by using polypropylene mesh instead of the ring which increased the incidence of dense adhesions and subsequently adding to the technical challenge of revisional surgery. The VBG eventually was abandoned in favor of other operations such as the Adjustable Gastric Banding⁽⁴⁾.

The Adjustable Gastric Banding is placed around the upper stomach to create a small proximal pouch. This produces a moderate restriction in the volume and type of foods the

patient is able to consume. The band also delays the emptying of the pouch, creating the sensation of satiety ⁽⁵⁾. The Laparoscopic Sleeve Gastrectomy (LSG) involves reduction of the size of the stomach by about 75% of its original size by surgically removing a large portion of it after insertion of a special bougie to calibrate a narrow tubular sleeve ⁽⁶⁾. It was found to be useful beyond its restrictive capabilities with physiologic alterations that affected metabolic processes and decreased hunger. This led to an increase of the popularity of the LSG in the late 2000s, and it rapidly replaced the LAGB as the preferred restrictive weight loss operation ⁽⁷⁾.

Laparoscopic greater curvature gastric plication (LGCP) is gaining ground in the treatment of morbid obesity, to avoid the placement of an implantable device or the irreversible resection of gastric tissue. Current technique of LGCP consists of infolding the greater curvature to reduce stomach volume by placement of rows of non-absorbable sutures ⁽⁸⁾. The reasons for revision after restrictive procedures have been reported to be unsatisfactory weight loss or intolerable adverse outcomes such as emesis, maladaptive eating syndrome, and gastroesophageal reflux disease (GERD) ⁽⁹⁾. Weight loss failure after bariatric surgery is defined as achieving or maintaining less than 50% of excess weight loss (EWL) over a period of 18 to 24 months or a body mass index (BMI) of greater than 35 over the same period ⁽¹⁰⁾.

There are specific causes of failure for each restrictive procedure, in VBG, most common causes of failure are pouch dilatation, stoma dilatation and disruption of staple line causing gastrogastic fistula ⁽¹¹⁾. In LAGB, causes include band slippage, erosion into the stomach, esophageal dilation, intolerance to the device, and ultimately a decreased quality of life ⁽¹²⁾. Failure after LSG and LGCP is mainly due to dilation of the gastric lumen, this may be attributed to a lack of adequate calibration at the time of the primary procedure or a natural process of stomach tissue to dilate and become more compliant over time ⁽¹³⁾. Various revisional options have been suggested, including biliopancreatic diversion (BPD), roux en-y gastric bypass (RYGB), minigastric bypass (MGB) or reestablishing restriction. In the latter category, the options include sleeve gastrectomy and repositioning of adjustable gastric band. Although RYGB is considered by many as the gold standard revisional procedure after failed restrictive surgery, MGB is

becoming an accepted alternative with satisfying outcome and similar long-term efficacy ⁽¹⁴⁾.

The minigastric bypass (MGB) consists of two components, including, first, a long sleeved gastric tube along the lesser curvature side and, second, a Billroth type II loop gastrojejunostomy with a 200 - 250 cm afferent limb. Some surgeons proposed placing an anchoring suture on the afferent limb as a valve to inhibit bile reflux ⁽¹⁵⁾. Short and long-term results after primary laparoscopic minigastric bypass (MGB) have been previously reported and MGB appears to be a simple, well tolerated, and effective alternative to RYGB. However, long-term outcomes for revisional MGB for a failed restrictive procedure (rMGB) have not yet been analyzed.

AIM OF THE WORK

Our objective in this study is to assess the effectiveness of revisional laparoscopic minigastric bypass for management of inadequate weight loss after different restrictive procedures.

PATIENTS AND METHODS

The present study is a prospective study that was conducted at Ain Shams University Hospitals in Egypt, between July 2016 and July 2018. It included **Forty (40) patients** who underwent a restrictive bariatric procedure 2 years ago or more with inadequate weight loss. **The study was approved by the Ethics Board of Ain Shams University and an informed written consent was taken from each participant in the study.**

Inclusion Criteria: Patients who underwent a restrictive bariatric surgery 2 years ago or more, with one or more of the following: Failure to achieve or maintain more than 50% of Excess Weight Loss (EWL) over a period of 2 years after restrictive bariatric surgery. Body mass index (BMI) of greater than 35 over a period of 2 years after restrictive bariatric surgery. Regain of 10% or more of the nadir weight in 1 year despite dietary counseling. Increase in Residual Gastric Volume (RGV), as assessed by radiology and/or gastroscopy.

Exclusion Criteria: Patients unfit for general anesthesia (ASA III, IV or V). Patients with failure due to intolerable adverse outcomes such as emesis, maladaptive eating syndrome, and gastroesophageal reflux disease (GERD). Patients with anatomical complications as band erosion, ring erosion, postoperative leakage. Patients with

inadequate weight loss who didn't have any dietary counseling yet. Patients with psychiatric illnesses receiving anti-psychotic medications.

Pre-operative assessment:

Full clinical history.

Outcome Assessment: Operative time, intraoperative complications, rate of conversion. Postoperative pain, consumption of analgesics, length of hospital stay, start of oral feeding. Weight

loss in term of percentage excess weight loss (%EWL), BMI at 1, 3, 6, 12, 18 and 24 month. Resolution of comorbidities.

Statistical Analysis

The collected data was revised, coded, tabulated and introduced to a PC using Statistical Package for Social Science (IBM Corp. Released 2011. IBM SPSS Statistics for Windows, Version 20.0. Armonk, NY: IBM Corp).

RESULTS

Table (1): Distribution of cases according to comorbidities

Parameter	No.	%
Hypertension	8	20.0
Type II Diabetes mellitus	7	17.5
Dyspnea on exertion	32	80
Knee pain	23	57.5
Back pain	29	72.5
Arthritis	3	7.5

The comorbidities (table 1) were diabetes mellitus (type II) in 7 patients (17.5%) all were on medical treatment, hypertension in 8 patients (20%) all were on antihypertensive drugs, dyspnea on excretion in 32 patients (80%), arthritis in 3 patients (7.5%), knee pain in 23 patients (57.5%) and back pain in 29 patients (72.5%) all of them had continuous pain that needed medications or physiotherapy and had no serious cause.

Table (2): Early postoperative complications

Parameter	No.	%
Port site infection	0	0.0
Bleeding	1	2.5
Stenosis at gastrojejunostomy	0	0.0
Leakage from gastric pouch staple line	1	2.5
Wound infection	1	2.5
Lung atelectasis/ chest infection	2	5.0
Port-site hernia	0	0.0
Deep venous thrombosis	0	0.0
Reoperation	1	2.5
Mortality	0	0.0

We had no anastomotic stenosis, port-site infection, port-site hernia or DVT in the early postoperative period. There was zero mortality in the early postoperative period (table 2).

Table (3): Late postoperative complications

Parameter	No.	%
Biliary reflux	3	7.5
Anemia	5	12.5
Hair loss	8	20.0
Gastrojejunostomy Stenosis/ulcer	0	0.0
Internal hernia	0	0.0
Neuropathy	3	7.5
Gall stones	4	10.0
Mortality	0	0.0

We had no anastomotic stenosis / ulceration or internal hernia in the late postoperative period. There was zero mortality in the late postoperative period (table 3).

Table (4): Patients' weight in different follow up measurements

	Weight		Test value*	P-value	Sig.
	Mean \pm SD	Range			
Pre-operative	117.08 \pm 16.67	93 – 167	NA	NA	–
1 month	112.39 \pm 16.01	89.28 – 160.32	44.412	0.000	HS
3 months	105.44 \pm 15.04	83.7 – 150.3	43.606	0.000	HS
6 months	100.88 \pm 14.44	80.35 – 144.29	38.549	0.000	HS
12 months	95.03 \pm 13.48	75.53 – 135.63	42.762	0.000	HS
18 months	91.20 \pm 12.93	72.51 – 130.21	42.708	0.000	HS
24 months	90.31 \pm 12.93	71.93 – 129.16	42.753	0.000	HS

*: Paired t-test; HS: Highly significant; S: Significant; NS: Non significant; P-value < 0.01 Highly significant; P-value < 0.05 Significant; P-value > 0.05 Non significant

Table (4) shows weight loss after the r-LMGB compared to the pre-revisonal weight throughout the 24 months follow up period; the mean weight decreased from 117.08 kg \pm 16.67 kg preoperatively to 112.39 kg \pm 16.01 kg at one month, 100.88 kg \pm 14.44 kg at six months, 95.03 kg \pm 13.48 kg at 12 months and reached as low as 90.31 kg \pm 12.93 kg at the end of the 2 years follow up period.

DISCUSSION

As morbid obesity is a chronic, life-long disease, an effective treatment should entail a multi-interventional approach with a lifetime follow-up of intensive consultations for nutritional support, physical and psychosocial support, and reoperation if necessary. The benefits of bariatric re-interventions have to dramatically outweigh the increased adverse outcomes and the higher complication rates of revision procedures ⁽¹⁶⁾.

Failure of adequate weight loss and weight regain has been reported to be as great as 25–30% after restrictive bariatric procedures. Weight regain after bariatric procedures is usually multifactorial and warrants psychological and dietary assessments. The reasons for revision after restrictive procedures have been reported to be due to insufficient weight loss, weight regain, and/or surgery-related complications ⁽¹⁰⁾.

Weight loss failure after bariatric surgery is defined as achieving or maintaining less than 50% of excess weight loss (EWL) over a period of 18 to 24 months or a body mass index (BMI) of greater than 35 over the same period ⁽¹⁰⁾.

Different factors such as lifestyle, mental health issues, hormonal/metabolic imbalance and technical issues after surgery may contribute to failure of different restrictive bariatric procedures. Thus, patients must be investigated and treated in the

specialized centers where multidisciplinary teams are available and knowledge about IWL is accumulated. The mini-gastric bypass (MGB) was introduced by Rutledge in 1997 and reported some years later. Since then, thousands of patients have been treated with this approach by several authors in different countries ⁽¹⁷⁾.

Short and long-term results from primary laparoscopic Mini-Gastric Bypasses (LMGB) have been previously reported, and LMGB appears to be a simple, well tolerated, and effective alternative to LRYGB. However, long-term outcomes for revisional LMGB for a failed restrictive procedure (r-LMGB) have not yet been heavily analyzed ⁽¹⁸⁾.

The aim of our study was to assess the effectiveness of revisional laparoscopic mini-gastric bypass for management of inadequate weight loss after different restrictive procedures. It is a prospective study which was done between July 2016 – July 2018, at Ain Shams University Hospitals, Cairo, Egypt. The study included 40 patients who underwent a restrictive bariatric procedure 2 years ago or more with inadequate weight loss. The mean time interval between the initial restrictive surgery and the revisional LMGB was 41.63 \pm 16.92 months. 18 patients (45%) had a VBG as their primary restrictive surgery, 10 patients (25%) had LAGB, 8 patients (20%) had LSG and 4 patients (10%) had LGCP. 28 patients (70%) underwent LMGB for IWL compared to 12 patients (30%) for WR.

Regarding the operative time and conversion to open surgery; in our study, the mean operative time was 118 minutes ranging between 77-188 minutes and all operations were performed laparoscopically except one case which was converted to open due to increased airway pressure with abdominal insufflation. The mean BMI of our patients was 41.59 kg/m².

Compared to the published results, a study conducted by Bruzzi and his colleagues (to assess 5-year outcomes of r-LMGB after failure of restrictive surgery) included 30 patients with mean BMI of 45.5 kg/m², the mean operative time was 140 minutes with no conversion to open⁽¹⁹⁾. Another study done by Almalki and his colleague (which compared between LMGB and LRYGB as a revision for failed restrictive procedures) included 81 patients in the r-LMGB group with a mean BMI of 37.8 kg/m², the mean operative time with r-LMGB was 168 with no conversion to open⁽²⁰⁾.

Our outcome relatively matches the results of previous studies such as *Bruzzi et al.*⁽¹⁹⁾ who reported a BMI decrease of 15 kg/m² and a %EWL of 65% after 24 months of follow-up of 30 patients who underwent r-LMGB, after failure of restrictive procedure, with a pre-revisional mean BMI of 45.5 kg/m².

Similarly, *Almalki et al.*⁽²⁰⁾ study showed, 81 patients with a mean BMI of 37.8 kg/m² who underwent r-LMGB as a group in a comparative study between LMGB and LRYGB as a revisional procedure after failure of restrictive procedure. At 24 months postoperative, the mean BMI reached was 27.2 kg/m² with a total BMI decrease of 10.6 kg/m² and a %EWL of 76.8%.

The total rate of complications in our study was comparative to other studies. Our study showed a rate of 12.5% compared to 10% in *Bruzzi et al.*⁽¹⁹⁾ study and 16% in *Almalki et al.*⁽²⁰⁾ study.

As regard the late postoperative complications; 3 cases (7.5%) with biliary reflux were reported in our study and were managed successfully with conservative treatment, in contrast to *Bruzzi et al.*⁽¹⁹⁾ study that reported 2 cases (6.66%) with intractable bile reflux that were converted to RYGB. No late complications were reported in *Almalki et al.*⁽²⁰⁾ study.

CONCLUSION

The revisional Laparoscopic Mini-Gastric Bypass (r-LMGB) appears to be a feasible and safe option after failed restrictive bariatric surgery.

However, additional studies with larger population and longer follow up period are required to evaluate longer-term success.

REFERENCES

1. **Kellogg TA (2011):** Revisional bariatric surgery. *Surg Clin North Am.*, 91(6):1353–71.
2. **Saheba I (2012):** Manual of Perioperative Care, chapter 29: Bariatric Surgery, John Wiley & Sons.
3. **Moises J, Eddie G, Roderick R et al. (2011):** Failed Restrictive Surgery: Is Sleeve Gastrectomy a Good Revisional Procedure? *Obes Surg.*, 21:157–160.
4. **Adrian G and Rebecca L (2015):** Minimally Invasive Bariatric Surgery, chapter 4: History of Bariatric and Metabolic Surgery, Springer, second edition.
5. **O'Brien PE, McDonald L, Anderson M and Brown WA (2013):** Long term outcomes after bariatric surgery: Fifteen year follow up of adjustable gastric banding and a systematic review of the bariatric surgical literature. *Ann Surg.*, 257(1):87–94
6. **Stephenson JA, Al-Ta'an O, Wiggins T, Sutton CD et al. (2010):** The surgical management of obesity. *InnovAiT.*, 3(10): 598–605.
7. **Braghetto I, Csendes A, Lanzarini E et al. (2012):** Is laparoscopic sleeve gastrectomy an acceptable primary bariatric procedure in obese patients? Early and 5-year postoperative results. *Surg Laparosc Endosc Percutan Tech.*, 22: 479–486.
8. **Almino R, Lyz Silva, Manoel N and Josemberg C (2015):** Laparoscopic Gastric Plication, Minimally Invasive Bariatric Surgery, Springer Science, 179-180.
9. **Schouten R, Wiryasaputra DC, van Dielen FM et al. (2010):** Long-term results of bariatric restrictive procedures: a prospective study. *Obes Surg.*, 20:1617–26.
10. **Chang VW and Lauderdale DS (2014):** Income disparity in body mass index and obesity in the United States: 1971-2002. *Arch Intern Med.*, 165(18):2122-2128.
11. **Samir A, Eran S, Ilanit M et al. (2015):** Laparoscopic Conversion of Failed Silastic Ring Vertical Gastroplasty (SRVG) and Vertical Banded Gastroplasty (VBG) Into Biliopancreatic Diversion (BPD); *J Gastrointest Surg.*, 19:625–630.

12. **Shen X, Zhang X, Bi J and Yin K (2015):** Long-term complications requiring reoperations after laparoscopic adjustable gastric banding: a systematic review. *Surg Obes Relat Dis.*, 11(4):956–64.
13. **Cesana G, Uccelli M, Ciccarese F et al. (2014):** Laparoscopic re-sleeve gastrectomy as a treatment of weight regain after sleeve gastrectomy. *World J Gastrointest Surg.*, 6(6):101–6.
14. **Elnahas A, Graybiel K, Farrokhyar F, Gmora S et al. (2013):** Revisional surgery after failed laparoscopic adjustable gastric banding: A systematic review. *Surg Endosc.*, 27(3):740–5.
15. **Kim Z and Hur KY (2011):** Laparoscopic mini-gastric bypass for type 2 diabetes: the preliminary report. *World J Surg.*, 35(3):631–6.
16. **Steffen R (2008):** The history and role of gastric banding. *Surg Obes Relat Dis.*, 4(3):S7–13.
17. **Hamdi A, Julien C, Brown P et al. (2014):** Midterm Outcomes of Revisional Surgery for Gastric Pouch and Gastrojejunal Anastomotic Enlargement in Patients with Weight Regain After Gastric Bypass for Morbid Obesity. *Obes Surg.*, 24: 1386- 1390.
18. **Lee WJ, Yu PJ, Wang W, Chen TC et al. (2005):** Laparoscopic Roux-en-Y versus mini-gastric bypass for the treatment of morbid obesity: a prospective randomized controlled clinical trial. *Ann Surg.*, 242(1):20–8.
19. **Bruzzi M, Voron T, Zinzindohoue F, Chevallier JM et al. (2016):** Revisional single-anastomosis gastric bypass for a failed restrictive procedure: 5-year results. *Surg Obes Relat Dis.*, 12:240–5.
20. **Almalki WJL, Jung-Chien C, Kong-Han S et al. (2018):** Revisional Gastric Bypass for Failed Restrictive Procedures: Comparison of Single-Anastomosis (Mini-) and Roux-en-Y Gastric Bypass, *Obes Surg.*, 28:970–975.