



## Original article

# Linear regression analysis showing the predictors of mortality in COVID-19

Hamzullah khan<sup>1\*</sup>, Khalida Rehman<sup>2</sup>

1- Associate Professor Hematology, Nowshera Medical College, MTI Nowshera, Pakistan.

2- Medical Officer, Department of Pathology, Nowshera Medical College, Nowshera, Pakistan.

### REPORT INFO

#### Report history:

Received 3 August 2020

Received in revised form 17 August 2020

Accepted 23 August 2020

#### Keywords:

Linear Regression

Predictor

Mortality

COVID-19

### ABSTRACT

To quantify the role of age, gender, exposure and rate of infection as predictors of mortality in corona virus disease 2019 (COVID-19), a cross sectional study was conducted covering 219 patients who were tested positive out of 554 suspects in Qazi Hussain Ahmed Medical Complex Nowshera, Pakistan. Linear regression analysis was used as a statistical tool. We observed that the positive contact history/exposure has been shown to have strong predictive value for mortality due to COVID-19 ( $\beta=0.394^{**}$ ,  $\Delta R^2=0.011$ ,  $p=0.001$ ). Similarly, age>60 years and rate of infection were also strong predictors of mortality with interaction values of ( $\beta=0.389^{**}$ ,  $\Delta R^2=0.046$ ,  $p=0.001$ ) and ( $\beta=0.431^{**}$ ,  $\Delta R^2=0.005$ ,  $p=0.03$ ), respectively. Gender as such is not a statistically significant predictor in mortality due to COVID-19. We concluded that using this regression analysis: the age of patients and positive exposure irrespective of gender statistically predicts the mortality in COVID-19.

### Introduction

Corona virus disease 2019 (COVID-19) was first reported from metropolitan city, Wuhan, Hubei province of China in Dec 2019, that causes severe respiratory disease/pneumonia. The etiology of COVID-19 is yet to be confirmed, but majority of the scientists agree that it most likely originated from the zoonotic corona virus, Severe Acute Respiratory Syndrome coronavirus (SARS-CoV) that emerged in 2002 [1].

Corona Virus disease termed as COVID-19, is an emerging highly contagious respiratory disease that is caused by novel corona virus. Its main clinical symptoms are fever, dry cough, fatigue, myalgia and dyspnea. Case fatality rate of 2.3% has been reported from china that is lower than SARS (9.5%), Middle East Respiratory

Syndrome (MERS) (34.4%) and Hemophilus Influenza (39%) [2].

In Pakistan the literature so far covering the prevalence and incidence is deficient. In Pakistan the so far (24<sup>th</sup> July 2020) reported data from government sources declares 270,400 confirmed cases with 5763 deaths. 219783 cases are recovered so far, and 1316 cases are under treatment in high dependency units countrywide. Sindh is the province with highest number of corona cases crossing 115883 [3].

Qiu H et al [4] reported from the infected zone of China that strong suspects from among the family members and other close contacts of COVID-19 patients, the infectivity ratio was as higher as 89% while it was (33%) for history of exposure to epidemic areas. Regarding mortality in age groups a study from China

reported 80% of the casualties (deaths) due to COVID-19 were in the adults aged >60 years [5].

### Objectives of the study and Methods

To quantify the role of age, gender, exposure and rate of infection as predictors of mortality in COVID-19, we did a cross sectional study covering 219 patients who were tested positive out of 554 suspects in Qazi Hussain Ahmed Medical Complex Nowshera, Pakistan from 10<sup>th</sup> March to 20<sup>th</sup> July 2020. Linear regression test was used using SPSS version 25 with task to predict a positive link of dependent variable, input (like infection (test positivity, exposure) with independent variable, output (Mortality) in COVID-19.

### Results and Discussion

We observed that the positive contact history/exposure has been shown to have strong predictive value for mortality due to COVID-19 as supported by the regression analysis. Beta value, i.e., ( $\beta=0.394^{**}$ ), showed that there was a positive link between exposure and mortality due to COVID-19 and one unit increment in the exposure will cause a 39.4% increment in the rate of mortality due to COVID-19. The value of  $R^2$ , i.e. ( $R^2=0.011$ ), shows that exposure is accounted for the variability of 1.1% in the cases of mortality. The p-value which is ( $p=0.001$ ), supported the statistically significant relationship (**table 1**).

Similarly, **Anzai A et al** [6] reported 33% of the infectivity in suspects with strong contacts but not with a contact directly COVID-19 patient. **Shi Y et al** [7] reported from Wuhan city, a significant relation of close contacts gets more infected from family members ( $p=0.031$ ) with COVID-19 that strongly supports our findings.

Mortality due to COVID-19 positively moderates with rate of infection supported by the results of an interaction term with ( $\beta=0.431^{**}$ ;  $\Delta R^2=0.005$ ,  $p\text{-value}=0.03$ ). The stated results showed that rate of infection positively associates with case fatality due to COVID-19 (**table 1**). Italy has been worst hit by the COVID-19, measuring 185000 cases mostly reported from the northern cities of the country. By April 2020, 45% increase in deaths were registered with a higher number being reported from the north of the country (76%) of the total. Most of them were men with an increasing trend by age with a mortality rate of 7.2% [8,9]. The severity of the infection has an impact on mortality rate. A study

reported from Wuhan, the epicenter of the first outbreak reported mortality in positive patients as higher as 20% [10].

Mortality due to COVID-19 positively moderates with age>60 supported by the results of an interaction term with ( $\beta=0.389^{**}$ ;  $\Delta R^2=0.046$  and  $P\text{ value}=0.001$ ). We concluded in one other intervention published in Microbes and infectious disease (MID) that the rate of infectivity and exposure to come across in contacts with positive patients with a probability of 2 times more than in people with age<55 years ( $p=0.005$ ,  $OR=2.01$ ) while the case fatality was three times more in patients with age >55 years ( $p=0.001$ ,  $OR=3.16$ ). The mortality rate was 5.41% [11].

Regarding the gender effect we observed that one unit increase in male gender exposure will cause 1.1% increase in rate of mortality as compared to female gender, which shows no clinically and statistically significant difference in gender groups ( $\beta=0.011$ ;  $\Delta R^2=0.001$ ,  $p=0.11$ ) (**table 1**). However, the literature from Italy reporting the infection being more common in male gender as compared to female gender with a higher mortality in the patient with age> 60 years [9]. The age vs mortality graph shows majority of the deaths occurred at age> 50 years (**figure 1**).

Hence we concluded that using this regression analysis we observed that age of patients, and positive exposure irrespective of gender statistically predicts the worst outcome in term of death in COVID-19 and this must be considered by the treating physician and treatment strategies should be age and risk exposure based to have hope for better outcome in COVID-19.

Therefore, in future it is the need of time to better understand the impact of gender and age in COVID-19 and to tailor the treatment according to the age and gender perspective and the therapeutic trials must include gender sensitive analysis.

**Conflict of interest:** None declared


**Authorship:** Both authors had read and approved the final version of the manuscript.

**Financial disclosure:** None declared

**Table 1.** Linear regression analysis to predict mortality in COVID-19 against different variables like age, gender, exposure and rate of infection.

Name of predictor	B ( $\beta$ -value)	$\Delta R^2$ (Variability)	Significance
Positive Contact History	0.394**	0.011	0.001
PCR Positivity (Rate of infection)	0.431**	0.005	0.03
Age >60 years	0.389**	0.046	0.001
Male gender	0.011	0.001	0.11

Significance at  $p=0.01$ \*\*, Significance at  $p=0.05$ \*

**Figure 1.** Age vs mortality graph.

## References

- Ahn DG, Shin HJ, Kim MH, Lee S, Kim HS, Myoung J, et al. Current Status of Epidemiology, Diagnosis, Therapeutics, and Vaccines for Novel Coronavirus Disease 2019 (COVID-19). J Microbiol Biotechnol 2020; 30(3): 313-324.
- Chen N, Zhou M, Dong X, Qu J, Gong F, Han Y, et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. Lancet 2020; 395: 507-13.
- Government of Pakistan. Coronavirus in Pakistan Confirmed Cases. Available at: [www.COVID.gov.pk/](http://www.COVID.gov.pk/). Retrieved 24 July 2020.
- Qiu H, Wu J, Hong L, Luo Y, Song Q, Chen D. Clinical and epidemiological features of 36 children with coronavirus disease 2019 (COVID). The Lancet 2020; 20(6): 689-696.
- Centers for disease control and prevention (CDC). COVID-19 Response Team. Severe Outcomes Among Patients with Coronavirus Disease 2019 (COVID-19) - United States,

- February 12-March 16, 2020. MMWR Morb Mortal Wkly Rep 2020; 69(12): 343-346.
- 6-**Anzai A, Kobayashi T, Linton NM, Kinoshita R, Hayashi K, Suzuki A, et al.** Assessing the Impact of Reduced Travel on Exportation Dynamics of Novel Coronavirus Infection (COVID-19). *J Clin Med* 2020; 9(2):601.
- 7-**Shi Y, Yu X, Zhao H, Wang H, Zhao R, Sheng J.** Host susceptibility to severe COVID-19 and establishment of a host risk score: findings of 487 cases outside Wuhan. *Crit Care* 2020; 24(1): 108.
- 8-**Presidenza del Consiglio dei Ministri - Dipartimento della Protezione Civile.** COVID-19 dati-regioni. [COVID-19 Italy: regional data digital repository]. Rome: Department for Civil Protection. Available at: <https://github.com/pcm-dpc/COVID-19/tree/master/dati-regioni>. Accessed Apr 13, 2020].
- 9-**Michelozzi P, de'Donato F, Scortichini M, De Sario M, Noccioli F, Rossi P, et al.** Mortality impacts of the coronavirus disease (COVID-19) outbreak by sex and age: rapid mortality surveillance system, Italy, 1 February to 18 April 2020. *Euro Surveill* 2020; 25(19): 2000620.
- 10-**Mizumoto K, Chowell G.** Estimating the risk of 2019 novel coronavirus death during the course of the outbreak in China, 2020. medRxiv. 2020.DOI: <https://doi.org/10.1101/2020.02.19.20025163>
- 11-**Khan H, Hussain S, Zar A, Fawad, S.** Analysis of demographic variables in acquiring infection and mortality due to COVID-19. *Microbes and Infectious Diseases* 2020; 1(2): 70-76.