

INFLUENCE OF NITROGEN SOURCES ON Yield AND ANATOMICAL CHARACTERS OF DAMSISA (*Ambrosia maritime*, L). PLANTS UNDER SANDY SOIL CONDITIONS

Matter, F. M. A.*; R. A. Agamy** and S.S. El-Ganaini**

* Dept. of Horticulture, Fac. of Agric. Fayoum University. Egypt.

** Dept. of Botany, Fac. of Agric. Fayoum University. Egypt.

ABSTRACT

Effects of the three nitrogen sources (namely, ammonium nitrate, ammonium sulphate and urea) each at the rates of 0, 50, 100 and 200 kg/fed., on vegetative, yield, anatomical characters and chemical constituents of damsisa plants were studied. The experiment was carried out as a pot experiment in Faculty of Agriculture, Fayoum University during the two successive seasons (2006/2007 and 2007/2008). The results obtained showed that, all sources of nitrogen fertilizer greatly increased growth parameters. All treatments (mainly, urea and ammonium sulphate) greatly increased chlorophyll a, b, carotenoids in the leaves and total carbohydrates, N, P and K contents of herb.

All treatments (especially, ammonium sulphate) increased stem section diameter, due to the increase in paraenchymatous and vascular tissues. All N sources greatly increased section diameter of leaf petiole, midvein thickness, blade thickness and dimensions of the vascular bundle as well as diameter of xylem vessels. Treating plants by ammonium sulphate greatly increased total sesquiterpene lactones. The greatest increase was observed on most previous botanical characters by ammonium sulphate. Thus, it can be recommended to use ammonium sulphate as nitrogen fertilizer for improving a yield and quality of damsisa plants under sandy soil conditions.

Keywords: Damsisa, Nitrogen fertilizer, Growth and Yield.

INTRODUCTION

Damsisa (*Ambrosia maritime* L). is one of the medicinal plants used colic and to expel renal stones (Mahran, 1967). In addition, damsisa usually grown near the bank of canals, where the snails of bilharziasis escape far from these plants and even it can kill some of these snails. On the other side, *Ambrosia maritime* is known to have molluscicidal properties, it is also exhibit an insecticidal activity against the larvae and pupae of *Anopheles pharoensis* (Teriest *et al.*, 1989).

Many investigators reported the importance of nitrogen fertilization for medicinal and aromatic plants. However, fertilization treatments at suitable level and source showed primitive effects on the plant growth, root system characters as well as active ingredients.

In this respect, Moa and Craker (1991) stated that an adequate supply of nutrients particularly nitrogen is one of the several factors responsible for increasing essential oil yield as well as vegetative growth. Also, this fact reported by several researchers El-Saeid *et al.*, (1996) on *Tagetes patula*, Hassanein *et al.*, (1998) on *Pelargonium graveolens*, Mansour *et al.*, (1999) on spearmint and major am plants and Mohamed and

Matter (2001) on marigold. Also, Jimenez *et al.* (2006) they found that the use of different nitrogen (N) forms ($\text{NH}_4^+\text{-N}$ or $\text{NO}_3^-\text{-N}$) shows benefits and limitations. For example, one of the advantages of ($\text{NH}_4^+\text{-N}$) is that it is fixed to the soil or substrate particles thanks to the cation exchange capacity of these materials and, therefore, it is not leached, decreasing ($\text{NO}_3^-\text{-N}$) pollution. Most plants can use both N forms, but the use efficiency and the preference for each form depend on the species, variety and plant age (Abbes *et al.*, 1995). Nitrogen and P uptake rates are higher in the plants supplied with NH_4^+ or $\text{NO}_3^- + \text{NH}_4^+$ than in plants provided NO_3^- alone (Jimenez *et al.*, 2007).

The present investigation was undertaken to study the effect of nitrogen source on the vegetative growth, chemical composition and anatomical characters of damsisa plants under sandy soil conditions.

MATERIALS AND METHODS

This investigation was conducted as a pot experiment during the two successive growing seasons of 2006/2007 and 2007/2008 at the Experimental Station, Faculty of Agriculture, Fayoum University to study the effect of nitrogen source on the vegetative growth, chemical composition and anatomical characters of damsisa (*Ambrosia maritime*, L) plants under sandy soil conditions.

Pots (30cm diameter and 50cm height) were filled with air dried sandy soil. The physical and chemical analysis of the tested soil used was carried out according to Olsen and Sommers (1982) and Page *et al.*, (1982).

Seeds of damsisa were sown in prepared seed beds on September 15th for both seasons. Uniform seedlings of 10-15cm length were transplanted into pots after 50 days from sowing.

The study included three sources of nitrogen; ammonium nitrate (33.5%N), ammonium sulphate (20%N) and urea (46.5%N). Four nitrogen levels were used for each (0, 50, 100 and 200 kg/fed.).The amounts of nitrogen fertilizers were divided into two equal doses and they were added twice during the growing seasons. The first one was one month after transplanting and the second was one month later. Control treatment received no nitrogen fertilizer. All the agricultural practices cultural practices for growing damsisa plant prevailing in the locality were followed.

Table (1): Some physico-chemical characteristics of the tested soil.

Physical properties												
Coarse sand			Fine sand		Silt	Clay	Soil texture					
70.98			22.99		2.33	3.70	Sandy					
Chemical properties												
O. matter%	EC dsm ⁻¹	pH	N %	P %	CaCO ₃ %	Ca ⁺⁺	Mg ⁺⁺	N ^{a+}	K ⁺	HCO ₃ ⁻	Cl ⁻	SO ₄ ⁻
0.67	4.5	7.5	0.06	0.07	10	0.52	0.12	0.04	0.02	0.05	0.48	0.55

The present experiment contained 10 treatments were distributed in a complete randomized block design with three replicates, each replicate contained five pots (one plant pot⁻¹). At the flowering stage, the following data were recorded as follows:

a) Vegetative growth characters per plant, i.e., plant height (cm), number of branches plant⁻¹, number of inflorescences plant⁻¹, inflorescences fresh weight /plant (g), plant fresh and dry weight (g) were estimated.

b) Chemical composition:

Chlorophylls (a&b) and carotenoids concentrations (mg g⁻¹ fresh weight of leaf) were determined using colorimetric method as described by Welburn and Lichtenthaler (1984). Total carbohydrates (%) were colorimetrically determined according to the method described by Herbert *et al.*, (1971). Nitrogen% was colorimetrically determined by using the orange G dye according to the method of Hafez and Mikkelsen (1981). Phosphorus% was determined according to method described by A.O.A.C. (1995). Potassium % was determined by Flame-Photometer Parkin-Elmer model 52 with acetylene burner according to Page *et al.*, (1982). Total sesquiterpene lactones content (mg/g) was determined colorimetrically in dry herb according to the method described by El-Sawy *et al.*, (1987).

c) Anatomical studies:

At 90 days after transplanting, samples for anatomical study were taken during the second season from the 5th internodes of the apex and its leaf (petiole and blade). Samples were killed and fixed, for two days, in FAA. solution (10 ml formalin+5 ml glacial acetic acid+35 ml distilled water+50 ml ethyl alcohol 95%). Samples were dehydrated and cleared in normal butyl alcohol series and embedded in paraffin wax (56-58°C m.p.). Cross and longitudinal sections 15µ thick were cut using a rotary microtome, adhesive with Haupts and stained with crystal violet erythrosine combination, cleared in carol xylene and mounted in Canada balsam (Willey, 1971).

The results were statistically analyzed using the LSD at probability level of 5% for comparison (Gomez and Gomez, 1983).

RESULTS AND DISCUSSION

a) Vegetative growth characters:

Plant height:

The results obtained in Table (2) revealed that, the most sources of N-fertilizers significantly increased plant height in comparison to the control. The greatest increase (20.48% and 37.40 %) of plant height was attained by addition of urea at 200 kg/fed. in the first and second season, respectively. These results are in accordance with the findings of many investigators as Mohamed (1999) on *Mentha longifolia* L. who found that increasing N level (urea) from 100 to 200 kg/fed., significantly increased plant height and Mohamed (2000) on *Nigella sativa* L. who concluded that the highest nitrogen level urea significantly increased plant height.

Number of branches plant⁻¹:

Data presented in Table (2) show that all sources and levels of N-fertilizers significantly increased number of branches plant⁻¹ as compared to control in the two seasons. However, the plants treated with ammonium sulphate gave the highest values for number of branches plant⁻¹ while, ammonium nitrate treatment gave the lowest values in comparison with the control in the first season. These results are in accordance with the findings of Kandeel *et al.*, (1992) on rosemary plant who found that adding (ammonium sulphate) at any level resulted in a significant increase in the number of branches plant⁻¹ and Faid (1993) on *Carum carvi* plants, who stated that number of branches plant⁻¹ was increased by increasing nitrogen rate from ammonium sulphate.

Table (2): Effect of the different nitrogen sources on vegetative characters of damsisa plants at the flowering stage during the two seasons

Characters Treatments Kg/ fed.	Plant height (cm)	No. of branches plant ⁻¹	No. of inflorescences plant ⁻¹ (g)	F.W. of inflorescences plant ⁻¹ (g)	F.W. of herb plant ⁻¹ (g)	D.W. of herb plant ⁻¹ (g)
First season (2006/2007)						
Control	41.5	26.6	23.0	11.6	83.0	35.6
Urea	50	47.3	35.3	28.6	12.1	111.0
	100	45.6	34.6	26.6	12.2	115.3
	200	50.0	38.0	34.0	14.2	134.0
ammonium sulphate	50	47.0	35.0	35.0	13.5	111.3
	100	44.3	38.3	28.6	14.1	129.0
	200	46.0	40.0	35.1	17.2	136.0
ammonium nitrate	50	46.0	32.0	25.3	11.4	96.0
	100	42.0	33.0	25.0	11.7	109.6
	200	42.3	36.0	26.1	14.1	93.0
LSD 5%	4.1	4.6	3.9	0.2	8.9	3.5
Second season (2007/2008)						
Control	39.3	29.8	24.5	11.8	85.2	35.5
Urea	50	43.7	35.5	29.1	12.0	105.3
	100	45.7	36.6	31.4	13.3	105.8
	200	54.0	37.7	35.3	14.1	128.4
ammonium sulphate	50	50.7	37.3	34.0	16.4	110.8
	100	50.6	40.5	35.6	15.1	118.3
	200	49.7	46.7	36.2	15.6	131.2
ammonium nitrate	50	49.7	36.3	25.7	11.4	97.3
	100	44.6	37.3	25.7	12.0	94.6
	200	43.7	38.0	27.5	12.1	89.3
LSD 5%	4.3	3.2	3.8	0.2	8.4	3.6

Number and fresh weight of inflorescences plant⁻¹:

Concerning the effect of N-sources fertilizers on number and fresh weight of inflorescences plant⁻¹, the data in Table (2) show that ammonium sulphate was more effective in enhancing number and fresh weight of inflorescences plant⁻¹ than urea and ammonium nitrate fertilizers. The high rate of ammonium sulphate (200 kg/fed.) gave the best results with increase as

(55.61% and 47.67%) and (48.28% and 32.20%) during the first and second season, respectively as compared to the control.

Fresh and dry weights of herb plant⁻¹:

The data obtained in the same Table (2) indicate that both fresh and dry weights of damsisa herb were greatly increased by application of different sources of nitrogen fertilizers. The best results were attained by ammonium sulphate, especially at the high level (200 kg/fed) that recorded the highest increase were (63.86 and 53.99%) and (54.93% and 53.93%) followed by urea treatment at the same level with increases (61.45% and 50.70%) and (51.55% and 50.00%) in both seasons as compared to the control treatment. Several species of ornamental plants have produced greater dry weight when supplied with 50% or more ammonium compared to nitrate (Aiello and Graves, 1997).

Attia and Ahmed (1997) treated *Chrysanthemum millefoliatum* with three N-fertilizer sources, namely urea, ammonium nitrate and ammonium sulphate. They found that, all nitrogen fertilizer sources significantly increased plant height, number of branches plant⁻¹, fresh and dry weight of the herb, number of flowers plant⁻¹ and total flowers weight plant⁻¹.

The positive effects of N on morphological parameters may be due to the important role of these elements in protoplasm formation and all proteins e.g. amino acids, nucleic acid, many enzymes and energy transfer materials (ADP and ATP, Russel, 1973). Therefore, the increase of plant growth with adding N fertilizers is attributed to the increase of the synthesized metabolites. The high amount of synthesized metabolites caused an increment of cell number and cell size (as indicated by anatomical observations) in different plant organs which resulted in more plant height, length of peduncle, number of branches plant⁻¹ and number of inflorescences plant⁻¹. Consequently, the fresh and dry weight of herb and inflorescences plant⁻¹ were increased. The present results are similar to those obtained by Kandeel *et al.*, (1992) on rosmary, El-Sherbeny *et al.*, (1997) on *Lavandula officinalis*, Medani (1998) on damsisa, Mohamed (2000) on *Coriandrum sativum* and Mohamed and Matter (2001) on marigold plants.

b) Chemical constituents:

Pigments and total carbohydrates concentration:

In both seasons chlorophylls and carotenoids as well as total carbohydrates increased as a result of treating plants with used N-sources at all levels Table (3) indictable. The highest values were obtained by treated plants with (100kg/fed.) with increases reached to 27.9%, 55.3% and 16.0% for chlorophyll a, b and carotenoids, respectively in the first season as compared to control. The same trend was found during the second season.

Total carbohydrates (%) were increased due to the three N sources. The greatest increases (65.6% and 59.1%) were obtained from ammonium sulphate at the medium rate of 100kg/fed. in the first and second seasons, respectively, as compared to control.

N, P and K concentrations:

Data in Table (3) show that nitrogen% increased as a result of all treatments. The highest values in this respect were recorded by ammonium sulphate at the rate of 200kg/fed.

Table (3): Effect of the different nitrogen sources on chemical constituents of damsisa plants at the flowering stage during the two seasons

Characters Treatments Kg/ fed.	Characters								
	Chlorophyll a mg/g	Chlorophyll b mg/g	Carotenoids mg/g	T. carbohydrates mg/g	N %	P %	K %	Total sesquiterpenes mg/g	
First season (2006/2007)									
Control	1.29	0.85	0.31	11.85	2.31	0.07	1.48	5.15	
Urea	50	1.62	1.29	0.33	15.53	3.14	0.16	1.70	6.07
	100	1.65	1.32	0.36	14.69	3.85	0.22	1.70	9.17
	200	1.33	0.90	0.33	12.34	3.76	0.15	1.59	6.08
ammonium sulphate	50	1.36	0.89	0.34	14.86	3.45	0.29	1.58	8.15
	100	1.34	0.92	0.35	18.62	3.25	0.22	1.55	6.77
	200	1.47	0.98	0.33	15.59	3.98	0.36	1.69	9.80
ammonium nitrate	50	1.33	0.90	0.35	13.89	3.71	0.15	1.66	9.53
	100	1.42	0.95	0.33	13.52	3.77	0.29	1.98	6.60
	200	1.39	0.94	0.34	12.38	3.97	0.22	1.44	7.60
Second season (2007/2008)									
Control	1.02	0.71	0.25	11.56	2.95	0.08	1.43	8.65	
Urea	50	1.52	0.83	0.28	15.68	2.96	0.18	1.67	8.79
	100	1.70	1.15	0.40	14.18	3.16	0.21	1.65	11.35
	200	1.61	0.88	0.33	11.45	3.09	0.16	1.62	8.87
ammonium sulphate	50	1.22	0.90	0.36	14.94	3.04	0.28	1.61	10.82
	100	1.44	0.90	0.37	18.35	3.08	0.28	1.60	10.26
	200	1.62	0.98	0.37	15.79	3.27	0.34	1.67	11.61
ammonium nitrate	50	1.34	0.71	0.35	14.20	3.09	0.16	1.67	11.05
	100	1.43	0.77	0.34	13.17	3.12	0.23	1.96	9.74
	200	1.44	0.97	0.34	11.67	3.26	0.24	1.50	9.55

Phosphorus and potassium were generally increased by the three N sources application. The greatest increase in phosphorus was observed by ammonium sulphate at the rate of 200kg/fed. While, the lowest K% was recorded by ammonium nitrate at the rate of 100kg/fed. in the first season. The same trend was also observed during the second season.

The high P uptake observed in the $\text{NH}_4^+\text{-N}$ treatments may be the result of a stimulation of the H_2PO_4 uptake when the NO_3^- uptake is strongly depressed (Mengel and Kirkby, 2001). Also, it was observed that plants treated with $\text{NH}_4^+\text{-N}$ source contained higher concentrations of inorganic anions (PO_4^{2-} , H_2PO_4^- , Cl^-), whereas cation (Ca^{2+} , Mg^{2+} , K^+) were found in lower concentrations. In contrast, the plants supplied with the $\text{NO}_3^-\text{-N}$ form contained higher cation content (Bar-Tal *et al.*, 2001). Nitrogen and P uptake rates are higher in the plant supplied with NH_4^+ or $\text{NO}_3^+\text{-NH}_4^+$ than in plants provided with NO_3^- alone (Jimenez *et al.*, 2007).

Sesquiterpene lactones:

Data tabulated in Table (3) indicate that total sesquiterpene lactones were greatly increased by application of all treatments. The maximum increase was obtained by the treatment 200kg/fed. of ammonium sulphate. This revealed increments of 90.9% in the first season. The same trend was also obtained in the second one in comparison to the control.

Collectively, the increase of leaf pigments to N may be attributed to the role of N in the pigment synthesis, especially N which is a component of chlorophyll molecule. The increase in total carbohydrates concentration resulting from N may be directly due to their effect on chlorophyll and indirectly activation of the anabolic processes of carbohydrates metabolism. Increases of N, P and K by N attributed to that nitrogen increases the capacity of the plants to absorb nutrients by increasing the size of plant root surface/ unit of soil volume, as well as the high capacity of the plants supplied with nitrogen in building metabolites, which in turn, contributes much to the increase of nutrient uptake (Mandour *et al.*, 1986). The increased in total sesquiterpene lactones by nitrogen fertilization was expected and agrees with Sidky and El-Mergawi (1997). They found on damsisia plant that good production of total sesquiterpene lactones was observed in plants fertilized by nitrogen.

c) Anatomical studies:


1- Stem stricture:

Microscopical counts and measurements of certain characters in transverse sections of damsisia stem treated with the three N-sources at all levels are given in Table (4). Likewise, microphotographs illustrating these treatments are shown in Figures (1 and 2).

Table (4): Effect of the different nitrogen sources on anatomical structures of damsisia stem.

Characters	Treatments	Control	Urea			Amm. sulphate			Amm. nitrate		
			Kg/fed.								
			50	100	200	50	100	200	50	100	200
Section diameter μ		2425	2950	3213	3088	3250	3388	3675	2975	3025	3225
Cortex	Thickness μ	173	205	177	183	176	195	195	180	190	205
	No. of layers	7.7	8.5	8.3	8.3	8.0	8.5	8.5	7.0	8.7	8.0
	Length of cells μ	67	103	93.3	107	90	87	90	90	93	93
No. of vascular bundles		18	20	22	20	21	22	22	21	19	22
Vascular bundle dimensions	Length μ	407	540	600	610	575	590	567	650	557	685
	Width μ	307	317	300	433	360	390	410	375	387	377
Av. Xylem vessel diameter		33	40	40	39	50	39	38	41	45	39
Medullary rays	Width μ	125	75	80	115	110	100	90	100	115	100
	No. of layers	6	7	6	8	7	6	7	6	7	7
Pith	Diameter μ	1250	1625	1675	1688	1625	1813	2000	1625	1713	1838
	No. of layers	29	35	34	34	29	37	39	35	37	36
	Length of cells μ	47	90	73	100	73	73	77	87	83	77

width of modularly rays (40%) was achieved by urea at 50kg /fed. Swaefy Hend *et al.* (2007) on peppermint found that treating the plants with 100%N+P (ammonium sulphate) increased dimensions of the stem.



Line = 1mm.

Fig (2): Longitudinal sections in damsisia stem under different sources of nitrogen:

a: Control

b: Urea at 200 kg/fed.

c: Ammonium sulphate at 200 kg/fed.

d: Ammonium nitrate at 200 kg/fed.

(cx: Cortex)

2- Leaf petiole:

It is clear from Table (5) and Figure (3) that, treating with the three N-sources at all levels greatly increased leaf petiole section diameter by increasing its tissues. The highest increase of leaf petiole section diameter was recorded in response to ammonium nitrate at 100kg/fed. (42.07%) with a great increase of length and width of vascular bundles as well as diameter of xylem vessels (31.11%, 8.89% and 53.19%, respectively).


3-Leaf blade:

Data tabulated in Table (5) and illustrated in Figure (4) show that, application of all N-forms greatly increased midvein thickness, mainly ammonium nitrate at 200 kg/fed. that caused the highest value (76%) followed by ammonium sulphate at the same level. Also, addition the application of ammonium sulphate at 50kg/fed. caused an increase in leaf blade thickness (40%) as compared to the control. This increment was due to the increase in thickness of both upper and lower palisade tissue as well as spongy tissue. Moreover, ammonium sulphate mainly at 200kg/fed. showed the highest increase in length and width of vascular bundles (250% and 100%) in comparison to the control.

Collectively, the increase in stem, leaf petiole and leaf blade tissues due to N-sources treatments could be attributed to the role of nitrogen in increasing cell division and expansion and this, in turn, was reflected in an increase of number and diameter of parenchyma cells in cortex and pith of stem. Also, stimulating cell division in cambial zone and consequently increased dimensions of the vascular bundles and xylem vessels. This stimulate effect was reported by Mohamed (1989) on chamomile, Medani (1998) on damsisa, Agamy (2000) on *Ocimum basilicum*, Mohamed and Matter (2001) on marigold plants and Swaefy Hend *et al.* (2007) on peppermint.

Table (5): Effect of the different nitrogen sources on anatomical structures of damsisa leaf petiole and blade

Characters	Treatments	Kg/fed.									
		Urea			Ammonium sulphate			Ammonium nitrate			
		Control	50	100	200	50	100	200	50	100	200
Leaf petiole											
Av. Section diameter μ		1338	1638	1563	1500	1588	1788	1575	1438	1900	1863
Vascular bundle dimensions μ	Length μ	450	460	480	530	470	600	500	510	590	580
	Width μ	450	360	370	360	340	430	400	400	470	520
Av. xylem vessel diameters μ		19	24	26	24	26	21	26	26	29	26
Leaf blade											
Midvein thickness μ		500	450	720	620	770	620	830	650	600	880
Blade thickness μ		250	220	250	250	350	250	250	280	290	330
Upper palisade thickness μ		100	88	98	93	113	100	105	113	120	118
Lower palisade thickness μ		50	40	50	38	43	43	38	80	40	43
Spongy thickness μ		50	63	75	78	105	88	75	43	75	93
Vascular bundles dimensions μ	Length μ	100	120	300	250	200	230	250	250	250	300
	Width μ	100	100	180	150	140	200	200	200	150	170



Line = 1mm.

Fig (4): Trans sections in damsisa leaf blade at different sources of nitrogen:

a: Control

c: Ammonium sulphate at 200 kg/fed.
(lb : Leaf blade

b: Urea at 200 kg/fed.

d: Ammonium nitrate at 200 kg/fed.
mv : Midvein)

REFERENCES

- Abbes, C.; L.E. Parent, A. Karam and D. Isfan (1995). Effect of $\text{NH}_4^+:\text{NO}_3^-$ ratios on growth and nitrogen uptake by onions. *Plant Soil* 171: 289-296.
- Agamy, R.A (2000). The effect of some fertilizers and growth regulators on botanical characters of *Ocimum basilicum* L. and *Cotula cinerea* Del. Ph.D., Thesis Fac. of Agric., Fayoum Cairo Univ., Egypt.
- A.O.A.C. (1995). Official Methods of Analysis, Published by the A.O.A.C. Washington, D. C., U. S. A.

- Aiello, A.S. and W.R. Graves (1997). Container medium and nitrogen from affect production of Amur maackia (*Maackia amurensis* Rupr.& Maxim.). HortScience, 32 (7): 1200-1203.
- Attia, F.A. and Ahmed (1997). Effect of some nitrogen fertilization forms and two growth regulators on *Chrysanthemum millefoliatum*, Ramat Cv. Icecap plants. J. Agric. Sci. Mansoura Univ., 22 (4): 1141-1154.
- Bar-Tel, A.; B. Alon; L. Karni and R. Rosenberg (2001). Nitrogen nutrition of greenhouse pepper. II- Effect of nitrogen concentration and NO_3^- : NH_4^+ ratio on Growth, transpiration and nutrient uptake. HortScience 36:1252-1259.
- El-Saeid, H.M.; M.S. Hussein; S.E. El-Sherbeny and E.A. Omer (1996). Effect of nitrogen on yield and active constituents of *Tagetes patula* L. Egypt. J. Hort. 23 (1): 101-112.
- El-Sawy, M.F.; J. Duncan; S. Amer; H. El-Ruweini; N. Brown and M. Hills (1987). The molluscicidal properties of *Ambrosia maritima* L. (Asteraceae). 3- A comparative field trail using dry and freshly harvested plant material. Trop. Med. Parasit., 38, 101-105.
- El-Sherbeny, S.E.; H.M. El-Saeid and M.S. Hussein (1997). Response of Lavender, *Lavandula officinalis* L. plants to different nitrogen sources. Egypt J. Hort. 24, 1, 7-17.
- Faid, A.M. (1993). Physiological study on caraway. M.Sc. Thesis, Fac. Agric., Zagazig Univ.
- Gomez, K.A. and Gomez, A.A. (1983). Statistical Procedure for Agricultural Research. A Wiley Inter-Science Publication. John Wiley&Sone Inc. New York.
- Hafez, A. and D.S. Hikkelsen (1981). Colorimetric determination of nitrogen for evaluating the nutritional status of rice. Commnu. Soil Sci. and Plant Analysis, 12 (1): 16-69.
- Hassanein, M.A.; S.M. Selim; T.S. Abd El-Aal and S.A. Mahmoud (1998). Effect of some nitrogenous fertilizer rates and soil properties on growth and essential oil of *Pelargonium graveolens* L. Annals of Agric. Sci. Moshtohor, Vol. 36 (1) p.539- 551.
- Herbert, D.; P.J. Phipps and R.E. Strange (1971). Methods in Microbiology, 5 B, Academic Press, London, 209-344.
- Jimenez, S.; J.I. Ales; M.T. Lao; B.M. Plaza and M. Perez (2006). Evaluation of N concentration quick test for field determination in a soil solution, drainage solution and nutrient solution in fustigation. Comman. Soil Sci. Plant Anal. 37:2461-2469.
- Jimenez, S.; M. Perez; B.M. Plaza; R. SAlinas and M.T. Lao (2007). Empirical Models of Phosphorus Uptake under Different Sources in *Dieffenbachia amoena* 'Tropic Snow'. HortScience 42: 412-416.
- Kandeel, A.M.; S.M. Mohamed and A.A. Sadek (1992). Effect of some agricultural treatments on growth and chemical composition of rosmary (*Rosmarinus officinalis* L.). 1- Effect of nitrogen fertilization. Egypt J. App. Sci., 7 (6): 340-352.
- Mahran, G.H. (1967). Medicinal Plants. 1st ed. Anglo, egypt. Book Shop. Cairo pp.465.

- Mandour, M.S.; S. El-Sherbeiny; N.B. Botros and S.H. El-Nagar (1986). Effect of nitrogen application upon growth, oil and nutrient content of Citronella grass. Bull. Egypt, Soc. Physiol. Sci. 145 (3) 6.
- Mansour, H.A.; E.I. El-Maadawy and E.A. El-Ghadban (1999). Organic and inorganic fertilization of spearmint and marjoram. Bull. Fac. Agric. Cairo Univ., Vol. 1, p. 226-239.
- Medani, R.A. (1998). Effect of N, P and K on growth, anatomical characters and some chemical constituents of damsisa plant. J. Aric. Sci. Mansoura Univ., 23 (11): 4869-4881.
- Mengel, K. and E.A. Kirkby (2001). Principles of plant nutrition. 5th ed. Intl. Potash Inst. Switzerland.
- Moa, Z. and I. Craker (1991). Photosynthesis and the production of herbs. Spice and Medicine Plant Digest. 9 (2): 15-18.
- Mohamed, M.I. (2000). Physiological studies on *Coriandrum sativum* L. and *Carum carvi* L. plants. Ph.D. Thesis, Fac. Agric. Cairo Univ. Egypt.
- Mohamed, S.A. (1989). Effect of mineral nutrition on some morphological and anatomical characters and yield of chamomile and fennel plants. M.Sc. Thesis, Fac. of Agric., Fayoum, Cairo Univ., Egypt.
- Mohamed, S. Faten (1999). Botanical studies on two wild medicinal plant (damsisa and wild mint) in Fayoum. M. Sc. Thesis, Fac. Agric., Fayoum, Egypt.
- Mohamed, S.A. and F.M.A. Matter (2001). Effect of ammonium nitrate and organic fertilizers on growth, volatile oil yield and chemical constituents of marigold (*Tagetes minuta* L.) plant. Fayoum J. Agric., Res. & Dev., Vol. 15, No. 1, pp. 95-107.
- Mohamed, S.A.; R.A. Medani and E.R. Khafaga (2000). Effect of nitrogen and phosphorus applications with or without micronutrients on black cumin (*Nigella sativa* L.) plants. 8th Conf. Agric. Dev. Res., Fac. Agric. Ain Shams Univ., Cairo, November 20-22, 2000. Annals of Agric. Sci., Sp. Issue, 3, 1323-1338.
- Olsen, S.R. and Sommers, L.E. (1982). Phosphorus, In Methods of Soil Analysis. Part2, pp.403-430. Page, A.I., Miller, R.H. and Keeny, T.R. (eds.), Am. Soc. of Agron. Madison WI.
- Page, A.I.; R.H. Miller and T.R. Keeny (Eds.) (1982). Methods of Soil Analysis. Part 2: 595. Am. Soc. of Agron., Madison WI 9.USA.
- Russel, E.W. (1973). "Soil Conditions and Plant Growth" Language Book Soc. Longman, London, 30-37.
- Sidky, M.M.A. and R.A. El-Mergawi (1997). Relation between, growth, biosynthesis and accumulation of major active constituents of *Ambrosia maritima* and some exogenous and endogenous factors. Bulletin of Fac. Agric., Cairo univ., 4(48): 631-654.
- Swaefy Hend, M.F.; Weaam R.A. Sakr; A.Z. Sabh and A.A. Ragab (2007). Effect of some chemical and bio-fertilizers on peppermint plants grown in sandy soil. 2. Effect on essential oil production, chemical composition and anatomical features. Annals Agric. Sci., Ain Shams Univ., Cairo, 52 (2): 465-484.

- Teriest, L.M.V. Vijver and J.J. Symoens (1989). Polymorphism and electrophoretic patterns in *Ambrosia maritime* L., *A. artemisiifolia* L. and *A. trifida* L. (Asteraceae). Plant Sci., 59: 211-218.
- Welburn, A.R. and H. Lichtenthaler (1984). Formula and program to determine total carotenoids and chlorophyll a and b of leaf extracts in different solvents. "Advances in Photosynthesis Research" (Sybesma C. Ed.) Vol.Π, pp. 9-12.
- Willey, R.L. (1971). Microtechnique. A Laboratory Guide. Mac Millan Publishing Co. Inc. New York., U.S.A.

تأثير مصادر النيتروجين على المحصول والصفات التشريحية لنباتات الدمسيسة تحت ظروف الأرض الرملية

فيصل محمود عبد المجيد مطر*، رمضان عبد العظيم عجمي** و سمير سيد الجنائني**

* قسم البساتين- كلية الزراعة – جامعة الفيوم- مصر

** قسم النبات الزراعي- كلية الزراعة – جامعة الفيوم- مصر

درس تأثير ثلاثة مصادر من السماد النيتروجيني (نترات الأمونيوم-سلفات الأمونيوم- اليوريا) بالمعدلات صفر-50-100&200 كيلو جرام للفدان على الصفات المورفولوجية والتشريحية والمحصول والمكونات الكيماوية لنباتات الدمسيسة فى تجربة أصص بكلية الزراعة جامعة الفيوم خلال موسمى 2006-2007 و 2007-2008 .

وتشير النتائج المتحصل عليها أن كل مصادر التسميد النيتروجيني أدت الى حدوث زيادة واضحة فى صفات النمو مثل ارتفاع النبات وعدد الفروع وعدد النورات ووزنها الطازج والوزن الطازج والجاف للعشب وكانت الزيادة السابقة على النحو التالى: سلفات الأمونيوم ثم اليوريا وأخيرا نترات الأمونيوم .

أظهرت كل المعاملات خاصة اليوريا وسلفات الأمونيوم زيادة واضحة فى محتوى الأوراق من كلوروفيلات أ و ب والكاروتينويدات وكذلك زاد محتوى العشب من الكربوهيدرات الكلية والنيتروجين والفوسفور والبوتاسيوم .

أيضا كل المعاملات خاصة سلفات الأمونيوم أدت إلى زيادة واضحة فى قطر قطاعات الساق وكانت هذه الزيادة ناشئة من حدوث زيادة فى كل من الأنسجة البرانشيمية والوعائية هذا بالإضافة إلى حدوث زياده فى قطر قطاعات عنق الورقه كنتيجة للزياده فى أنسجته, كما أظهرت كل مصادر السماد النيتروجيني زيادة واضحة فى سمك العرق الوسطى ونصل الورقة وأبعاد الحزم الوعائية وقطر أوعية الخشب.

أدت معاملة نباتات الدمسيسة بالمصادر المختلفة للسماد النيتروجيني (خاصة سلفات الأمونيوم) إلى حدوث زيادة واضحة فى محتوى العشب من السيسكوترينينات الكلية. لوحظ أن أعلى زيادة فى معظم الصفات السابقة لنباتات الدمسيسة تم الحصول عليها نتيجة استخدام سلفات الأمونيوم لذا توصى الدراسة بإستخدامة كمصدر للسماد النيتروجيني للحصول على نباتات دمسيسة ذات محصول وجودة عالية.