

Comparative Study of Esophageal Varices Grading by Multi-Detector Computed Tomography and Endoscopy

FATMA A. EL-SHARAWY, M.D.* and MOHAMED M. MABROUK, M.D.**

The Departments of Radio-Diagnosis & Medical Imaging and Internal Medicine**, Faculty of Medicine, Tanta University, Egypt*

Abstract

Background: Cirrhosis is often complicated by Esophageal Varices (EV) and portal hypertension. The use of upper GI endoscopy (GIE) as a screening method is limited regarding invasiveness, expensive, needs sedation as well as patient's poor acceptance of the procedure. In contrast; Multidetector Computed Tomography (MDCT) imaging is non-invasive, does not necessitate sedation, and allows accurate assessment of variceal site and size, also better tolerated by patients than upper GI Endoscopy (GIE).

Aim of Study: To prove that MDCT is a non-invasive alternative diagnostic tool to EGD in grading of Esophageal Varices.

Patients and Methods: 50 patients with liver cirrhosis were included. Evaluation of Multidetector Computed Tomography (MDCT) in the diagnosis of esophageal varices and its grading was done by comparing the grades of Esophageal Varices at Multidetector Computed Tomography (MDCT) and upper GI endoscopy independently. Extra-esophageal findings were also assessed by Multidetector Computed Tomography (MDCT) that cannot by (GIE).

Results: At upper GI endoscopy, 3 patients had grade 0 Esophageal Varices, 25 patients had grade 1, 16 patients had grade 2 and 6 had grade 3. The sensitivity, specificity, positive and negative predictive values and accuracy of Multidetector Computed Tomography (MDCT) for defining Esophageal Varices in all grades were 99.5%, 99.6%, 99.4%, 99.5% and 99.5% respectively. Important extra-esophageal findings were determined by Multi-Detector Computed Tomography (MDCT) only. The acceptance of patients for Multidetector Computed Tomography (MDCT) was significantly more than that for upper GI Endoscopy (GIE).

Conclusion: Multi-Detector Computed Tomography (MDCT) is a fast, non-invasive procedure for diagnosis and grading of Esophageal Varices.

Key Words: Esophageal Varices (EV) – Upper GI endoscopy (GIE) – Computed Tomography (CT) – Multidetector Computed Tomography (MDCT) – Esophago-Gastro-Duodenoscopy (EGD) – Intra-Venous (IV).

Introduction

Development of significant complications for portal hypertension as ascites and/or Esophageal and Gastric Varices occurs when hepatic venous pressure gradient increases above 10mmHg [1].

Esophageal varices is one of the major complications of liver cirrhosis, with an estimated prevalence of approximately 80% with decompensated patients and 50% in compensated cirrhosis [2].

Esophagoscopy (EGD) is the gold standard in the diagnosis of esophageal Varices; however, the use of endoscopy as a method of screening is limited, as it is invasive, expensive, needs sedation together with patient's poor acceptance of the procedure [3].

Several studies have searched for alternatives to conventional endoscopy including biochemical, clinical and ultrasound parameters, transient elastography and video capsule endoscopy for the non-invasive or minimally invasive diagnosis of esophageal Varices. These methods have shown promising performance characteristics for overcoming bleeding and were perceived by patients as preferable to endoscopy; yet they have limited sensitivity and specificity and none of them had been proved as an alternative for upper GI endoscopy [4].

CT imaging as better tolerated by most of the patients, non-invasive, does not need sedation, and allows accurate assessment of variceal site and size than endoscopy is considered a good alternative to upper GI endoscopy [5].

With the introduction of MDCT imaging with its multi-planar capabilities, esophageal, para-esophageal and Gastric Varices as well as other porto-systemic shunts was progressively recognized in patients with liver cirrhosis [6].

Aim of the study: To prove that MDCT is a non-invasive alternative diagnostic tool to EGD in grading of Esophageal Varices.

Patients and Methods

This prospective study was carried out on 50 cirrhotic patients (30 males, 20 females) their age ranged from 31 to 75 years with a mean age of 49.9 years. They were referred to Diagnostic Radiology and Medical Imaging Department, from Internal Medicine and Tropical Medicine Departments; Tanta University Hospital throughout the period from April 2018 to November 2019.

Inclusion criteria:

- Cirrhotic patients diagnosed by clinical, laboratory and radiological parameters.
- Patients suspicious to have hepatocellular carcinoma.
- Patients with portal hypertension diagnosed by abdominal ultrasonography.

Exclusion criteria:

- Active gastrointestinal bleeding.
- History of endoscopic variceal ligation.
- History of adverse reactions to iodinated contrast agent.

Methods:

The procedure was carried out over two separate studies; MDCT study with IV contrast injection and EGD study. MDCT images were evaluated for diagnosis and grading of Esophageal Varices and then correlated with EGD.

1-Multidetector computed tomography (MDCT):

All studies were obtained using a 128 multi-detector CT scanner (GE Optima CT model 660) installed in Diagnostic Radiology and Medical Imaging Department. Before examination the patients swallow two 5g packets of effervescent powder containing sodium bicarbonate, tartaric acid and citric acid. Patients lied supine with head first. Scout films were taken routinely in all patients.

MDCT was performed before and after the injection of non-ionic contrast medium. No positive oral contrast material was administered. After completion of the nonenhanced scans, an IV bolus injection of non-ionic contrast material was administered at a dose of 1.5-2mg/kg at a rate of 4ml/s through a 22-gauge IV catheter inserted into an ante-cubital vein using an automatic injector.

After injection of intravenous contrast material, the liver was scanned in arterial (scanning delay, 20-25 seconds), portal (scanning delay, 60 seconds), and delayed (scanning delay, 3-5 minutes) phases. Images were performed from the lower thorax and abdomen to the iliac wing during a single breath-hold at the end of inspiration.

After completion of the study, a high-quality data set with a 0.7-mm reconstruction interval was transferred to a picture archiving and communication workstation equipped with dedicated 2D and 3D software tools which allowed processing of Multiplanar Reconstructions (MPRs) and Volume Renderings (VRs), by using surface-shaded transparent and endo-luminal virtual endoscopy modes.

2- Esophagoscopy :

All studies were obtained using Pentax EG-2985. *Esophagoscopy* was done within 1 week from the MDCT study. Patients were fasting overnight. Before the procedure, Midazolam 3-5mg IV was given with Xylocaine local spray above the tongue and nasopharynx. Using Pentax EG-2985, the patient lied on the examination trolley on the left side with the IV access line preferably in the right arm. The endoscope was inserted under direct vision. Examination of the entire esophagus was done.

Results

Our study included 50 cirrhotic patients 30 of them were males and 20 females. Their age ranged from 31 to 75 years with a mean of 49.9 years. (Table 1) shows the different age groups.

Table (1): Age and sex distribution in the studied patients (n=50 patients).

Age group (years)	Sex		Total
	Male	Female	
30-39Y	0	1	1
40-49Y	11	9	20
50-59Y	16	9	25
60-69Y	1	1	2
70-80Y	2	0	2
Total	30	20	50

The causes of liver cirrhosis were different among the studied patients, 13 had hepatitis B (26%), 26 had hepatitis C (52%), 11 had Bilharzia- sis & other causes of cirrhosis (22%). The different causes of cirrhosis are shown in Fig. (1).




Fig. (1): Causes of liver cirrhosis in the studied patients (n=50 patients).

In our study, Esophageal Varices were classified into four groups by MDCT:

Grade 0: No Varices visualization on the inner surface of the Esophagus.

Grade 1: One varix less than 5mm in diameter detected on the inner surface from the esophagus.

Grade 2: Several Varices less than 5mm detected on the inner surface from the esophagus.

Grade 3: One varix 5mm or greater or several Varices occupy more than half of the circumference of the Esophagus. The results are listed in (Table 2).

Table (2): MDCT grades of Esophageal Varices in the studied patients (n=50 patients).

Grade	No. of patients	Percentage
Grade 0	3	6
Grade 1	25	50
Grade 2	16	32
Grade 3	6	12
Total	50	100

In our study, Esophageal Varices were classified into four groups by EGD:

Grade 0: No Varices visualization on the inner surface of the esophagus.

Grade 1: Small Varices were defined as those that flatten with insufflations or protrude minimally into the esophageal lumen.

Grade 2: Moderate-sized Varices were defined as those that protrude into the esophageal lumen but did not touch each other.

Grade 3: Large Esophageal Varices were defined as those that protrude into the Esophageal lumen and touch each other. The results are listed in (Table 3).

Table (3): Endoscopic grades of Esophageal Varices in the studied patients (n=50 patients).

Grade	No. of patients	Percentage
Grade 0	3	6
Grade 1	25	50
Grade 2	17	34
Grade 3	5	10
Total	50	100

The ability of MDCT and Virtual Endoscopic CT to identify Esophageal Varices (EV) as compared to EGD was listed in (Table 4).

Table (4): MDCT and virtual endoscopic CT compared to endoscopy to identify Esophageal Varices.

Grades	Grade0	GradeI	Grade II	GradeIII
True positive	3	24	17	5
True negative	47	26	33	45
False positive	0	0	1	0
False negative	0	1	0	0

Table (5): sensitivity, specificity and accuracy of MDCT and Virtual Endoscopic CT to identify Esophageal Varices as referred to endoscopy.

	Sensitivity	Specificity	PPV	NPV	Accuracy
Grade 0	100	100	100	100	100
Grade 1	98	100	100	98.1	99
Grade 2	100	98.5	97.1	100	99
Grade 3	100	100	100	100	100
Total	99.5	99.6	99.4	99.5	99.5

The sensitivity, specificity, positive and negative predictive values and accuracy of CT for defining EV in all grades were 99.5%, 99.6%, 99.4%, 99.5% and 99.5% respectively.


In our study, we identified extra-esophageal pathology by multi-detector MDCT only. The different findings are showed in (Table 6).

Table (6): Extra-esophageal findings identified by MDCT in the studied patients (n=50 patients).

CT extra-esophageal findings	No. of patients	Percentage
HCC	4	8
Splenomegaly	37	74
Ascites	31	62
Para-esophageal varices	29	58
Gastric varices	23	46
Perisplenic collateral	17	34

In our study, when we compare the acceptance of MDCT and upper GI endoscopy techniques from the patient side, 47 patients (94%) out of 50 found that MDCT is more preferable and accepted than upper GI endoscopy. Patient preference is shown in Fig. (2).

Fig. (2): Comparison between MDCT and endoscopy as regards patient preference.







Fig. (3): Male patient aged 68 years old with hemoptysis: MDCT porto-venous phase sagittal (A), coronal (B) axial (C) and virtual CT (D) images show enhanced vessels protruded into the esophageal lumen-white arrow-(esophageal varices grade 2 by CT) measuring between 2.8-4.3mm confirmed by EGD image (E).



Upper endoscopy 3 days later done by professional internal medicine doctor revealed grade 3 esophageal varices.







Fig. (4): Female patient aged 49 years old with recurrent hemoptysis & U/S revealed hepatic focal lesion & splenomegally: MDCT porto-venous phase axial coronal (A), sagittal (B), axial (C) and endoscopic (D) images show multiple enhanced vessels involving the whole circumference of the esophageal lumen-white arrow-(esophageal varices grade 3 by CT) confirmed by EGD image (E).



☆

Upper endoscopy a week later done by professional internal medicine doctor revealed grade 2 esophageal varices.

☆





Fig. (5): Male patient aged 61 years old with recurrent hemoptysis & previous variceal ligation: MDCT Porto-venous phase sagittal (A), axial (B) coronal (C) and endoscopic (D) images show multiple enhanced vessels involving the whole circumference of the esophageal lumen-white arrow-(esophageal Varices score 3 by CT) confirmed by GED image (E). The liver is cirrhotic.



Upper endoscopy done by professional internal medicine doctor revealed grade 3 esophageal varices.



Discussion

Approximately 60-80% of patients with liver cirrhosis develop Esophageal Varices during their life at a rate of 5% per year, the progression from small to large Varices occurs in 5-10% of patients after the first year [8] .

Esophagoscopy is the gold standard in the diagnosis of oesophageal Varices in cirrhotic patients [9] .

Screening for esophageal Varices using upper GI endoscopy is always done for patients with cirrhosis to avoid significant morbidity and mor-

tality associated with bleeding from Varices [10]. It is recommended every year in patients with small Varices and every 2-3 years in patients without Varices to allow initiation of primary preventive treatment, this means that a large number of cirrhotic patients undergo unnecessary endoscopic examination [11].

The use of upper GI endoscopy as a method of screening is limited, as it is invasive, expensive, needs sedation, and the patient's poor acceptance of the procedure [12].

MDCT imaging is non-invasive, does not necessitate sedation, and allows accurate assessment of variceal site and size, and it is also better tolerated by most of the patients than upper GI endoscopy [13].

In this study we compare the ability of MDCT in the diagnosis of esophageal Varices as a non-invasive procedure and its acceptance to the patients to upper GI endoscopy. It included (50) patients with cirrhosis (30) males (60%) and (20) females (40%). So male affection is more predominant than females in agreement with Moftah et al. study [14] where male patients were 74% and female patients were 26%.

The causes of liver cirrhosis were different among the studied patients, 13 had hepatitis B (26%), 26 had hepatitis C (52%), 11 had Bilharziasis (22%). So hepatitis C is a major cause of liver cirrhosis in agreement with Dessouky et al. study [15] where the most common cause of liver cirrhosis was hepatitis C (68%).

In our study, CT scan had a sensitivity of (99.5%), a specificity of (99.6%). So close to that found by Dessouky et al. study [15] where the overall CT sensitivity was (99%), and specificity (98%), in comparison to the upper GI endoscopy.

As regards conventional CT in diagnosis and grading of esophageal Varices; the study of Perri et al. [16] reported that CT has approximately 90% sensitivity in the detection of esophageal Varices, but only about 50% specificity. Also, Kim et al. Study [17] got results with sensitivity (92%), specificity (84%), Dessouky et al. study [15] also reported (99%) sensitivity, specificity (98%) while ours recorded sensitivity (99.5%), specificity (99.6%).

A well optimized protocol for the evaluation of Esophageal Varices with contrast, very

small slice thickness, bolus tracking technique that allows more accurate timing of arterial and portal venous phases and acquisition of high-resolution images associated with 2D and 3D reconstructions and endo-luminal virtual images that provide accurate results about different grades of Esophageal Varices. Moreover, the higher sensitivity and specificity in our study may be due to the fact that we use 128 slice MDCT while in the study of Perri et al., [16] they used 4 detectors and in Also Kim et al., [17] they used 16 detectors in their studies.

An efficient and well-tolerated technique for distending the esophagus was done by the administration of effervescent powder. The slow passage of the effervescent powder through the esophagus and the supine position of the patient save the developed gas within the esophageal lumen longer, thus causing more efficient wall distension. This was in agreement with Cansu et al. study [18] who reported that using the effervescent powder in the detection and grading of esophageal Varices by MDCT increased the success rate of MDCT. Thus, small Varices, which are difficult to be detected via MDCT, can be detected and upper GI endoscopy may not be required to evaluate these low risk bleeding Varices.

In our study, when we compare the acceptance of MDCT and upper GI endoscopy techniques from the patient side, 47 patients (94%) out of 50 found that MDCT is more preferable and accepted than upper GI endoscopy, only 3 patients (6%) found upper GI endoscopy more preferable. This was in agreement with Dessouky et al. [15] who reported MDCT more tolerable compared to upper GI endoscopy and patients are more willing to utilize it for follow-up.

MDCT has high performance of in the detection and grading of Esophageal Varices in our study yet there is an increased risk of radiation hazard. with some limitations of MDCT when compared to upper GI endoscopy for the screening of Esophageal Varices and also; therapeutic intervention cannot be performed during MDCT, whereas this is possible during upper GI endoscopy. A dose-modulation program to reduce the dose of radiation.

Our study provided an opportunity for cirrhotic patients for dual screening and evaluation strategy of two crucial pathological conditions, which are HCC and Esophageal Varices, without any added cost, effort, time or risk of radiation. Considering the high cost of performing

multiple tests and the relative invasiveness of upper GI endoscopy, a single non-invasive surveillance tool for both Varices and HCC may be important. These factors constitute a major advantage of MDCT over upper GI endoscopy [19].

In our study; the performance characteristics of MDCT have been proved to be superior to upper GI endoscopy for the detection and evaluation of other important extra- esophageal abnormalities, which were also considered other risk factors that needed rapid and selective management and may give further information regarding the propensity of Esophageal Varices and variceal bleeding such as splenomegaly, ascites, HCC and gastric submucosal Varices. This goes with Dessouky et al., [15].

Conclusion:

MDCT with IV contrast is an excellent, safe, non-invasive alternative diagnostic tool for detection and grading of esophageal varices as compared to conventional upper GI endoscopy.

References

- 1- GARCIA-TSAO G. and BOSCH J.: Management of Varic- es and variceal hemorrhage in cirrhosis. *N. Engl. J. Med.*, 362 (9): 823-32, 2010.
- 2- GOMEZ E.V., RODRIGUEZ Y.S., BERTOT L.C., et al.: The natural history of compensated HCV-related cirrhosis: A prospective long-term study. *J. Hepatol.*, 58 (3): 434-44, 2013.
- 3- FANTI L. and TESTONI P.A.: Sedation and analgesia in gastrointestinal endoscopy: What's new? *World J. Gastroenterol.*, 16 (20): 2451-7, 2010.
- 4- CHERIAN J.V., DEEPAK N., PONNUSAMY R.P., et al.: Non-invasive predictors of esophageal varices. *Saudi J. Gastroenterol.*, 17 (1): 64-8, 2011.
- 5- ELKAMMASH T., ELFIKY I., ZAITON F., et al.: Diagnostic performance of multidetector computed tomography in the evaluation of esophageal Varices. *The Egyptian Journal of Radiology and Nuclear Medicine*, 47 (1): 43-51, 2016.
- 6- ZHAO L.Q., HE W., JI M., et al.: 64-row multidetector computed tomography portal venography of gastric variceal collateral circulation. *World J. Gastroenterol.*, 16 (8): 1003-7, 2010.
- 7- HASHIZUME M., AKAHOSHI T. and TOMIKAWA M.: Management of gastric varices. *J. Gastroenterol. Hepatol.*, 26 Suppl 1: 102-8, 2011.
- 8- SARANGAPANI A., SHANMUGAM C., KALYANASUNDARAM M., et al.: Noninvasive prediction of large esophageal varices in chronic liver disease patients. *Saudi J. Gastroenterol.*, 16 (1): 38-42, 2010.
- 9- HWANG J.H., SHERGILL A.K., ACOSTA R.D., et al.: The role of endoscopy in the management of variceal hemorrhage. *Gastrointest. Endosc.*, 80 (2): 221-7, 2014.
- 10- MORISAKA H., MOTOSUGI U., ICHIKAWA S., et al.: Association of splenic MR elastographic findings with gastroesophageal varices in patients with chronic liver disease. *J. Magn. Reson. Imaging*, 41 (1): 117-24, 2015.
- 11- POZA CORDON J., FROILAN TORRES C., BURGOS GARCÍA A., et al.: Endoscopic management of esophageal varices. *World J. Gastrointest. Endosc.*, 4 (7): 312-22, 2012.
- 12- SINGEAP A.M., STANCIU C. and TRIFAN A.: Capsule endoscopy: The road ahead. *World J. Gastroenterol.*, 22 (1): 369-78, 2016.
- 13- WELLS M.L., HANSEL S.L., BRUINING D.H., et al.: CT for Evaluation of Acute Gastrointestinal Bleeding. *Radiographics*, 38 (4): 1089-107, 2018.
- 14- MOFTAH S.G., KAMAL S. and HANNA A.: CT esophagography: Non-invasive screening and grading of esophageal Varices in cirrhosis. *The Egyptian Journal of Radiology and Nuclear Medicine*, 45 (2): 263-70, 2014.
- 15- DESSOUKY B.A. and ABDEL AAL EL S.M.: Multidetector CT oesophagography: An alternative screening method for endoscopic diagnosis of oesophageal varices and bleeding risk. *Arab J. Gastroenterol.*, 14 (3): 99-108, 2013.
- 16- PERRI R.E., CHIOREAN M.V., FIDLER J.L., et al.: A prospective evaluation of computerized tomographic (CT) scanning as a screening modality for esophageal varices. *Hepatology*, 47 (5): 1587-94, 2008.
- 17- KIM Y.J., RAMAN S.S., YU N.C., et al.: Esophageal varices in cirrhotic patients: Evaluation with liver CT. *AJR Am. J. Roentgenol.*, 188 (1): 139-44, 2007.
- 18- CANSU A., AHMETOGLU A., KUL S., et al.: Diagnostic performance of using effervescent powder for detection and grading of esophageal varices by multidetector computed tomography. *Eur. J. Radiol.*, 83 (3): 497-502, 2014.
- 19- DENG H., QI X., ZHANG Y., et al.: Diagnostic accuracy of contrast-enhanced computed tomography for esophageal varices in liver cirrhosis: A retrospective observational study. *J. Evid. Based Med.*, 10 (1): 46-52, 2017.

الملخص العربي

دراسة مقارنة بين الأشعة المقطعية متعددة الكواشف ومنظار المرئ في تقييم درجات دوالي المرئ

إن ارتفاع ضغط الوريد الباطني و دوالي المرئ من مضاعفات تليف الكبد. تشخيص دوالي المرئ باستخدام المنظار كوسيلة للفحص محدود، نافذ ؛ مكلف ويحتاج الي تخدير كما أنه قد يتم رفضه من المريض. على العكس فالتصوير بالأشعة المقطعية غير نافذ ، لا يحتاج إلى تخدير ومتقبل من المرضى كما يتيح تشخيص أدق لمكان وحجم الدوالي من المنظار.

الهدف من البحث: إثبات أن الأشعة المقطعية متعددة الكواشف يمكن أن تستخدم كبديل غير نافذ لمنظار المرئ في تشخيص درجات دوالي المرئ.

المرضى: تشتمل هذه الدراسة على خمسين مريض لديهم تليف بالكبد تم فحصهم بالأشعة المقطعية متعددة الكواشف لتشخيص دوالي المرئ ودرجاتها مع مقارنتها بمنظار المرئ وتجديد درجاتها كل على حده. كما أتاحت المقطعية تشخيص إصابات ما خارج المرئ والتي لا تمكن بواسطة المنظار.

معايير الاشتمال: المرضى الذين يعانون من:

- ارتفاع ضغط الوريد الباطني.
- تليف بالكبد.
- سرطان الكبد.

معايير الاستبعاد: المرضى الذين لديهم :

- حدوث مضاعفات لاستخدام صبغة المقطعية.
- نزيف نشط بالجهاز الهضمي.
- ربط دوالي المرئ بالمنظار.

الطرق:

- أخذ التاريخ المرضي للمريض من حيث إصابة الكبد أو قئ دموي أو حقن دوالي مرئ سابق وغيرها.
- الفحص اكلينيكي كامل وتقييم العلامات الحيوية.
- فحص التحاليل الطبية للمريض.
- مراجعة الاشعات السابقة اذا توافرت مع المريض.
- عمل اشعة مقطعية متعددة الكواشف على المرئ ومنظار مرئ .

تم مقارنة نتائج الأشعة المقطعية متعددة الكواشف على المرئ مع نتائج فحص منظار المرئ وتسجيلها وحساب درجات حساسياتها ودقة كل منها في التشخيص.

وعلى ذلك فإن الاشعة المقطعية بالصبغة هي طريقة تشخيصيه بديله متطورة وغير نافذه لتحديد درجات دوالي المريء مقارنة بمنظار المريء وتعطي معلومات اضافية عما خارج المريء من إصابات مما يسمح بنتائج أفضل في تقييم درجات دوالي المريء.