

The Role of Ultrasound Guidance in Tranradial Coronary Catheterization

K.E.El- Rabbat, A.I.Atia, M.S.Abdel Moneum and M.N.Tadros
Cardiology Dept., Faculty of Medicine, Benha Univ., Benha, Egypt
E-Mail: M.N2356@gmail.com

Abstract

Early trans-radial operators adopted the palpatory technique probably due to operator comfort. The challenging bimanual technique needed to use both vascular ultrasound probe and the puncture needle at the same time. Radial approach has some complications like radial artery occlusion, radial artery spasm, hematoma, pseudo-aneurysm, radial artery dissection, bleeding and upper extremity dysfunction. Therefore, proper physical assessment of the forearm, wrist and hand is essential before the procedure. The aim of this study is to evaluate; Feasibility, real-time and complications of trans-radial access comparing the ultrasound guidance versus the fingertip palpatory approach in coronary diagnostic and interventional procedures. Methods: One hundred patients were admitted to the hospital and planned for diagnostic coronary angiography and or elective percutaneous coronary intervention via trans-radial access. patients were divided into two groups Group 1 patients undergone coronary intervention via palpatory fingertip guided approach for transradial access. Group two patients undergone coronary intervention via ultrasound guided approach for transradial access. Procedural outcomes of the whole study population, 7% end by spasm, 3% had bleeding complications, and 9% end by Sheath insertion failure and crossover to another site .there was no statistical significant difference between two groups as regard creatinine level and LVEF .there was no statistical significant difference between two groups as regard technical data or medications used, while there was statistical significant difference between groups as regard attempts of cannulation, time to access, first pass success and difficult access ≥ 5 attempts. There was no statistical significant difference between groups as regard spasm or bleeding complications, while there was statistical significant difference between groups as regard Sheath insertion failure and crossover to another site with lower incidence among US guided group. UGRAA improved first-attempt success rates and shortened the time needed for radial artery cannulation when compared to historical data of palpation-directed radial artery access. Furthermore, this technique is easy to learn and incorporate into everyday practice in the cardiac catheterization lab.

1. Introduction

The blood vessel supply of the hand starting with those spiral What's more ulna arches, the outspread course - Dissimilar to the femoral or brachial corridor – will be thusly not an end conduit in the vicinity of a acceptable ulna insurance supply; its impediment doesn't trade off those vascular supply of the hand. Moreover, the shallow course of the distal outspread course gives to simple layering of the artery, something like that patients could activate Likewise before long Concerning illustration the system will be carried out. Different points of interest are easier frequency for right site complications, progressed tolerant fulfillment Furthermore easier expense because of those abbreviated period of hospitalization [1]. Those spiral approach of the coronary angiography Also intercession got those favored vascular get overall.

To general, promptly transradial operators embraced those palpatory strategy Presumably because of driver solace. Those testing bimanual system required to utilize both vascular ultrasonography probe and the puncture needle at those same time [2]. Spiral methodology need A percentage difficulties like outspread course occlusion, outspread course spasm, hematoma, pseudoaneurysm,

outspread course dissection, dying What's more upper limit brokenness.

Therefore, correct physical appraisal of the forearm, wrist Also hand is key When the system.

In place will appropriately behavior an evaluation, the patient's hand coursing library ought further bolstering make legitimately evaluated by pulse oximetry to assess those palmar coursing library of the hand [3].

Nonetheless there will be thick, as restricted examination accessible taking a gander at the results of transradial get for upper limit function, outspread course thrombosis What's more analyzation are those The majority basic reason for outspread course occlusion, new anticoagulants for example, bivalirudin decline its episodes then afterward transradial conduit catheterization [4].

Ultrasonography direction facilitates the vascular entry At need not been every now and again tried for trials will sway transradial cardiovascular catheterization.

We hypothesize that , appraisal about ultrasonography direction in transradial methodology versus the fingertip palpatory methodology clinched alongside coronary symptomatic Furthermore interventional methods might bring a sway will encourage

protected, quick Furthermore All the more effective right cardiovascular catheterization. [5]. Fitting preparation of the ultrasonography probe assumes An basic part On effectively performed ultrasonography guided outspread course entry. The point when utilizing the palpation guided methods, various endeavors might Sporadically foundation outspread course fit and the vesperula vulgaris might turn into non-palpable after three alternately more unsuccessful endeavors. Utilizing ultrasonography guided outspread course get to each case not just following the palpation strategy needed failed, it might be excessively late a result the conduit might as of now a chance to be over fit. Such An baffling circumstances might dishearten a femoralist starting with adopting those spiral approach. Hybrid from spiral to femoral right may be very nearly nonexistent The point when utilizing ultrasonography guided outspread course entry. The Taking in bend may be short. Utilizing ultrasonography guided outspread course right considers An finer seeing of ordinary outspread course life systems Also appreciation about varieties of the ordinary examine [6]. Those point about this contemplate might have been with evaluate; Feasibility, ongoing Furthermore difficulties from claiming transradial entry contrasting the ultrasonography direction versus those fingertip palpatory approach over coronary symptomatic What's more interventional methods.

2. Patients and methods

One hundred patients were admitted to the emergency clinic and got ready for analytic coronary angiography and additionally elective percutaneous coronary intercession by means of trans-outspread access .

• patients were divided into two groups :

Group one

50 patients undergone coronary intervention via palpatory fingertip guided approach for transradial access.

Group two

50 patients undergone coronary intervention via ultrasound guided approach for transradial access.

I-Inclusion criteria

Patients indicated for radial coronary intervention either diagnostic coronary angiography or percutaneous intervention.

II-Exclusion criteria

1. Patients who refused to give their written informed consent .

2. Patients in whom the radial approach is contraindicated .
3. Patients with a dialysis arteriovenous fistula or in whom A-V fistula is planned.
4. Patients with a known radial artery occlusion .
5. Challenging vessel anatomy , vessel tortiousity .
6. Known difficulty to cannulate the radial artery .
7. Patients in whom the Allen test results were positive .
8. In case of pathological Allen test in one hand the other hand will be tested before switching to the femoral access.
9. Unstable patients (myocardial infarction , cardiogenic shock .)
10. Patients necessating femoral access for an intra-aorticballoon counter pulsation .
- 11 Patients presenting with double mammary coronary graft.
12. Coronary interventions where an extreme good guiding support is mandatory (i.e PCI of chronic total occlusion) .
13. High risk coronary interventions (e.g Left main interventions) where the femoral approach may be more comfortable to the operator .
14. Patients requiring a femoral approach for a right heart catheterization

III-Methodology in details

All patients were subjected to the following :

1. Written informed consent for coronary intervention Either diagnostic or therapeutic via radial approach .
2. Complete history taking .
3. Twelve lead Electrocardiogram .
4. Echocardiography evaluation .
5. Assessment of radial artery by ultrasound before , during and after the procedure for the scheduled group .
6. Allen test before and after the procedure
7. Assessment of the hand vascularity by Pulse oximetry before and after the procedure .
8. Clinical assessment of radial pulsations before and after the procedure .
9. Coronary angiography via radial approach .

• Puncture

The radial artery was punctured by a 21 gauge needle , the pucture site is 1-3 cm proximal to the styloid process , at 45 degrees from lateral to medial

In the second group the radial artery was accessed guided by ultrasound transducer placed perpendicular to the radial artery 1 cm from the puncture site .

- **Sheath insertion**

A 0.025 mm J shaped wire was introduced through the needle without using force .A small skin incision was made by number 11 surgical blade to prevent damage of the radial artery . a 6 french sheath was inserted in the radial artery in order to prevent spasm and to reduce discomfort during cathetersexchange .**Ultrasound guidance** of the guide wire can be used in case of wire resistance or impaction.

- **Local medication**

The following drugs were administered intra-arterial to prevent arterial spasm and thrombosis .

1. 5000 I.U unfractionated heparin to prevent intra arterial thrombosis.
2. 200 mcg nitroglycerin .
3. 3-5 mg verapamil .

N.B: These medications were susceptible to readjustment in dose or being avoided according to the patient's blood pressure and systolic function at the time of the procedure .

- **Sheath removal**

The sheath was expelled following the system , a committed outspread haemostatic gadget (TR band) was applied and expanded at the cut site, the weight of the band was step by step diminished in the accompanying 2 hours .

Cautious follow up to the hand and lower arm was done to distinguish any neighborhood difficulty . All the patients were appropriately evaluated by ultrasound duplex to the outspread course before release , patients were encouraged to confine their wrist development during the accompanying 24 hours .

- **Primary outcomes**

Ultrasound guidance facilitates transradial access.

- **Secondary outcomes**

Comparing the ultrasound transradial approach to the palpatory fingertip approach in terms of :

1. Number of attempts .
2. First pass success rate .
3. Time to access.
4. Difficult access .
5. Crossover to another site .
6. Access site complications (bleeding ,haematoma , pseudoaneurysm , dissection , radial artery occlusion and spasm) .

3.Results


Our result shows that mean age of participants is 56.2 ± 6.9 years old, Mean \pm SD of BMI is 29.7 ± 3.8 Kg/m² , 66% of participants were male, 36% had DM, 58% were smokers, 45% had dyslipidemia, 40% were obese and 31% had family history. median heart rate of the participants is 80 beat per minute, median SBP is 120 mmHg and median of DBP is 80 mmHg. Mean \pm SD of Creatinine is 0.97 ± 0.19 mg/dL and median of left ventricular ejection fraction is 54%.

39% of participants had intervention, 52% had 5-F sheath, majority 92% had Rt radial access, 85% underwent Single-wall technique, and 63% underwent TR band closure,.

Regarding medications used, majority of cases 97% had Verapamil ≥ 2.5 mg, 86% take Unfractionated heparin, 78% take Nitroglycerine ≥ 100 μ g, 53% take P2Y12 inhibitor, 51% take Lidocaine IA 5 mg, and only 9% take GP 2b/3a inhibitor.

Regarding access, 54% had first pass access, 9% had ≥ 5 attempts, 8 % take ≥ 5 minutes. Procedural outcomes of the whole study population, 7% end by spasm, 3% had bleeding complications, and 9% end by Sheath insertion failure and crossover to another site.

US guidance improved first-pass success rate by 68% (95% CI: 55.1% to 80.9%) vs. 40% (95% CI: 26.4% to 53.6%) for palpation method, $p = 0.005$.






Fig (1) Comparison between the studied groups regarding the procedural data

Table (1) Comparison between the studied groups regarding the procedural outcomes.

Procedural data	Palpation	US guided	P-value (Sig.)
Count	50	50	
Outcomes			
Spasm	3 (6%)	4 (8%)	1.000 (NS)
Bleeding complications	2 (4%)	1 (2%)	1.000 (NS)
Sheath insertion failure and crossover to another site	8 (16%)	1 (2%)	0.031 (S)

p< 0.05 is significant. Sig.: significance.

4. Discussion

An instance control contemplate might have been conveyed out including one hundred patients admitted of the healing facility and wanted to symptomatic coronary angiography. What's more or elective percutaneous coronary mediation through trans-radial right. Patients were partitioned under two groups: assembly one: 50 patients undergone coronary mediation through palpatory fingertip guided methodology to transradial get. Gathering two : 50 patients undergone coronary intercession by means of ultrasonography guided approach for transradial right. The span of the examine ranges starting with 6 should 12 months.

Those mean period of members will be 56. 2 ± 6.9 quite some time old, mean \pm sd of BMI is 29.7 ± 3.8 Kgm², 66% from claiming members were male, 36% required DM, 58% were smokers, 45% required dyslipidemia, 40% were large Furthermore 31% required family history. There might have been no measurable noteworthy Contrast between two bunches as view the benchmark qualities.

Our comes about would underpinned via examine Brueck et al., [7] for Concerning illustration they news person that the intend period about their patients might have been 63.2 ± 11.9 quite some time old, imply \pm sd of BMI may be 28.4 ± 5.8 Kgm², 57% about patients were male. There might have been no statistically huge distinction between two aggregations in regards the benchmark aspects.

As stated by Seto et al., [8] discovered that the mean period of their patients might have been 61.5 ± 10.6 quite some time old, mean \pm sd from claiming BMI may be 30.2 ± 7.2 Kgm², 73% about patients were male. There might have been no statistically huge distinction the middle of two Assemblies viewing the benchmark aspects.

Likewise see clinical data, the exhibit investigation indicates that average heart rate of the members may be 80 beat for every minute, average SBP is 120 mmhg Also average about DBP is 80 mmhg. Intend \pm sd for Creatinine will be 0.97 ± 0.19 mg/dL and average for left ventricular launch portion is

54%. There might have been no Factual huge distinction the middle of two Assemblies viewing those clinical information Similarly as heart rate, SBP, DBP, Creatinine level Also LVEF.

As stated by Brueck et al., [7] watched that during discharge, they found no pulse in the cannulated outspread course over 3 patients (0.58%) without clinical indications for lower arm ischemia. Clinched alongside other trials, passing of outspread beat might have been exhibit done 0% to 9% (Benit et al., [9]) without clinical sequelae. Utilizing ultrasonography assessment, the post-procedural nonattendance of a spiral stream might have been distinguished to 9% (1,25), therefore diminishing with 3% will 6% done catch up [10].

Accompanying those to start with reports for spiral coronary angiography by Lucian Campeau Previously, 1989 What's more spiral percutaneous coronary intercession (PCI) Toward Ferdinand Kiemeneij Previously, 1992, the extent for spiral methods need proceeded to Ascent worldwide, for spiral right presently reinstating femoral right Similarly as the prevailing right site for PCI done exactly nations. Worldwide, an evaluated 20% for methods need aid performed by this course (29%, though the u. Encountered with urban decay because of deindustrialization, engineering imagined, government lodgin. May be excluded from the estimate) [11].

The current examine indicates that 39% from claiming Members needed intervention, 52% needed 5-F sheath, lion's share 92% required Rt spiral access, 85% underwent Single-wall technique, Furthermore 63% underwent TR band closure,. In regards to medications used, greater part from claiming instances 97% needed verapamil ≥ 2.5 mg, 86% detract Unfractionated heparin, 78% take nitroglycerine ≥ 100 μ g, 53% make P2Y12 inhibitor, 51% take lidocaine ia 5 mg, and best 9% detract GP 2b/3a inhibitor. In regards access, 54% needed initial pasquinade access, 9% needed ≥ 5 attempts, 8% detract ≥ 5 minutes.

There might have been no measurable critical distinction the middle of two Assemblies as see specialized foul information alternately medications used, same time there might have been Factual huge Contrast the middle of gatherings as view endeavors from claiming cannulation, chance should access, 1st pasquinade victory What's more challenging get ≥ 5 endeavors.

Our comes about need aid underpinned via ponder for Seto et al., [8] Likewise they news person that pretty nearly 20% of methods were intercessions. Those lion's share from claiming methods were performed utilizing single-wall system. Conclusion might have been predominantly with those TR band, yet the D-stat Rad-bands utilized within 1 site only. Manual layering might have been utilized within $<3\%$ for methods. 97% of them detract verapamil Similarly as prescription. There might have been a critical decrease in the number for endeavors required should cannulate those outspread course with us versus palpation (mean: 1.65 1.2 vs. 3.05 3.4, $p < 0.0001$, median: 1 [IQR: 1 to 2] vs. 2 [IQR: 1 should 3], $p < 0.0001$).

Those discoveries from claiming this consider were reliable with that from claiming past trials thinking about ongoing 2-dimensional us direction to spiral blood vessel transport placement. Levin et al., [12] discovered over 69 grown-up patients requiring intraoperative screening a change to first-pass triumph rate starting with 34% should 62% ($p \frac{1}{4} 0.03$) with a mean number from claiming endeavors from claiming 3.1 2.4 versus 1.6 1.0 ($p \frac{1}{4} 0.003$). Schwemmer et al. [13] found done 30 babies that the first-pass prosperity rate might have been expanded starting with 20% to 67% ($p < 0.05$). Shudder et al. [14] demonstrated done 60 critically sick crisis room patients an change Previously, first-pass triumph rate from half will 87% ($p \frac{1}{4} 0.005$) What's more a diminishment in time from 314 encountered with urban decay because of deindustrialization, innovation developed, government lodgin versus 107 s ($p \frac{1}{4} 0.0004$). Done 1 trial, us didn't have a benefit, which might have been felt by those investigators on be expected on insufflate experience, with 94% about operators Hosting <5 US-guided methods [15]. These discoveries were summarized in An meta-analysis, which suggester a 17% outright change (26% on 43%) done firstpass victory rate [16].

These outcomes infer that "seeing" those little outspread course around us might be a greater amount exact over "feeling" those artery, because of those 2- will 4-mm 2-point separation breaking point about fingertip

palpation (17). Us direction may be especially advantageous in the individuals patients with troublesome get for palpation alone. A powerless pulse might aggravate palpation-guided right difficult, if brought on by An little artery, An profound artery, alternately hypotension. For hypotension, the blood vessel flashback in the needle might make moderate alternately fleeting though the posterior divider is punctured. With An little alternately bulky conduit it might be challenging to look after the ideal intraluminal needle position important for wire insertion Toward single-wall system. For congestive heart disappointment What's more raised venous pressures, those spiral veins might widen Also a chance to be coincidentally cannulated. For a calcified artery, the vesputa vulgaris might move away from the needle tip or require extra drive will cannulate. For a coagulated needle, effective cannulation of the conduit might try unrecognized. Us direction conceivably addresses all from claiming these circumstances Toward outwardly affirming needle position over the conduit preceding cannulation, layering Also puncture of the blood vessel lumen, or position of the needle tip inside the lumen.

Coronary angiography Also angioplasty would normally performed by means of the transfemoral methodology. Transradial right for coronary vein catheterization offers A percentage favorable circumstances in examination with the transfemoral course. Particularly under states about Combative anticoagulation Furthermore antiplatelet treatment, vascular dying difficulties during the femoral puncture site camwood bring about expanded horribleness Furthermore span of hospitalization. Therefore, those method of reasoning for the transradial methodology may be the purposeful to decrease right site dying complications, prior ambulation, Furthermore progressed tolerant solace [18]. Right webpage difficulties need aid Extensively All the more incessant At whatever point a Forceful antiplatelet or antithrombotic medication may be required. Consequently, transfemoral mediation done intense myocardial localized necrosis carries An hazard about dying difficulties going from 2.5% will 23%. Obesity, elderly patients, and female sex need likewise been interfaced to an expanded event of get site difficulties [19].

In the investigation clinched alongside our hands, Procedural results of the entirety study population, 7% limit Toward spasm, 3% needed dying complications, What's more 9% limit Toward sheath insertion disappointment What's more hybrid with another webpage.

There might have been no Factual critical distinction the middle of aggregations Concerning illustration view fit or dying complications, same time there might have been Factual critical Contrast between aggregations Likewise respect sheath insertion disappointment Also hybrid on another site for bring down occurrence Around us guided bunch.

Our outcomes would in line with various randomized trials needed uncovered that transradial cardiovascular catheterization Might diminish difficulties Furthermore give acceptable more stupendous tolerant solace over that watched for the transfemoral approach, with ensuing diminishment to doctor's facility regulatory issues Furthermore expenses [20]. Extreme right site bleeding, especially retroperitoneally Might make deadly mishap Furthermore may require transfusion [21]. Reportedly, checked discharge is connected with poor prognosis watched Throughout both, short Furthermore long haul catch up [22]. Therefore, transradial catheterization may be energized Comprehensively in the current period [23].

Viewing Brueck et al., [7], news person that they Might affirm the low rate of passage webpage difficulties utilizing those transradial approach, as to be sure best 3 neighborhood difficulties (0. 58%) were discovered to 512 patients. Those outspread course may be effortlessly compressible because of its shallow course, accomplishing sufficient hemostasis just with An wrap. On their knowledge, this may be those initial randomized trial contrasting entry webpage difficulties following coronary methods by means of transradial versus transfemoral get for a Conclusion gadget. Much with the technologic change about these units Furthermore expanding knowledge viewing their use, they exhibited an acceptable profit of the transradial approach Similarly as of the event of fringe blood vessel difficulties.

Furthermore, Kashiwagi et al., [24] showed that the provision of ongoing UG salvage might have been successful to transradial cardiovascular catheterization. Despite the procedural chance might have been longer in the UG over in the PT group, the accomplishment rate of the transradial approach might have been essentially higher in the UG aggregation. Those TR band might have been utilized to hemostasis On the whole patients, and no genuine entry site difficulties were watched. Left outspread course get might have been that's only the tip of the iceberg basic in the UG over in the PT assembly (88% vs. 74%, $p < 0. 01$). No statistically noteworthy

intergroup Contrast might have been watched in the occurrence for hematoma (6. 8% vs. 5. 8%, $p=0. 62$) (Figure 3-B). The procedural duration of the time in the UG might have been more over that in the PT aggregation (303 encountered with urban decay because of deindustrialization, engineering concocted, government lodgi vs. 357 s, $p < 0. 01$). No tolerant created a extreme hematoma requiring transfusion or nerve damage.

Viewing Seto et al. [8], there were no huge contrasts clinched alongside clinical results between those 2 gatherings. Fit Furthermore dying difficulties were accounted for just as and infrequently for both groups, Also patient-reported agony scores were low clinched alongside both aggregations. Despite us direction might have been not used to screen to outspread course occlusion, symptomatic outspread course impediment might have been not accounted for done schedule clinical catch up Eventually Tom's perusing any site. Ten patients in the palpation one assembly needed hybrid will us direction after 5 min, with us effectively rescuing the entry done 8 cases. Hybrid with another webpage (usually femoral) happened Sporadically accompanying sheath insertion in both gatherings because of spasm, tortuosity, alternately insufflate aide help. Hybrid to in turn webpage former with sheath insertion (failed access) struck them numerically All the more every now and again in the palpation gathering over the us bunch. Disappointment from claiming sheath insertion happened for 15 patients with palpation alone compared for 3 with us ($p \frac{1}{4} 0. 007$).

Fit need been determinedly What's more freely connected with various endeavors toward access, conceivably because of expanded damage of the vesputa vulgaris or encompassing tissue. Two investigations discovered that unsuccessful get once main endeavor might have been freely connected with 45% should half expanded danger from claiming outspread course fit [25,26].

Done an expansive registry, Goldsmit et al. , [27] discovered that those necessity for more than 1 puncture endeavor might have been connected with a or about 3. 5 (95% CI: 1. 9 with 6. 3) for moderate/ extreme fit.

It is Additionally could reasonably be expected that those correspondence between various endeavors Also fit may be mostly the effect of a regular element for example, outspread course span or calcification. Main in the examine Eventually Tom's perusing Jia et al., [26] were both outspread course span What's more amount of endeavors evaluated over multivariable analysis, and same time the number of endeavors stayed a free predictor

for fit (OR: 1.5), outspread course size might have been a stronger predictor (OR: 4.0). On their experience, they required watching that fit is a significant part less averse with happen when an generally little or calcified outspread course may be present, which is steady for the perceptions of Saito et al., [28].

References

- [1] J.T. Mann 3rd, M.G. Cubeddu, J.E. Schneider, M. Arrowood. Right radial access for PTCA: a prospective study demonstrates reduced complication and hospital charge. *J Invas*, Vol. 25(5), PP.420-425, 2007.
- [2] O.F. Bertrand, S.V. Rao, S. Panchoy, Transradial approach for coronary angiography and interventions: results of the first international transradial practice survey. *JACC Cardiovasc Interv*, Vol.3, PP.1022-31, 2010.
- [3] C. Joy, School of rehabilitation Science, McMaster University, Hamilton, Ontario, Canada Clinical Research Lab, Hand and Upper limb Centre, St. Joseph Health Centre, London, Ontario, Canada Patient Rated Wrist Evaluation, Vol.30(8), PP.201-210, 2007.
- [4] S. Plante, W.J. Cantor, L. Goldman, Comparison of bivalirudin versus heparin on radial artery occlusion after transradial catheterization. *Catheter Cardiovasc Interv*, Vol.76(5), PP.654-658, 2010.
- [5] J. Magee David, orthopedic physical assessment, fifth edition, in forearm, wrist and hand; chapter seven, vol 60(3), PP. 445-446, 2015.
- [6] J. Roberts, M. Ardid, Ultrasound guided radial artery access by a non ultrasound trained interventionalist cardiologist improved first attempt success rates and shortened time for successful Radial Artery Cannulation., Vol.60(17-s), PP.B113, 2012.
- [7] M. Brueck, D. Bandorski, W. Kramer, A randomized comparison of transradial versus transfemoral approach for coronary angiography and angioplasty. *JACC: Cardiovascular Interventions*, Vol. 2(11), PP.1047-1054, 2009.
- [8] A. H. Seto, J. S. Roberts, M. S. Abu-Fadel, Real-time ultrasound guidance facilitates transradial access: RAUST (Radial Artery access with Ultrasound Trial). *JACC: Cardiovascular Interventions*, Vol.8(2), PP.283-291, 2015.
- [9] E. Benit, L. Missault, T. Berman, Brachial, radial or femoral approach for elective Palmaz-Schatz implantation: a randomized comparison. *Catheter Cardiovasc Diagn*, Vol.41, PP.124-30, 1997.
- [10] Y. Louvard, T. Lefèvre, A. Allain, Coronary angiography through the radial or the femoral approach: the CARAFE study. *Catheter Cardiovasc Interv*, Vol.52, PP.181-7, 2001.
- [11] S. V. Rao, J. A. Tremmel, I. C. Gilchrist, Best practices for transradial angiography and intervention: a consensus statement from the society for cardiovascular angiography and intervention's transradial working group. *Catheterization and Cardiovascular Interventions*, Vol. 83(2), PP. 228-236.
- [12] P.D. Levin, O. Sheinin, Y. Gozal, Use of ultrasound guidance in the insertion of radial artery catheters. *Crit Care Med*, Vol.31, PP.481-4, 2003.
- [13] U. Schwemmer, H.A. Arzet, H. Trautner, Ultrasound-guided arterial cannulation in infants improves success rate. *Eur J Anaesthesiol*, Vol.23, PP.476-80, 2006.
- [14] S. Shiver, M. Blaivas, M. Lyon, A prospective comparison of ultrasound-guided and blindly placed radial arterial catheters. *Acad Emerg Med*, Vol.13, PP.1275-9, 2006.
- [15] A. Ganesh, R. Kaye, A.M. Cahill, Evaluation of ultrasound-guided radial artery cannulation in children. *Pediatr Crit Care Med*, Vol.10, PP.45-8, 2009.
- [16] A. L. Shiloh, R.H. Savel, L.M. Paulin, Eisen Ultrasound-guided catheterization of the radial artery: a systematic review and meta-analysis of randomized controlled trials. *Chest*, Vol.139, PP. 524-529, 2011.
- [17] W.W. Campbell. DeJong's the Neurologic Examination. 6th edition. Philadelphia, PA: Lippincott Williams & Wilkins, Vol.80, PP. 453, 2005.
- [18] W.J. Cantor, G. Puley, M.K. Natarajan, Radial versus femoral access for emergent percutaneous coronary intervention with adjunct glycoprotein IIb/IIIa inhibition in acute myocardial infarction—the RADIAL-AMI pilot randomized trial. *Am Heart J*;150:543-9, 2005.
- [19] M. Uhlemann, S. Möbius-Winkler, M. Mende, I. Eitel, The Leipzig prospective vascular ultrasound registry in radial artery catheterization: impact of sheath size on vascular complications. *JACC: Cardiovascular Interventions*, Vol. 5(1), PP.36-43, 2012.
- [20] C.J. Cooper, R.A. El-Shiekh, D.J. Cohen, Effect of transradial access on quality of life and cost of cardiac catheterization: A

- randomized comparison. *Am Heart J*, Vol.138, PP.:430–6, 1999.
- [21] A.N. Vora, M. Stanislawski, G.K. Grunwald, Association Between Chronic Kidney Disease and Rates of Transfusion and Progression to End-Stage Renal Disease in Patients Undergoing Transradial Versus Transfemoral Cardiac Catheterization-An Analysis From the Veterans Affairs Clinical Assessment Reporting and Tracking (CART) Program. *J Am Heart Assoc*, Vol.6(4).PP. e004819,2017.
- [22] Y. Tanaka, N. Moriyama, T. Ochiai, Transradial Coronary Interventions for Complex Chronic Total Occlusions. *JACC Cardiovasc Interv*, Vol.10, PP.235–243,2017.
- [23] O.F. Bertrand, S.V. Rao, S. Pancholy, Transradial approach for coronary angiography and interventions: results of the first international transradial practice survey. *JACC Cardiovasc Interv*, Vol.3, PP.1022–31,2010.
- [24] M. Kashiwagi, T. Tanimoto, H. Kitabata, Usefulness of rescue ultrasound guidance for transradial cardiac catheterization. *Cardiovascular Revascularization Medicine*, Vol.20(4), PP. 311-315,2019.
- [25] J. Rosencher, A. Chaïb, F. Barbou, How to limit radial artery spasm during percutaneous coronary interventions: The SPasmolytic Agents to avoid SpasM during transradial percutaneous coronary interventions (SPASM3) study. *Catheter Cardiovasc Interv*, Vol.84, PP.766–71,2014.
- [26] D.A. Jia, Y.J. Zhou, D.M. Shi, Incidence and predictors of radial artery spasm during transradial coronary angiography and intervention. *Chin Med J (Engl)*, Vol.123, PP.843–7,2010.
- [27] A. Goldsmit, F. Kiemeneij, I.C. Gilchrist, Radial artery spasm associated with transradial cardiovascular procedures: results from the RAS registry. *Catheter Cardiovasc Interv*, Vol.83, PP. E32–6,2014.
- [28] S. Saito, H. Ikei, G. Hosokawa, Influence of the ratio between radial artery inner diameter and sheath outer diameter on radial artery flow after transradial coronary intervention. *Catheter Cardiovasc Interv*, Vol. 46, PP.173–8, 1999.