



DJS Vol. 36 (2013) 179-185		
	<b>Delta Journal of Science</b> Available online at <a href="https://djs.journals.ekb.eg/">https://djs.journals.ekb.eg/</a>	
Research Article		ZOOLOGY

## Characterization of ITS1 gene of *Leishmania infantum* isolated from Iraqi patients with visceral leishmaniasis by PCR- RFLP and sequencing methods.

**Ilham A. Majeed<sup>1</sup>**

**Fatemeh Gaffarifar<sup>2</sup>**

**Nada Al-Bashier<sup>3</sup>**

1. Medical Laboratory Dept. College of Health & Medical Technology/ Baghdad/ Iraq.

2 Parasitology and Entomology Dept., Faculty of Medical Sciences, Tarbiat Modares University,  
P.O.Box 14115-331.

3. Medical Research Center/ College of Medicine/ Al-Nahrain University/ Baghdad/ Iraq.

### ABSTRACT

For the best of our knowledge, there is no information about molecular characterization of Iraqi isolates of visceral leishmaniasis, the present work aimed to characterize three different Iraqi isolates of *Leishmania infantum* by polymerase chain reaction (PCR), restriction fragment length polymorphism (RFLP) and sequencing methods.

Three isolates from bone marrow of Iraqi patients infected with kala azar were used in this study. The isolates were already diagnosed by isoenzyme as *Leishmania infantum*. Patients were inhabiting different parts of Baghdad. The samples after microscopic examination were cultured on modified NNN media. Then DNA was extracted for amplifying ITS1 (internal transcribed spacer 1) gene by PCR. Identification of samples was studied using RFLP (digestion with Apo1 restriction enzyme) and sequencing of PCR products.

The PCR of all samples showed a band under about 500 bp. The results by using PCR-RFLP method showed no restriction with digestion with Apo1 restriction enzyme. The results of sequencing showed differences with all separated gene from *Leishmania infantum* in the gene bank.

In this study we found that the sequences of ITS1 gene of *Leishmania infantum* separated from Iraqi patients are different from other samples, as there is no similarity with *Leishmania infantum* (MHOM/TN/80/IP11). The more similarity is with the Iranian isolate of *Leishmania infantum* (MCAN/IR/97/LON) with 41%, whereas the similarity with *Crithidia luciliae* internal transcribed spacer 1, ITS1 is 96%.

**Key words: Visceral leishmaniasis, PCR-RFLP, sequencing, Iraq**

### INTRODUCTION

Leishmaniasis is endemic in 88 countries and 12 million people per year have been estimated to be at risk of infection. Leishmaniasis forms a spectrum of diseases ranging from benign, self-healing skin ulcer to overwhelming lethal systemic infection (1).

The leishmaniasis comprise a group of diseases that display widely different clinical manifestations in humans, which depend not only on the species initiating infection but also on

the general health and genetic make-up of the infected individual (2).

Visceral leishmaniasis is a serious health problem and an important endemic disease in Iraq. During the last years there was an increase in visceral leishmaniasis cases in Iraq (3).

Majeed *et al.* (2012) reported that in Iraq, visceral leishmaniasis incidence per 100 000 population was 2.6 in 2007, 3.1 in 2008, and 4.8 in 2009, mostly in children aged <5 years (4).

In Iraq visceral leishmaniasis is usually detected in infants and children (5, 6). According to Zukerman & Lainson at 1977,

visceral leishmaniasis in Iraq is considered to be of infantile type (7).

Correct identification of the etiologic agent of leishmaniasis is important for clinical, epidemiologic reasons, prevention, and treatment. Therefore, molecular characterization based on partial sequences of *Leishmania* species is important.

ITS1 is a suitable target for detection, identification and characterization of *Leishmania* spp for this reason, this gene was used in this study. In the present study, PCR and sequencing of ITS1 gene were employed to study visceral leishmaniasis of the Iraqi isolates.

## Materials and Methods

### 1) Isolates:

Three Iraqi isolates of *L. infantum* was used in this study, they were collected by bone marrow aspirate from three patients living in different parts of rural areas of Baghdad.

Those patients were all less than five years old, one girl and two boys.

The isolates were cultivated on liquid NNN medium containing RPMI

so that large number of promastigote can be harvested for DNA extraction,

also *Leishmania infantum* strain (MHOM/TN/80/IP11) was cultivated and used as positive control.

### 2) DNA extraction:

About  $2 \times 10^6$  of promastigotes of each isolates and MHOM/TN/80/IP11 strain were initially washed with phosphate buffered saline (PBS).

The procedure of the extraction of DNA was followed according to the manufacturer instruction of the kit (Bioneer Corporation).

The DNA extraction products were detected in 0.8% agarose gel and photographed. The results showed that concentration of the extracted DNA was high.

### 3) PCR amplification and gel electrophoresis

Genomic DNA from promastigotes of all isolates and *Leishmania infantum* strain (MHOM/TN/80/IP11) were used as templates to amplify the ITS1 gene by PCR. The reaction was performed in 25 $\mu$ l of the solution containing, 3 $\mu$ l of template DNA, 0.5 $\mu$ l of dNTP (with concentration 10mM and final concentration of 200  $\mu$ M), 0.5 $\mu$ l of Taq DNA polymerase (with concentration 5 Unit/ml), 2.5 $\mu$ l of 10X PCR buffer, 0.75 $\mu$ l of MgCl<sub>2</sub> (50 mM), 15.75 $\mu$ l of distilled water and 1 $\mu$ l of each of primers (10Pmol/ $\mu$ l).

The primer sequences that we used in this study were as follows (Spanakos et al., 2008) (8):

LeF, 5'-TCCGCCCCGAAAGTTACCGATA-3and

LeR, 5'-CCAAGTCATCCATCGCGACACG-3';

These primers could detect 18S ribosomal RNA gene, partial sequence; internal transcribed spacer 1, complete sequence; and 5.8S ribosomal RNA gene, partial sequence of *Leishmania* spp.

The PCR program was as follow: 5 min at 94oC for initial denaturation step, then 30 cycles of denaturation at 94oC for 1 min, annealing at 54oC for 30 s, an extension at 72oC for 45 s, and a final extension at 72oC for 5 min.

The PCR products were analyzed by electrophoresis on 1.5% agarose gels, visualized by UV illumination and finally photographed.

### 4) Digestion with restriction endonuclease enzyme (RFLP)

Based on the sequences of *L. infantum* retrieved from Gen Bank, we expected that ApoI could digest the PCR product, then ApoI (Fermentase co.) was selected as a suitable enzyme used for all *L. donovani* complex (Spanakos et al., 2008)(8).

RFLP was performed according to the methods described by (Marfurt et al., 2003) (9). We provided total 31  $\mu$ l volume, include 10  $\mu$ l of PCR products (0.1-0.5  $\mu$ g of DNA), 18  $\mu$ l of nuclease-free water, 2  $\mu$ l of 10X buffer Tango, and 1  $\mu$ l of ApoI enzyme (Fermentase) at 37°C water bath overnight (16 h). The digest was used for electrophoresis on 3.5% agarose gel. Fragment sizes were estimated by their comparison with bands of 100bp DNA length standard ladder.

### 5) DNA sequencing

The PCR products were excised from a 3.5% agarose gel and DNA was isolated using the QIA quick Gel Extraction Kit (Qiagen), according to the manufacturer's instructions.

All PCR products were sequenced with dideoxy method (Fazapajouh Company) using the PCR primers. Using the Clustal W 1.8 software package.

## RESULTS

### 1. Evaluation of the PCR Method

Based on the results of PCR product in the gel electrophoresis, all isolates showed DNA size 477 bp, whereas for MHOM/TN/80/IP11 was about 380 bp (Fig.1).

### 2. RFLP

The resultant restriction fragments of samples were separated on a 3.5% agarose gel and difference was observed in amplicon size between isolates and MHOM/TN/80/IP11 strain.

The PCR product of all isolates didn't digested by ApoI (Fig.2) but MHOM/TN/80/IP11 was digested to two fragments (Fig.3).

### 3. Sequencing

PCR products containing clinical samples and strain MHOM/TN/80/IP11 were sequenced according to F and R primers. All isolates in this study showed similarity sequences (Fig.4).

The results of sequencing indicate that the length of fragments of all clinical isolates were 477 bp. It was additionally found out that there is no restriction site of ApoI (AATT) in ITS1 sequences for all clinical isolates, whereas restriction enzyme site was shown in MHOM/TN/80/IP11 ITS1 sequence.

## Discussion

One of our findings in this study is that the sequences of ITS1 gene of the isolates separated from Iraqi patients are different from other *Leishmania infantum* strains that registered in gene bank.

ITS1 gene is a target to study the characterization of visceral leishmaniasis because ITS1 sequences are adequately polymorphic to detect different strains (Al-Jawabreh *et al.*, 2006; Fryauff *et al.*, 2006) (10,11) and so we can compare our data with many other study on *Leishmania* ITS1 sequences that are saved in sequences data banks (Schonian *et al.*, 2003)(12).

In this study it was found that the sequences of ITS1 gene of *Leishmania infantum* separated from Iraqi patients shows no similarity with *Leishmania infantum* strain (MHOM/TN/80/IP11), and moderate similarity with Iranian isolate of *Leishmania infantum* (MCAN/IR/97/LON) with 41% (Fig. 5), whereas the similarity with *Crithidia luciliae* internal transcribed spacer 1, ITS1 is 96% (Fig.6). We need more molecular studies on other genes of the Iraqi *L. infantum* to explain the similarity with the genus *Crithidia luciliae* and we may explain this similarity as these isolates are new strain of *L. infantum*. The sequence variants of ITS1 might be due to the presence of multiple strains/clones or the existence of intragenomic variations in the multicopy ITS1, or a combination of both. Finally, we may need to revise the taxonomy of the Iraqi strain of *Leishmania infantum*.

### Acknowledgment

This study was performed as part of research activity of staff member of the College of Health & Medical Technology/ Baghdad/ Iraq and supported by the Dept. of Parasitology & Entomology at Tarbiat Modares University/ Tehran. Thank is due to the staff of the department of parasitology at Tarbiat Modares University.

The authors declare there is no conflict of interest.

### References

- Desjeux, P. Leishmaniasis: current situation and new perspectives. *Comp. Immunol., Microbiol. Infect. Dis.* (2004); 27: 305-318.
- Alexander, J. & Russell, D. G. (1992). The interaction of *Leishmania* species with macrophages. *Advan. Parasitol.* (1992); 31: 175-254.
- Rashed, A. Study of Kala-azar in Iraq for the period 1999-2003[F.I.C.M.S dissertation]. Scientific Council of Community and Family Medicine: Baghdad; (2005).
- Majeed, B.; Sobel, J.; Nawar, A.; Badri, S.; Muslim, H. The persisting burden of visceral leishmaniasis in Iraq: data of the National Surveillance System, 1990-2009. *Epidemiol Infect.* (2012); 4:1- 4.
- Hussein, A. Clinical and Laboratories Study for Kala-azar in Thi-Qar City [High Diploma. Thesis]. College of Med. and Heal. Tech. Found. Of Tech. Educ. Baghdad; (2004).
- Rifaat, L. Kh. Comparison of serodiagnostic method in Kala-azar with reference to immunoglobulin changes [M.Sc. Thesis]. Baghdad Univ.; (1988).
- Zuckerman, A. & Lainson, R. *Leishmania*. In: Julius P, Kreier E, editors. *Parasitic protozoa* Vol. I. New York, San Francisco, London: Academic press; (1977); p. 57-133.
8. Spanakos G. & Piperaki, E.T. Detection and species identification of Old World *Leishmania* in clinical samples


using a PCR- based method. *Trans Roy Soc Trop Med Hyg.* (2008); 102:46-53.

Marfurt, J.; Niederwieser, I.; Makia, N.D.; Beck H.P. & Felger, I. Diagnostic genotyping of Old and New World *Leishmania* species by PCR-RFLP. *Diagn. Microbiol. Infect. Dis.* (2003); 46:115-124.


Al-Jawabreh, A.; Schoenian, G.; Hamarsheh, O. & Presber, W. Clinical diagnosis of cutaneous leishmaniasis: a comparison study between standardized graded direct microscopy and ITS1-PCR of Giemsa-stained smears. *Acta Trop.* (2006); 99:55-61.

Fryauff, D.J.; Hanafi, H.A.; Klena, J.D.; Hoel, D.F.; Appawu, M.; Rogers, W.; Pupilampu, N.; Odoom, S.; Kweku, M.; Koram, K. Wilson, M.D.; Racznik, G. & Boakye, D. Short report: ITS-1 DNA sequence confirmation of *Leishmania major* as a cause of cutaneous leishmaniasis from an outbreak focus in the Ho district, Southeastern Ghana. *Am J. Trop. Med. Hyg.* (2006); 75: 502-504.


Schonian, G.; Nasereddin, A.; Dinse, N.; Schweynoch, C.; Schilling, H.D.; Presber, W. & Jaffe, C.L. PCR diagnosis and characterization of *Leishmania* in local and imported clinical samples. *Diagn. Microbiol. Infect. Dis.* (2003); 47:349-358.



**Fig. 1.** Agarose gel electrophoresis of PCR products. Lanes 1 to 3: *Leishmania infantum* isolates from Iraqi patients, (Lane M) is a 100 bp ladder. Lanes 4 : *Leishmania infantum* as standard strain (MHOM/TN/80/IP11).



**Fig.2:** Agarose gel electrophoresis of PCR and RFLP products ( with *ApoI* enzyme). The marker (lane M) is a 100 bp ladder. Lanes 1&3 ITS1 from patients isolates before digestion , Lanes 2 and 4 of after digestion with *APOI* enzyme.



**Fig.3:** Agarose gel electrophoresis of PCR products and after digestion with *ApoI*. The marker (lane M) is a 100 bp ladder. Lane 1 PCR product of *Leishmania infantum* strain (MHOM/TN/80/IP11) after digestion with *APOI* enzyme , Lanes 2 ITS1 gene before digestion with *APOI* enzyme.

TCCGCCCGAAAGTTCACCGATATTTCTTCAATAGAGGAAGCAAAAGTCGTAACAAGGTAGCTGT  
AGGTGAACCTGCAGCTGGATCATTTTCCGATGATACCATACACAAAAACAAAAACCGGAGGGT  
 TTGGGTGTGGCGTGATGTGTGTATGTGTGTGCGTGTA AAAAGCGCATGCGCATATATGCATGC  
 ATAGTAGTGCCCGGCTCTCTACGTTGGGAGGAGCGGAACTAAACATTTCCGTTTCTCTCTAAC  
 ACATAAACAAACACAACATAGCCCAGCGCCGTTGCGTGCTTTCTCTCTCTCAACTCTCTCTCT  
 TGTGGGGGGTGTGTGTGTGGGGGTTTGTGCGCGCGTGTGCCGAACAAGGCCAATCGATGCAC  
 GTGTGTGTATTGTATTGTTCTTTCTTAGAGAACGATATAAAAACCGCGTGCATGGATGACGGCTC  
AAATAACGTGTCGCGATGGATGACTTGG

**Fig. 4 :** Result of PCR product contain 18S ribosomal RNA gene, partial sequence; internal transcribed spacer 1, complete sequence; and 5.8S ribosomal RNA gene, partial sequence of visceral leishmaniasis isolate.

CCGATGATACCATACACAAAAACAAAACCGGAGGGTTTGGGTGTGGCGTGTATGTGTGTA  
TGTGTGTGCGTGTAAAAAGCGCATGCGCATATATGCATGCATAGTAGTGCCCGGCTCTCTA  
 CGTTGGGAGGAGCGGAACTAAACATTTCCGTTTCTCTCTAACACATAAAACAAACACAACA  
 TAGCCAGCGCCGTTGCGTGCTTTCTCTCTCTCAACTCTCTCTCTTGTGGGGGGTGTG  
 TGTGTGGGGGGTTTGTGCGCGCGTGTGCCGAACAAGGCCAATCGATGCACGTGTGTGTATT  
 GTATTGTTCTTTCTTAGAGAACGATATAAAAACCGCGTGCATGGATGACGGCTCAAA

**Fig.5:** The results of sequencing of ITS1 for Iraqi isolates, showed 41% similarity with Iranian isolates that this similarity is shown in the beginning and the end of ITS1 gene (underline regions).

Query	13	TACACAAAAACAAAACCGGAGGGTTTGGGgtgtggcgtgtatgtgtgtatgtgtgtgcgt	72
Crithidia	1	TACACAAAAAC-AAAACCGGAGGGTTTGGGTGTGGCGTGTATGTGTATGTGTGTACGT	59
Query	73	gtAAAAAGCGCATGCGCATATATGCATGCATAGTAGTGCCCGGCTCTCTACGTTGGGAGG	132
Crithidia	60	GTAAAAAGCGCATGCGCATATATACATATATAGTAGTGCCAGCTCTCTACGTTGGGAGG	119
Query	133	AGCGGAACTAAACATTTCC-GTTTCTCTCTAACACATAAAACAAACACATAGCCCAG	191
Crithidia	120	AGCGGAACTAAACATTTCCGTTTCTCTCTAACACATAAAACAAACACATAGCCCAG	179
Query	192	CGCCGTTGCGTGCTTtctctctctctcaactctctctcttgggggggtgtgtgtgtg	251
Crithidia	180	CGCCGTTGCGTGCTTTCTCTCTCTCAACTCTCTCTTTTGTGGGGGGTGTGTGTGTG	239
Query	252	gggtttgtgcgcgcgtgtgCCGGAACAAGGCCAATCGATGCACGTGTGTATTGTATTG	311
Crithidia	240	GGGTTTGTGCGCGCGTGTGCCGAACAAGGCCAATCGATGCACGTGTGTATTGTATTG	299
Query	312	TTCTTTCTTAGAGAACGATATAAAAACCGCGTGCATGGATGACGGCTCAAA	362
Crithidia	300	TTCTTTCTTAGAGAACGATATAAAAACCGCGTGCATGGATGACGGCTCAAA	350;

**Fig.6** : Alignment of the Iraqi isolate and *Crithidia luciliae* with 96% similarity.

## الملخص العربي

توصيف جين مفساح الاستنساخ الداخلي 1 (ITS1) للشمانيا الطفلية والمعزولة من مرضى عراقيين مصابين بالشمانيا الحشوية وذلك باستخدام تقنيات تفاعل انظيم البلمرة التسلسلي (PCR) , تباين اطوال قطع التقييد (RFLP) والتتابع (Sequencing)

ندى البشير

فاطمة غفاري فار

الهام عبدالواحد مجيد

على حد علمنا لا تتوفر معلومات حول التوصيف الجزيئي للعزلة العراقية للشمانيا الحشوية لذا هدفت هذه الدراسة توصيف العزلة العراقية للشمانيا الطفلية (*Leishmania infantum*) باستخدام تقنيات تفاعل انظيم البلمرة التسلسلي (PCR) , تباين اطوال قطع التقييد (RFLP) والتتابع.

تم جمع ثلاثة عزلات من نخاع العظم لمرضى مصابين بمرض الكلا ازار ويقطنون مناطق مختلفة من بغداد. بعد التعرف على الطفيلي تحت المجهر تم انماءها على الوسط الزرعي المحور. استخلص الدنا (DNA) لتضخيم جين مفساح الاستنساخ الداخلي -1 (ITS 1) بواسطة تفاعل انظيم البلمرة التسلسلي (PCR). التعرف على العينات تمت دراسته بتقنية تباين اطوال قطع التقييد (RFLP) (هضم باستخدام انظيم الاقترع Apo 1) ثم اجراء التتابع لناتج البلمرة.

اظهرت نتائج تفاعل انظيم البلمرة التسلسلي للعينات ان هناك حزمة تحت حوالي 500 زوج القواعد. اما نتائج تفاعل انظيم البلمرة التسلسلي مع تباين اطوال قطع التقييد فلم تظهر اي اقترع باستخدام الانظيم الهاضم كما كانت نتائج التتابع مختلفة عن الجين المفصول من الشمانيا الطفلية في بنك الجينات.

يستنتج من ذلك ان تتابع هذا الجين في الشمانيا الطفلية المعزولة من المرضى العراقيين تختلف تماما عن العزلة القياسية (IPII / 80 / MHOM/ TN) ولكن ظهر بعض التشابه مع العزلة الايرانية للشمانيا الطفلية (MCAN/ IR/ 97/ LON) بنسبة 40% بينما كانت نسبة التشابه مع احد انواع الكرايندية (*Crithidia luciliae*) هي 96%.