



EVALUATION OF PERFORMANCE OF DAMASCUS GOAT AND ITS CROSSES WITH EGYPTIAN BALADI AND BARKI

[4]

Khallouf, N.¹; Manal Elsayed² and Mona, A. Osman³

1-Department of Animal Production, Faculty of Agriculture, Aleppo University, Aleppo, Syria.

2-Department of Animal Production, Faculty of Agriculture, Ain Shams University, Shoubra El-Kheima, Cairo, Egypt.

3-Animal Production Research Institute, Ministry of Agriculture and Land Reclamation, Dokki, Giza Egypt.

Keywords: Egyptian Baladi goats, Barki, Damascus, Crossbreeding, Heritability, Conception rate, Litter size

ABSTRACT

A total of 3108 records on conception rate (CR) and 2123 on litter size (LS) were used in this study. CR data represented 1600 Damascus (D) does, 201 Barki (BR) does, 229 Baladi (BL) does and 1070 crossbred does between D and each of BR and BL, while LS represented 1070 D does, 128 BR does, 145 BL does and 775 crossbred does. These records were used to study the effect of crossing and other fixed effects and to estimate heritability of these two traits in the Damascus goats. Statistical models included varying combination of the fixed effects of station, genotypes within station, year of mating, month of mating, and parity and the random effect of animal nested within station, genotype, year of mating and month of mating. Variance and covariance components for Damascus data were estimated through the Gibbs Sampling technique using a model that included the fixed effects of station, year and month of mating and parity, and the random effects of doe additive genetics and permanent environmental.

CR mean in different analyses ranged from 0.39 to 0.63. Damascus, with a range of CR of 0.48 to 0.63, surpassed both local breeds (ranging from 0.33 to 0.58) and was close to the crossbreds that ranged between 0.34 and 0.68. LS mean ranged from 1.26 to 1.50 in different analyses.

Baladi scored the highest in LS (1.49). Damascus ranged from 1.15 to 1.24 in different locations. LS ranged from 1.37 to 1.67 for crossbreds. For both traits there was no clear evidence of the merit of Damascus goats or their crosses over the Barki and Baladi goats. More data taken under sound herd management are needed to clarify the utility of the Damascus and its crosses.

Heritability estimates \pm SE for CR and LT in Damascus goats were 0.03 ± 0.003 and 0.04 ± 0.003 , respectively.

INTRODUCTION

Reproductive traits, such as fertility, prolificacy, fecundity and lamb survival have been recognized as major factors influencing profitability as reported by **Rao (1997)**. **Dickerson (1970)** stated that the cost of animal products depends upon the efficiency of three basic functions, female production, reproduction, and growth of the young. The female production conceptually contributes to all the outputs directly and indirectly from those breeding females that are kept for reproducing next generation. Improving female reproduction performance is an important target for increasing profitability of sheep (**Abdulkhalik et al 1989**) especially in lamb production. Litter size and number of lambs born are considered as two important traits in sheep and goats. Such two traits usually have low heritability and could be affected by many environmental factors such as year and month of mating, season and parity. Such environmental influences can mask genetic differences

(Received December 10, 2007)

(Accepted December 10, 2007)

when comparing between animals as reported by **Kennedy (1992)** and they must be controlled and/or corrected to elucidate genetic differences between animals (**Khallouf et al 2004**).

Dickerson (1970) stated that there is much greater potential for increasing both biological and economic efficiency of lamb production through genetic improvement in reproductive rate than through improvement in growth rate or body composition. The same can be said for goats. **Rao (1997)** reported that improving fertility could be quite important to successful accelerated breeding systems.

Improving the performance of local breeds of goats through crossing with Damascus goat has been tried in Egypt. Preliminary study by **Abdulla (1990)** indicated that crossing Damascus bucks to Barki does to improve their kid production seemed to be better than using the local Zairaibi bucks.

This work aimed at studying environmental factors affecting conception and litter size at birth in the pure breeds of Baladi, Barki and Damascus goats and to evaluate crosses between the latter and former two in three different locations in Egypt.

MATERIALS AND METHODS

Data

Data in the present study were collected from three experimental stations, Borgelarab, Sakha and Elgemmiza, of the Sheep and Goat Research Department, Animal Production Research Institute (APRI), Egyptian Ministry of Agriculture and Land Reclamation during the period 1982-2006. Damascus goats were imported from Cyprus and raised at Borgelarab station and then some were moved to Sakha station. Damascus bucks were mated to the local Barki does at Borgelarab and to Baladi does at Sakha in an effort to improve the performance of these local breeds. Also a Damascus herd was raised at Elgemmiza station as a pure breed. A total of 3108 records on conception rate (CR) and 2123 on litter size (LS) were used in this study. CR data represented 1600 Damascus (D) does, 201 Barki (BR) does, 229 Baladi (BL) does and 307 crossbred does between D and each of BR and BL, while LS represented 1070 D does, 129 BR does, 144 BL does and 780 crossbred does.

Management

Natural mating was practiced once a year mainly during autumn and winter months for all does but sometime empty does were exposed to bucks out of this period in an effort to maximize kid production. Does were divided into groups of 25-30 each joined with a fertile buck. Kids were weaned at three months of age. Animals were fed on Egyptian clover (*Trifolium alexandrinum*) from December to May. From June to November they were fed on crop stubbles and green fodder if available, besides a concentrate mixture, clover hay and rice straw.

Statistical analysis

Data on CR and LS of does were analyzed using Proc Mixed of **SAS (1996)** with repeated measures. Various models were followed for four alternative sets of data: all Damascus and crosses data, Damascus data, Damascus and crosses data in Sakha station and Damascus and crosses data in Borgelarab station). Gibbs Sampling (GS) techniques were applied to estimate the variance components and genetic parameters of the studied categorical traits for Damascus data. GS Program referring to Monte Carlo Markov Chain Methods is based on Bayesian methods for the estimation of variance covariance components (**Van Tassell and Van Vleck, 1996**).

The following model was applied for all Damascus and crosses data to obtain estimates for the investigated traits:

$$y_{ijklmno} = \mu + s_i + b_{j(i)} + r_k + m_l + a_{m(ijkl)} + p_n + e_{ijklmno}$$

Where

- y_{ijklmn} = record of conception rate (CR) measured as 1 if the doe gave birth and 0 otherwise,
- μ = the overall mean,
- s_i = the fixed effect of station, $i = 1, 2, 3$ for Borgelarab, Elgemmiza and Sakha station, respectively,
- $b_{j(i)}$ = the fixed effect of genotypes within station, $\{j(i) = D, \frac{1}{2}D.\frac{1}{2}BK, \frac{1}{4}D.\frac{3}{4}BK, \frac{3}{4}D.\frac{1}{4}BK, \frac{7}{8}D. \frac{1}{8}BK, \text{ for Borgelarab}; D, \frac{3}{4}D.\frac{1}{4}BL, \frac{1}{4} D.\frac{3}{4}BL, \frac{1}{2}D.\frac{1}{2}BL, \frac{3}{8}D. \frac{5}{8}BL, \frac{7}{8}D. \frac{1}{8}BL, \frac{5}{8}D. \frac{3}{8}BL \text{ for Sakha}; \text{ and } D \text{ for Elgemeza}\}$,
- r_k = the fixed effect of year of mating,
- k = 1982,.....,2006,
- m_l = the fixed effect of month of mating, $l =$

1,2,.....,12,
 $a_{m(ijkl)}$ = the random effect of animal m, nested within station i, genotype j, year of mating k and month of mating l,
 p_n = the fixed effect of parity, n =1,2,.....,10,
 $e_{ijklmno}$ = random error.

The following model was applied for Damascus data:

$$y_{ijklmno} = \mu + s_i + r_k + m_l + a_{m(ijkl)} + p_n + e_{ijklmno}$$

The definition of terms in the model is as those in the previous model with the absence of the effect of genotypes and some of subclasses for different fixed effects. Mean separations was done using **Duncan (1955)**, to evaluate the differences between locations, parities and month of mating.

The following model was applied for Damascus and cross data in Borgelarab station once and Sakha once more:

$$y_{ijklmn} = \mu + b_j + r_k + m_l + a_{m(ikl)} + p_n + e_{ijklmn}$$

The definition of terms is the same as those above.

The following threshold multiple trait model was fitted to estimate variance-covariance components and genetic and phenotypic parameters for CR using BS:

$$Y = X\beta + Zu + e,$$

where,

Y is N.1 matrix of observations of CR,

X is the incidence matrix for fixed effects,

β is the vector of an overall mean and fixed effects of station, year of mating, month of mating and parity,

Z is the incidence matrix for random effects,

U is the vector of random effects of does additive genetic effect and permanent environmental effect for CR of doe and

e is a vector of random errors normally and independently distributed with zero mean and variance $\sigma_e^2 I$.

The variance-covariance matrix was as follows:

$$Var \begin{bmatrix} u \\ e \end{bmatrix} = \begin{bmatrix} G & 0 \\ 0 & R \end{bmatrix}$$

where,

G is the additive genetic and permanent environmental variance-covariance matrix,

R is the residual variance-covariance matrix.

The same pervious models were used to analyze litter size (LS) after deleting records with zero conception.

LS and CR were transformed by ARCSIN transformation before the analysis of variance but means and SE were decoded to the original scale shown in different tables.

RESULTS AND DISCUSSION

Least squares means \pm standard errors of the studied traits in different levels of fixed effects except year of mating for all Damascus and crosses data are given in **Table (1)**.

Tables (1) and (2) show that no significant difference was observed for CR between different stations. The effects of genotype, parity, month and year of mating on CR were significant (**Table 1**). Damascus showed higher CR than Barki and Baladi goats. In contrast to these results **Abdelsalam et al (2000)** showed that conception rate was the highest in Barki does followed by Damascus and then by Zaraibi. In the present study $\frac{7}{8}D.\frac{1}{8}BK$ and $\frac{3}{4}D.\frac{1}{4}BK$ does had much higher CR (0.62 for both) than all purebreds while, $\frac{1}{2}D.\frac{1}{2}BK$ were equal (0.47) to Barki. CR tended to increase from the first to the third parity (**Tables 1 and 2**). Guney et al (2005) reported CR in Damascus in North Cyprus as 70.2% and 80.5% and litter size as 1.62 and 1.56. The highest estimates of CR were recorded in the matings of January, August and September. Damascus crossbreds with Baladi had higher CR than that Damascus crossbreds with Barki (**Tables 3 and 4**).

Least squares means for CR and LS through years 1983 – 2006 for all Damascus and crosses data are presented in **Figure (1)**. LS shows no specific year trend but CR showed a slump during the late eighties and early nineties possibly due to poor management in that period.

Table 1. Least squares means, standard errors (\pm SE) and probability of type I error (P) across the factor for CR and LS Damascus and all crossbred data.

Factor	No	CR		No	LS	
		Mean \pm SE	P		Mean \pm SE	P
Station:			0.16			0.00
Borgelarab	1387	0.54 \pm 0.04		984	1.36 \pm 0.04 ^a	
Elgemmiza	427	0.50 \pm 0.04		305	1.53 \pm 0.04 ^b	
Sakha	1294	0.43 \pm 0.11		834	1.54 \pm 0.11 ^c	
Genotype(Station):			0.00			0.00
Barki(BK)(Borgelarab)	201	0.47 \pm 0.05 ^b		128	1.41 \pm 0.09 ^b	
Damascus(D)(Borgelarab)	280	0.56 \pm 0.04 ^a		198	1.39 \pm 0.08 ^c	
7/8D*1/8BK(Borgelarab)	69	0.62 \pm 0.06 ^a		61	1.44 \pm 0.10 ^{bc}	
1/2D*1/2BK(Borgelarab)	211	0.47 \pm 0.05 ^b		135	1.37 \pm 0.09 ^c	
1/4D*3/4BK(Borgelarab)	435	0.52 \pm 0.04 ^{ab}		298	1.38 \pm 0.08 ^c	
3/4D*1/4BK(Borgelarab)	191	0.62 \pm 0.05 ^a		164	1.37 \pm 0.09 ^c	
Damascus(D)(Elgemmiza)	427	0.50 \pm 0.04 ^{ab}		305	1.49 \pm 0.00 ^b	
Baladi(BL)(Sakha)	229	0.36 \pm 0.05 ^c		145	1.76 \pm 0.09 ^a	
Damascus(D)(Sakha)	893	0.44 \pm 0.04 ^b		567	1.42 \pm 0.08 ^{bc}	
1/2D*1/2BL(Sakha)	164	0.49 \pm 0.05 ^b		117	1.67 \pm 0.09 ^{ab}	
Parity:			0.05			0.00
1	975	0.51 \pm 0.05 ^a		690	1.34 \pm 0.09 ^c	
2	718	0.52 \pm 0.05 ^a		490	1.38 \pm 0.09 ^{bc}	
3	509	0.53 \pm 0.05 ^a		358	1.46 \pm 0.09 ^{abc}	
4	369	0.44 \pm 0.05 ^b		236	1.45 \pm 0.09 ^{bc}	
5	228	0.47 \pm 0.06 ^a		139	1.42 \pm 0.10 ^{bc}	
6	135	0.48 \pm 0.06 ^a		94	1.56 \pm 0.10 ^{ab}	
7	81	0.45 \pm 0.07 ^b		51	1.34 \pm 0.11 ^c	
8	47	0.58 \pm 0.08 ^a		34	1.52 \pm 0.13 ^{abc}	
9	23	0.44 \pm 0.10 ^b		15	1.67 \pm 0.15 ^a	
≥ 10	23	0.50 \pm 0.10 ^a		16	1.62 \pm 0.16 ^{abc}	
Month of mating:			0.00			0.10
1	15	0.77 \pm 0.11 ^{ab}		11	1.33 \pm 0.15	
2	4	0.13 \pm 0.22 ^c		1	1.00 \pm 0.46	
3	13	0.00 \pm 0.0 ^c		2	1.74 \pm 0.33	
5	37	0.42 \pm 0.08 ^b		18	1.46 \pm 0.13	
7	114	0.59 \pm 0.06 ^{ab}		83	1.60 \pm 0.09	
8	41	0.75 \pm 0.09 ^a		39	1.71 \pm 0.11	
9	2653	0.70 \pm 0.04 ^{ab}		1803	1.49 \pm 0.06	
10	226	0.56 \pm 0.05 ^{ab}		166	1.49 \pm 0.10	

Means followed by different superscripts differ significantly at $P < 0.05$.

Table 2. Least squares means, standard error (\pm SE) and probability of type I error (P) across the factor for CR and LS for Damascus data.

Factor	No	CR		No	LS	
		Mean \pm SE	P		Mean \pm SE	P
Station			0.06			0.00
Borgelarab	1036	0.47 \pm 0.01 ^a		984	1.48 \pm 0.08 ^a	
Elgemmiza	426	0.46 \pm 0.01 ^a		305	1.62 \pm 0.08 ^a	
Sakha	1296	0.40 \pm 0.01 ^a		834	1.44 \pm 0.13 ^a	
Parity			0.06			0.00
1	975	0.44 \pm 0.05 ^{ab}		690	1.37 \pm 0.10 ^{ab}	
2	717	0.47 \pm 0.06 ^{ab}		490	1.41 \pm 0.10 ^{ab}	
3	510	0.49 \pm 0.06 ^{ab}		3581	1.50 \pm 0.10 ^{ab}	
4	369	0.41 \pm 0.06 ^{ab}		2361	1.49 \pm 0.10 ^{ab}	
5	228	0.47 \pm 0.06 ^{ab}		139	1.51 \pm 0.10 ^{ab}	
6	135	0.39 \pm 0.07 ^{ab}		94	1.61 \pm 0.12 ^{ab}	
7	81	0.35 \pm 0.08 ^b		51	1.37 \pm 0.13 ^b	
8	47	0.62 \pm 0.09 ^a		34	1.56 \pm 0.14 ^a	
9	23	0.34 \pm 0.12 ^b		15	1.69 \pm 0.18 ^b	
\geq 10	23	0.45 \pm 0.14 ^{ab}		16	1.62 \pm 0.22 ^{ab}	
Month of mating			0.00			0.03
1	15	0.46 \pm 0.16 ^{abc}		11	2.24 \pm 0.27 ^{abc}	
2	4	0.17 \pm 0.21 ^c		1	0.89 \pm 0.47 ^c	
3	13	0.28 \pm 0.24 ^{bc}		2	2.13 \pm 0.47 ^{bc}	
5	37	0.20 \pm 0.09 ^{bc}		18	1.11 \pm 0.24 ^{bc}	
7	114	0.34 \pm 0.13 ^{bc}		83	1.59 \pm 0.20 ^{bc}	
8	41	0.89 \pm 0.19 ^a		39	1.28 \pm 0.22 ^a	
9	2657	0.74 \pm 0.02 ^{ab}		1803	1.42 \pm 0.03 ^{ab}	
10	227	0.47 \pm 0.05 ^{abc}		166	1.45 \pm 0.08 ^{abc}	

Means followed by different superscripts differ significantly at P<05




Figure 1. Least squares means of CR and LS for all Damascus and crossbreds data

Least squares means \pm standard errors for CR and LS in different levels of fixed effects except year of mating for Damascus data are given in **Table (2)**.

Least squares means for CR and LS through years 1983 – 2006 for Damascus data are presented in **Figure (2)**. As the case for the whole set of data, Damascus data show a slump during the early nineties for CR but no specific trend is observed for LS.




Figure 2. Least squares means of CR and LS for Damascus data

Table (1) shows significant differences in LS of does attributed to station, genotype and parity. LS ranged from 1.38 for Barki to 1.68 for Baladi with majority of their crosses with Damascus scoring higher LS than the pure Damascus. The effect of breed on LS was in good agreement with the findings of **Amoah et al (1996)** and **Guney et al (2005)**. LS increased with the order of parity with a drop at the fifth and seventh parity. The influence of genotype and parity was non significant on LS in Borgelarab station, while it was highly significant in Sakha station (**Tables 3 and 4**). LS was not significantly affected by month of mating in both stations. LS tended to increase in ninth and tenth parities (**Tables 1, 2, 3 and 4**).

Least squares means \pm standard errors for CR and LS in different levels of fixed effects for Damascus and crossbreds in Borgelarab data are given in **Table (3)**.

Table 3. Least squares means, standard error (\pm SE) and probability of type I error (P) for CR and LS for Damascus and crosses data in Borgelarab station.

Factor	No	CR		No	LS	
		Mean \pm SE	P		Mean \pm SE	P
Genotype:			0.09		0.92	
Barki(BK)	201	0.33 \pm 0.08		128	1.26 \pm 0.11	
Damascus (D)	279	0.48 \pm 0.07		198	1.24 \pm 0.11	
$\frac{7}{8}$ D . $\frac{1}{8}$ BK	69	0.40 \pm 0.09		61	1.32 \pm 0.13	
$\frac{1}{2}$ D . $\frac{1}{2}$ BK	211	0.34 \pm 0.07		135	1.26 \pm 0.11	
$\frac{1}{4}$ D . $\frac{3}{4}$ BK	435	0.37 \pm 0.07		298	1.24 \pm 0.11	
$\frac{3}{4}$ D . $\frac{1}{4}$ BK	191	0.41 \pm 0.08		164	1.24 \pm 0.12	
Parity:			0.00		0.16	
1	445	0.47 \pm 0.07 ^{ab}		352	1.14 \pm 0.10	
2	316	0.45 \pm 0.07 ^{abc}		231	1.19 \pm 0.10	
3	227	0.46 \pm 0.07 ^{abc}		127	1.22 \pm 0.11	
4	187	0.41 \pm 0.07 ^{abc}		119	1.22 \pm 0.11	
5	95	0.36 \pm 0.08 ^{bc}		53	1.15 \pm 0.12	
6	49	0.35 \pm 0.09 ^{bc}		29	1.35 \pm 0.13	
7	28	0.32 \pm 0.10 ^{bc}		16	1.23 \pm 0.15	
8	15	0.25 \pm 0.12 ^c		8	1.22 \pm 0.19	
9	8	0.41 \pm 0.15 ^{abc}		6	1.42 \pm 0.21	
≥ 10	16	0.41 \pm 0.12 ^a		13	1.45 \pm 0.17	
Month of mating:			0.00		0.54	
1	1	0.07 \pm 0.40 ^c		na.	na.	
2	4	0.38 \pm 0.21 ^{bc}		1	0.97 \pm 0.46	
3	13	0.00 \pm 0.0 ^c		2	0.99 \pm 0.35	
5	14	0.70 \pm 0.12 ^a		14	1.40 \pm 0.14	
7	114	0.52 \pm 0.08 ^{ab}		83	1.51 \pm 0.09	
8	41	0.65 \pm 0.16 ^a		39	1.31 \pm 0.19	
9	1102	0.70 \pm 0.03 ^{ab}		753	1.35 \pm 0.05	
10	47	0.16 \pm 0.16 ^a		92	1.27 \pm 0.43	

Least squares means for CR and LS through years 1983 – 2006 for Borgelarab station data presented in **Figure (3)** show no specific trend.

Least squares means \pm standard errors for Damascus and crossbred in Sakha data are given in **Table (4)**.

Least squares means for CR and LS through years 1983 – 2006 for Sakha station data presented in **Figure (4)** show no specific trend.




Figure 3. Least squares means of CR and LS for Borgelarab station




Figure 4. Least squares means of CR and LS for Sakha station.

Table 4. Least squares means, standard error (\pm SE) and probability of type I error (P) across the factor for CR and LS Damascus and all crossbred in Sakha station.

Factor	No	CR		No	LS	
		Mean \pm SE	P		Mean \pm SE	P
Genotype			0.20			0.00
Baladi(BL) (Sakha)	230	0.58 \pm 0.01		144	1.49 \pm 0.18 ^{ab}	
Damascus (D) (Sakha)	893	0.63 \pm 0.01		567	1.15 \pm 0.17 ^b	
½D . ½BL (Sakha)	164	0.68 \pm 0.01		117	1.41 \pm 0.18 ^{ab}	
Parity			0.29			0.00
1	415	0.62 \pm 0.11		258	1.22 \pm 0.01 ^{ac}	
2	303	0.68 \pm 0.11		186	1.16 \pm 0.01 ^c	
3	217	0.72 \pm 0.11		152	1.33 \pm 0.01 ^{abc}	
4	142	0.63 \pm 0.12		90	1.38 \pm 0.01 ^{abc}	
5	101	0.70 \pm 0.12		64	1.35 \pm 0.01 ^{abc}	
6	64	0.65 \pm 0.12		47	1.45 \pm 0.01 ^{ab}	
7	42	0.57 \pm 0.13		26	1.19 \pm 0.01 ^{bc}	
8	22	0.80 \pm 0.14		19	1.59 \pm 0.01 ^a	
9	10	0.72 \pm 0.17		7	1.47 \pm 0.01 ^{ab}	
≥ 10	1	0.19 \pm 0.43			na.	
Month of mating			0.00			0.32
1	11	1.07 \pm 0.17 ^a		11	1.43 \pm 0.18	
2	na	na		na	na	
3	na	na		na	na	
5	17	0.15 \pm 0.15 ^c		1	1.03 \pm 0.52	
7	na	na		na	na	
8	na	na		na	na	
9	1289	0.68 \pm 0.11 ^b		837	1.60 \pm 0.11	
10	na	na		na	Na	

.na= not available.

Means followed by different super scripts differ significantly at $P < 0.05$.

As an approximate indication of heterotic effects, when the means weighed by their respective proportion of genetics from the two breeds was subtracted from actual mean of crossbreds, for CR, differences were all positive but that for 1/2D*1/2BK when all data considered with station included in the model and all negative but that for 1/4D*3/4BK in the analysis of Borgelarab data. As for LS, all values were negative except that for 7/8D * 1/8BK and 1/2D*1/2BL in the analysis

of the full data set and that of 7/8D*1/8BK at Borgelarab and Damascus halfbreds at Sakha and Borgelarab.

Heritability estimates for LS and CR in Damascus in the present data are quite low, being 0.04 ± 0.003 and 0.03 ± 0.003 , respectively. These results are in disagreement with the higher estimates reported by **Bagnicka *et al* (2006)** and **Odubote (1989)** of 0.35 for litter size, while in agreement with **Constantinou (1988)**.

CONCLUSION

The rather low mean conception rate in four runs ranging from 0.39 to 0.63 reflects serious defects in the management of these herds which could render breed/cross comparisons valueless. However, litter size was generally what is expected. In conception rate, Damascus surpassed both local breeds and was close to the crossbreds but generally the opposite was true for litter size. In view of the varying results of the merit of the crossbreds in both traits and the possible presence of genotype by station interaction, no conclusion could be reached regarding the use of Damascus in crossing Baladi and Barki goats.

ACKNOWLEDGEMENT

Authors thank Dr. Salah Galal, Professor of Animal Breeding and Genetics, Ain Shams University for assisting in modeling and in reviewing the manuscript.

REFERENCES

- Abdelsalam, M.M.; M. Eissa; G. Maharem and A.I. Heider (2000). Improving the productivity of the Barki goat by crossbreeding with Damascus or Zaraibi breeds. *Alex. J. Agric. Res.* **45**:33-42.
- Abdulkhalik, A.M.; W.W.R. Harvey and C.F. Parker (1989). Genetic parameters for ewe productivity traits in the Columbia, Suffolk and Targhee breeds. *J. Anim. Sci.* **67**: 3250.
- Abdulla, M.E. (1990). **Study on the Productive and Reproductive Performance of Desert Barki Goats and their Crosses with Damascus and Zaraibi Goats.** pp. 71-75. M.Sc. Thesis. The Institute of Animal Production Research (APRI)
- Amoah, E.A.; S. Gelaye; P. Guthrie and C.E. Rexroad, Jr. (1996). Breeding season and aspects of reproduction of female goats. *J. Anim. Sci.* **74**:723-728.
- Bagnicka, Emilia; Ewa Wallin; Marek Łukaszewicz and Tormod Ådnøy (2006). Heritability for reproduction traits in Polish and Norwegian populations of dairy goat. *Small Ruminant Research*, **68**:256-262.
- Constantinou, A. (1988). Genetic and environmental relationships of body weight, milk yield and litter size in Damascus goats. *Small Ruminant Research* **2**:163-174.
- Dickerson, G.E. (1970). Animal size and efficiency. *Anim. Prod.*, **27**:367.
- Duncan, D.B. (1955). Multiple range and multiple F test. *Biometrics*, **11**:1-24.
- Güney, O.; O. Torun and O. Özüyanık (2005). Milk production, reproductive and growth performances of Damascus goats under northern Cyprus conditions. *Small Ruminant Research* **65**: 176-179.
- Kennedy, B.W. (1992). **Genetic of Milk and Type.** Doc. No. 27 pp. 1-5. RV. Ontario Agr. College Guelph, Canada. Collection at the National Agricultural Library. D.Van Vleck; U., Cornell Ithaca, NY.
- Khallouf, N.; Manal Elsayed; G.H. Ghadry and K. Fatal (2004). Environmental and genetic factors affecting milk production and lactation period in Damascus goats in Syria. **8th Conf. Agric. Dev. Res., Fac. Agric., Ain Shams Univ., Cairo, Egypt, 2004, Annals Agric. Sci., Sp. Issue, 1:237-246, 2004.**
- Odubote, I.K. (1989). Genetic parameters for litter size at birth and kidding interval in the West African Dwarf goats. *Small Ruminant Research* **2**:163-164.
- Rao, S. (1997). **Genetic Analysis of Sheep Discrete Reproductive Traits Using Simulation and Field Data.** Dissertation submitted to the Faculty of the Virginia Polytechnic Institute and State University, in partial fulfillment of requirements for the degree of Doctor of Philosophy in Animal Science.
- SAS (1996). **SAS User's Guide, Statistics.** Version 8 Edition SAS Institute INC. Cary NC.
- Van Tassell, C.P. and L.D. Van Vleck (1996). Multiple-trait Gibbs sampler for animal models: Flexible programs for Bayesian and likelihood-based (co) variance component inference. *J. Anim. Sci.* **74**: 2567-2586.