

## BROOMRAPE AND OTHER WEED CONTROL IN CARROT (*DAUCUS CAROTA* L.)

\*GHALWASH, A. M., I. E. SOLIMAN\* and AZZA E. KHAFFAGY\*

\* Weed Research Central Laboratory, ARC, Giza, Egypt.

(Manuscript received 13 January 2014)

---

### **Abstract**

Four field experiments were conducted at Sakha Agricultural Research Station, Kafer El-Sheikh Governorate in the two successive winter seasons 2011/12 and 2012/13, to study the efficacy of some weed control treatments on broomrape and annual weeds control, as well as, on carrot yield productivity and its quality. The first study, used the herbicides glyphosate twice, imazapic twice, as post-emergence herbicides and hand pulling twice to broomrape control as a parasite weed. The second study, used the herbicides pendimethalin, butralin, oxyfluorfen, metribuzin as pre-emergence herbicides and hand weeding twice for controlling annual weeds. The results indicated that herbicide: imazapic twice, gave the best broomrape control and the highest increase in carrot roots yield and its quality, followed by glyphosate, and hand pulling twice of broomrape spikes, respectively. The herbicides imazapic and glyphosate decreased number of broomrape spikes/ m<sup>2</sup>, weight of broomrape spikes/m<sup>2</sup>, spikes length and number of capsules/spike. The previous pre-herbicides exceeded in a great extent the unweeded treatment in controlling annual broad-leaved weeds and annual grassy weeds. The maximum reduction values of dry weight of broad-leaved, grassy and total annual weeds were obtained by using oxyfluorfen, pendimethalin, metribuzin, butralin and hand weeding twice in two seasons, respectively, compared with the control treatment. The herbicides oxyfluorfen, imazapic, pendimethalin, metribuzin, glyphosate and butralin gave the highest increases in yield and its components in both seasons, compared with the control treatment. All tested herbicides gave the highest significant increase in the quality characters in carrot roots, followed by hand pulling of broomrape or hand weeding of the annual weeds. Also, the highest net return was obtained by herbicides used and hand weeding twice. Thus, these herbicides treatments can replace hand weeding for the control of weeds and broomrape in carrot crop.

**Key words:** *Orobanche*, Carrot, Broomrape, Weed Control.

### **INTRODUCTION**

Carrot has a few herbicides that could be used as pre- or post-emergence in Egypt. The use of herbicides in carrot fields plays an important role in improving the growth of carrot plants and consequently increase the productivity of land unit

area and lowering the cost of production as compared to hand weeding. Carl *et. al.* (2000) reported that weed control options are very limited in carrots. Carrots don't compete well against weeds, and herbicides are important tools for producing high-yield and high-quality carrots. Boydston *et. al.* (2003) reported that carrot is a small seeded, initially slow growing crop that can suffer severe yield losses from weed competition. Dittmar and Stall (2012) indicated that weeds reduce carrot yields by reducing the size of carrot roots through direct competition for nutrients, space and water. Weeds also deform carrot roots, making them unmarketable. Furthermore, weeds which grow late in the season may also cause severe harvesting problems. Karaliauskaite *et. al.* (2008) stated that the efficiency of metribuzin herbicide was higher when it was sprayed single (0.5 l/ha) at carrot 1-2 leaves stage and the amount of weeds was reduced by 84%. On the other hand, Jacobsohn *et. al.* (1980) mentioned that (*Orobancha aegyptica* L.) is a phanerogamic holoparasite without chlorophyll that may cause severe damage to many broad-leaved vegetables and field crops. Carrot plants without control become stunted due to heavy parasitization with broomrape and they were completely destroyed by the end of the season. Aviv *et. al.* (2002) stated that parasitic (*Orobancha spp.*) are major constraints to vegetable crop production in the Mediterranean basin to Eastern Europe. Schaffer *et. al.* (1991) proved that (*Orobancha aegyptica* Pers) and (*O. crenata*, Forsk) decreased the total sugar content similarly in carrot roots. Sucrose was the primary sugar in non infected roots and its level was greatly reduced in infected roots. So, broomrape infection can reduce carrot roots quality even when the visual appearance of the roots are not affected. Jacobsohn and Kelman (1982) reported that broomrape (*Orobancha spp*) control and high yields of carrots were obtained by spray application of 1.0 or 1.5 kg glyphosate /ha in January with 1-3 additional sprays applied at equal intervals (15 days) up to application. Nandula *et. al.* (2002) mentioned that the changes in the composition of both free and bound amine acids in carrot are associated with broomrape parasitism.

Farag *et. al.* (1994) stated that weeds associated carrot plants caused a 57-67% reduction in carrot root yield. In weed control plots, carrot root yield, root/shoot ratio and total carotenoids were increased, while, total soluble solids and dry matter content were decreased. Some selective herbicides gave the hope of solving this problem. Further, Gesagared herbicide gave the highest total sugar content and hand weeding twice can be used for weed control and improvement of yield and quality in carrots crops.

So, the present investigation was conducted to study the effect of some weed control treatments on broomrape and annual weeds control, as well as to study their effects on growth, and quality characters of carrot roots.

## MATERIALS AND METHODS

Four field experiments were carried out at Sakha Agricultural Research Station, during the two 2011/12 and 2012/13 successive winter seasons, to study the effectiveness of some weed control treatments on controlling broomrape (*Orobancha crenata*, Forsk.) and some annual weeds in carrot (*Daucus carota* L.), as well as carrot yield and its components. Carrot cultivar, Chantenay Red cored, was sown on the 5<sup>th</sup> and 10<sup>th</sup> of October in the first and second seasons, respectively, and yield uprooted harvested at the 5<sup>th</sup> and 10<sup>th</sup> of February) 2012. The soils of the four experiments were clay textured.

Table 1. Soil mechanical analysis of the experimental sites.

oil depth (cm)	Particle size distribution %			Texture	Bulk density, (g/cm <sup>3</sup> )	eld capacity %
	Sand	Silt	Clay %			
0-15	15.6	19.35	64.97	Clay	1-1	44.80
15-30	20.4	14.3	65.30	Clay	1-21	41.45
30-45	17.09	17.0	65.01	clay	1-28	39.27

The experiments were carried out in a randomized complete black design, with four replicates. plot area was 10.5 m<sup>2</sup>. The present study divided into two parts as follows:-

### Part I-Effect of *Orobancha* control treatments on broomrape.

In these experiments, the plots were naturally infested with broomrape seeds. Other weeds were removed by hand weeding. Each experiment consisted of four *Orobancha* control treatments as follows:

- 1- Glyphosate [(N – phosphonomethyl ) glycine] , known commercially as Round up 48% WSC, was applied twice with equal rate at 36.0 g a.i./fed. after 50 days from sowing , followed with 21 days interval.
- 2- Imazapic [(±)-2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-5-methyl-3-pyridinecarboxylic acid], known commercially as Oroban 10 % EC, was applied twice and equal rate at 20 g a.i./fed. after 50 days from sowing , followed with 21 days interval.
- 3- Hand pulling (twice), beginning after 15 days from emergence broomrape spikes with 21 days intervals between them.
- 4- Untreated check.

**Part II- Effect of weed control treatments on annual weeds.**

Two experiments were carried out, each experiment consisted of six weed control treatments, as follows:-

1. Butralin [4-(1,1-dimethylethyl)-*N*-(1-methylpropyl)-2,6-dinitrobenzenamine], known commercially as Amex 48% EC was applied once with a rate of 96.0 g a.i./fed., surface application after sowing and before irrigation.
2. Oxyfluorfen [2-chloro-1-(3-ethoxy-4-nitrophenoxy)-4-(trifluoromethyl)benzene], known commercially as Goel 24% EC was applied once and equal rate at 120 g a.i./fed., after 21 days from sowing.
3. Metribuzin [4-amino-6-*tert*-butyl-4,5-dihydro-3-methylthio-1,2,4-triazin-5-one,4-amino-6-*tert*-butyl-3-methylthio-1,2,4-triazin-5(4*H*)-one], known commercially as Sencor 70% WP was applied twice and equal rate of 36.0 g a.i./fed., after 21 days from sowing, followed with 21 days interval.
4. Pendimethalin [*N*-(1-ethylpropyl)-3,4-dimethyl-2,6-dinitrobenzenamine], known commercially as Stomp 50% EC was applied once and equal rate of 850 g a.i./fed., surface application after sowing and before irrigation.
5. Hand weeding (twice) was carried out at 18 and 32 days from sowing.
6. untreated check.

Herbicides in both field experiments were sprayed by knapsack sprayer CP3 in water volume of 200 liters per faddan. The preceding summer crop in the two seasons was maize (*Zea mays* L.). Plot area was 10.5m<sup>2</sup> (3.5m length and 3.0m width). Each plot contained four rows. During the two seasons, calcium super phosphate (15.5%) at a rate of 100 kg/fed was added before planting and nitrate (33.5%N) at a rate of 100 kg/fed.) was added before planting. Other cultural practices for carrot production were applied as recommended. The collected data were as follows:

**In experiments of controlling broomrape.****1. broomrape characteristics.**

Prior to carrot harvesting, number of broomrape spike/m<sup>2</sup>, weight broomrape/m<sup>2</sup>, spike length (cm) and number of capsules/ spike were estimated.

**In experiments for controlling the annual weeds.****1. On susceptibility rating of weeds.**

The susceptibility of weeds to herbicides was measured after 28 days from application of the herbicides by the reduction percentage of the dry weight of each species compared to the un-weeded check according to Frans and Talbert (1977) as follows:

- Susceptible (S) = > 90 % reduction.
- Moderately susceptible (MS) = 80 - 89 % reduction.

- Moderately tolerant (MT) = 60 - 79 % reduction.
- Tolerant (T) = < 60% reduction.

## **2- On dry weight of annual weeds.**

Annual weeds were hand weeding after 45 and 75 days from sowing and classified into three categories (broad-leaved, grassy and total weeds). The dry weight of weeds was determined in g/m<sup>2</sup>, after drying in a forced draft oven at 70 C° for 48 hours.

### **- Carrot yield and its components.**

At harvest time, samples of ten carrot plants were collected randomly from the central rows of each plot to assess the following criteria: root length (cm), root diameter (cm), number of roots / m<sup>2</sup>. Carrot plants were harvested from each plot, carrot root yield per plot was weighed and estimated as ton per feddan (fed).

### **2. Carrot quality characters.**

At harvest, the following data were recorded from ten carrot plants from each plot:

- a) Total soluble solids % (T.S.S) in carrot roots was measured by a Carl Zeiss handrefractometer.
- b) Dry matter % in roots.
- c) Total sugar was determined in carrot roots on fresh weight basis, according to the methods outlined by Poschenok (1976).
- d) Total carotenoid content (mg/100gm fresh weight) was determined according the method described by Ranganna(1977).

### **3. Economic evaluation.**

Net return was calculated by expressing the cost and yield of the unit area in monetary. The retail price used in computing cash returns was L.E 5 (Egyptian pounds) for carrot/kg for both seasons. The costs were negated from the overall cash returns as the resulted cash was considered to be the net return.

### **Statistical analysis.**

The obtained data were subjected to proper statistical analysis of variance, according to **Snedecor and Cochran (1980)** and the least significant differences (LSD) at the 5 % level of probability were calculated.

## RESULTS AND DISCUSSION

### Part I- Effect of broomrape control treatments.

#### - On broomrape.

Data presented in Table (2) show that the nature broomrape infestation in untreated check treatment was 20.02 and 22.93 number of spikes/m<sup>2</sup> or 113.35 and 106.55g/m<sup>2</sup> in both seasons, respectively. Imazapic applied twice at 20 g a.i./fed (post-emergence) and hand pulling twice gave approximately equal reduction effect, where reduction in number of broomrape spikes/m<sup>2</sup>, dry weight of spikes (g/m<sup>2</sup>), spike length (cm) and number of capsules / spike was (83.1, 86.6, 80.9 and 80.5%) (80.1, 85.1, 75.5 and 79.2%) in the first season, and (81.6, 82.1, 80.3 and 81.1%) and (78.9, 81.7, 77.3 and 76.5%) in the second season, respectively, compared to untreated check. While, glyphosate applied twice at 36.0 g a.i./fed as post-emergence herbicide gave the following reduction percentage in the previous respective characteristics (75.7, 82.7, 61.4 and 70.7%), in first season and (75.0, 79.1, 70.6 and 71.1%), in the second season, respectively, compared to untreated check. The obtained results were in agreement with those findings of Hassanein and Kholosy (1997) and Ghalwash *et. al.* (2008) who reported that the action of imazapic and glyphosate on broomrape is attributable to its selective accumulation in the young parasite plant up to four times as high as that in faba bean host root, three days after spraying.

Table 2. Effect of broomrape control treatments on broomrape infestation in 2011/12 and 2012/13 winter seasons.

Broomrape control treatments	2011/12 season			
	Number of spikes/m <sup>2</sup>	Dry weight of spikes g/m <sup>2</sup>	Spike length (cm)	Number of capsules/ spike
Glyphosate (36.0 g a.i./fed)	4.86	19.62	15.73	6.93
Imazapic (20.0 g a.i./fed)	3.38	15.23	7.78	4.61
Hand pulling twice	3.98	16.93	9.98	4.92
Untreated check	20.02	113.35	40.79	23.68
LSD at 5%	1.15	3.43	2.08	1.28
2012/13 season				
Glyphosate (36.0 g a.i./fed)	5.72	22.32	14.53	7.76
Imazapic (20.0 g a.i./fed)	4.23	19.05	9.73	5.08
Hand pulling twice	4.83	19.47	11.24	6.29
Untreated check	22.93	106.55	49.42	26.82
LSD at 5%	1.13	2.26	1.79	1.39

#### - On carrot yield components and quality.

Data in Table (3) indicated that imazapic at 20 g a.i./fed and glyphosate at 36 g a.i./fed. gave the highest significant increase percentage in weight of carrot

roots/fed by (52.8 and 52.19 %) , respectively compared to untreated check in first season and (50.27 and 49.21%), respectively in second season. While, hand pulling twice gave an increases percentage by (31.43 and 37.64%), respectively in 2011/12 and 2012/13 seasons, respectively. Similar results were obtained in case of carrot components and its quality in both seasons. In the first season, imazapic at 20 g a.i. /fed and glyphosate at 36 g a.i. /fed gave the highest values of root length (16.7 and 15.6 cm), root diameter (3.7 and 3.1 cm), number of roots/m<sup>2</sup> (175.7 and 158.4), TSS by (9.9 and 8.9 %) total carotene (81.7 and 76.1 mg), and total sugar (10.2 and 9.5 g), respectively. The obtained results in the first season confirmed to a great extent those observed in second season with minor differences. In the second season, imazapic at 20 g a.i. and glyphosate at 36 g a.i. /fed. gave the highest values of root length (15.96 and 14.0 cm), root diameter (3.85 and 3.3 cm), number of roots/m<sup>2</sup> (152 and 148), TSS (8.7 and 9.4 %), total carotene (86.7 and 79.5 mg) and total sugar (10.0 and 9.2 g), respectively.

Table 3. Effect of bromrape control treatments on yield components and its quality at harvest in carrot during 2011/12 and 2012/13 winter seasons.

Bromrape control treatments	2011/12 season							
	Root length (cm)	Root diameter (cm)	Number of root/m <sup>2</sup>	Weight of root (ton/fed)	TSS %	Dry matter %	Total carotenoides (M/g)	Total sugar g/(F.w)
Glyphosate (36.0 g, a.i./fed)	15.60	3.10	158.4	11.86	8.90	10.80	76.10	9.50
Imazapic (20.0 g, a.i./fed)	16.70	3.70	175.7	12.02	9.90	11.5	81.70	10.20
Hand pulling twice	11.20	2.01	135.7	8.27	7.90	7.80	59.70	8.60
Untreated check	8.30	1.51	95.7	5.67	5.50	5.70	42.30	5.90
LSD at 5%	1.29	0.851	31.5	2.31	1.97	2.50	12.40	1.61
	2012/13 season							
Glyphosate (36.0 g, a.i./fed)	14.00	3.30	148.2	10.83	9.40	11.80	79.50	9.20
Imazapic (20.0 g, a.i./fed)	15.46	3.85	152.7	11.06	8.70	12.20	86.70	10.00
Hand pulling twice	10.10	2.12	126.5	8.82	7.51	8.10	60.10	8.20
Untreated check	7.30	1.60	88.3	5.50	5.70	5.90	39.30	5.50
LSD at 5%	1.53	0.821	27.90	3.31	2.03	2.97	15.9	1.67

#### - Economic evaluation (net return):

Data in Table (4) showed that the net profit of carrot roots yield by weed control treatments in the first season could be arranged in descending order as follows: 10678 LE by imazapic at 20 g a.i./fed., 10554 LE by glyphosate at 36 g a.i. /fed, and 7979 LE by hand pulling twice, compared to untreated check 5850 LE. In the second season the same trend approximately for net profit was confirmed to those observed in first season. The net profit of carrot roots yield by the following treatments in a descending order was 9814 LE by imazapic at 20 g a.i./fed, 9627 LE by glyphosate at 36 g a.i. /fed, and 8231 LE by hand weeding twice, compared to untreated check 5373 LE. These results are in agreement with those reported by

(Sanjeev *et. al.* 2003, Dillared *et. al.* 2004 and Dixit *et. al.* 2005) who found that use of the herbicide pendimethalin and hand hoeing gave the highest yield and net return. This result may be due to the increase in leaf area /plant, dry weight of leaves % and gross head weight due to these treatments.

Table 4. Cost of weed control treatments, total root yield ton/fed and net return of carrot yield (L.E./fed) as affected by weed control treatments during 2011/12 and 2012/13 seasons.

Weed control treatments	2011/12			2012/13		
	Cost of W.C.T. (L.E./fed.)	T. root yield ton/fed.	N. ret. of yield (L.E./fed)*	Cost of W.C.T. (L.E./fed)	T. root yield ton/fed.	N. ret. of yield (L.E./fed)*
Glyphosate (36.0 g a.i./fed)	120	11.86	10554	120	10.83	9627
Imazapic (20.0 g a.i./fed)	140	12.02	10678	140	11.06	9814
Hand pulling twice	400	9.31	7979	400	9.59	8231
Untreated check	0	6.50	5850	0	5.97	5373
LSD : at 5% level		2.21			2.66	

\* N. ret = Net return  
L.E/person per day

Notice: The labor costs was assumed as 25

## Part II - Effect of weed control treatments on weeds and carrot:

### A - On annual weeds:

#### - Susceptibility of weeds:

Data in Table (5) illustrated that the susceptibility of weeds species was measured depending on the reduction of the dry weight of every species than untreated control after applied the soil-herbicides with 30 days interval. (*Medicago interterta*), *Sonchus oleraceas* and (*Chenopodium spp.*) as annual broad-leaved weeds were susceptible (> 90%) to butralin at 96 g a.i. /fed, oxyfluronfen at 120 g ai/fed, metribuzin at 40 g a.i./fed and pendimethalin at 850 g a.i. /fed. Meanwhile, (*Portulaca oleracea*), (*Beta vulgaris*), (*Ammi majus*) and (*Mililotus indicus*) as annual broad-leaved weeds and (*Phalaris spp.*) as annual grassy weed were moderately susceptible (80-89%) to oxyfluronfen at 120 g a.i./fed., and metribuzin at 36 g a.i./fed. Further, (*Portulaca oleracea*) and (*Phalaris spp.*), were susceptible (> 90%) to oxyfluronfen at 120 g a.i./fed, and pendimethalin at 850 g a.i./fed. (*Beta vulgaris*), (*Ammi majus*) and (*Mililotus indicus*) were fluctuated between susceptible and moderate susceptible to oxyfluronfen at 120 g a.i./fed, and pendimethalin at 850 g a.i./fed. These results obtained in the first season are confirmed with similar results in second season, that



mean the previous four soil-herbicides therefore exceeded in a great extent to unweeded treatment from view point of controlling (*Medicago interterta*), (*Sonchus oleraceas*) and (*Chenopodium spp.*), (*Portulaca oleracea*), (*Beta vulgaris*), (*Ammi majus*) and (*Mililotus indicus*) as annual broad-leaved weeds and (*Phalaris spp.*) as annual grassy weeds.

Table 5. Susceptibility of annual weed species to some herbicides treatments at 30 days from application during 2011/12 and 2012/13 winter seasons.

Weed control treatments	2011/12 season							Annual grassy weeds (g/m <sup>2</sup> )
	Annual broad-leaved weeds (g/m <sup>2</sup> )							
	<i>Medicago interterta</i>	<i>Sonchus oleraceus</i>	<i>Chenopodium spp</i>	<i>Portulaca oleracea</i>	<i>Beta vulgaris</i>	<i>Ammi majus</i>	<i>Mililotus indicus</i>	<i>Phalaris spp</i>
Butralin (96.0 g a.i./fed)	(18.6)	(11.4)	(17.3)	(32.4)	(37.4)	(35.2)	(42.6)	(34.6)
	94 S	96 S	90 S	87 MS	87 MS	86 MS	87 MS	89 MS
Oxyfluorfen (120.0 g a.i./fed)	(12.2)	(10.2)	(12.3)	(25.7)	(17.6)	(42.6)	(47.2)	(26.2)
	97 S	97 S	96 S	91 S	91 S	81 MS	86 MS	91 S
Metribuzin (36.0 g a.i./fed)	(12.4)	(19.3)	(26.6)	(28.5)	(33.2)	(41.8)	(32.4)	(30.70)
	97 S	95 S	91 S	89 MS	89 MS	81 MS	89 MS	88 MS
Pendimethalin (850.0 g a.i./fed)	(26.5)	(21.7)	(11.6)	(24.7)	(31.7)	(28.7)	(46.9)	(27.1)
	92 S	94 S	96 S	91 S	88 MS	91 S	86 MS	91 S
	2012/13 season							
Butralin (96.0 g a.i./fed)	(18.1)	(27.3)	(21.2)	(31.6)	(37.4)	(36.4)	(41.5)	(41.3)
	94 S	84 MS	95 S	89 MS	89 MS	88 MS	88 MS	81 MS
Oxyfluorfen (120.0 g a.i./fed)	(12.5)	(21.5)	(21.4)	(21.6)	(16.2)	(38.3)	(46.1)	(24.2)
	97 S	93 S	95 S	97 S	90 S	84 MS	86 MS	93 S
Metribuzin (36.0 g a.i./fed)	(16.3)	(25.3)	(18.3)	(29.4)	(42.6)	(34.0)	(31.0)	(28.4)
	95 S	89 MS	96 S	87 MS	83 MS	89 MS	88 MS	89 MS
Pendimethalin (850.0 g a.i./fed)	(23.6)	(23.4)	(17.9)	(26.2)	(43.3)	(31.9)	(43.2)	(25.7)
	91 S	92 S	96 S	93 S	83 MS	92 S	84 MS	92 S

S= > 90 %

MT= 60-79 %

MS= 80-89 %

T = < 60 %

#### - Dry of broad-leaved weeds:

Data in Table (6) indicated that dry weight of broad-leaved weeds (g/m<sup>2</sup>) was significantly influenced by weed control treatments. These findings were true at

the two evaluated samples and such effect was constant from one season to another. The tested herbicides (butralin, oxyfluorfen, metribuzin, pendimethalin) were effective in reducing the dry weight of broad-leaved and grassy weeds under natural heavy weed infestation, followed by hand weeding twice in the first and second seasons. These results are similar to those obtained by Sanjeev *et al* (2003) who found that the highest weed control efficiency were recorded from oxyfluorfen at 0.16 kg a.i./ha, pendimethalin at 0.75 kg a.i./ha and metribuzin at 0.70 kg a.i./ha, as compared to untreated check. The most effective treatments in reducing the dry weight of broad-leaved weeds up to 75 days after sowing were oxyfluorfen (90.0%), pendimethalin (88.3%), metribuzin (85.7%) and butralin (84.4%) in the first season, while in the second season, they were 88.0, 86.2, 85.1 and 84.2%, respectively. Hand weeding twice reduced the dry weight of broad-leaved weeds by 84.0 and 84.1%, relative to the control at both growing seasons, respectively.

**– Dry weight of grassy weeds:**

Data presented in Table (6) indicated that weed control treatments had a significant effect on dry weight of grassy weeds. These results were fairly true after 45 and 75 days from sowing and this effect was constant from season to another. It could be noticed that dry weight of grassy weeds reached to about ( 207.6 g/m<sup>2</sup>) in control plots at 75 days after sowing in the first seasons. All weed control treatments were significantly superior over the control plots. In this respect, the application of butralin, oxyfluorfen, metribuzin and pendimethalin gave about 86.8, 91.0, 89.1 and 90.5% reduction in dry weight of grassy weeds in the first season, respectively compared with the untreated check. The results of those treatments in the second season gave the same trend. These results were in harmony with those reported by Hegazy *et. al.* (1993), who revealed that using oxyfluorfen as post-emergence resulted in a good control of annual weeds.

**– Dry weight of total weeds:**

Data in Table (6) also revealed that dry weight of total weeds /m<sup>2</sup> was significantly affected by weed control treatments. However, it could be noticed that all herbicides were highly effective for reducing the dry weight of total weeds than that of untreated check. These findings were true after 45 and 75 days from sowing and this effect was constant from season to another. This means that applying was necessary to eliminate the weed plants that survived or escaped from the herbicides. Similar results were obtained by Hegazy *et. al.* (1993) and Sanjeev *et al* (2003).

Table 6. Effect of weed control treatments on dry weight of annual weeds ( $\text{g/m}^2$ ) after 45 and 75 days from sowing in 2011/12 and 2012/13 winter seasons.

Weed control treatments	2011/12 season					
	Dry weight of weeds in $\text{g/m}^2$ at 45 days from sowing			Dry weight of weeds in $\text{g/m}^2$ 75 days from sowing		
	Broad-leaved weeds ( $\text{g/m}^2$ )	Grassy weeds ( $\text{g/m}^2$ )	Total weeds ( $\text{g/m}^2$ )	Broad-leaved weeds ( $\text{g/m}^2$ )	Grassy weeds ( $\text{g/m}^2$ )	Total weeds ( $\text{g/m}^2$ )
Butralin (96.0 g a.i./fed)	80.8	17.8	98.6	306.9	28.4	335.3
Oxyfluorfen (120.0 g a.i./fed)	57.6	13.9	71.5	196.4	15.8	212.2
Metribuzin (36.0 g a.i./fed)	75.6	16.4	92.0	280.4	16.7	297.1
Pendimethalin (850.0 g a.i./fed)	65.3	14.8	80.1	229.3	16.4	245.7
Hand weeding twice	83.7	18.6	102.3	314.9	31.4	346.3
Untreated check	319.8	98.0	417.8	1964	207.6	2151
LSD at 5%	72.4	42.6	83.7	265.2	68.8	112.4
	2012/13 season					
Butralin (96.0 g a.i./fed)	85.4	19.4	104.8	251.7	29.6	281.3
Oxyfluorfen (120.0 g a.i./fed)	60.5	14.3	74.8	190.8	19.7	210.5
Metribuzin (36.0 g a.i./fed)	74.8	17.6	102.4	236.6	23.9	260.3
Pendimethalin (850.0 g a.i./fed)	80.3	16.8	97.1	219.6	20.8	230.4
Hand weeding twice	92.6	21.3	113.9	252.5	36.8	296.3
Untreated check	412.0	80.7	402.7	1592	219.4	1811
LSD at 5%	64.8	34.9	71.3	197.8	49.3	93.8

**B - On carrot yield and its components:**

Data in Table (7) indicated that weed control treatments had a significant effect on root length at harvest during the two growing seasons. Hand weeding twice application gave the highest values and significantly increased the root length of carrot than the control at the harvest by (31.5 and 26.8 %) in both seasons, respectively.

Butralin, oxyfluorfen, metribuzin and pendimethalin were significantly superior over hand weeding twice. Also, oxyfluorfen, pendimethalin and metribuzin superior in the length of carrot plants and recorded the root length (16.68, 16.46 and 15.70 cm), respectively as compared to hand weeding twice treatment (13.15 cm) at harvest in first season, this effect was constant at the second seasons. This reduction in root length under the control plots might be attributed to the negative effects of weeds on crop growth which may be occurred as a result of the competition between carrot and weed plant.

BROOMRAPE AND OTHER WEED CONTROL IN CARROT  
(*DAUCUS CAROTA* L.)

Table 7. Effect of weed control treatments on yield and its components at harvest in carrot during 2011/2012 and 2012/2013 winter seasons.

Weed control treatments	2011/12 season							
	Root length (cm)	Root diameter (cm)	Number of root/m <sup>2</sup>	Weight of root (ton/fed)	TSS %	Dry matter %	Total carotenoides (M/g)	Total sugar g/ (F.w)
Butralin (96.0 g. a.i./fed)	15.25	2.11	152.8	11.54	9.00	9.80	74.30	9.20
Oxyfluorfen (120. 0 g. a.i./fed)	16.68	2.80	175.7	12.54	9.90	11.50	84.60	10.4
Metribuzin (40.0 g a.i./fed)	15.7	2.60	168.3	11.86	9.20	10.8	80.20	9.70
Pendimethalin (850.0g a.i./fed)	16.46	2.70	171.2	12.06	9.50	11.20	82.00	10.20
Hand weeding twice	13.15	2.11	140.8	9.31	8.70	8.50	64.80	9.20
Untreated check	9.01	1.95	100.9	6.50	6.50	6.10	47.7	6.80
LSD at 5%	1.45	0.862	32.4	2.43	2.08	2.98	14.35	1.89
	2012/13 season							
Butralin (96.0 g. a.i./fed)	12.70	2.68	143.4	10.94	8.80	11.80	76.30	8.70
Oxyfluorfen (120. g. a.i./fed)	15.26	3.90	159.6	12.57	10.60	12.90	85.50	10.00
Metribuzin (40.0 g a.i./fed)	13.75	3.03	149.8	12.00	9.80	11.90	81.30	8.90
Pendimethalin (850.0g a.i./fed)	14.75	3.15	155.9	12.23	10.20	11.50	84.20	9.60
Hand weeding twice	10.95	2.18	132.6	9.59	8.00	8.40	63.20	8.90
Untreated check	8.02	1.86	97.4	5.97	6.20	6.70	44.30	6.10
LSD at 5%	1.68	0.928	29.30	3.86	2.04	3.11	16.43	1.98

Data revealed that root diameter was significantly affected by weed control treatments at harvest during the two growing seasons. Plots that were hand weeding twice produced the thickest stem as compared to herbicide treatments. It could be noticed that the results of this character had the same trend of that of root length under this study.

This reduction in the untreated check reflect the negative impacts of weeds on crop growth which may be occurred as a result of the competition between carrot and weed plants for the environmental resources (light, water and nutrients) which are necessary for plant growth. Chemical weed control treatments were superior in increasing root diameter of carrot that hand weeding twice and control treatments, during both growing seasons. These results were in complete agreement with these obtained by Farag *et. al.* (1994).

Data in Table (7) further indicated that the number of carrot root/m<sup>2</sup> was significantly affected by weed untreated check during the two growing seasons. Results illustrated that weed control treatments were correlated with an increase in number of carrot /m<sup>2</sup> in both seasons. This might be due to the eliminating treatment exposed to competition of annual weeds. The highest significant number of carrot root /m<sup>2</sup> was obtained from oxyfluofen, pendimethalin, metribuzin, and butralin. These treatments increased the number of carrot root/m<sup>2</sup> by (42.6, 41.1, 40.0 and 33.9 %) in the first season, respectively. The results of these treatments in the second season gave the same trend. Meanwhile, the hand weeding treatment gave the lowest increase in number of carrot root/m<sup>2</sup> (28.3 and 26.5 %) in the both seasons, respectively, compared to the untreated check.

Data revealed that weed control treatments had a significant effect on final carrot root yield (ton/fed) in both growing seasons. Dense weeds growing with carrot plants all over the growing seasons in control plots resulted in the lowest yield ( 6.50 and 5.97 ton/fed). The significant increases in carrot yield and its components accompanied with the quality were obtained by all herbicides used and hand weeding treatment. Further, the herbicides while were more efficient in controlling broomrape or annual weeds caused the highest increases in quantity and quality of carrot. This was true in the both seasons.

Data showed that all tested herbicides were significantly superior over the treatments in carrot root yield/ fed in both seasons. In this respect, the highest carrot root yield/ fed (12.54 and 12.57 ton/fed) was achieved from oxyfluorfen, followed by pendimethalin ( 12.06 and 12.23 ton/fed), respectively in the both seasons. This may be due to that applying the herbicides were necessary to eliminate the weed plants,

which survived or escaped from the herbicides and assure the importance of using the suitable herbicides due to the expected problem of weed flora.

Results presented in Table (7) showed the effect of weed control treatments had a significant effect on total soluble solids in carrot root and all exceeded significantly untreated check in both seasons. Oxyfluorfen, pendimethalin, metribuzin and butralin treatments were the potent treatments. These results are similar to those reported by Farag *et. al.* (1994).

All herbicidal treatments (oxyfluorfen, pendimethalin, metribuzin and butralin) were exceeded in dry matter in carrot content roots in the two seasons, compared to the untreated check. Meanwhile, the hand weeding treatment gave the lowest increase in dry matter in carrot roots (28.2 and 20.2 %) in both seasons, respectively, compared to the untreated check.

Data in Table (7) also, revealed that all herbicides treatments gave higher total carotene content in both seasons and were not significantly different in most cases. On the other hand, the untreated check was the lowest in total carotene content. The reduction in carotene content values, under hand weeding and untreated check treatments, reflected the negative impacts of annual weeds on carrot growth, which might occur as a result of competition between carrot and weeds. Also, the results showed that, using the tested herbicides was necessary to eliminate weeds and to avoid its negative impacts on carrot. Similar results had been reported by Farag *et. al.* (1994).

Regarding the effect of weed control treatments on total sugars in carrot roots, data indicated that all tested herbicides gave the highest total sugars in carrot roots. Hand weeding gave the lowest total sugars, compared to all tested herbicides. This result showed that a single hand weeding was insufficient to provide the desired weed control level and this was reflected on the limited increases in carrot growth. Such results, presented in Table (6) agreed with those obtained by Farag *et. al.* (1994). These effects might be attributed to the dominant weeds in the hand weeding treatment, and this assured the importance of using suitable herbicides.

#### **-Economic analysis.**

Data in Table (8) showed that the net profit of carrot roots yield by weed control treatments in the first season could be arranged in a descending order as follows: 11026 LE by oxyfluronfen at 120 g a.i./fed, 10554 LE by pendimethalin at 850 g a.i. /fed, 10466 LE by metribuzin at 40 g a.i./fed, (10116 LE) by butralin at 96 g a.i. /fed. and (8748 LE) by hand weeding twice, compared to untreated check (6570 LE). In second season the same trend approximately for net profit was confirmed to those observed in the first season. The net benefit of carrot roots yield by the following

treatments in a descending order was 11053 LE by oxyfluronfen at 120 g a.i./fed., (10807 LE) by pendimethalin at 850 g a.i. /fed, (10610 LE) by metribuzin at 40 g a.i./fed, (9198 LE) by hand weeding twice and (9576 LE) by butralin at 96 g a.i. /fed. compared to untreated check (6156 LE). These results are agreement with those reported by (Sanjeev *et. al.* 2003,) who found that use of the herbicide pendimethalin and hand hoeing gave the highest yield and net return.

Table 8. Cost of weed control treatments, total root yield (t/fed) and net return of carrot yield (L.E./fed.) as affected by weed control treatments during 2011/12 and 2012/13 seasons.

Weed control treatments	2011/12			2012/13		
	Cost of W.C.T. (L.E /fed)	T. root yield (ton/fed.)	N. ret. of yield (L.E /fed)*	Cost of W.C.T. (L.E /fed)	T. root yield (ton/fed.)	N. ret. of yield (L.E /fed)*
Butralin (96.0 g, a.i./fed)	270	11.54	10116	270	10.94	9576
Oxyfluorfen (120.0 g, a.i./fed)	260	12.54	11026	260	12.57	11053
Metribuzin (40.0 g, a.i./fed)	190	11.84	10466	190	12.00	10610
Pendimethalin (850.0 g, a.i./fed)	300	12.06	10554	300	12.23	10807
Hand weeding twice	450	10.22	8748	450	10.72	9198
Untreated check	0	7.30	6570	0	6.84	6156
LSD : at 5% level		2.43			3.86	

\* N. ret = Net return

Notice: The labor costs was assumed as 25 L.E/person per day

## REFERENCES

1. Aviv, D., Z. Amsellem and J. Gessel. 2002. Transformation of carrots with mutant acetolactate synthase for *Orobanche* (broomrape) control. *Pest Management Sci.*, 58 :12, 1187-1193.
2. Boydston, Rich, W. Martin, P. G. Prosser and S. Dollars. 2003. Carrot weed control – Research with caparol and notron *Proceedings of the Pacific northwest vegetable association, PASCO*, P. 81-88.

3. Carl, E., B. Brent, E. B. Outwell, E. J. Ogbuchickwe, E. Milt and J. Mcgiffen. 2000. Weed control in carrots the efficacy and economic value of linuron. Hort Sci. 35 (6):1089- 1091.
4. Dillard, H. R., R. R. Bellinder and D. A. Shah. 2004. Integrated management of weeds and diseases in a cabbage cropping system. Crop-protection. 23(2): 163-168.
5. Dittmar, P.J. and W. M. Stall. 2012. Weed control in carrot. This document is HS201, one of
6. a series of the Horticultural Sciences Department, Florida Cooperative Extension vice, Institute of Food and Agricultural Sciences, University of Florida.
7. Dixit, A. J. , J. R. Ramteke, S. T. Thorate and N. D. Jambhale. 2005. Efficacy of herbicides in controlling weeds in cabbage. Maharashtra J.Agric. Univ. 30 (3): 362-363.
8. Farag, I. A., H. A. Hussein and M. A. Farghali. 1994. Effect of chemical weed control on growth of weeds, yield and quality of carrot. Assiut J. of Agric. Sci., 25: 3, 1-12.
9. Frans, R.E. and R. Talbet. 1977. Design of field experiment and the measurement and analysis of plant response. Res. Methods in Weed Sci. Soc. Field. South. USA, Auburn, Alabama.
10. Ghalwash, A. M., I. E. Soliman and Azza E. Khaffagy. 2008. Performance of some faba bean (*Vicia faba*, L.) cultivars under numerous broomrape (*Orobanche crenata*, Forsk.) control treatments. J. Agric. Sci. Mansoura Univ., 33 (4): 2439-2448.
11. Hegazy, R. T., H. M. El-Sheakh and A. K. I. Kafoury. 1993. Effect of planting methods and some herbicides on the growth of onion seedlings and weeds in nursery. Egypt. J. Appl. Sci., 8 (9): 388 – 397.
12. Hassanein, E. E. and A. S. Kholosy. 1997. Demonstration plots of faba bean for broomrape control in fayoum Governorate. NVRP for wild oats control in wheat and other winter crops. 5th Annual Meeting, Cairo, 11-15 Sept: 109-111.
13. Jacobsohn, R. and Y. Kelman. 1982. Proper timing of glyphosate application for broomrape control in carrot and celery. Phytoparasitica, 10: (4), 268.
14. Jacobsohn, R. Geenberger, A. , Katan, J., Levi, M. and H. Alon. 1980. Control of Egyptian broomrape (*Orobanche aegyptica* ) and other weeds by means of solar heating of the soil by polyethylene mulching. Weed Sci. 28 (3): 312- 316.
15. Karaliauskaite, D., Starkute, R., Bundiniene O., J. Jankauskiene. 2008. Chemical weed control in carrot crop ISHA Acta Horticulturae 830: iv Balkan Symposium on vegetable and potatoes.



16. Nandula, V. K., J. G. Foster and C. L. Fay. 2002. Impact of Egyptian broomrape (*Orobanche aegyptica*, Pers) parasitism on amino acid composition of carrot (*Daucus carota*, L.) . *J. of Agric., and Food Chemistry*, 48 (9): 3930-3934.
17. Poschenok, N. 1976. *Biochemical analytical methods of plants*. Publishers, Naokova Damka Kiev, 276 PP. (In Russian).
18. Ranganna, S. 1977. *Manual of analysis of fruit and vegetable products*. Tata Mcgraw – Hill Publishong company Limited, New Delhi, PP. 634.
19. Sanjeev, A., K. S. Sandhu and S. Ahuja. 2003. Efficiency of weed control in cabbage onion cropping system. *Annual of Biology*. 19 (1) : 31- 34.
20. Schaffer, A.A., R. Jacobsohn, D. M. Joel, E. Eliassi and M. Fogelman. 1991. Effect of broomrape (*Orobanche* spp.) infection on sugar content of carrot roots. *Hort. Sci.*, 26 (7): 892- 903.
21. Snedecor, G. W. and W. G. Cachran. 1980. " *Statistical Methods* "6 th ed., Iowa State Univ. Press, Ames. Igrawowa, U.S.A, 325-330.
22. Topps, J. H. and R.L. Wain. 1957. Investigation of fungi toxicity of 3- and 5-alkyl salicylinlide and para- chloroaniline. *Ann. Appli. Biol.* 45 (3): 506-511.

## مكافحة الهالوك والحشائش الأخرى في الجزر

عادل مصطفى غلوش\* ، إبراهيم السيد سليمان\* ، وعزة السيد خفاجى\*

\*المعمل المركزى لبحوث الحشائش - مركز البحوث الزراعية - الجيزة - مصر

اجريت أربعة تجارب حقلية فى المزرعة البحثية بمحطة البحوث الزراعية بسخا محافظة كفر الشيخ فى موسمين متتاليين ٢٠١٢/٢٠١١ و ٢٠١٣/٢٠١٢م بهدف دراسة تأثير فاعلية بعض معاملات مكافحة الحشائش على حشيشة الهالوك و الحشائش الحولية وانعكاس ذلك على محصول الجزر وجودته. التجربة الأولى (مبيد جليفوسيت مرتين ومبيد ايمازبيك مرتين) لمكافحة الهالوك فى محصول الجزر بينما شملت التجربة الثانية استخدام مبيد بنديمتالين والبيوترالين والاكسيفلورفين والمتربيوزين بعد انبات نباتات الجزر) لمكافحة بعض الحشائش الحولية فى محصول الجزر بالإضافة الى النقاوة اليدوية مرتين . وقد أشارت النتائج المتحصل عليها التجربة الاولى أعطى مبيد ايمازبيك مرتين أفضل مكافحة للهالوك وانعكس ذلك على زيادة محصول الجزر وجودته تليه معاملة النقاوة اليدوية مرتين و مبيد جليفوسيت مرتين على الترتيب. وقد أكدت هذه النتائج تحت العدوى الطبيعية العالية للهالوك بالتربة أنه يمكن استخدام ايمازبيك ، جليفوسيت كلاهما مرتين وهذه المعاملات أدت الى نقص عدد شماريخ الهالوك / م٢ ووزن شماريخ الهالوك /م٢ وطول شمراخ الهالوك وأيضاً عدد الكبسولات / شمراخ هالوك خلال موسمى النمو . التجربة الثانية أعطت مبيدات الحشائش (بنديمتالين والبيوترالين والاكسيفلورفين قبل الانبات والمتربيوزين بعد الانبات كفاءة عالية ضد الحشائش العريضة و النجيلية . أدى استخدام مبيدات الحشائش الى نقص كبير فى الوزن الجاف للحشائش عريضة الأوراق وضيقة الأوراق والحشائش الكلية حيث ظهر ذلك من استخدام مبيدات الحشائش ( اكسيفلورفين ، بنديمتالين، متربيوزين و بيوترالين والنقاوة اليدوية مرتين) فى كلا الموسمين وذلك مقارنة بمعاملة الكنترول. أدى استخدام مبيدات اكسيفلورفين ، بنديمتالين، متربيوزين و بيوترالين الى زيادة محصول الجذر و مكوناته فى كلا الموسمين مقارنة بمعاملة الكنترول. جميع مبيدات الحشائش السابقة أعطت زيادة معنوية لصفات الجودة فى الجزر بينما معاملة النقاوة اليدوية مرتين أعطت أقل قيم لهذه الصفات خلال موسمى الزراعة. أيضاً أظهرت النتائج أن استخدام هذه المبيدات والنقاوة اليدوية مرتين أدت الى زيادة صافى الربح . ولذلك توصى هذه الدراسة بأنه يمكن استخدام المبيدات السابقة لتحل محل النقاوة اليدوية فى مكافحة الحشائش الكلية والهالوك فى محصول الجزر.