

CERTAIN ASPECTS ON REPRODUCTIVE POTENTIAL AND FEEDING CAPACITY OF THE SPIDER *CHIRACANTHIUM INCLUSUM* HENTZ (ARACHNIDA: MITURGIDAE)

ABU-ZAED, AMAL E., MERVAT A. KANDEL AND A. A. A. EL-SAYED

Plant Protection Research Institute, ARC, Dokki Giza

(Manuscript received 14 March 2010)

Abstract

Studies were carried out under laboratory conditions at 25 ± 1 °C and 65-70% R.H. to study the effect of two species of Lepidopterous insects, *Pectinophora gossypiella* and *Spodoptera littoralis* on biological aspects of the spider *Chiracanthium inclusum*. *P. gossypiella* was the best as it reduced in duration of different predator stages, whereas, *P. gossypiella* increased the fecundity, as well as increasing consumption prey. While, female deposited average 134.3 eggs/sac when fed on *P. gossypiella* and 97.6 eggs/sac on *S. littoralis*, respectively. On the contrary, female longevity lasted 114.0 days, while, male adult hood lasted 75.0 days on *P. gossypiella*, and 135.0 days and 98.3 days for both sexes on *S. littoralis*, respectively. The female and male predator consumed 348.3 and 190.6 prey of *P. gossypiella* and 411.66 and 217.33 prey of *S. littoralis*, respectively.

INTRODUCTION

Winter spider (*Chiracanthium inclusum* Hentz) is a nocturnal collected running spider that is often one of the most numerous spiders collected in crops such as, cotton and citrus Whitcomb *et al.* 1963 and Amalin *et al.* 2001. Nyffeler *et al.* 1990 and Miliczky and Calkins (2002) reported that it is known to prey the eggs of *Helicoverpa zea* and the eggs of other crop pests. On the other hand, Taylor and Pfannenstid (2008) demonstrated that a large proportion of the spider *C. inclusum* collected at night from cotton fields showed that it had fed on extra floral nectar. In addition, other species of *Chiracanthium* were reported in significant numbers in cotton fields in the USA (Whitcomb and Bell 1964). However, Ruiz (2006) reared *Chiracanthium inclusum* on eggs of *H. zea* and scale insects as a prey and observed it to be with cotton bollworm. In 2008 Pfannenstiel *et al.* studied the effect of nectar and egg of *H. zea* on feeding, immatures duration and adult longevity of the spider *C. inclusum*.

Therefore the aim this research was to examine the effect of two cotton pests *P. gossypiella* and *S. littoralis* on some biological developmental stages, feeding capacity and fecundity of the spider *C. inclusum* in order to better understand its role, as a natural enemy of insect pests in cotton fields.

MATERIALS AND METHODS

1. Prey used

The two insect species of prey used in this study were *Pectinophora gossypiella* (PBW) and *Spodoptera littoralis*. Eggs and larvae of pink bollworm, *P. gossypiella* were reared under controlled conditions ($25\pm 1^\circ\text{C}$ and 65-70 % R.H. with complete dark all day time) for several generations on a semi- artificial diet according to Rashad and Ammar (1985). On the other hand, larvae of *S. littoralis* were reared on the castor bean leaves.

2. The spider *C. inclusum*

Laboratory culture of *C. inclusum* began with adults and egg sacs collected at the end of August and mid September from cotton fields at Mansoura region Dakahalyia Governorate and kept under the laboratory conditions. The predator was reared for one generation on eggs + 1st instar larvae of PBW and 2nd instar larvae of *S. littoralis*.

Feeding capacity

To rearing the winter spider 90 individuals spiderling were reared from eggs to maturity on different prey. 90 newly emerged *C. inclusum* spiderling were placed individually in glass tubes (15 cm high X1.5cm wide) with sufficient numbers of the two spotted spider mite *T. urticae* were offered daily until the end of 1st instar spiderling. After that, these tubes were divided in two groups, each of 45 tubes. Sufficient numbers of eggs and newly hatched larvae of PBW were offered daily to first group, while, the second group was offered the 2nd instar larvae of *S. littoralis*. The numbers offered from each prey were increased as the predator's spiderling grew older. The numbers of consumed prey from each PBW and *S. littoralis* were recorded daily.

Also, the numbers of deposited egg sacs and the number of deposited eggs inside sacs were counted as well as pre oviposition, oviposition, post oviposition, longevity of females and male estimated. The total consumptions were assessed. The relation between duration of different stages and adult longevity, also, total of consumption for each prey were examined by analysis of variance (ANOVA) using Costat program.

RESULTS AND DISCUSSION

Developmental time of spiderling stage

Development of spiderling *Chiracanthium inclusum* passes through seven instars. Duration of these instars were affected by different foods. As shown in Table (1), the

total developmental periods were highly significant shorter for spiderlings reared on eggs + first instar larvae of *Pectinophora gossypiella* (127.93 days) while, being longer (146.39 days) when reared on *S. littoralis* for completion of development, respectively. Obtained results indicated that rearing on *P. gossypiella* decreased developmental time by 12.5% than rearing on *S. littoralis*.

Feeding Capacity

The results in Table (2) showed that the highly significant difference occurred in feeding capacity of *C. inclusum* between spiderlings instars 2nd – 7th when fed on mixed of (eggs + 1st instar larvae) of *P. gossypiella* and 2nd instar larvae of *S. littoralis*. Obtained data cleared that the mean consumption rate increased with the progressive of spiderling age. Also, data analysis in Table (2) revealed that the average of total consumption of the tested predator *C. inclusum* from (2nd –7th) instars fed on each of the two preys *P. gossypiella* & *S. littoralis* were 514.0 and 325.0 prey, respectively.

Under standard laboratory conditions, adult female and male of *C. inclusum* lived an average of 114 & 75 days, when reared on *P. gossypiella* while, the averaged of 135 and 98.3 when feeding on *S. littoralis*, respectively (table, 3).

Mansour *et al.* (1980) reared the spider *Chiracanthium mildi* on the prey *S. littoralis* larvae. Females required a mean of 231.0 days from hatching to mature, reached adult after 9-10 months, and lived for an average of 240 days as adult. Male required 185.0 days from hatching to reach maturity, becoming adult after 7-8 months, and lived for an average of 73.0 days as adult. When rearing *C. jsvium* on *Ceratitis capitata*, Sallam (1996) stated that eggs incubation period averaged 14.06 days and the total life span was 387.62 and 234.73 days for female and male, respectively.

Ovipositional period

Table (3) showed that high significant differences occurred between pre-oviposition, oviposition and post oviposition periods for *C. inclusum* when reared on (eggs+ larvae) of *P. gossypiella*, these periods averaged 19.1, 30.3 and 64.3 days, respectively, while, prolonged to 24.6, 38.3 and 72.0, respectively when adult female fed on 2nd instar larvae of *S. littoralis* at the same pattern.

Number of deposited eggs/female

Data in Table (3) revealed that the number of deposited eggs/sac was affected by prey. This average was higher for female fed on eggs+larvae of PBW. Also, data analyses showed a highly significant differences between the average number of deposited eggs, whereas, female deposited 134.3 and 97.6 eggs when it fed on PBW and *S. littoralis*, respectively. Also, high significant differences occurred between eggs/sac hatchability percentages when female fed on PBW and *S. littoralis*,

where by, hatching percentages were 93.96 and 82.16%, when female fed on the above mentioned prey, respectively.

Edwards (1958) recorded 112 eggs in a single egg/sac produced by *C. inclusum*, while, Peck and Whitcomb (1970) reported the occurrence a range of 17-86 eggs per sac. However they stated that *Chiracanthium* spiderlings appeared to feed on the infertile eggs in their egg mass. The 1st instar spiderling fed on larvae of *T. cinnabarinus* and 1-day old *S. littoralis* larvae. Amalin *et al.* (2001) reared *C. inclusum* on citrus leaf miner larvae, female produced 1-3 egg masses with an average of 57 eggs per egg mass. Oviposition took place 2-7 days after mating.

Incubation periods of eggs

At 25±1°C and 65-70% R.H. egg incubation averaged 13.23 days and 18.61 days when females reared on PBW and *S. littoralis*, respectively.

Data indicated that rearing females on *P. gossypiella* shortened adult longevity by 15.6% time and increased number of eggs per sac eggs by 27.4% than that females reared on *S. littoralis* Table (3).

Table 1. Development of immature stages of *Chiracanthium inclusum* when fed on different prey at 25±1°C and 65-70 % R.H.

prey	1 st instar	prey	2 nd instar	3 rd instar	4 th instar	5 th instar	6 th instar	7 th instar	Total immature stage
Tetraneus	9.30±0.3	PBW	15.67±0.1	11.33±0.13b	23.67±0.2b	19.33±0.12	26.33±0.31	22.3±0.61	127.93±2.9b
		<i>S. littoralis</i>	17.1±0.2	15.67±0.2a	27.67±0.5a	28.0±0.4	25.6±0.7	25.0±0.3	146.39±1.79a
		p	Ns	0.01*	0.009**	Ns	Ns	0.0048**	0.0005***
		LSD		2.61	2.28			1.49	4.84

Table 2. food consumption of the spider *S. littoralis C. inclusum* during its immature stages at 25±1°C and 65-70 % R.H.

Prey	1 st instar	prey	2 nd	3 rd	4 th	5 th	6 th	7 th	Total immature stage
Tetraneus	43.3±4.49	PBW	42.66±2.4	57.0±1.5a	75.6±a	79.7±0.5a	110.7±1.6a	148.6±1.15a	514.0±3.05a
		<i>S. littoralis</i>	38.3±2.6	40.0±2.3b	40.6±1.8b	56.0±3.1b	65.6±1.9b	74.3±2.3b	315.0±5.2b
		p	Ns	0.003**	0.0022**	0.0015**	0.000***	0.000***	0.000***
		LSD		7.68	13.94	8.53	8.43	6.73	25.51

Table 3. Effect of different prey on fecundity and longevity of the spider *Chiracanthium inclusum* at 25±1°C and 65-70 % R.H.

Preys	Ovipositional period			Total eggs	Hatching %	Incubation period	Adult longevity	
	Pre-oviposition	Oviposition	Post oviposition				♀♀	♂♂
PBW	19.1±0.5b	30.3±0.3b	64.3±0.6b	134.3±3.9a	93.96±0.5a	13.23±0.5b	113.7±1.7b	75.0±0.5b
<i>S. littoralis</i>	24.6±0.3a	38.3±0.30a	72.0±1.7a	97.6±2.2b	82.16±0.13b	18.61±0.1a	134.9±1.7a	98.3±0.4a
p	0.001**	0.0001***	0.014*	0.0008***	0.000***	0.002**	0.0003***	0.000***
LSD	1.85	1.30	5.15	11.18	1.64	2.26	5.15	1.01

REFERENCES

1. Amalin, D.M., J.E., Pena, J. Reiskind, and R. McSorley. 2001. Comparison of the survival of three species of sac spiders on natural and artificial diets. *J. Arachnology*, 29:253-263.
2. Amalin, D.M., J., Reisk, J.E. Pena and R.McSorley. 2001. Predatory behavior of three species of sac spider attacking citrus leaf miner. *J. of Acarology*, 29: 72-81.
3. Carrol, D.P. 1980. Biological notes on the spiders of some citrus groves in central and southern California. *Entomol. News*, 91: 147-154.
4. Greenstone, M.H. and K.D. Sunderland. 1999. Why a symposium on spiders in agro ecosystems now? PP. 267-267. In *Proceeding, 15th International Congress of Arachnology and Symposium on Spiders in Agro ecosystems, 27 June-3July, 1998 Chicago II*.
5. Mansour, F., D. Rosen and A. Shulov. 1980. Biology of the spider *Chiracanthium mileid* (Arachnida : Clubionidae). *Entomophaga*, 25 (3) : 237-248.
6. Miliczky, E.R. and C.O. Calkins. 2002. Spiders (Araneae) as potential predators of leaf roller larvae and egg masses (Lepidoptera: Tortricidae) in central Washington apple and pear orchards. *Pan-Pacific Entomology*, 78:140-150.
7. Nyffeler, M. and G. Benz. 1987. Spiders in natural pest control a review. *J. of Applied Entomology*, 103: 321-329.
8. Nyffeler, M., R.G, Breene, D.A. Dean and W.L. Sterling. 1990. Spiders as predators of arthropod eggs. *J. of Applied Entomology*, 109:490-501.

9. Peck, W.B. and W.H. Whitcomb. 1970. Studies on the biology of spider *Chiracanthium inclusum* (Hentz). Univ. of Arkansas (Fayetteville) Agricultural Experiment Station Bulletin 753: 76 pages.
10. Pfannenstiel, R.S. 2008. Spider predators of Lepidopteran eggs in South Texas field crops. *Biological Control*, 46: 202-208.
11. Rashad, A.M. and E.D. Ammar. 1985. Mass rearing of the spiny bollworm, *Earias insulana* on semi- artificial diet. *Bull. Soc. ent. Egypte*, 65: 239- 244.
12. Ruiz, R.I. 2006. Response of *Chiracanthium inclusum* spiderlings to kairomones associated with the eggs and scales of their prey, the cotton bollworm. M.Sc. Thesis, University of Texas at Pan American, Edinburg, Texas, U.S.A.
13. Sallam , G.M.E. 1996. Studies on true spiders in Giza Governorate. M. Sc. Thesis, Fac. Agric., Cairo Univ., Egypt, PP.139.
14. Taylor, R.M. 2008. Plant nectar contributes to the survival, activity, growth and fecundity of the nectar-feeding wandering spider *Chiracanthium inclusum* (Hentz) (Miturgidae). Ph.D. Dissertation, Ohio State Univ., Columbus Ohio, U.S.A.
15. Turnbull, A.L. 1973. Ecology of the true spiders (Arancomorphae). *Ann. Rev. Entomol.*, 18:305-348.
16. Whitcomb, W.H. and K. Bell. 1964. Predaceous insects, Spiders and mites of Arkansas cotton fields. Univ. of Arkansas (Fayetteville) Agricultural Experiment Station Bulletin 690: 84 pages.
17. Whitcomb, W.H., H. Exline and R. C. Hunter. 1963. Spiders of the Arkansas cotton field. *Annals of the Entomol. Soci. America*, 56:653- 660.

تأثير الفرائس المختلفة علي المظاهر البيولوجية ومعدل التكاثر و الكفاءة الأفتراسية للعنكبوت *Chiracanthium inclusum* Hentz

آمال إبراهيم أبو زيد - ميرفت عبد السميع قنديل - علي أحمد السيد

معهد بحوث وقاية النباتات - مركز البحوث الزراعية - الدقي - الجيزة

تم دراسة تأثير نوعين من الغذاء (الفرائس) وهما دودة اللوز القرنفلية و دودة ورق القطن علي بعض المظاهر البيولوجية لدورة الحياة للاطوار الغير كاملة والأطوار الكاملة و الكفاءة التناسلية وكذلك الكفاءة الأفتراسية للعنكبوت *C. inclusum* تحت الظروف المعملية من درجة حرارة 25 ± 1 °م ورطوبة نسبية 65-70%.

وقد اوضحت النتائج: قصر دورة الحياة للاطوار غير كاملة من العمر الثاني للمفترس و حتي العمر السابع وخلال هذه المراحل استهلكت الأفراد كمية اكبر من دودة اللوز القرنفلية مقارنة بدودة ورق القطن. حيث تشير النتائج ان افضلية دودة اللوز القرنفلية للمفترس حيث اعطت اعلي معدل لوضع البيض للانثي داخل كل كيس منفرد 134.3 بيضة/كيس مقارنة بـ 97 بيضة/كيس منفرد عند التغذية علي دودة ورق القطن الا ان متوسط حياة الطور الكامل كانت 114 يوم للانثي ، 75 يوم للذكر عند التغذية علي دودة اللوز القرنفلية مقارنة بـ 135 يوم للانثي و 98.3 يوم للذكر عند التغذية علي العمر اليرقي الثاني لدودة ورق القطن.