

Animal Reproduction Research Institute,  
AL-Haram, Giza, Egypt.

## **PREGNANCY RATE IN EGYPTIAN BUFFALOES AFTER SYNCHRONIZATION OF ESTRUS WITH PGF<sub>2α</sub> OR OVULATION BY OVSYNCH PROGRAM**

(With 2 Tables and 2 Figures)

By

**F.A. EL ZOHERY; A.A. NASRA  
and S.E.M. AL GHANDOUR**

(Received at 20/9/2006)

تقييم معدل الحمل في الجاموس المصري في فترة ما بعد الولادة باستخدام  
الحقن المزدوج للبروستاجلاندين أو نظام تزامن التبويض

فوزى الزهيري ، عبد الخالق نصره ، صلاح الغندور

تحول استطالة الفترة بين الولادتين في الجاموس المصري من زيادة الكفاءة الانتاجية له مما يؤثر سلبا على اقتصاديات انتاجه الأمر الذي حدى بالبحث عن وسائل متعددة لزيادة الخصوبة به وعلية تم اختيار عدد ٤٨ جاموسة حلابة، قسمت الى ثلاث مجموعات: المجموعة الأولى: حقنت في العضل بهرمون البروستاجلاندين مرتين بينهم ١٤ يوم، ثم تم حقن الحيوانات التي لم تأتى في الشياح بجرعة تالثة ثم التلقيح الاصطناعى بعد ٧٢-٨٠ ساعة. المجموعة الثانية: استخدم معها الحقن المحرر للحائة المنسلية- البروستاجلاندين - المحرر للحائة أما المجموعة التالثة فكانت عينة ضابطة. ولقد كانت نتائج المجموعة الثانية أعلى في نسبة الحمل من المجموعتين الأخرتين.

### **SUMMARY**

Postpartum in buffaloes plays a great role in delay of the cyclicity in buffaloes that seemed to be anestrous. So, the use of hormones to induct estrus or induct ovulation has an importance to increase pregnancy rate. The present study was designed to evaluate the efficacy of the two methods in inducing high pregnancy rate. The experimental animals (48 lactating buffaloes) were divided into three groups: PGF<sub>2α</sub> group 1(n=16); Ovsynch group 2 (n=16) and control group (n=16). Pregnancy rate was significantly (p<0.001)increased in group 2 than group 1 and control group due to higher level of progesterone and more accurate synchronization for ovulation., It is more suitable to use the regimen of ovsynch in postpartum buffaloes especially in multiparous females with good condition score.

### **C-Blood Samples:**

Blood samples were collected on days 0 and 25 after insemination from the jugular vein by venipuncture into 10 ml vacutainer tubes. Serum was separated and stored at  $-20^{\circ}\text{C}$  until Progesterone hormone assay.

### **D- Hormonal assay:**

Progesterone was measured by direct radio immunoassay (RIA) using coat A count kit (Diagnostic Products Corporation) DPC.

### **E- Pregnancy diagnosis:**

Progesterone  $>1.0$  ng/ml on day 25 post insemination was considered as pregnant and confirmed 60 days after the insemination by rectal palpation.

### **F- Statistical analysis:**

The data were statistically analyzed according to Snedecor and Cochran (1980). A Chi- square test was used to compare pregnancy rates to TAI (ovsynch) versus induced estrus with  $\text{PGF}_{2\alpha}$ .

## **RESULTS**

**Group I** ( $\text{PGF}_{2\alpha}$  group), 4/8(50%) primiparous buffaloes responded and became pregnant (2, 1 and 1 pregnant buffalo after the 1<sup>st</sup>, 2<sup>nd</sup> and 3<sup>rd</sup> injections respectively), 5/8 (62.5%) multiparous buffaloes became pregnant (2, 2 and 1 after the 1<sup>st</sup>, 2<sup>nd</sup> and 3<sup>rd</sup> injections, respectively) with high significance ( $P < 0.001$ ) as shown in Table 1.

On the other hand pregnancy rates in (Group II, Ovsynch group), were 4/8 (50%) primiparous buffaloes (1 and 3 after injection of  $\text{PGF}_{2\alpha}$  and 2<sup>nd</sup> injection of GnRH, respectively), 6/8 (75%) multiparous buffaloes became pregnant (1, 2 and 3 after injection of 1<sup>st</sup> injection of GnRH,  $\text{PGF}_{2\alpha}$  injection and 2<sup>nd</sup> injection of GnRH, respectively) with high significant ( $P < 0.001$ ).

**Group III** (control group) for primiparous and multiparous buffaloes showed pregnancy rate 3/8 (37.5%) and 4/8 (50%) respectively as shown in Table 1.

The overall of pregnancy rates were 9/16 (56.25%) in  $\text{PGF}_{2\alpha}$  group, 10 /16 (62.5%) in ovsynch group and 7/16 (43.75%) in control group with a high significant ( $P < 0.001$ ) increase in ovsynch group rather than control group and  $\text{PGF}_{2\alpha}$  group as shown in Table 1.

**Table 1:** Pregnancy rates in different groups (%)

Animal Group	Primiparous	Multiparous	tal
Group I ( $\text{PGF}_{2\alpha}$ )	50.0 (4/8) <sup>a</sup>	62.5 (5/8) <sup>b</sup>	56.25 (9/16) <sup>B</sup>
Group II ( Ovsynch )	50.0 (4/8) <sup>a</sup>	75.0 (6/8) <sup>a</sup>	62.5 (10/16) <sup>A</sup>
Group III ( Control )	37.5 (3/8) <sup>b</sup>	50.0 (4/8) <sup>c</sup>	43.75 (7/16) <sup>C</sup>

Total rates with different capital letters superscript in the last column differs significantly ( $P \leq 0.001$ ).

Pregnancy rates with different small letters superscripts in the same column differs significantly ( $P \leq 0.001$ ).

**Progesterone concentration:**

As presented in Table (2) serum concentration of progesterone were not different for primiparous buffaloes in Groups I, II and III just before treatment ( $0.686 \pm 0.115$ ,  $0.755 \pm 0.076$  and  $0.676 \pm 0.045$  ng/ml respectively,) while, in multiparous buffaloes in Groups I, II and III were ( $0.927 \pm 0.157$ ,  $0.722 \pm 0.058$  and  $0.767 \pm 0.085$  ng/ml, respectively).

Regarding progesterone level after 25 days post treatment, it is found that the group II has a higher significant increase ( $P \leq 0.001$ ) ( $4.886 \pm 0.333$  ng/ml) for primiparous and ( $5.083 \pm 0.100$  ng/ml) for multiparous buffaloes; than group I ( $4.684 \pm 0.126$ ,  $4.734 \pm 0.230$  ng/ml); and group III ( $4.837 \pm 0.353$ ,  $4.239 \pm 0.124$  ng/ml) primiparous and multiparous pregnant animals, respectively.

**Table 2:** Serum progesterone profile for different groups (ng/ml) (Mean  $\pm$  SEM)

Animal groups	Pretreatment (on day 0)		Post insemination (on day 25)			
	Primiparous	Multiparous	Non- Pregnant		Pregnant	
			Primiparous	Multiparous	Primiparous	Multiparous
G I (PG - PG)	N=8 0.686 <sup>b</sup> $\pm$ 0.115	N=8 0.927 <sup>a</sup> $\pm$ 0.157	N=4 1.118 <sup>Ba</sup> $\pm$ 0.076	N=3 0.865 <sup>Cb</sup> $\pm$ 0.288	N=4 4.684 <sup>Bb</sup> $\pm$ 0.126	N=5 4.734 <sup>Ba</sup> $\pm$ 0.230
	N=8 0.755 <sup>b</sup> $\pm$ 0.076	N=8 0.722 <sup>a</sup> $\pm$ 0.058	N=4 2.473 <sup>Aa</sup> $\pm$ 0.448	N=2 1.139 <sup>Bb</sup> $\pm$ 0.044	N=4 4.886 <sup>Aa</sup> $\pm$ 0.333	N=6 5.083 <sup>Aa</sup> $\pm$ 0.100
	N=8 0.676 <sup>b</sup> $\pm$ 0.045	N=8 0.767 <sup>a</sup> $\pm$ 0.085	N=5 0.903 <sup>Cb</sup> $\pm$ 0.042	N=4 1.385 <sup>Aa</sup> $\pm$ 0.096	N=3 4.837 <sup>Aa</sup> $\pm$ 0.353	N=4 4.239 <sup>Cb</sup> $\pm$ 0.124

Progesterone levels (on day 0) with different small alphabetical superscripts in the same column differs significantly ( $P \leq 0.001$ ).

Progesterone levels with different capital alphabetical superscript in the same column differ significantly at least at  $p \leq 0.001$ . while, progesterone level (either in pregnant or in the non- pregnant group in the same row) with different small alphabetical superscript differ significantly at least at ( $p \leq 0.01$ ).

### Pregnancy rate




Figure 1: Pregnancy rates (%) in different groups.

### Progesterone profile




Figure 2: Serum progesterone profile (ng/ml) in different groups.

As shown in figure 1 pregnancy rate was higher in group II than other groups, while Figure 2 showed progesterone profile was higher comparatively in group II than other groups.

## DISCUSSION

The main purpose of this study was to evaluate the effect of  $\text{PGF}_{2\alpha}$  as a way of synchronization of estrus and the Ovsynch program for ovulation synchronization with pregnancy rates in Egyptian buffaloes.  $\text{PGF}_{2\alpha}$  group resulted in 56.25% pregnancy rate in lactating buffaloes these results corresponded to the results in cows by Lucy *et al.*, (1986); Stevenson *et al.* (1987) and Archbald *et al.* (1992) that possibly due to the variability in time from AI to ovulation. This variability in time from injection of  $\text{PGF}_{2\alpha}$  to estrus and subsequent ovulation may be directly related to the number and size of the ovulatory follicles at the time of  $\text{PGF}_{2\alpha}$  injection. In contrast to the Ovsynch protocol that showed pregnancy rate 62.5% which is considered high significance ( $P < 0.001$ ) rather than other groups ( $\text{PGF}_{2\alpha}$  group and control group), this result agreed with those applied on Brazilian buffaloes by Roy *et al.* (1996); Burke *et al.* (1996); Momcilovic *et al.* (1998); Berber *et al.* (2002); Bartolomeu *et al.* (2002) and Paul and prakash (2005). A high percentage of pregnancy rate in lactating buffaloes (75%) at a random stage of the estrus cycle ovulated a follicle after first injection of GnRH specially in multiparous rather than primiparous that was in accordance with that results in cows revealed by Silcox *et al.* (1993). The difference between pregnancy rates of primiparous (50%, 50%, 37.5%) and multiparous (62.5%, 75%, 50%) in the different groups respectively agreed with Pietro *et al.* (2003) and demonstrated that the parity is a decisive factor in the efficiency of the protocols. Also, they reported that primiparous spend energy for the continuity of the corporal growth and multiparous converge that energy for the reproductive processes.

In regarding to plasma progesterone concentrations of pregnant buffaloes in group I and II tended to be higher in multiparous buffaloes  $4.734 \pm 0.230$  ng/ml and  $5.083 \pm 0.100$  ng/ml than in multiparous of group 3 ( $4.239 \pm 0.124$  ng/ml while they are not significant to the corresponding levels of the primiparous buffaloes  $4.684 \pm 0.126$  ng/ml and  $4.886 \pm 0.333$  ng/ml in group I and II respectively that agreed with Mee, *et al.* (1993). This may be attributed to high level of progesterone concentrations of GPG buffaloes were probably due to the stimulatory effects of GnRH, or to prolonged effects of GnRH on the proportion of large luteal cells in the developing corpus luteum during pregnancy. So, estrus expression is limited with  $\text{PGF}_{2\alpha}$  than GPG protocol based on fixed-time insemination protocols in lactating buffaloes and necessitates

minimal amounts of estrus detection in order to reach maximal pregnancy rates.

## REFERENCES

- Archbald, L.F.; Tran, T.; Massey, R. and Klapstein, E. (1992):* Conception rates in dairy cow after timed-insemination and simultaneous treatment with gonadotrophin-releasing hormone and/or prostaglandin F<sub>2α</sub>. *Theriogenology*. 37: 723-226.
- Bartolomeu, C.C.; Del Rei, A.J.M.; Madureira, E.H.; Souza, A.J.; Silva, A.U. and Baruselli, P.S. (2002):* Timed insemination using synchronization of ovulation in buffaloes using CIDR-B, CRESTAR and Ovsynch. *Anim. Breed Abstr.* 70: 332 -235.
- Baruselli, P.S.; Madureira, E.H.; Barnabe, V.H.; Barnabe, R.C.; Berber, R.C.A. and Amaral, R. (2000):* Timed insemination using synchronization of ovulation in buffalo. 14<sup>th</sup> International Congress of Animal Reproduction, Barazelu, V. 14, P. 39 -43.
- Baruselli, P.S.; Amaral, R.; Barufi, F.B.; Valentim, R. and Marques, M.O. (2001):* Lecirelin and Buserelin (gonadotrophin-releasing hormone agonists) are equally effective for fixed timed insemination in buffalo. *Braz. J. Vet. Res. Anim. Sci.* Vol.38 No.3 P.215-218, Sao. Paulo.
- Baruselli, P.S.; Madureira, E.H.; Banarbe, V.H.; Barnabe, R.C. and Berber, R.C.A. (2003):* Evaluation of synchronization of ovulation for fixed timed insemination in buffalo (*Bubalus bubalis*). *Braz. J. Vet. Res. Anim. Sci.* Vol.40 No.6 P. 433-437, Sao. Paulo.
- Berber, R.C.; de A.; Madureira, E.H. and Baruselli, P.S. (2002):* Comparison of two Ovsynch protocols (GnRH versus LH) for fixed timed insemination in buffalo (*Bubalus bubalis*) *Theriogenology*. 57: 1421-1430.
- Britt, J.S. and Gaska, J. (1998):* Comparison of two estrus synchronization programs in a large, confinement-housed dairy herd. *JAVMA*.212:210-212.
- Burke, J.M.; De la Sota, R.L.; Risco, C.A.; Staples, C.R.; Schmitt, E.J-P. and Thatcher, W.W. (1996):* Evaluation of timed insemination using a gonadotrophin-releasing hormone agonist in lactating dairy cows. *J. Dairy Sci.* 79: 1385-1393.