

Dept. Theriogenology
Fac. Vet. Med., Assiut University.

**MORPHOMETRIC AND IMMUNOHISTOCHEMICAL
VARIATIONS IN THE CAMEL (*CAMELUS
DROMEDARIUS*) TESTIS IN RELATION TO SOME
ENDOCRINOLOGICAL ASPECTS DURING
DIFFERENT SEASONS OF THE YEAR**
(With 2 Tables and 3 Figures)

By

D.R.I. DERAR; H.A. HUSSEIN and A.M. SALEH*

*Dept. of Anatomy and Histology, Fac. Vet. Med., Assiut University

(Received at 16/12/2004)

**التغيرات في القياس الظاهري وكيمياء النسيج المناعي في خصية
الجمال وحيد السنام نسبة إلى بعض النواحي الهرمونية
خلال المواسم المختلفة من السنة**

ضرار رفعت ضرار ، حسن عبد الصبور حسين ، عبد المهيمن مصطفى صالح

تمت دراسة التغيرات الموسمية التي تطرأ علي مستوي هرمون التستستيرون في المصل مدعومة باستكشاف التفاعل المناعي لإنزيم ٣ بيتا هيدروكسي ستيرويد ديهيدروجينيز في خلايا لايدج لعدد ٥٤ جمال وحيد السنام وناضج جنسيا وسليم ظاهريا. أظهرت النتائج أن خلايا لايدج كانت كثيرة ومنتشرة في النسيج البين خلوي مع تفاعل قوي وإيجابي لإنزيم ٣ بيتا هيدروكسي ستيرويد ديهيدروجينيز أثناء أشهر ديسمبر ويناير وفبراير ومارس. تفاعل هذا الإنزيم كان خافتا وضعيفا تدريجيا أثناء الأشهر من أبريل وحتى يوليو عندها لم يكن ممكنا ملاحظة التفاعل المناعي لهذا الإنزيم. وفي ذات الوقت -ومصاحبا لهذه التغيرات-كانت هناك زيادة كبيرة في هرمون التستستيرون في المصل ابتداءً من شهر ديسمبر وانتهاءً بشهر أبريل مع ملاحظة أن أعلى مستويات له كانت في شهر يناير. لوحظ أن مستوي هرمون الثيروكسين في المصل يمر تقريبا بنفس التغيرات من زيادة ونقصان متوازيا مع التي شهدها هرمون التستستيرون. من خلال نتائج الدراسة الحالية يمكن القول أن إنزيم ٣ بيتا هيدروكسي ستيرويد ديهيدروجينيز له أهمية قصوى في تنظيم عملية تخليق هرمون التستستيرون من خلايا لايدج في ذكور الجمال وحيدة السنام وأن هرمون الثيروكسين هام جدا للنشاط التناسلي أثناء موسم التزاوج في ذكور الجمال وحيدة السنام ويمر بنفس التغيرات التي يمر بها هرمون الذكورة.

SUMMARY

Seasonal variation in serum testosterone, thyroxin and the testicular morphology were studied in 54 sexually mature and apparently healthy one-humped camels during the different seasons of the year. The

testosterone and thyroxin serum levels were measured and 3 β -hydroxysteroid dehydrogenase activity of Leydig cells was assessed immunohistochemically to aid in the interpretation of results. The activity of 3 β -HSD was high during cold months and severely depressed to the minimum activity in hot months. Concomitantly, serum testosterone and thyroxin levels increased during the winter and early spring and decreased thereafter. Their levels reached the peak during the months of January till April. These results suggested that 3 β -HSD is a key enzyme in the regulation of the testosterone production in Leydig cells of the male dromedary. Thyroxin is a crucial hormone for the male reproductive activity during the breeding season in the dromedary and fluctuated in the same pattern as serum male androgen.

Key words: *Camel, immunohistochemical, testis, thyroxin 3 β -HSD.*

INTRODUCTION

Seasonal changes in the camels could be clarified through studying morphology of the testis, histochemical observation of the testes and studies on the male accessory sex glands (Abdel-Raouf and Owaida 1974, Abdel-Raouf *et al.* 1975). It has been found that in seasonal breeders the mating and nonmating seasons are clearly related to different levels of testosterone in the plasma and testes (Racey, 1978). Clear correlation between testicular steroidogenesis and reproduction is well exemplified in the dromedary that is not a typical seasonal breeder. The rate of synthesis of testosterone is high during the mating season and low during the nonmating season (Friedlander *et al.* 1984).

Seasonality in the male is evidenced by changes in sexual behaviour, morphology and function of the genital organs, as well as changes in endocrinological profiles. In seasonal breeders, the effect of photoperiod is undeniable. In this regard, Vaughan *et al.* (1982) explained that chronic exposure of female Syrian hamsters (long day breeders) for 9 weeks to a short photoperiod (10L:14D) depressed the pituitary-thyroid axis as indicated by a drop in circulating titers of thyroid stimulating hormone (TSH), thyroxin (T₄), triiodothyronine (T₃) and the free thyroxin index (FT₄) compared to animals maintained under long photoperiodic conditions. 90% of iodine circulating in the animal's blood is in the form of T₄ (Wilson, 1975). Lack of iodine prevents production of both T₄ and T₃ (Guyton, 1991). The enzyme hydroxysteroid dehydrogenase (3 β -HSD) plays a central role in the

biosynthesis of steroid hormones, including androgens (Conley and Bird 1997; Penning 1997). 3β -HSD is present in the testis, ovary and placenta, adrenal gland as well as in a large number of peripheral intracrine tissues, including the prostate, breast, liver and skin (Ferre *et al.* 1975; Lacoste *et al.* 1990). It catalyzes the final step in progesterone biosynthesis in the ovary and is required for testosterone production in the testis. Different 3β -HSD isoforms have been cloned from various tissues from humans, rats and mice (Simard *et al.* 1993 and 1995 and Penning 1997).

The purpose of the present work was to investigate the correlation between seasonal changes in serum testosterone and thyroxin hormones from one side and the 3β -HSD activity and testicular morphology on the other side.

MATERIALS and METHODS

The testes of 48 sexually mature (5-12 years old) and apparently healthy one-humped camels were obtained from Bany-Ady (Assiut governorate) and Cairo slaughterhouses. The materials were collected at regular monthly intervals over a period of twelve months. Within one hour after slaughter, the scrotum was incised, the testes were removed and the volume was taken. For the histological morphometric study, the testes were cut in slices and small cubes from the testicular parenchyma were taken and fixed in neutral buffer formalin contained 1% glutaraldehyde. Then, processed for paraffin embedding, and sections of 5 μ m thick were cut and stained with H&E.

Morphometric studies: The weight of testes was taken and their volume was measured by water displacement method (Willett and Ohms, 1957 and Scherle, 1970). The testicular parenchymal volume was calculated by subtraction of 11% from the testicular volume. These 11% represent the volume of tunica albuginea and rete testis (Wrobel 1990). Different histological morphometric values were carried out on H&E stained sections using Leica Q 500 MC Image analyzer.

3β -Hydroxysteroid dehydrogenase (3β -HSD):

Fixation: Immediately after slaughter, the testes were removed from their envelopes and samples of suitable size (about 1.0 x 0.5 x 0.5 cm) were taken from different regions testis every month. Immersion fixation was carried out in two steps. Fixative I (30 min) contained 4% paraformaldehyde; 15% v/v saturated picric acid; 0.1% glutaraldehyde in 0.1 M phosphate buffer, pH 7.4. Fixative II (several hours) had the same composition as fixative I but without glutaraldehyde. Following fixation,

the blocks were washed in 0.1 M phosphate buffer, transferred into a graded series (10%, 20%, 30%) of saccharose-containing rinsing buffer and shipped by air to Regensburg. Here, the samples were immersed in Tissue Tek OCT compound (Miles, Elkhardt, Ind., USA) and snap-frozen in liquid nitrogen. Cryostat sections (12 μm thick) were mounted on gelatin/chrome alum-coated slides and air-dried for 2 - 3 min before further treatment.

Immunohistochemistry: All subsequent steps were carried out in a moist chamber on slides with sections surrounded by water-repellent PAP-PEN (SCI Science Services, München, Germany). Sections were rinsed (3 x 10 min) in TBS: 0.1 M TRIS (pH 7.4); 0.8 % NaCl; 0.0015 % Triton X-100 between the consecutive steps of the test. (1) Non-specific bindings were blocked by preincubation (60 min) with blocking buffer containing 0.1 M TRIS (pH 7.4); 0.15 % Thimerosal; 0.8 % Triton X-100; 0.8 % NaCl; 20 % normal goat serum; 20 % fetal calf serum. (2) Incubation (overnight) with primary antiserum (rabbit anti-mouse adrenal/gonadal 3β -HSD) at a dilution in blocking buffer of 1:512 overnight at 4°C. (3) Incubation (60 min) in secondary antibody/biotinylated in blocking buffer. (4) Blocking of endogeneous peroxidase with phenylhydrazine. (5) Incubation (60 min) in AB complex (ABC). (6) Developing with 0.5 mg/ml DAB; in 0.1 M TRIS (pH 7.4) containing 0.002 % $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$; 0.4 % $\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$ and 0.012 % H_2O_2 . (7) Rinsing once in TBS, dehydration, mounting in DPX.

Controls for immunohistochemistry: Controls included: (a) Omission of the primary antiserum. (b) Substitution of primary antiserum by non-immune serum 1: 500 in blocking buffer. (c) Blocking of the primary antibody by preincubation with the matching antigen in excess. No immunostaining was obtained after any of these control procedures (a-c).

Blood sampling and hormonal assay: Samples were collected by Venipuncture at a monthly interval from six mature male camels housed in the veterinary teaching clinic, Assiut Univ. and its neighborhood. Samples were kept at 4°C for 30 minutes and the serum was obtained after centrifugation at 3000 rpm for 20 minutes and stored at -20°C till assay. Serum testosterone and Thyroxin were determined using ELISA kit (Biosource, Europe, S.A., Code, 40 17000). Inter- and intra-assay coefficients were 6.2, 6.4% for testosterone and 11.4 and 11.7 % for Thyroxin, respectively.

Statistical analysis: Statistical analysis of the collected data was carried out according to procedures of completely random design, SAS (1995).

RESULTS

The weight, volume densities (mean \pm SE) of testes and testicular parenchyma throughout the year are shown in (Table 1, Fig 1A). There was one fold difference between the highest and lowest mean testicular volume during different reproductive cycle. The testicular weight and volume showed similar peak during the breeding phases. The testes began to increase in weight and volume from quiescence (September) and attained a peak during breeding season (December, January and February). Then, testes declined in volume and weight in March and April to reach the lowest values in July.

The epithelial height and diameter of seminiferous tubules (table 1, Fig 1B) displays statistically significant annual changes. The tubular diameter showed significant increase in February and March and lowest values were recorded in July. The epithelial height showed significantly higher values during the breeding season in December and January. The Leydig cells started to increase in number in September to reach the maximum number in December and showed significant seasonal variations (table 1, Fig 1B).


Table 1: seasonal and monthly variation in testicular volume, diameter of seminiferous tubules and epithelial height of seminiferous tubules in the male dromedary (n = 48).

Mon.	Day length (hrs)	Testicular weight (g)	Testicular Volume (c.c.)	Parenchyma volume of (c.c.)	Leydig cells number/Crosse section 250X	Diameter of seminiferous tubules (μ m)	Epithelial height (μ m)
Jun July (Summer) Aug.	14.15	64.0 \pm 3.0 ^a	65.8 \pm 3.3 ^a	47.1 \pm 2.2 ^a	55 \pm 4.4 ^a	169.4 \pm 2.5 ^a	77.5 \pm 1.6 ^{ab}
	13.50	64.3 \pm 3.5 ^a	65.3 \pm 3.9 ^a	47.1 \pm 2.6 ^a	40 \pm 3.7 ^a	163.5 \pm 1.6 ^a	77.5 \pm 1.7 ^{ab}
	13.15	64.8 \pm 4.0 ^a	66.8 \pm 4.3 ^a	47.2 \pm 3.2 ^a	40 \pm 4.1 ^a	170.3 \pm 1.7 ^a	74.6 \pm 2.3 ^{ab}
Sept. Oct. (Autumn) Nov.	12.30	66.5 \pm 5.1 ^a	68.9 \pm 5.3 ^a	50.8 \pm 3.2 ^a	90 \pm 5.2 ^{ab}	187.5 \pm 3.8 ^{ab}	63.4 \pm 2.6 ^a
	11.30	71.0 \pm 3.3 ^{ab}	73.8 \pm 3.3 ^{ab}	58.0 \pm 2.2 ^{ab}	95 \pm 5.0 ^{ab}	190.5 \pm 4.6 ^{ab}	78.4 \pm 1.4 ^{ab}
	10.15	78.0 \pm 3.4 ^{ab}	79.3 \pm 3.6 ^{ab}	63.0 \pm 3.5 ^{ab}	110 \pm 6.7 ^{ab}	198.8 \pm 2.8 ^b	86.1 \pm 5.2 ^b
Dec. Jan. (Winter) Feb.	10.45	94.8 \pm 4.3 ^b	95.8 \pm 4.3 ^b	78.0 \pm 2.1 ^b	182 \pm 8.0 ^b	200.4 \pm 4.3 ^b	90.3 \pm 2.1 ^b
	11.0	96.6 \pm 5.6 ^b	97.6 \pm 6.6 ^b	79.6 \pm 2.4 ^b	280 \pm 8.7 ^b	210.1 \pm 4.8 ^{bc}	91.4 \pm 3.1 ^b
	11.45	93.9 \pm 5.3 ^b	94.9 \pm 4.3 ^b	77.2 \pm 3.6 ^b	265 \pm 10 ^b	210.4 \pm 5.1 ^{bc}	90.5 \pm 4.6 ^b
March April (Spring) May	12.0	90.0 \pm 5.7 ^b	90.5 \pm 5.7 ^b	73.2 \pm 4.8 ^b	180 \pm 9.6 ^b	220.6 \pm 5.2 ^c	90.3 \pm 2.1 ^b
	12.50	71.2 \pm 4.3 ^{ab}	73.8 \pm 5.3 ^{ab}	67.1 \pm 4.5 ^b	160 \pm 9.0 ^b	190.5 \pm 6.1 ^{ab}	86.4 \pm 2.5 ^b
	13.50	60.1 \pm 4.1 ^{ab}	61.1 \pm 4.2 ^{ab}	55.5 \pm 3.3 ^a	90 \pm 4.1 ^a	180.7 \pm 4.2 ^{ab}	81.5 \pm 1.9 ^{ab}


- Values in mean \pm mean standard error.
- means in the same column with the same letter were not significantly different.
- a, b, c means with different superscripts were significantly different (p<0.05).

Figure 1: seasonal variation in: (A) testicular weight, testicular volume and Parenchymal volume B) Leydig cells number, Tubular diameter and Epithelial height of the testicular tubule.

Testicular weight (W), Testicular volume and parenchymal volume (V)






Leydig cells number (No.), diameter of semi. tub. and epith. height (um)



3 β -HSD immunohistochemistry

The hormonal activity of the Leydig cell was assessed by 3 β -HSD enzyme immunohistochemistry. This reaction revealed dark gray granules localized in the cytoplasm of the Leydig cells. The intensity of the reaction and the population of Leydig cells showed marked annual variations. In June, July and August, a weak reaction in and low number Leydig cells were observed. September and October, intertubular tissue showed few number of Leydig cells with strong 3 β -HSD reaction. The number of the Leydig cells increased to reach their maximum population and strongest 3 β -HSD reaction in December, January and February (qualitative and quantitative). In March April and May, the intertubular tissues contained abundant number of Leydig cells with relatively weak 3 β -HSD reaction (Figure 2).

Figure 2: Changes in 3 β -HSD immunoreactivity throughout different seasons of the year in the male camel. 250x.



Hormonal concentration


In the beginning of the year, during January to April months, serum testosterone increased significantly and reached a peak level during these months As shown in table (2), Figure (3).

Table 2: Serum concentrations of testosterone and thyroxin all over the year in male camels (n = 6).

Month	Season	Testosterone ng/ml	Thyroxin ng/dL
June	Summer	2.58 ± 0.72 ^c	66.50 ± 7.81 ^{bc}
July		1.30 ± 0.75 ^c	52.16 ± 9.05 ^c
August		1.11 ± 0.56 ^c	50.33 ± 7.70 ^c
September	Autumn	2.54 ± 0.44 ^c	41.33 ± 16.06 ^c
October		1.43 ± 0.26 ^c	4.42 ± 4.49 ^{cd}
November		1.49 ± 0.40 ^c	26.50 ± 2.57 ^d
December	Winter	15.94 ± 3.40 ^b	64.50 ± 8.34 ^{bc}
January		30.86 ± 2.96 ^a	75.83 ± 12.08 ^b
February		30.37 ± 3.00 ^a	96.50 ± 19.20 ^{ab}
March	Spring	29.93 ± 3.88 ^a	91.00 ± 17.39 ^b
April		30.99 ± 5.88 ^a	116.00 ± 18.50 ^a
May		13.59 ± 2.23 ^b	87.83 ± 6.16 ^b

- Values in mean ± mean standard error.
- Means in the same column with the same letter were not significantly different.
- a, b, c means with different superscripts were significantly different(p<0.05).

Figure 3: annual variation in serum testosterone and thyroxin levels in the male dromedary.



DISCUSSION

The reproductive activity of the camel builds up during September and October, and the animal is in actual rut during November, December, January and February, with a drop in March and thereafter (Abdel-Raouf and owaida, 1974, Abdel-Raouf *et al.*, 1975 and Tingari *et al.*, 1984). In the present study, the testes began to increase in volume from quiescence (September) and attained a peak during breeding season (December, January and February). Then, testes declined in volume in March and April to reach the lowest values in July. The present findings are coincident with those of Singh and Bharadwaj (1978) and Zayed *et al.* (1995).

Thornton *et al.* (2002) suggested that plasma androgen and/or IGF-1 levels may be important in modulating the expression of some s in the regulation of the testosterone production in Leydig cells. In this study, a steroidogenic enzymes like 3beta-HSD. These results suggest that 3beta HSD is a key enzyme strong positive relation between serum androgen and histochemistry reaction of 3β-HSD was noticed. In June and July, The Leydig cells showed low population and a very weak 3β-HSD activity. The activity of the 3β-HSD increased steadily in September and October and reached the maximum activity in December, January, February and March (qualitative and quantitative). The Leydig cells are in highest population and highest morphological differentiation and their smooth ER (SER) is highly developed (Zayed *et al.* 1995). In April and May, the Leydig cells decreased steadily in population and 3β-HSD activity to reach the minimum state in Jun and July. The cells were small in size, SER is reduced and many Leydig cells are degenerating (Zayed *et al.* 1995). These findings supported the previous results reported for camel (Yagil and Etzion, 1980), stallion (Johnson and Thompson, 1987) and Japanese black bear (Komatsu *et al.* 1997).

Serum testosterone level was 15.94 ± 3.40 ng/ml in December, reached a plateau in April (30.99 ± 5.88 ng/ml) and decreased to 13.59 ± 2.23 ng/ml in May and dropped to its lowest level at June (2.58 ± 0.72 ng/ml) and continued at this nadir through the summer and autumn seasons. It has been previously shown that the differential testosterone synthesis between the seasons in the dromedary is not quantitative. However, during the mating season, the synthesis of testosterone synthesis is through both 4-ene and 5-ene pathways, whereas during the non mating season, the synthesis occurs mainly through the 5-ene pathway and at a lower rate than that of this pathway during the mating

season (Bedrak *et al.*, 1983). Delgadillo *et al.* (2004) reported that Short days enhanced testosterone secretion and long days inhibited it in seasonal breeder males. During the mating season, Leydig cells were highly packed and larger than during the nonmating season (Friedlander *et al.*, 1984).

Abdel-Raouf *et al* (1975) claimed that the largest seminiferous tubule diameters and the greatest numbers of spermatogonia, spermatids and spermatozoa were found in the spring. The numbers of mature Leydig cells, compared to the numbers of pre-Leydig and immature Leydig cells, increased by the end of winter so that, during the spring, the interstitial cells were mainly of the mature type. Degenerative changes with diminished numbers of mature cells were seen in the summer and this trend continued into early and mid-autumn. In the stallion (long day breeder), Johnson and Thompson (1987) found that the volume of Smooth Endoplasmic Reticulum (SER)/g and testosterone/g tended to be higher in the breeding than non-breeding season. Leydig cell number/g, volume of SER/testis, testosterone/testis, and Leydig cell number/testis were significantly greater in the breeding than in the non-breeding season. Volume of SER/testis and testosterone/testis were related significantly to the cell number/testis, and SER/testis was related ($P < 0.05$) to testosterone/testis.

Our findings support the concept that the thyroid gland plays a fundamental role in seasonal reproduction in the male camel. An annual cycle of serum thyroxin was detected; values reached a peak in winter (late breeding season) and a minimal level in summer (late anestrus). Significant increase in serum thyroxin concentration was found during the period from December till June. Maximum level of thyroxin was found in April (116.00 ± 18.5 ng/dL). The increase in serum thyroxin concentrations was coincident with an increase in serum androgen concentration. Moreover, a simultaneous significant decline in serum thyroxin and androgen levels were found nearly in the same time during the nonmating season. Wasfi, *et al.* (1987) reported values of 9.33 ± 1.15 ng/ml (1.43 ± 0.18 nmol/l) for T_3 concentration in normal Saudi Arabian camels. Nazifi *et al.*, (1999) found that the concentrations of T_3 and T_4 were higher in the breeding season compared with the rest of the year ($p < 0.05$). Thyroidal hormones (T_3 and T_4) showed significant correlations with serum total protein and glucose.

A significant positive correlation between plasma levels of cholesterol and both serum thyroxin and testosterone in males was reported (Heller *et al* 1981). Webster *et al* (1991), Anderson *et al.*

(2003) and Hernandez *et al* (2003) mentioned that thyroid hormones did not alter onset of the breeding season but they were permissive for various species to enter seasonal anestrus. Responsiveness to T₄ is lost gradually during the mid to late anestrus season and thyroid hormones can influence the timing of the breeding season and thus may be required for the maintenance or entrainment of the endogenous reproductive rhythm (Anderson *et al*, 2002 Billings *et al*, 2002). Viguié *et al* (1999) provided strong evidence that thyroid hormones can act directly within the brain to promote seasonal inhibition of neuroendocrine reproductive function in the ewe. Further, the reproductive neuroendocrine axis is not equally responsive to thyroid hormone at all times of the year (Thrun *et al*, 1997). It is concluded that the decline of thyroid function, as gauged by hormone secretion in summer, aids in preservation of body water by decreasing pulmonary water loss and dropping basic metabolism (Yagil *et al*, 1978).

Form the present findings, it seems that, Thyroid hormones are necessary only during a limited interval early in the breeding season to promote seasonal reproductive activation in the male dromedary. There is a critical period of responsiveness during which thyroid hormones must be present for rut to develop. In conclusion, the present findings indicated that there is a concomitant rise and fall in 3 β -HSD, serum testosterone and thyroxin level in the male dromedary which is indicative and diagnostic for the onset of reproductive seasonality. Thyroxin is a key hormone for the resumption of sexual activity after the non rut period in the dromedary.

REFERENCES

- Abdel-Raouf, M. and Owaida, M.M. (1974):* Studies on reproduction in camels (*Camelus dromedarius*). IV. Gross changes in the morphology of the testis in relation to age and season. *Ass. Vet. Med. J.*, 1 :215-223.
- Abdel-Raouf, M.; Fateh El-Bab, M.R. and Owaida, M.M. (1975):* Studies on reproduction in the camel (*Camelus dromedarius*). V. Morphology of the testis in relation to age and season. *J Reprod. Fertil.* 43: 109-116.
- Anderson, G.M.; Connors, J.M.; Hardy, S.L.; Valent, M. and Goodman, R.L. (2002)* Thyroid hormones mediate steroid-independent seasonal changes in Luteinizing Hormone pulsatility in the ewe. *Biol. Reprod.* 66, 701–706.

- Anderson, G.M.; Hardy, S.L.; Valent, M.; Billings, H.J.; Connors, J.M. and Goodman, R.L. (2003):* Evidence that thyroid hormones act in the ventromedial preoptic area and the premammillary region of the brain to allow the termination of the breeding season in the ewe. *Endocrinol.* 144, 2892-2901.
- Bedrak, E.; Rosenstrauch, A.; Kafka, M. and Friedlander, M. (1983):* Testicular steroidogenesis in the camel (*Camelus dromedarius*) during the mating and the nonmating seasons. *Gen. Comp. Endocrinol.* 52, 255-64.
- Billings, H.J.; Viguié, C.; Karsch, F.J.; Goodman, R.L.; Connors, J.M. and Anderson, G.M. (2002):* Temporal requirements of thyroid hormones for seasonal changes in LH secretion. *Endocrinol.*, 143, 2618-2625.
- Conley, AJ and Bird, IM (1997):* The role of cytochrome P450 17 alpha-hydroxylase and 3 beta-hydroxysteroid dehydrogenase in the integration of gonadal and adrenal steroidogenesis via the delta 5 and delta 4 pathways of steroidogenesis in mammals. *Biol. Reprod* 56:789–799.
- Delgadillo, J.A.; Cortez, M.E.; Duarte, G.; Chemineau, P. and Malpoux, B. (2004):* Evidence that the photoperiod controls the annual changes in testosterone secretion, testicular and body weight in subtropical male goats. *Reprod. Nutr. Dev.* 44: 183-193.
- Ferre, F.; Breuiller, M.; Cedarol, L.; Duchesne, M.J.; Saintot, M.; Descomps, B. and Crustes de Paulet, A. (1975):* Human placental D5–3 β hydroxysteroid dehydrogenase activity D5–3 β _HSD; intracellular distribution, kinetic properties, retransinhibition and influence of membrane delipidation. *Steroids* 26:551–570.
- Friedlander, M.; Rosenstrauch, A. and Bedrak, E. (1984):* Leydig cell differentiation during the reproductive cycle of the seasonal breeder *Camelus dromedarius*: An ultrastructural analysis. *Gen. Comp. Endocrinol.*, 55, 1-11.
- Guyton, A.C. (1991):* The thyroid metabolic hormones in: *Textbook of medical physiology*, 8th edn: W.B. Saunders Company, Philadelphia.
- Heller, R.F.; Miller, N.E.; Lewis, B.; Vermeulen, A.; Fairney, A.; James, V.H. and Swan, A.V. (1981):* Associations between sex hormones, thyroid hormones and lipoproteins *Clin. Sci.* 61 (5) 649-51.

- Hernandez, J.A.; Hallford, D.M. and Wells, N.H. (2003):* Ovarian cyclicity in thyroid-suppressed ewes treated with propylthiouracil immediately before onset of seasonal anestrus. *J. Anim. Sci.* 81:29–34.
- Johnson, L. and Thompson, D.L. Jr. (1987):* Effect of seasonal changes in Leydig cell number on the volume of smooth endoplasmic reticulum in Leydig cells and intratesticular testosterone content in stallions. *J. Reprod. Fertil.* 81, 227-32.
- Komatsu, T.; Tsubota, T.; Yamamoto, Y.; Atoji, Y. and Suzuki, Y. (1997):* Seasonal changes in the immunolocalization of steroidogenic enzymes in the testes of the Japanese black bear (*Ursus thibetanus japonicus*). *J. Vet. Med. Sci.* 59, 521-529.
- Lacoste, D; Bélanger, A. and Labrie, F. (1990):* Biosynthesis and degradation of androgen in human prostatic cancer cell lines. In: Castagretta L, D'Aquino S, Labrie F, Bradlow H (eds) Steroid formation, degradation and action in peripheral tissues. *Ann N Y Acad Sci.* 595:389–392.
- Nazifi, S.; Gheisari, H.R. and Poorabbas, H. (1999):* The influences of thermal stress on serum biochemical parameters of dromedary camels and their correlation with thyroid activity. *Comp. Clinic. Pathol.* 9, 49–54.
- Penning, TM. (1997):* Molecular endocrinology of hydroxysteroid dehydrogenases. *Endocr Rev.* 18:281–305.
- Racey, P.A. (1978):* Seasonal changes in testosterone levels and androgen-dependent organs in male moles (*Talpa europea*). *J. Reprod. Fert.* 52, 195-200.
- SAS. (1995):* Statistical Analysis System. SAS/ STAT Users's Guide, Release 6.12 Edition. Cary, NC. SAS inst., Inc.
- Scherle, W.E. (1970):* A simple method for volumetry of organs in quantitative stereology *Mikroskopie.* 26: 57-60.
- Simard, J.; Couet, J.; Durocher, F.; Labrie, Y.; Sanchez, R.; Breton, N.; Turgeon, C. and Labrie, F. (1993):* Structure and tissue-specific expression of a novel member of the rat 3 beta-hydroxysteroid dehydrogenase/delta5-delta 4 isomerase (3 beta-HSD) family. The exclusive 3 beta-HSD gene expression in the skin. *J. Biol. Chem.* 268:19659–19668.

- Simard, J.; Sanchez, R.; Durocher, F.; Rheaume, E.; Turgeon, C.; Labrie, Y.; Luu-The, V.; Mebark, F.; Morel, Y. and De Launoit, Y. (1995):* Structure-function relationship and molecular genetics of the 3 beta-hydroxysteroid dehydrogenase gene family. *J. Steroid Biochem Mol Biol* 55:489–505.
- Singh, U.B. and Bharadwaj, M.B (1978):* Histological and histochemical studies on the testis of camel (*Camelus dromedarius*) during the various seasons and ages. Part II, *Acta Anat.* 101, 280-288.
- Thornton, M.J.; El-Alfy, M. and Labrie, F. (2002):* Seasonal changes in the expression of some steroidogenic enzymes in male red deer skin. *Endocrine (Abstract)* 4, 90.
- Thrun, L.A.; Dahl, G.E.; Evans, N.P. and Karsch, F.J. (1997):* A Critical period for thyroid hormone action on seasonal changes in reproductive neuroendocrine function in the ewe. *Endocrinol.* 138, 3402-3409.
- Tingari, M.D.; Ramos, A.S.; Gaili, E.S.; Rahma, B.A. and Saad, A.H. (1984):* Morphology of the testis of the one-humped camel in relation to reproductive activity. *J. Anat.* 139, 133-43.
- Vaughan, M.K.; Powanda, M.C.; Richardson, B.A.; King, T.S.; Johnson, L.Y. and Reiter, R.J. (1982):* Chronic exposure to short photoperiod inhibits free thyroxin index and plasma levels of TSH, T4, triiodothyronine (T3) and cholesterol in female Syrian hamsters. *Comp. Biochem. Physiol. A.* 71, 615-8.
- Viguié, C.; Battaglia, D.F.; Krasa, H.B.; Thrun, L.A. and Karsch, F.J. (1999):* Thyroid hormones act primarily within the brain to promote the seasonal inhibition of Luteinizing Hormone secretion in the ewe. *Endocrinol.* 140, 1111-1117.
- Wasfi, I.A.; Hafez, A.M.; El-Tayeb, F.A.A. and El-Taher, A.Y. (1987):* Thyroid hormones, cholesterol and triglycerides level in the camel. *Res. Vet. Sci.* 42, 418.
- Webster, J.R.; Moenter, S.M.; Woodfill, C.J. and Karsch, F.J. (1991):* Role of the thyroid gland in seasonal reproduction. II. Thyroxine allows a season-specific suppression of gonadotropin secretion in sheep. *Endocrinol.* 129, 176-183.
- Willett, E.L. and Ohms, J.L. (1957):* Measurement of testicular size and its relation to production of spermatozoa by bulls. *J. Dairy Sci.* 12: 1559-1569.
- Wilson, J.G. (1975):* Hypothyroidism in ruminants with special reference to fetal goiter. *Vet. Rec.* 97, 161-164.

- Wrobel, K.H. (1990):* The postnatal development of the bovine Leydig cell population. *Reprod. Dom. Anim.* 25, 51 – 60.
- Yagil, R. and Etzion, Z. (1980):* Hormonal and behavioural patterns in the male camels (*Camelus dromedarius*). *J Reprod Fert*, 58, 61-65.
- Yagil, R.; Etzion, Z. and Ganani, J. (1978):* Camel thyroid metabolism: effect of season and dehydration. *J. Appl. Physiol.* 45, 540-4.
- Zayed, A.E.; Hifny, A.; Abou-Elmagd, A. and Wrobel, K.H. (1995):* Seasonal changes in the intertubular tissue of the camel testis (*Camelus dromedarius*). *Ann. Anat.*,177:199-212.