

Dept. of Poultry Disease,  
Fac. Vet. Med., Assiut Univ.

**STUDIES ON COMPYLOBACTER INFECTION  
IN LAYER CHICKENS**  
(With 5 Tables and 1 Figure)

By

**A.A. IBRAHIM and HEBAT-ALLAH A. MOHAMED\***

\* Animal Health Research Institute  
(Received at 26/3/2003)

دراسات عن الإصابة بالكامبيلوباكتر في قطعان الدجاج البياض

عوض عبد الحافظ إبراهيم ، هبة الله عبد الحليم محمد

تم فحص الأحشاء الداخلية لـ ٦٧ عينة من دجاج بياض عمر ٢٥-٣٦ أسبوعا حديث النفوق و ٢٥ عينة من قناة البيض و ٣٥ مسحة من فتحة المجمع من حالات حية . وقد تم عزل ميكروب الكامبيلوباكتر جو جيناي وكولاي في ٢١ حالة من ٦٧ بنسبة (٣٤,٣%) و ٣ حالات من ٢٥ بنسبة (١٢%) و ١١ حالة من ١٥ بنسبة (٣١,٤%) علي التوالي. وقد تم إجراء العدوى الصناعية في كتاكيت عمر يوم التي أدت إلى ظهور أعراض المرض ونسبة وفيات وصلت إلى ٧٠% ولكن كتاكيت عمر ٤ أسابيع كانت مقاومة لحد ما ويعمل اختبار الحساسية للعترات المعزولة معمليا وجد أن حمض النالاديكس والإريثروميسين والجنتاميسين هي الأدوية الأكثر تأثيرا .

**SUMMARY**

Sixty seven samples of visceral organs (liver, spleen), blood and bile of freshly dead layers, 25 samples of oviduct and 35 cloacal swabs from living layers 25-36 weeks old collected from farms at Assiut Governorate were examined. Isolation of Campylobacter (C.) revealed 23 positive cases of C. jejuni and C.coli out of 67 (34.3%), 3 out of 25 (12%) and 11 out of 35 (31.4%) respectively. Trials for reproducing the infection in 1 day old Chicks lead to 70% mortality while 4 weeks old chicks were somewhat resistant. In vitro sensitivity test showed that Naladixic acid, Erythromycin and Gentamycin were effective.

**Key words:** *Compylobacter infection, layer chickens.*



## INTRODUCTION

Campylobacteriosis is attributed to infection by thermophilic members of the genus *Campylobacter*, the three species of clinical significance *C.jejuni*, *C.coli* and *C. laridis* are microaerophilic, gram negative, spiral, uniflagellate organisms. (Sebald and Veron, 1963).

Commercial poultry serve as reservoirs of *Campylobacter* infection with isolation rates in feces reached 72% in chickens. (Simmons and Gibbs, 1979).

*Campylobacter jejuni* is the most frequently occurring member of the thermophilic triad (Munroe *et al.*, 1983).

*Campylobacter jejuni* has been identified as one of the major causes of diarrheal disease in humans throughout the world. (Skirrow 1990).

The organism induced infection of humans is epidemiologically linked to the consumption of improperly prepared poultry products or foods cross-contaminated by poultry products. (Shane, 1992)

Recent evidence suggests that *Campylobacter jejuni* can colonize the oviduct of laying hens but the source and role of this colonization are unknown. (Camarda *et al.*, 2000)

Modugno *et al.* (2000) found that *Campylobacter jejuni* biotypes I and II were common, type III was rare and type IV was absent.

### The aim of the present work is:

- Isolation and identification of *Campylobacter* spp. from laying hens.
- Biotyping of *Campylobacter jejuni*.
- Experimental infection of baby chicks with the isolated organism.
- Testing the isolate against several antibiotic discs to give a suitable treatment.

## MATERIALS and METHODS

### Samples:

Visceral organs (liver, spleen), bile and blood of freshly dead 67 layers 25 samples of oviduct and 35 cloacal swabs from living layers 25-36 weeks old were used for isolation.

### Media used were:

1. Skirrow selective media for isolation and purification (Skirrow, 1977).
2. Semisolid brucella medium for maintenance. (Park *et al.*; 1984 and Mossel, 1985).
3. Triple sugar iron agar (Park *et al.*; 1984).

4. Semisolid brucella medium with cysteine for hydrogen sulphide production by lead acetate strips (Park *et al.*; 1984).
5. Semisolid brucella medium with sodium selenite for selenite reduction test (Ullmann, 1979).
6. Semisolid brucella medium with potassium nitrate for nitrate reduction test. (Park *et al.*; 1934).

**Reagents and Indicators:**

1. 3% hydrogen peroxide for catalase test.
2. Sodium hippurate solution (Sigma) and ninhydrin solution for hippurate hydrolysis test.
3. Nitrate solution (A) (0.8% sulphamic acid in 5N acetic) and nitrate solution (B) 0.5%  $\gamma$ -naphthyl amine in 5N acetic acid) for nitrate reduction test (Diem *et al.*, 1957).
4. Tetramethyl paraphenylenediamine - 2HCl for oxidase test.
5. Lead acetate test paper strips.

**Stain used is:**

Gram's Stain.

**Pathogenicity test:**

Twenty, 1 day old baby chicks and twenty, 4 weeks old chicks were used in our experiment. They were obtained from the Faculty of Agriculture Assiut University poultry farm.

**Sensitivity discs used were:**

Chloramphenicol (30Mg), Naladixic acid (30Mg), Ampicillin (10Mg), Colistin sulphate (10Mg), Streptomycin (10Mg), Gentamycin (10Mg) Erythromycin (15 Mg), Tetracyclin (30 Mg), and lincomycin (2M).

**Methods:**

**1- Isolation:**

Dead laying chickens were subjected to post-mortem examination and swabs were taken from liver, spleen, bile, blood and oviduct. Cloacal swabs from living cases were taken. These swabs were streaked on skirrow selective media and incubated at 37C for 48h. under microaerophilic conditions by use of anaerobic Jar and campy-gas pack - generating packets. Suspected colonies were subcultured in brucella semisolid media and incubated aerobically at 37C for 24h, then maintained in a refrigerator. Further identification and subculture were done weekly.

## 2- Identification of the organism:

Specimens from suspected colonies were stained by Gram stain to show typical morphology of the organism.

Biochemical reactions to study: the motility, oxidase test, catalase production, susceptibility or resistance to naladixic acid, nitrate reduction test, sodium hippurate hydrolysis and hydrogen sulphide production by using lead acetate strips and T S I media.

The differentiation between *C. jejuni*, *C. coli* and *C. laridis* is based on naladixic acid sensitivity and hippurate hydrolysis (Table 1) (Skirrow and Benjamin, 1980, Varnam and Evans, 1991).

Table 1: the differentiation between *C. species*:

<i>C. species</i>	Naladixic acid sensitivity	Hippurate hydrolysis
<i>C. jejuni</i>	Sensitive	+
<i>C. coli</i>	Sensitive	-
<i>C. laridis</i>	Resistant	-

## 3- Biotyping:

According to Ilor scheme, (1984). Table 2

Table 2: Shows biotyping scheme for *C. jejuni*:

Test	<i>C. jejuni</i>			
	I	II	III	IV
Hippurate hydrolysis	+	+	+	+
H <sub>2</sub> S production on TSI	-	-	+	+

## Pathogenicity test:

Six groups of chicks were divided as follow:

- 1st group was ten-1day old chicks inoculated orally with  $9 \times 10^7$  CFU (Ruiz – palacios *et al.*; 1981).
- 2nd group was five-1day old chicks, left in contact with the 1st group.
- 3rd group was five-1day old chicks, left as control.
- 4th group was ten-4week old chicks inoculated orally with  $9 \times 10^7$  CFU (Ruiz – palacios *et al.*; 1981).
- 5th group was five-4week old chicks left in contact with 4th group.
- 6th group was five-4week old chicks and left as control.

## Sensitivity test:

Discs were placed on brucella blood agar (without antibiotic supplement) according to Fennel *et al.*, 1984

## RESULTS

Some of the naturally infected laying chickens showed diarrhea, while P.M examination of freshly dead cases revealed distension in the intestinal tract extending to the ceca and accumulation of mucus and petechial haemorrhages present in some cases. Other cases showed mottling of the parenchyma of liver and hydropericardium.

Bacteriological examination revealed isolation of small, round, moist, non haemolytic colourless to cream coloured colonies.

By Gram stain showed curved and gull winged forms, the isolates were motile; with a characteristic corkscrew kind of movement.

Results of biochemical tests are shown in Table 3.

Table 3: Shows Biochemical tests:

Test	Isolate
1- Catalase	+ ve
2- Oxidase	+ ve
3- Nitrate reduction	+ ve
4- Hippurate hydrolysis	+ ve
5- Sodium selenite reduction	+ve
6- Hydrogen sulphide production:	
a- TSI agar	- ve
b- Lead acetate stirps	+ve

Percentage of *C. jejuni* and coli in isolated positive cases present in Table 4.

Table 4: Shows percentage of *C.jejuni* and coli in positive cases:

Visceral Samples (67)				Oviduct samples (25)			Cloacal swabs (35)				
No. of positive	%	<i>C. jejuni</i>	<i>C. coli</i>	No. of positive	%	<i>C. jejuni</i>	<i>C. coli</i>	No. of positive	%	<i>C. jejuni</i>	<i>C. coli</i>
23	34.4	18	5	3	12	3	-	11	31.4	8	3

Biotyping of *C. jejuni* according to flor scheme(1984) were Biotype I and II

### The pathogenicity test:

The 1st group showed signs of diarrhea at the third day post inoculation with mortality reached 70% at the 10th day postinoculation and the gross lesions were enlargement of liver with red and yellow mottling and congestion enlargement of the heart, distention of the intestine extending to the ceca (Fig.1) with profuse mucus.

The 2<sup>nd</sup> group showed mild signs and lesions at the 5<sup>th</sup> day of inoculation.

The 4<sup>th</sup> and 5<sup>th</sup> groups showed mild lesions compared with 1<sup>st</sup> group at 8<sup>th</sup> day post inoculation without mortality.

There were no clinical signs in 3<sup>rd</sup> and 6<sup>th</sup> groups.

Reisolation of the organism in experimentally infected chicks revealed that the *Campylobacter* could be reisolated from 1<sup>st</sup> group.

The sensitivity of *C. jejuni* against antimicrobial discs illustrated in Table 5.

Table 5: illustrates the sensitivity of *C. Jejuni* against antimicrobial discs

Types of discs	Sensitivity
Naladixic acid	+++ ve
Erythromycin	++ ve
Gentamycin	++ ve
Chloramphenicol	+ ve
Tetracycline	+ ve
Ampicillin	-ve
Colistin sulphate	- ve
Streptomycin	- ve
Lincomycin	-ve

## DISCUSSION

In our study we recovered *C. jejuni* from reproductive tract beside the internal organs and cloacal swabs this result is in agreement with that reported by Camarda *et al.*, (2000).

The present study revealed isolation of *C. jejuni* from 31.4% of examined cloacal swabs this percent is less than that recorded by Modugno *et al.*, (2000) who detected *C. jejuni* from 73% of examined Cloacal swabs in laying hens. The authors also isolated this organism from the oviducts in percentage of (29.4%), this percent is higher than that we recorded in our isolation, where we found *C. jejuni* in 12% of examined oviducts. The comparatively low percent of isolation that recorded by our result may be due to the explanation of Wegmuller *et al.*, (1993) that the method of culturing with selective enrichment may lose sensitivity because of non optimal growth conditions and the polymerase chain reaction (PCR) has been more sensitive and now the

direct Colony hybridization method is sufficiently sensitive to detect small numbers of *C. jejuni* in chickens.

*C. jejuni* biotype I and II were common, this result is similar to that reported by Modugno *et al.* (2000).

The experimentally infected 1 day old chicks showed signs of depression and diarrhea, we agreed the result of Ruiz – palacios *et al.* (1981). On the other hand we differ with them in the mortality rate where they recorded low percentage of mortality (32%) compared with 70% in our experiment.

Gross lesions in inoculated 1 day old chicks revealed distention of intestine to the two cecai with accumulation of the mucus with areas of haemorrhage and hepatic changes were present, this result is in agreement with that observed by Sanyal *et al.* (1983) and Welkos (1984).

The experimental birds kept in contact with inoculated chick revealed signs and lesions similar to that in inoculated chick but in mild picture, this result is somewhat similar to that obtained by Clark and Bueschkens (1988) where they also found beside these lesions focal hepatic necrosis. The reason that contact chicks infected from inoculated one was explained by Lindblom *et al.* (1986) and Pakamunski *et al.* (1986) that the chicks are coprophagic by nature and *C. jejuni* readily colonizes in the chick, a rapid transmission through an entire flock could be expected. Also Evans (1992) noticed that once *C. jejuni* is present in a flock, the feed, water, litter and even the air rapidly become contaminated and help to disperse the organism.

The experimentally infected 4 week old chicks showed mild signs and lesions compared with the group I, We agreed the result of Engvall *et al.*, (1986), Hoop and Ehrsam (1987), where they observed that the flock usually become infected without clinical signs when the chicks are three to five week old, but infection has been observed as early as seven days old.

The sensitivity of the isolates to different antibiotics revealed that Naladixic acid, Erythromycin and Gentamycin were the most effective drugs, this result is similar with the result reported by Das *et al.* (1996). The isolated organism was resistant to Ampicillin as reported by Erdger and Diker (1995).



## REFERENCES

- Camarda, A., Newell, D.G., Nasti, R. And Modugno, G.Di (2000):* Genotyping *Campylobacter jejuni* Strains isolated from the gut and oviduct of laying hens. *Avian Diseases* 44 : 907-912.
- Clark, A.G. and Bueschkens, D.H. (1988):* Horizontal spread of human and poultry – derived strains of *Campylobacter jejuni* among broiler Chickens held in incubators and Shipping boxes. *Journal of food Protection*, 51, 438 – 441.
- Das, S.C., Mullick, S.G. and Biswas, G. (1996):* Isolation and identification of *Campylobacter* from poultry: biotyping and in vitro antimicrobial sensitivity. *Indian Journal of veterinary Research*. 5(1) 29-34.
- Diemair, W. (1957):* *laboratorium subchfur – lebensmittel chemiker.*, 7 Auflag, verlag von theador steimkoff, leipzig
- Engvall, A., Bergqvist, A., sandstedt, K. and Danielsson-Tham, M.L. (1986):* Colonization of broilers with *Campylobacter* in conventional broiler – chicken flocks. *Acta veterinaria Scandinavica*, 27, 540-547.
- Erdger, J. and Di Ker, K.S. (1995):* Multiple antibiotic resistance in Poultry isolotes of *Campylobacter*. *Veterinar fak Itesi Dergisi Ankara universitesi, Ankara, Turkey.*
- Evans, S.J. (1992):* Introduction and spread of thermophilic *Campylobacters* in broiler flocks. *Veterinary Record*. 131,574-576.
- Fnnell, C.L., Totten, F.A., Quinn, T.C., Patton, D.I., Molmes, K.K. and Stamm,W.E.(1984):* Characterization of *Campylobacter* – like organism isolated from homosexualmen *J. Inf. Dis.* 149: 58-66.
- Hoop, R. and Ehram, H. (1987):* Ein Beitrag zur Epidemiologie von *Campylobacter jejuni* and *Campylobacter Coli* in der Huhuermast. *Schweizer Archiv fur Tierheilkinde*, 129, 193-203.
- Lindblom, G.B., Sjogren, E. and kajser, B. (1986):* Natural *Campylobacter* Colonization in chickens raised under different environmental conditions. *Journal of Hygiene*, 96,385-391.
- Lior, H. (1984):* New extended biotyping Scheme for *Campylobacter jejuni*, *Campylobacter coli* and *Campylobacter laridis*. *Journal of Clinical Microbiology*, 20,636-640.
- Modugno, G.D, Camarda, A., Circella, E. and Ficarella, V. (2000):* Epidemiological features of *Campylobacter jejuni* infection among fowls. *Selezione veterinaria* 11,978-997.

- Mossel, D.A.A. (1985):* Media for *Campylobacter jejuni* and other *Campylobacters*. *International Journal of Food Microbiology*, 2,119-122.
- Munroe, D.I., Prescott, J.F and Penner, J.L. (1983):* *Campylobacter jejuni* and *Campylobacter coli* Serotypes isolated from Chickens, Cattle and Pigs. *Journal of Clinical Microbiology*, 18,877-881.
- Pakamunski, S., Kass, N., Borochovich, E., Marantz, B. and Rogol, M. (1986):* Incidence of *Campylobacter* Spp. In broiler flocks monitored from hatching to slaughter. *Avian Pathology*, 15,83-92.
- Park, C.E., Smibert, R.M., Bluser, M.J. Vanderzant, C. and Stern, N.J. (1984):* *Campylobacter* in: " Compendium of methods for the microbiological examination of foods" 2nd Ed. Speck, M. I. (ed) American Public Health Association Washington, D.C.
- Ruiz-Palacios, G.M., Escamilla, E. and Torres, N. (1981):* Experimental *Campylobacter* diarrhea in chickens. *Infection and Immunity*, 34,250-255.
- Sanyal, S.C., Neogi, P.K.B., Islam, K.M.N. and Huq, M.I. (1983):* Pathogenesis of *Campylobacter jejuni* diarrhoea in an experimental model in: Pearson, A. D., Skirrow, M. B., Rowe, B., Davics, J. R. and Jones, D. M. (eds) *Campylobacter II* P. 122 (London, Public Health Laboratory Service).
- Sebald, M., and M. Veron (1963):* Teneur en bases de l'ADN et Classification des vitriens. *Ann Inst Pasteur (Paris)* 105:897-910.
- Shane, S.M. (1992):* The significance of *Campylobacter jejuni* infection in poultry: a review. *Avian pathology*. 21: 189-213
- Simmons, N.A. and F.J. Gibbs (1979):* *Campylobacter* Spp. in oven-ready Poultry. *Journal Infection*. 1:159-162.
- Skirrow, M.B. (1977):* *Campylobacter* enteritis : a "new" disease. *British medical Journal* 2,9-11
- Skirrow, M.B. (1990):* *Campylobacter* *Lancet*. 336, 921-923.
- Skirrow, M.B. and Benjamin, J. (1980):* "1001" *Campylobacters* : Cultural characteristics of intestinal *Campylobacters* from man and animals. *Journal of Hygiene*, 85,427-442.
- Ullmann, U. (1979):* Methods in *compyloacter*, In " Methods in Microbiology" 13th Ed. Bergan, T. and Norris, J.R. Academic Press, New York: 435- 452.

*Varnam, A.H. and Evans, M.G. (1991):* Campylobacter in : " food borne Pathogens" Wolfe Publishing Ltd. London England.

*Wegmuller, B., Luthy, J. and candrian U. (1993):* Applied and Environmental Microbiology. 59:2161-2165.

*Welkos, S.L. (1984):* Experimental gastroenteritis in newlyhatched chicks infected with Campylobacter jejuni. Journal Medical Microbiology 18:233-248.



**Fig. 1:**  
**Distension of the intestinal tract in experimentally infected chick  
by C. jejuni**