

Animal Health Research Institute,
Dokki, Egypt.

ROLE OF IBIS IN TRANSMITTING REOVIRUS INFECTION AMONG POULTRY FARMS

(With 6 Figures)

By

AHLAM, A. FARGHLI and LEILA, A. TANTAWY

(Received at 8/5/2004)

فيروس الريو في أبو قردان

أهلام عبد الحميد فرغلي ، ليلى طنطاوي

تم فحص عدد ١٣ عينة أظهرت ثلاثة منها ورم في الركبة، ووجد في حالة واحدة بطش حمراء في البنكرياس. بالفحص الفيروسي تم عزل فيروس الريو. تم العزل في بيض مخضب عن طريق حقن البيض بالحقن على الغشاء. ظهرت بثرات بيضاء على الغشاء. وتجمعات نووية داخل الخلايا على الزرع النسيجي. الفيروس مقاوم للتكوير وفورم والاشعير ودرجة حرارة ٥٦ م لمدة ساعة. تم حقن ٢٠ كتكوت تحت جلد القدم عمر ٣ أسابيع وتم العزل والفحص الباثولوجي لهذه الكتاكيت وظهر ورم في الرجل بعد ٤،٤،٣، ١٢، ١٥ يوم بعد الحقن. بالفحص الباثولوجي أظهر عتامة في الغشاء المحيط بالقلب والتهابات في عضلة القلب. والفحص الهستوباثولوجي تتركز في خلايا الكبد وتفسيرات في البنكرياس والغدة الهضمية.

SUMMARY

Out of thirteen Ibis examined, 3 cases showed swelling of the digital flexor and metatarsal extensor tendon, and one case has red patches on the surface of pancreas. Virological examination revealed reovirus resembles that in cage bird. Experimental infection of 3 weeks old 20 chickens through foot pad inoculation was done. Histopathological and virological examination of the inoculated chickens were described. The results revealed diarrhoea at 3rd day PI and swollen legs at 4th, 12th and 15th day PI. Isolation by egg inoculation (CAM) showed pock lesions in one case and tissue culture (vera) revealed syncytia. Virus resisted ether, chloroform and it was heat stable. Pathological lesions were pericarditis, myositis, hepatic vacuolar degeneration and necrosis in addition to lesions in pancreas and proventriculus.

Key words: Ibis, reovirus, poultry farms.

INTRODUCTION

Ibis in Egypt has a close relation with agricultural land and farmer, it is called farmer's friend. Giza zoo is considered as a permanent habitat for it. So research works are needed to know the role of Ibis as a carrier or reservoir of infection for domestic poultry and zoo birds. Soad and Wafaa (2003) isolated some bacterial microorganisms (*Pseudomonas aeruginosa*, *E. coli*, *Salmonella enterica typhimurium*, *Proteus vulgaris* and *P. haemolytica*) from Ibis.

Avian Reoviruses are members of the Orthoreovirus genus in the Reoviridae family. They can be differentiated in commercial poultry by antigenic configuration, pathotype, relative pathogenicity, growth in cell culture, sensitivity to trypsin and host specificity (Jones *et al.*, 1994) and (Jones and Georgian, 1984). Concerning the economic importance of Reovirus, Pass *et al.* (1982) reported the economic significance of viral arthritis (VA) and malabsorption syndrome (MAS) in broiler breeder pullets and hens.

In young meat type chickens, economic losses related to reovirus infections are frequently associated with increased mortality, viral arthritis/tenosynovitis (Rosenberg and Olson, 1997), loss in weight gain, poor feed conversion, uneven growth rates and reduced marketability of affected birds (Dobson and Gilsson, 1992). Reoviruses are prevalent world wide in chickens, turkeys and other avian species. Viral arthritis is observed primarily in meat type chickens but can be found in lighter breeder (Jones and Onunkwo, 1978 and Schwartz *et al.*, 1976) and in turkeys (Alafaleg and Jones, 1989, Back and Nagaraja, 1996 and Nersessian *et al.*, 1986) in the digestive and respiratory tract of clinically normal chickens and turkeys Kawamura *et al.* (1965) and Wooley *et al.* (1972). Van der Heide (1996) estimated that greater than 80% of reoviruses isolated from chickens were apathogenic. Reovirus can be isolated also from pigeons (McFerran *et al.*, 1976) and from diseased pigeons, pheasants, parrots and exotic avian species (Gough *et al.*, 1988). Reovirus was also isolated from the liver of an African grey parrot (Graham, 1987), from American woodcock (Docherty *et al.*, 1994), from Bob white quail (Ritter *et al.*, 1986) and from pheasants (Mutlu *et al.*, 1998). Cadman *et al.* (1994) detected antibodies to avian reovirus by ELISA in Ostriches from Zimbabwe. The purpose of this study was to investigate the epidemiology of reovirus in Ibis and the probability of being a reservoir and carrier of the disease.

MATERIALS and METHODS

1- History and PM examination:

Thirteen Ibis Farm Abu Rawash, Giza. were examined, 3 cases only showed swelling of the digital flexor and metatarsal extensor tendon. One case has red patches on the pancreas.

2- Egg inoculation:

Collected pancreas, liver and synovial fluid of each case was inoculated in 5 eggs per each 10 day old embryonated chicken eggs via chorioallantoic membrane.

3- Agar gel precipitodion test:

Harvested and allantoic fluid were added. Frozen thawed 3 times and put in agar gel made by 1.2 purified agar in phosphate buffer pH 7.2. Reference sera which taken from Dr. Attia Sami, Virology Department Cairo, Univ. put in the center allantoic membrane in prepheral well and examined for 3 days for precipitation line.

4- Tissue Culture:

Vero cell line were inoculated with inoculum (synovial fluid and pancreatic emulsion) and examined for 5 days for cytopathic effect and syncytia.

5- Heat stability:

Viral isolates were divided into 5 tubes each then incubated in water bath at 56°C for 5, 10, 15, 30 min. and 1 hour. Control tube kept in room temperature. Treated and control samples were checked for infectivity in tissue culture.

6- Chemical test:

a- Ether sensitivity:

It was carried according to the method described by Andraws and Horsman (1949). Two ml of the virus was prewarmed in water bath 37°C and 0.8 ml of chemically pure diethylether was added. The tube was agellated and incubated at 37°C for 30 minutes then the aqueous layer was collected for the residual virus to be titrated in tissue culture.

b- Chloroform sensitivity:

The technique used was that described by Feldman and Wang (1967). One part of chemical purified chloroform was fixed with 19 parts of tissue culture containing the virus to have final dilution of 5% chloroform. The mixture was left at room temperature for 10 min. with continuous vigorous shaking. The mixture was centrifuged at 3000rpm for 5 min. and aqueous layer was collected and the residual virus was collected and titrated in tissue culture.

7- Virus neutralization:

Virus was titrated 4 well per each dilution 0.0025 serum (Virology Department, Cairo University), the same amount was added to each well which contained 0.0025 of virus. Incubation for 1 hour and MEM was added 0.050 in each well (alphaprocedure). The difference between virus titration only and virus serum titration was calculated as neutralization index.

8- Experimental design:

Three weeks old 20 chickens were inoculated via foot pad, other 8 chickens kept as control. Four sacrificed birds were examined at 2,4,12 and 15 days post inoculation (PI) for:

- 1- Postmortem examination.
- 2- Diagnosis by agar gel precipitation test.
- 3- Histopathological examination.

9- Histopathological examination:

Samples from tendon, pancreas, liver and heart were fixed in 10% neutral buffered formalin. The specimens were treated chemically then sectioned at 5µm and stained with haematoxylin and Eosin (Culling 1963).

RESULTS

Viral isolation:

One case of Ibis from three cases of swelling legs showed whitish pock lesions on CAM. Death of embryos occurred within 3-5 day PI. One line of precipitation appeared against the positive serum by agar gel.

Tissue culture:

In case of the first passage, cell rounding occurred. By the second passage syncytia appeared. Third passage showed necrobiotic changes.

Virus titration	Ether	Chloroform
$10^{-4.75}$	$10^{-4.75}$	$10^{-4.75}$

Thermostability:

Virus was stable at 56°C for 1 hour.

Neutralization index:

Index	2
-------	---

Clinical signs & Post-mortem findings:

At 2nd day PI one dead bird showed diarrhoea, at 4th day PI 3 birds showed tenosynovitis, swollen legs (Fig. 6) with greenish diarrhoea appeared at 12 and 15 day PI. Liver appeared pale at 3rd day PI, other cases showed congestion at 4th day PI.

Virus reisolation:

100% of pancreas was positive until 9th day PI but 25% of liver was positive.

3-Histopathological findings:

At 2nd and 4th day PI, the liver showed severe congested blood vessels (Fig. 1) in addition to severe vacuolar degeneration (Fig. 2) of the hepatocytes. Focal coagulative necrosis was observed. The proventriculus showed hyperplasia and metaplasia of the lining epithelium with cellular degenerative changes. While the heart at 2nd day PI revealed focal pericardial and myocardial oedema with degeneration in the myocardial muscle fibers. Heart lesions became severe at 4th day PI as pericarditis (Fig. 3), pericardial oedema with myositis and heterophilic infiltration between the myocardial fibers (Fig. 4). At the 15 day PI the lesions became less in heart, liver and proventriculus while the pancreas showed vacuolar degeneration of the acinar cells of the exocrine tissue (Fig. 5) in addition to cellular necrotic debris.

DISCUSSION

Reovirus is very common among domestic poultry and other avian species. Inoculum was prepared from synovial fluid, pancreas and liver of infected Ibis and inoculated in embryonated chicken eggs via chorioallantoic membrane. Pock lesions were developed in addition to embryo mortality within 3-5 day PI and stunting of survival embryo. These results agreed with Mustaffa and Spradbrow (1971) and Bains, *et al.* (1974).

The infected CAM homogenate were tested by AGPT using Reovirus S1133 antiserum. Isolates that produced pathological lesions in chicken and gave precipitation line were propagated for the passage in vero cell. Syncytia formation followed by detachment from monolayer cell sheet. The choice of vero cells for reovirus isolation was manipulated with Sahu and Olson (1975), Tantawi *et al.* (1984), Shapouri *et al.* (1996) and Samah (1997).

Reovirus in Ibis resemble that in cage bird. Trials of reisolation proved that the best organ for reisolation is the pancreas.

Symptoms and postmortem lesions resembled that described by Sterner (1989) and Van der heid (1996). The histopathological lesions involved myocarditis and heterophilic infiltration between the myocardial fibers as mentioned by Kerr and Olson (1969) and Olson and Solomon (1968). Lesions in liver as vacuolar degeneration and necrosis were also observed by Gourvca and Schnitzer (1982).

These lesions might be due to the replication of the virus in the cytoplasm consuming the cellular elements and cause irritation and cell damage.

In conclusion, Ibis could be a carrier and a reservoir for avian reovirus, so control measures are needed to prevent spreading of infection among poultry farms and zoo birds.

REFERENCES

- Al Afaleq, A. and Jones, R.C. (1989):* Pathogenicity of three turkeys and three chickens reoviruses with particular reference to arthritis/tenosynovitis. *Avian Pathol* 18:433-440.
- Andrews and Horstman (1949):* The susceptibility of viruses to ethyl ether. *J.G. Microbiol* 3, 29,297.
- Back, A. and Nagaraja, K.V. (1996):* Pathogenicity of chicken reovirus strain S3311 and 1733 for turkeys. *Proc.international symposium on Adeno virus and reovirus infection in poultry. Rauschholzhausen, Germany, 245-250.*
- Bains, B.S.; Mackenzi, M. and spradbrow (1974):* Reovirus associated mortality in broiler chickens in victoria. *Avian Dis.*18: 472-476.
- Cadman, H.F.; Kelly, P.J.; Zhou, R.; Davelaar, F. and Mason, P.R. (1994):* A sero-survey using enzym linked immunosorbant assay for antibodies against poultry pathogen in Ostrich (*Struthio Camelus*) from zimbabwe. *Avian Dis.* 38: 621-625.
- Culling, C.F.A. (1963):* Handbook of histopathological techniques. 2nd ed., Butterworths, London.
- Dobson, K.N. and Glisson, J.R. (1992):* Economic impact of a documented case of reovirus infection in broiler breeders. *Avian Dis.* 36: 788-791.
- Docherty, D.E.K.A.; Converse, W.R.; Hansen and Norman, G.W. (1994):* American Woodcock (*scelopox minor*) mortality associated with a reovirus. *Avian Dis.* 38: 899-904.
- Feldman and Stephan Wong, D.S. (1961):* Sensitivity of various viruses to chloroform. *Proc., Sci, Exp. Biol.* 106,136.

- Gough, R.E.; Alexander, D.J.; Collins, M.S.; Lister, S.A. and Cox, W.J. (1988): Routine virus isolation or detection in the diagnosis of diseases in birds. Avian Pathol. 17:893-907.*
- Gouvea, V. and Schmitzer, T.J. (1982): Pathogenicity of avian reoviruses: examination of six isolates and a vaccine strain. Infect. Immun 38: 731-738.*
- Graham, D.L. (1987): Characterisation of avian reo like virus from and pathogenicity for parrots. Avian Dis. 31: 411-419.*
- Jones, R.C. and Omunkwo, O. (1978): Studies on experimental tenosynovitis in light hybrid chickens. Avian Pathol. 7:171-181.*
- Jones, R.C. and Georgiou, K. (1984): Reovirus induced tenosynovitis in chickens: The influence of age in infection. Avian Pathol. 13: 441-457.*
- Jones, R.C.; Afaleq, A.I.; Savage, C.E. and Islam, M.R. (1994): Early pathogenesis in chicks of infection with a trypsin sensitive avian reovirus. Avian Pathol 23: 683-692.*
- Kamamura, H.F.; Shimizu, M.; Maeda and Tsubahara, H. (1965): Avian reovirus: its properties and serological classification. Natl Inst Anim Health Q (Tokyo) 5:115-124.*
- Kerr, K.M. and Olson, N.O. (1969): Pathology of chickens experimentally inoculated or contact infected with an arthritis producing virus. Avian Dis. 13: 729-745.*
- Mc Ferran, J.B.; Connor, T.J. and MackCraken, R.M. (1976): Isolation of reoviruses and Adenoviruses from avian species other than domestic fowl. Avian Dis. 39: 32-745.537.*
- Mustafa Bajee, and Spradbrow, P.B. (1971): The isolation avian Reovirus. Aust. Vet. J. 47-284.*
- Mutlu, O.F.; Grund, C. and Cuvén, F. (1998): Reovirus infections of pheasants (*Pheasianus Colchicus*), Tierärztliche Praxis Ausgabe Grobtiere Nutztiere 26: 104-107.*
- Nersissian, B.N.; Goodwin, M.A.; Rage, R.K.; Kleven, S.H. and Brown, J. (1986): Studies on orthoreovirus isolated from young turkeys. III Pathogenic effect on chicken embryos, chicks poults and suckling mice. Avian Dis. 30: 585-592.*
- Olson, N.O. and Solomon, D.P. (1968): A natural outbreak of synovitis caused by the viral arthritis agent, Avian Dis. 12: 311-316.*
- Pass, D.A.; Robertone, D.M. and Wilcox, G.E. (1982): Runting syndrome in broiler chicken in Australia. Vet. Rec. 110, 386-387.*

- Ritter, D.G.; Levy, D.H.; Guy, J.J. and Barnes, H.J. (1986): Intestinal cryptosporidiosis and reovirus isolation from bobwhite quail (*colinus virginianus*) with enteritis. *Avian Dis.* 30: 603-608.
- Rosenberg, J.K. and Olson, N.O. (1997): Viral arthritis. In B.W. nek, H.J. Barnes, C.W. Beard, I, R. McDougald, and Y.M.Saif (eds). *Diseases of poultry* 10th ed. Iowa state University press; Ames, J.A, 711-718.
- Sahu, S.P. and Olson, N.O. (1975): Comparison of the characteristics of avian reoviruses isolated from the digestive and respiratory tract, with viruses isolated from the synovia. *Am.J. Vet. Res.* 36: 847-850.
- Samah, H.M. (1997): Studies on avian reoviruses in Egypt. M.V.Sc. Thesis, Fac. of Vet. Med. Benisuef branch. Cairo University.
- Schwartz, L.D.; Gentry, R.F.; Rothenbacher, H. and Van der Heid, L. (1976): Infectious tenosynovitis in commercial white leghorn chickens. *Avian Dis.* 20: 769-773.
- Shapouri, M.R.S.; Freneth Larochalle, R.; Azella, M. and Silim, A. (1996): Characterisation of monoclonal antibodies against avian reovirus strain S1133. *Avian Pathol.*, 25: (1), 57.
- Soad, A.; Nasif and Wafee, M.M. Hassan (2003): The role of Ibis in transmission of avian bacterial infection. *Egypt. Vet. Med. Asso.* 63: 159-163.
- Sterner, F.J.; Rosenberger, J.K.; Margolin, A. and Ruff, M.D. (1989): In vitro and in vivo characterisation of avian reovirus. II clinical evaluation of chickens infected with two reovirus pathotypes. *Avian Dis.* 22: 545-554.
- Tantawi, H.H.; Amina, N.; Youssef, Y.I.; Fawzia, Al-Abdulla, J.M.; El-Batravi, A.; El Ghawas, A.; Nasser, A.A. and Reda, I.M. (1984): Infectious tenosynovitis in broilers and broiler breeders in Egypt. *Vet. Res. Communication*, 8, 229-235.
- Van der Heid, L. (1996): Introduction on avian reovirus. *Proc. international symposium on Adeno virus and reo virus infection in poultry*, Rauischholzhausen, Germany, 138-142.
- Wooley, R.E.; Dees, T.A.; Cromack, A.S. and Gratzek, J.B. (1972): Infectious enteritis in turkeys: Characterisation of two reoviruses isolated by sucrose density gradient centrifugation from turkeys with infectious enteritis. In proceeding 5th scientific conference, Egypt. *Am J.Vet. Res.* 33: 157-164.




Fig. 1: Liver of chicken at 2nd day PI showed congested blood vessels (H&E x 100).




Fig. 2: Liver of chicken at 2nd day PI showed vacuolar degeneration and necrosis (H&E x 250).




Fig. 3: Heart of chicken at 4th day PI showed pericarditis (H&E x 250).



Fig. 4: Heart of chicken at 4th day PI showed myositis with heterophilic infiltration (H&E x 250).



Fig. 5: Pancreas of chicken at 15 day PI showed vacuolar degeneration of acinar cells (H&E x 400).




Fig. 6: Leg of inoculated chicken at 4th day PI swelling.