

## Role of Eosinophils Apoptosis and Phagocytosis in Sinonasal Polyps: An Immunohistochemical and Ultrastructural Study

Ahmed H. Abdel-Rahman\* – Abdel- Salam Hashiem\*\* – Ahmed S.Gad\*\* -  
Ahmed Moh. Nassar\*\* – Ahmed Abdel – Basset\*\* – Aly Abdalla\*\* –  
Atif I. M. Ali\*\*\*

From Departments of Pathology\*, Otorhinolaryngology\*\* and Histology\*\*\* Al-Azhar University (Assuit & Cairo).

### Abstract

**Background:** Inflammatory cells disappear from airway tissue through apoptosis and phagocytosis. However, the cells may also be cleared through primary cytolysis, necrosis secondary to apoptosis or transepithelial migration.

**Material and Methods:** This study evaluated the role of apoptosis and phagocytosis of tissue eosinophils in the sinonasal polyps received from fifty patients. Immunohistochemically, the amount of CD 95 — and CD 68 – positive cells was determined by counting the average number in ten randomly chosen high power fields. The inflammatory infiltrate consisted of numerous eosinophils as well as a considerable amount of lymphocytes and macrophages. Also, we detected a frequent expression of CD 95 on the eosinophils and numerous other inflammatory cells.

**Results:** CD 68 – positive cells (most probably macrophages) approximately equalled the number of eosinophils. Ultrastructurally, seventy three percent of the examined eosinophils (365 / 500) were apoptotic; about one third of which (89 / 365, 24.3 %) were secondarily necrosed. However, only four percent (20 / 500) were cytolytic. The phagocytic action of macrophages engulfing eosinophils was confirmed by electron microscopy.

**Conclusion:** Thus, our study concluded the elimination of tissue eosinophils predominantly through apoptosis and phagocytosis that might be important for clearance of eosinophils from the respiratory tissue and could improve the allergic manifestations.

**Key Words:** Sinonasal polyps – Apoptosis – Phagocytosis – Eosinophils – Macrophages – CD 95 (Fas receptor) – CD68 – Immunohistochemistry – Electron Microscopy.

### Introduction

Sinonasal polyps are non-neoplastic swellings usually arising from the ethmoid sinuses and prolapse into the nasal cavity. Previous studies had been concerned the etiology, the relation to allergy, histopathology and treatment of these lesions (Hellquist, 1996), however, a little attention about apoptosis and resolution of the eosinophilic inflammatory infiltrate in these polyps had been performed (Davidsson *et al.*, 2000).

The polyps characteristically contain a leucocyte infiltrate in which eosinophils are often remarkably prominent (Jahnsen *et al.*, 1995). The later cells are bone marrow derived granulocytes with a richness of surface receptors for immunologic ligands

(Rothenberg, 1998). They harbor specific granules which, beside chemokines, cytokines, and growth factors, contain large amounts of tissue – toxic cationic proteins namely major basic protein, eosinophil – derived neurotoxin, eosinophil cationic protein (ECP) and eosinophil peroxidase (Martin *et al.*, 1996). Moreover, eosinophils produce tumor necrosis factor – alpha (TNF –  $\alpha$ ), transforming growth factor – alpha (TGF –  $\alpha$ ) and transforming growth factor – beta 1 (TGF –  $\beta$ 1). Thus, these cells actively contribute to mucosal inflammation and may cause several morphological changes occurring in sinonasal polyps. The demonstration of mRNA for interleukin (IL) -6 & IL - 12 can

imply T-helper 1 cell response ( Davidsson *et al.*, 1996). Endothelial adhesion molecules particularly vascular cell adhesion molecule - 1 (VCAM -1) and P-selectin are thought to play a crucial role in human eosinophil extravasation at inflammatory sites and nasal polyps ( Karlsson and Hellquist, 1996, Desreumaux and Capron, 1996 & Westergren *et al.*, 1997 ). In addition, eosinophils abounding and degranulation in airway mucosal and pulmonary tissues are believed to have pathogenic roles in asthma, nasal polyposis, allergic rhinitis and eosinophilic pneumonia ( Erjefalt and Persson, 2000, Erjefalt *et al.*, 2001 & Emanuelsson *et al.*, 2004 ). During active seasonal allergic rhinitis eosinophils may exhibit a more pronounced degranulation than in many other common diseases characterized by tissue eosinophilia e.g. inflammatory bowel disease and asthma ( Erjefalt *et al.*, 1998 & Erjefalt *et al.*, 2001 ). However, a little information is known about the dynamics of eosinophilic degranulation in allergic rhinitis especially when the disease progresses from an asymptomatic baseline situation into an active symptomatic condition ( Emanuelsson *et al.*, 2004 ).

Infiltration of leucocytes in diseased airway tissues is counterbalanced by their elimination through apoptosis (Woolley *et al.*, 1996 ) and prompt engulfment by macrophages ( Haslett, 1999 & Henson *et al.*, 2001 ). In vitro study has generated a detailed information regarding molecular and pharmacologic regulation of apoptosis (Nutku *et al.*, 2003). Human purified blood eosinophils seem to be prone to massive apoptosis particularly when cultured in the absence of specific growth factors or in the presence of glucocorticoids in the cell medium ( Gardai *et al.*, 2003 & Bochner, 2004 ). Steroid administration in vitro has been shown to significantly inhibit the survival rate of eosinophils from patients with atopic dermatitis and has suggested that steroid - induced apoptosis decreases the eosinophil count, in vivo, in these patients (Matsukura *et al.*, 1997 ).

Further support for the role of apoptosis is provided by demonstration of Fas-receptor ( CD 95 ; APO -1 ) on lung eosinophils ( Tsuyuki *et al.*, 1995 ) and by

the delay of neutrophils apoptosis mediated by interleukin-6 ( Biffi *et al.*, 1996 ).

Moreover, the production in nasal polyps of ILs-3, -5 & -8 as well as granulocyte -macrophage colony - stimulating factor ( GM - CSF ) & TNF -  $\infty$  indicates a possible involvement of these cytokines in the promotion of eosinophil survival probably by preventing or delaying apoptosis ( Bachert *et al.*, 1997 ). While the biological activities of IL-3 and GM - CSF extend to affect several cell types, IL-5 acts specifically on eosinophils to promote their maturation, activation and survival (Pretolani and Goldman, 1997 & Till *et al.*, 1997 ).

Monocytes and macrophages are certainly different regarding the induction of apoptosis; monocytes spontaneously undergo apoptosis after cytokine withdrawal whileas macrophages have no great potential to undergo apoptosis (Davidsson *et al.*, 2000 ). The later cells from different sources and at various stages of activation may change in their ability to engulf apoptotic cells suggesting the possibility that the dead cells would not be immediately eradicated from all tissues (Uller *et al.*, 2004). Furthermore, the occurrence of apoptotic eosinophils and neutrophils has been reported in human skin indicating that the apoptotic cells in vivo should be well detectible by careful histologic examination of tissue (Ying *et al.*, 1997 & Erjefalt and Persson, 2000 ).

A clear identification of morphologic characteristics, achievable exclusively by transmission electron microscopy (TEM) analysis, is considered crucial for the assessment of occurrence of apoptosis of cells in vivo by specifically appearance of eosinophil cell debris within the macrophages as would be the case if apoptotic eosinophils had been engulfed in vivo ( Stadelmann *et al.*, 2000 & Uller *et al.*, 2004 ). In addition, TEM analysis further involves the detection of other features of the tissue eosinophils such as occurrence of cytolysis as well as necrosis secondary to apoptosis ( Erjefalt *et al.*, 1999 & Uller *et al.*, 2004 ).

We, thus, aimed to investigate the role of apoptosis and phagocytosis of tissue eosinophils in the sinonasal polyps.

## Material and methods

The material of this work is consisted of fifty sinonasal polyps obtained by routine polypectomy from Otorhinolaryngology Department, Al-Husseini University Hospital during the period from August 2006 to August 2008. All patients were males with a ges ranged from 28 to 51 years (mean,  $42.8 \pm 1.7$ ). All patients had received a local steroid treatment.

Small pieces of the specimens were used for ultrastructural study while the remaining tissue specimens were immediately placed in 4% neutral buffered formaldehyde and cut up within the next 24 hours. The following techniques were carried out:

### 1) Routine histopathologic technique:

Formalin fixed and paraffin embedded blocks were sectioned at 6  $\mu$ m thickness and examined microscopically using hematoxylin and eosin (Hx & E) stain to ensure the integrity of specimens before the immunoperoxidase technique and to confirm the clinical diagnosis.

### 2) Immunoperoxidase (IP) technique:

According to Hellquist *et al.*, (1997) & Sundelin *et al.*, (1997) the sections were deparaffinized, hydrated and incubated with 3% H<sub>2</sub>O<sub>2</sub> for 5 minutes to block the endogenous peroxidase activity and then immersed in boiling citrate buffer (PH6) in a microwave oven, twice, for 5 minutes (600 W). After rinsing in Tris buffered saline (TBS), the sections were incubated with monoclonal primary antibodies for CD 95 (Anti - Fas, Clone UB-2, Oncor, Gaithersburg, MD) at a dilution of 1:25 and for CD68 (Anti-CD68, Clone PG - M1, Dako, Copenhagen, Denmark) at a dilution of 1:150 for 1 hour at room temperature. Then after rinsing in distilled water & TBS buffer, secondary antimouse antibody using peroxidase labelled bition streptavidin complex detection system (Strept AB complex / AP, Dako, Copenhagen, Denmark) for CD 95 and CD 68 was applied. Diaminobenzidine was added as a chromogen. Then, the sections were

counterstained with hematoxylin and coversliped.

### Immunohistochemical analysis:

CD 68 - positive cells (most probably macrophages) and CD 95 - positive eosinophils count were evaluated by counting their average numbers in ten randomly selected high power fields (HPF; X 400 magnification) per polyp. The average number of either positive cells was scored as follows; 0: negative, 1+: < 5 positive cells, 2+: 5 - < 20, and 3+:  $\geq 20$  cells per field. CD68 immunopositivity was detected as a brownish nuclear staining while CD95 immunoreactivity appeared as a brownish membranous and cytoplasmic signalling.

### 3) Transmission electron microscopy (TEM):

According to Uller *et al.* (2001) and Erjefalt *et al.* (2003), samples for TEM study were prepared. Small pieces of biopsies were put in 2.5% gluteraldehyde, then in cacodylate buffer for 20 minutes and fixed in a fume cupboard with osmium tetroxide for 2 hours at 4°C. The specimens were dehydrated, cleared in propylene oxide for 30 minutes at room temperature and embedded in Beem capsule filled completely with epoxyresin and polymerized at 60°C for 24 hours. The capsule was then trimmed and sectioned at 1  $\mu$ m (Semithin section) and 90 nm (Ultrathin sections). The sections were placed on a 200 - mesh, thin - bar, copper grid and contrasted by staining in uranyl acetate (4% in distilled water for 30 minutes at 4°C) and with lead citrate (0.5% 1M sodium hydroxide for 3 minutes at room temperature). The ultrastructural analysis was carried out using a Philips CM10 TEM (Philips, Eindhoven, Netherlands).

### Ultrastructural analysis:

The biopsies that contained sufficient numbers of eosinophils and had a well - preserved morphology without mechanical artifacts were thirty five out of the fifty cases (35 / 50, 70%). In these appropriate biopsies, each individual eosinophil was carefully analysed at x

4,000 – x 12,000 magnification . In total , we examined 500 eosinophils .

**Results**

All biopsies displayed a typical morphology in the routinely stained sections including oedema , excessive inflammatory cell infiltrate , basement membrane thickening and goblet cell proliferation ( Fig . 1 ) . The inflammatory infiltrate consisted of numerous eosinophils as well as considerable number of lymphocytes and macrophages that are almost entirely found towards the periphery of the polyps (Fig.2 )

Fas receptor ( CD 95 ) - positive eosinophils were found in most cases ( 46 out of 50 ; 92 % ) . Regarding the average number of CD 95 + ve eosinophils , 29 cases ( 29 / 50 ; 58 % ) were of grade 3 + (Fig . 3 ) , 8 ( 16 % ) were of grade 2 + and 9 ( 18 % ) were of grade 1 + ( Fig 4 ) . The remaining cases ( 4 / 50 ; 8 % ) didn't express CD95 on eosinophils ( Table 1 ) . In addition , CD95 molecule was expressed on numerous other inflammatory cells which had not been counted in this study .

CD 68 - positive cells approximately equalled the number of eosinophils . They are detected in all cases although their incidence differed from one polyp to another . Their average number was graded as 3 + in 28 ( 56 % ; Fig . 5 ) , 2 + in 12 ( 24% ) and 1 + in 10 ( 20 % ; Fig . 6 ) cases ( Table 2 ) . In most cases ( 46 out of 50 ;

92 % ) , CD 95 - positive eosinophils & CD 95 + ve other inflammatory cells appeared more frequent towards the periphery compared to the central part of the polyp ( Fig . 7 ) . The immunohistochemical results , regarding the susceptible CD 95 positive eosinophils for programmed cell death , were confirmed by their apoptotic ultrastructural morphology ( Fig . 8A-D ) . Eosinophils ultrastructurally were distributed in the epithelium and subepithelial layer of the polyp . Four percent of TEM – examined tissue eosinophils ( 20 / 500 ) were cytolytic as detected by incomplete chromatolysis , cell membrane disruption and partly dissolved cytoplasm with spilling of protein – containing free granules . On the other hand , seventy three percent ( 365 / 500 ) of the tissue eosinophils were clearly apoptotic with appearance of electron – dense granules , condensed chromatin , nuclear bleb formation , preserved plasma membrane and non – dilated organelles . Secondary necrosis , as a final stage of apoptosis , was encountered in 24 . 3 % of the apoptotic eosinophils ( 89 / 365 ) and was distinguished from cytolysis as shown by a condensed dark nucleus as well as plasma and organellar membranes rupture . Thus , most of tissue eosinophils were died by apoptosis rather than by cytolysis ( Table 3 ) . In addition , macrophages exhibited signs of engulfment of cell material definitely eosinophils as shown by their specific granules

**Table 1 : Immunoreactivity of eosinophils for CD 95 in studied polyps ( n = 50):**

CD95 - positive eosinophils			CD95 - negative eosinophils		
No , of cases	%	grade	No , of cases	%	grade
29	58	3 +	4	8	0
8	16	2 +			
9	18	1 +			
Total= 46	92		4	8	

**Table 2 : Immunoreactivity of macrophages for CD 68 in studied polyps ( n = 50) :**

CD95 - positive cells			CD95 - negative cells		
No , of cases	%	grade	No , of cases	%	grade
28	56	3 +	0	0	0
12	24	2 +			
10	20	1 +			
Total= 50	100		0	0	

**Table 3 : Ultrastructural findings in TEM – examined eosinophils (n = 50):**

Finding	No ; of cells	%
Cytolysis	20	4
Apoptosis only	276	48.7
Apoptosis with secondarynecrosis	89	24.3
	} 365	} 73



**Fig 1 : A case of sinonasal polyp showing oedema , excessive inflammatory cell infiltrate & basement membrane thickening . Fig 2 : A sinonasal polyp showing an inflammatory infiltrate almost found towards the periphery ( H x & E x 200 ) .**





Fig 3 : A sinonasal polyp showing grade 3 CD95 - positive cells ( brownish cytoplasmic staining ) .

Fig 4 : A sinonasal polyp showing grade 1 CD95 - positive cells ( brownish cytoplasmic staining ) .

Fig 5 : A sinonasal polyp showing grade 3 CD68 - positive macrophages ( brownish nuclear staining ) .

Fig 6 : A case of sinonasal polyp showing grade 1 CD68 - positive macrophages . (Immunoperoxidase [ IP ] x 400 ) .

Fig 7 : A case of sinonasal polyp showing CD95 - positive inflammatory cells more frequently towards the periphery than in the central part of the lesion ( IP x 100 ) .



**Fig . 8 : Electron micrographs showing cytolitic eosinophils detected by incomplete chromatolysis of the nucleus , disrupted cell membrane and spilling of protein – containing granules ( SP ) ; 8A . Apototic eosinophils revealed condensed chromatin ( apoptotic body ; Ab ; 8A-D ) and disrupted cell membranes ( CM ; blue arrow head ; 8B & C ) . Secundarily necrotic eosinophils displayed a condensed dark nucleus and damaged cell membrane ( CM ; 8A-C ) . The apoptotic eosinophils revealed electron – dense granules ( G ; 8E ) .  
( Double fixation & Double stain : 8A-D x 10,000 and 8E x 12,000 ) .**



## Discussion

Sinonasal polyp is an upper airway inflammatory disease in which the structural modifications of epithelium (secretory hyperplasia and squamous metaplasia) and lamina propria (basement membrane thickening, extracellular matrix accumulation and collagen deposition) are associated with inflammatory cell infiltration (Molet *et al.*, 2003).

The uniform handling of the tissue and the immediate fixation minimized the risk that the number of cells which might enter apoptosis after polypectomy would differ between cases. Compared to electrophoresis that demonstrates the apoptotic process after its point of no return (180 – bp fragmentation), terminal – d – UTP nick end labeling assay (TUNEL) is less specific in this respect. The incorporation of nucleotides made by the TUNEL technique indicates DNA breaks of any size with free 3 – OH ends. Such breaks may occur early in the apoptotic process and at a stage when DNA repair is still possible. Therefore, TUNEL does not necessarily indicate that the individual cell will definitively die (Davidsson *et al.*, 2000).

Our study revealed that the inflammatory infiltrate in the sinonasal polyps consisted of numerous eosinophils as well as a considerable number of lymphocytes and macrophages. Moreover, we displayed a variable grades of CD 95 expression on tissue eosinophils (regarding their average number) in 92 % of the cases as well as on numerous other inflammatory cells. These positive cells were seen more frequently towards the periphery compared to the central part of the polyps.

In addition, CD68 – positive cells (most probably macrophages) approximately equalled the number of eosinophils although their incidence varied from one polyp to another. All polyps contained CD 68 – positive cells in various grades regarding their average number.

Both CD 95 – and CD 68 – positive cells were numerous in 58% & 56 % of cases respectively. Our findings were in agreement with those observed by Davidsson *et al.* (2000) and Henson *et al.* (2001) who stated that the Annexin – V

positivity of eosinophils indicates that these cells signal their entry into apoptosis and the TUNEL positivity of eosinophils indicates that these cells have DNA breaks, which occur during the substantial period of the apoptotic process, and thus point to the apoptotic morphology *in vivo*. In addition, they reported that Annexin – V positivity demonstrates the switch of the membrane – bound phosphatidylserine molecule that occurs early in the apoptotic process. They also demonstrated that many eosinophils are actively phagocytosed within the tissue and these cells signal, as well, macrophage – like cells by the switch of phosphatidylserine molecule (binding of Annexin – V) which seems to be associated with the active phagocytosis by CD 68 – positive cells.

Davidsson *et al.* (2000) stated that the further out toward the periphery of the polyp, the more pronounced the apoptosis (TUNEL positivity) and also the phagocytosis.

Steroids are known to reduce the survival rate of human eosinophils. Eosinophil apoptosis is likely an indirect effect of the elimination of survival – inducing cytokines caused by steroids. Also Fas ligand / Fas receptor interactions are suggested to be involved in the regulation of apoptosis and defects in this system could contribute to the accumulation of these cells in diseases (Matsukura *et al.*, 1997). This statement may interpret the absence of eosinophil CD 95 immunoreactivity in 14 % of our cases.

Hebestreit *et al.* (1996) has been shown that activation of Fas with specific anti – Fas antibodies resolves chronic eosinophilic tissue inflammation, whereby apoptotic eosinophils were identified within the nasal polyps by *in – situ* detection of DNA strand breaks. As all our patients had received a local steroid treatment, some of the apoptotic eosinophils were probably died due to this therapy. This is supported by Nonaka *et al.*, (1999); Linden *et al.* (2000); Belvisi (2004) and Ochkur *et al.* (2007) observation that the airway steroids through inhibition of growth factors including IL – 5, both *in vitro* and *in vivo*,



should increase the incidence of eosinophil apoptosis. Lombardi and Passalacqua (2003) increasingly recognized that the harmful effects of tissue eosinophilia are either due to the inappropriate neutralization of their products or failure to quickly eliminate these cells from the tissues. Wooley *et al.* (1996) & Duncan *et al.* (2003) postulated that eosinophil apoptosis may occur in vivo, thus, promoting the removal of airway eosinophils and the resolution of inflammation in asthma.

Ultrastructurally, we demonstrated a distribution of eosinophils in the epithelium and subepithelial layer of sinonasal polyp. Our finding revealed that the most common mechanism for eosinophilic elimination was apoptosis that was encountered in 73% of TEM-examined eosinophils about third of which displayed features of secondary necrosis whileas cytolysis as another mechanism was found in only 4% of our TEM-examined cells. In addition, the macrophages were seen engulfing eosinophils. These ultrastructural findings confirmed the immunohistochemical ones and were in coinciding with those observed by Nutku *et al.* (2003); Gardai *et al.* (2003); Uller *et al.* (2004) and Erjefalt *et al.* (2004). In contrast, Erjefalt and Persson (2000) and Farahi *et al.* (2007) found that the apoptotic eosinophils are exceedingly rare in human nasal polypous tissues in vivo and suggested the possibility that turnover of human airway tissue eosinophils in vivo, irrespective of steroid therapy, largely involves other mechanisms than apoptosis. Alternative modes include non-injurious egression between epithelial cells into the airway luman and proinflammatory disintegration through cytolysis.

Uller *et al.* (2001) mentioned that cytolysis of eosinophils is a significant event that should not be confused with eosinophilic necrosis which occurs in vitro secondary to apoptosis and may also occur in vivo in airway tissues under the exceptional conditions caused by anti-Fas treatment of mouse airway.

Saunders *et al.* (1999) and Davidsson *et al.* (2000) reported that TUNEL technique or phosphatidylserine (Annexin) staining of the histologic sections would predictably produce false-positive results because any section-induced cell damage would make the intracellular phosphatidylserine

amenable for staining. Also, depending on the concentration and incubation time of involved enzymes and nucleotides, TUNEL method may fail to stain even truly apoptotic cells or, more commonly, may stain virtually all cells in the tissue irrespective of apoptosis.

Phagocytes taking up apoptotic cells may employ a range of possible recognition mechanisms (Davidsson *et al.* 2000). The apoptosis and phagocytosis of eosinophils by macrophages, as detected in our study, have also described in vivo in human allergen-induced cutaneous late-phase reactions (Ohta and Yamashita, 1999). In vitro studies have displayed that apoptosis of eosinophils is activated in a sequential manner and that eosinophils are phagocytosed as intact cells by macrophages (Simon and Alam, 1999).

Davidsson *et al.* (2000) mentioned that eosinophils are phagocytosed by macrophage-like cells (CD 68-positive cells) within the polypoid tissue presumably after the phosphatidylserine switch attracts the macrophages. Thrombospondin is another recognition pathway for phagocytosis although it had not been detected in any of their studied polyps; Thus, they believed that the phosphatidylserine switch was to occur early in the apoptotic process and probably before the thrombospondin bindings.

Human eosinophils, 100-fold more common in tissue than in blood, may therefore be the target for cytotoxic T-lymphocytes in tissue (Vignola *et al.*, 1999). The Fas receptor (CD 95; death receptor) is likely a universal receptor, present on many human cell types and upregulated by certain stimuli when needed (Sorkness *et al.*, 2007).

Presence of numerous inflammatory cells expressing CD95 within the polypous tissue in our as well as in Davidsson *et al.* (2000) and Henson *et al.* (2001) studies may well imply that the high number of apoptotic cells is due partly to Fas-induced apoptosis.

## Conclusion

This study added further information regarding tissue eosinophilia, apoptosis and engulfment of eosinophils by macrophages in sinonasal polyps. Moreover, it concluded that the removal of eosinophils

by cytolysis, apoptosis and phagocytosis in the sinonasal polyps might be important mechanisms, particularly the later two, for the clearance of these cells from the respiratory tissue and therefore could improve the allergic and hypersensitivity manifestations.

## References

- Bachert C, Wagenmann M, Hauser U, Rudack C (1997): IL-5 synthesis is upregulated in human nasal polyp tissue. *J. Allergy Clin. Immunol.*, 99: 837-42.
- Belvisi MG (2004): Regulation of inflammatory cell function by corticosteroids. *Proceedings of the ATS.*, 1(3): 207-14.
- Biffi WL, Moore EE, Moore FA, Barnett CC (1996): Interleukin-6 delays neutrophil apoptosis via a mechanism involving platelet-activating factor. *Trauma.*, 40:575-8.
- Bochner BS (2004): Verdict in the case of therapies versus eosinophils: the injury is still out. *J. Allergy Clin. Immunol.*, 113:3-9.
- Davidsson A, Andersson T, Hellquist BH (2000): Apoptosis and phagocytosis of tissue-dwelling eosinophils in sinonasal polyps. *Laryngoscope*, 110: 111-6.
- Davidsson A, Danielsen A, Viale G (1996): Positive identification in situ of mRNA expression of IL-6 and IL-12 and chemotactic cytokine RANTES in patients with chronic sinusitis and polypoid disease. *Acta Otolaryngol. (Stockh.)*, 116:604-10.
- Desreumaux P and Capron M (1996): Immunobiology of eosinophils in mucosal tissues. *Mucosal Immunol. Update*, 4:45-8.
- Duncan CJ, Lawrie A, Blaylock M G, Douglas JG (2003): Reduced eosinophil apoptosis in induced sputum correlates with asthma severity. *J. Eur. Respir.*, 22:484-90.
- Emanuelsson A C, Greiff L, Andersson M, Persson CG A, Erjefalt J S (2004): Eosinophil degranulation status in allergic rhinitis: observations before and during seasonal allergen exposure. *J. Eur. Respir.*, 24(5): 750-7.
- Erjefalt J S, Andersson M, Greiff L (1998): Cytolysis and piecemeal degranulation as distinct modes of activation of airway mucosal eosinophils. *J. Allergy Clin. Immunol.*, 102: 287-94.
- Erjefalt J S, Greiff L, Andersson M, Adelroth E, Jeffery P K, Persson C G (2001): Degranulation patterns of eosinophil granulocytes as determinants of eosinophil driven disease. *Thorax*, 56: 341-4.
- Erjefalt M G, Greiff L, Andersson M (2003): Marginal ultrastructural signs of degranulation in circulating eosinophils and neutrophils from symptomatic patients with asthma and allergic rhinitis. *Am. J. Respir. Crit. Care Med.*, 167: 757-61.
- Erjefalt J S, Malm G M K, Gleich G J (1999): Eosinophil cytolysis (ECL) as a major mechanism for sIgA-induced granule protein release. *Am. J. Respir. Crit. Care Med.*, 159: A99-A 104.
- Erjefalt J S and Persson C G (2000): New aspects of degranulation and fates of airway mucosal eosinophils. *Am. J. Respir. Crit. Care Med.*, 161: 2074-85.
- Erjefalt JS, Uller L, Erjefalt MM, Persson CG (2004): Rapid and efficient clearance of airway tissue granulocytes through transepithelial migration. *Thorax*, 59: 136-43.
- Farahi N, Cowburn A S, Upton P D, Deighton J (2007): Eotaxin-1 / cc chemokine Ligand 11: A novel eosinophil survival factor secreted by human pulmonary artery endothelial cells. *J. Immunol.*, 179(2): 1264-73.
- Gardai S J, Hoontrakoon R, Goddard C D, Day B J (2003): Oxidant-mediated mitochondrial injury in eosinophil apoptosis: enhancement by glucocorticoids and inhibition by granulocyte-macrophage colony-stimulating factor. *J. Immunol.*, 170: 556-66.
- Haslett C (1999): Granulocyte apoptosis and its role in the resolution and control of lung inflammation. *Am. J. Respir. Crit. Care Med.*, 160: S5-S11.
- Hebestreit H, Yousefi S, Balatti I (1996): Expression and function of the Fas receptor on human blood and tissue eosinophils. *Eur. J. Immunol.*, 26: 1775-80.
- Hellquist H B (1996): Histopathology of nasal polyps. *Allergy Asthma*, 17: 237-42.
- Hellquist H. B., Olejnicka B., Jadner M., Andersson T. (1997): Fas-receptor is expressed in human lung squamous cell carcinomas whereas bcl-2 and apoptosis are not pronounced, A preliminary report. *Br. J. Cancer.*, 76: 175-9.
- Henson P. M., Bratton D. L., Fadok V. A. (2001): The phosphatidyserine receptor: a crucial molecular switch? *Not. Rev. Mol. Cell Biol.*, 2: 627-33.
- Jahnsen F L, Haraldsen G, Aanesen J P, Haye R (1995): Eosinophil infiltration is related to increased expression of vascular cell adhesion molecule-1 in nasal polyps. *Am. J. Respir. Cell Mol. Biol.*, 12: 624-32.
- Karlesson M G and Hellquist H B (1996): Endothelial adhesion molecules for nasal-homing T-cells in allergy. *Virchows Arch.*, 429: 49-54.
- Linden M, Svensson C, Andersson E, Andersson M (2000): Immediate effect of topical budesonide on allergen challenge-induced nasal mucosal fluid levels of

- granulocyte - macrophage colony - stimulating factor and interleukin - 5 . *Am. J. Respir. Crit. Care Med.*, 162 : 1705 - 8 .
26. Lombardi C and Passalacqua G (2003) : Eosinophilia and diseases:clinical revision of 1862 cases.*Arch. Int.Med.*,163(11): 1371 - 3 .
  27. Martin L B, Kita H, Leiferman M, Gleich G J (1996): Eosinophils in allergy : role in disease , degranulation and cytokines. *Int.Arch. Allergy Immunol.*,109:207-15 .
  28. Matsukura M, Yamada H, Yudate T, Tezuka T (1997): Steroid - induced changes of eosinophils in atopic dermatitis. *Int. Arch. Allergy Immunol.*,114(suppl.1):51- 4
  29. Molet S. M., Hamid Q. A., Hamilos D. L. (2003):IL-11 and IL-17 expression in nasal polyps: relation to collagen deposition and suppression by intranasal fluticasone propionate. *Laryngoscope*,113:1803 - 11 .
  30. Nonaka R, Nonaka M, Takanashi S, Jordana M (1999) : Eosinophil activation in the tissue : synthetic steroid, budesonide, effectively inhibits the survival of eosinophils isolated from peripheral blood but not nasal polyp tissues . *J. Clin. Lab. Immunol.*, 51 : 39 - 53 .
  31. Nutku E, Aizawa H, Hudson SA, Bochner B S (2003): Ligation of Siglec-8: a selective mechanism for induction of human eosinophil apoptosis.*Blood.*,101: 5014 - 20 .
  32. Ochkur S I, Jacobsen E A, Protheroe CA, Biechele T L (2007): Coexpression of IL-5 and Eotaxin-2 in mice creates an eosinophil- dependant model of respiratory inflammation with characteristics of severe asthma.*J. Immunol.*,178 (12) : 7879 - 89 .
  33. Ohta K and Yamashita N (1999): Apoptosis of eosinophils and lymphocytes in allergic inflammation. *J. Allergy Clin. Immunol.* , 104 : 14 - 21 .
  34. Pertolani M and Goldman M (1997): Cytokines involved in the downregulation of allergic airway inflammation . *Res. Immunol.*, 148 : 33 - 8 .
  35. Rothenberg M E ( 1998 ) : Eosinophilia . *N. Engl. J. Med.* ; 338 : 1592 - 600 . .
  36. Saunders M W, Wheatley AH, George S J, Lai T (1999) : Do corticosteroids induce apoptosis in nasal polyp inflammatory cells ? : In vivo and in vitro studies . *Laryngoscope*, 109 : 785 - 90 .
  37. Simon H and Alam R (1999) : Regulation of eosinophil apoptosis : transduction of survival and death signals . *Int. Arch. Allergy Immunol.*, 118 : 7 - 14 .
  38. Sorkness R L, Herricks K M, Szakaly R J, Lemanske R F (2007): Altered allergen-induced eosinophil trafficking and physiological dysfunction in airways with preexisting virus- induced injury.*Am.J.Physiol. Lung Cell Mol. Physiol.*,292 (1):L85- L 91.
  39. Stadelmann C and Lassmann H (2000) : Detection of apoptosis in tissue sections . *Cell Tissue Res.*, 301 : 19 - 31 .
  40. Sundelin K, Jadner M, Norberg S L, Davidsson A (1997) : Metallothionein and Fas (CD 95) are expressed in squamous cell carcinoma of the tongue.*Eur.J. Cancer* 33: 1860 - 1864 .
  41. Till S, Walker S, Dickason R (1997) : IL - 5 production by allergen - stimulated T cells following grass pollen immunotherapy for seasonal allergic rhinitis . *Clin. Exp. Immunol.* , 110 : 114 - 21 .
  42. Tsuyuki S, Bertrand C, Erard F (1995) : Activation of the Fas receptor on lung eosinophils leads to apoptosis and the resolution of eosinophilic inflammation of the airways . *J. Clin. Invest.*, 96: 2924- 31 .
  43. Uller L, Andersson M, Greiff L, Persson C G A, Erjefalt J S (2004): Occurrence of apoptosis, secondary necrosis and cytolysis in eosinophilic nasal polyps . *Am. J. Respir. Crit. Care Med.*, 170 ( 7 ) : 742 - 7 .
  44. Uller L, Ottosson P, Erjefalt M M, Steven T (2001) : Anti - Fas mAb induces apoptosis of eosinophils, neutrophils and epithelial cells and may aggravate lung inflammation in allergic mice . *Am. J. Respir.Crit.Care Med.*,163: A322- A 325 .
  45. Uller L, Persson C G, Kallstrom L, Erjefalt J S (2001): Lung tissue eosinophils may be cleared through luminal entry rather than apoptosis : effects of steroid treatment . *Am J. Respir. Crit. Care Med.*, 164 :1948 - 56 .
  46. Vignola A M, Chanez P, Chiappara G, Siena L (1999) : Evaluation of apoptosis of eosinophils , macrophages and T- lymphocytes in mucosal biopsy specimens of patients with asthma and chronic bronchitis *J. Allergy Clin. Immunol.* , 103 : 563 - 73
  47. Westergren V., Viale G., Delliorto P., Pellegrini C. ( 1997 ) : RANTES is more prevalent in bacterial than non - bacterial maxillary sinusitis and P-selectin is preferentially upregulated in diseased mucosae . *Arch. Otolaryngol. Head Neck Surg.* , 123 : 1103 - 10 .
  48. Woolley K L, Gibson P G, Carty K, Wilson A J (1996) : Eosinophil apoptosis and the resolution of airway inflammation in asthma . *Am. J. Respir. Crit. Care Med.*, 154 : 237 - 43 .
  49. Ying S, Meng Q, Barata T L, Kay A B (1997):Association of apoptosis of neutrophils and eosinophils and their ingestion by macrophages with resolution of the allergen - induced cutaneous late - phase response in atopic human subjects . *Proc Assoc. Am. Physicians*, 109 : 42 - 50.

دور التساقط الخلوي والتهام خلايا الإيوسين في السليلة المخاطية للجيوب الأنفية :

دراسة هستوكيميائية مناعية وتركيبية عالية الدقة

أحمد حسن عبد الرحمن\* - عبد السلام هاشم\*\* - أحمد سيد جاد\*\* - أحمد محمد نصار\*\* -  
أحمد عبد الباسط\*\* - علي عبد الله\*\* - عاطف إبراهيم محمد علي\*\*\* -  
من أقسام الباثولوجي\* والأنف والأذن والحنجرة\*\* والهستولوجي\*\*\*  
بكلية طب الأزهر بنين فرعي أسبوط والقاهرة

تختفي خلايا الالتهاب من أنسجة المجري الهوائي عن طريق التساقط الخلوي والالتهام بالخلايا الآكلة ويمكن التخلص أيضاً من هذه الخلايا عن طريق التحلل الخلوي الأولي والموت الثانوي للتساقط الخلوي أو هجرة الخلايا عبر النسيج الظهاري . وقد قيمت هذه الدراسة دور التساقط الخلوي والتهام خلايا الإيوسين في السليلة المخاطية للجيوب الأنفية والتي استئصلت من خمسين مريضاً . وبالطريقة المناعية الهستوكيميائية تم تحديد كمية الخلايا الإيجابية لسي دي 95 وسي دي 68 عن طريق حساب متوسط العدد في عشر مجالات عشوائية عالية التكبير وكان النسيج الالتهابي المتسرب متضمناً العديد من خلايا الإيوسين وعدد لا بأس به من الخلايا الليمفاوية والعديد من الخلايا الآكلة . وقد تم أيضاً إظهار سي دي 95 بصورة متكررة علي خلايا الإيوسين وكذلك إظهاره علي العديد من الخلايا الالتهابية الأخرى أما الخلايا الإيجابية لسي دي 68 ( وهي غالباً الخلايا الآكلة ) فقد تساوت تقريباً في عددها مع خلايا الإيوسين ووجد بالطريقة التركيبية عالية الدقة أن 73 % من عدد خلايا الإيوسين المفحوصة تعاني من تساقط خلوي ( 365 من 500 خلية ) كان ثلثها تقريباً ( 89 من 365 بنسبة 24.3 % ) يعاني من الموت الثانوي للتساقط الخلوي وفي المقابل كان 4 % فقط من خلايا الإيوسين ( بعدد 20 من 500 خلية ) متحللة ذاتياً وقد تأكد بهذه الطريقة حدوث التهام خلايا الإيوسين بالخلايا الآكلة . وعلي ذلك نستنتج من هذه الدراسة أن التخلص من خلايا الإيوسين يتم غالباً عن طريق التساقط الخلوي والتهام الخلايا وهذا قد يكون مهماً في تطهير النسيج التنفسي من خلايا الإيوسين وبذلك قد تتحسن مظاهر الحساسية .