

SOIL TAXONOMY AND EVALUATION MODEL FOR SPECIFIC AGRICULTURE LAND USE OF PHYSIOGRAPHIC UNITS IN THE DESERT ZONE LOCATED ON THE EAST-NORTHERN RIM OF QARUN LAKE, FAYOUM DEPRESSION, EGYPT

Abdel Aal T.S.*, Soliman Y.R.A. and Hassanein A.E.

*Soils and Water Dept., Fac. of Agric., El Fayoum Univ., Egypt
Soils, Water and Environ. Res. Institute, Agric. Res. Center, Giza, Egypt

ABSTRACT:

The desert zone located on the east-northern rim of Qarun lake represents one of the promising desert outskirts of El Fayoum Governorate whether be under demand for agriculture use or to be future planned projects for later on use. That is true, since it is considered a good model for representing many of the landscape features in El Fayoum depression. So, it selected to be identified within the context of physiography, soil classification and land evaluation for specific agriculture land use. This area includes both the continental alluvium of the Nile and desert sediments that were derived from local parent rocks. The physiographic features were identified, using visual interpretation of Landsat data ETM7 (Enhanced Thematic Mapper 7), according the applied physiographic approach, and found to be as the Nile alluvial terraces, desert alluvial terraces, dissected slopes, aeolian plain, pediplain, bajada, cuesta fronts, cuesta summits and wadis. The later landforms whether are almost flat or gently slopping. The relatively high tableland structures were delineated as dissected cuesta of summits and fronts. Forty-five mini pits were located and studied for setting up a characteristic map legend. The differences were represented by nine soil profiles to be fully described and soil samples were selected for laboratory analyses.

Soil taxa were categorized according to the **Key of Soil Taxonomy (USDA, 2006)** till the soil family level into:

- i) The *Aridisols*, soil families are a) *Typic Haplocalcids*, coarse loamy, mixed, hyperthermic in pediplain unit b) *Typic Haplogypsids*, coarse loamy, mixed, hyperthermic in bajada unit; c) Soils of wadis unit are found in a complex pattern of *Typic Haplogypsids*, fine loamy, mixed, hyperthermic and d) *Typic Calcigypsids*, fine loamy, mixed, hyperthermic in the desert alluvial terraces unit.
- ii) The *Vertisols* include a) *Chromic Gypsite*, fine clayey, semectitic, hyperthermic in the Nile alluvial terraces unit.
- iii) *Entisols* include a) *Lithic Torriorthents*, fine loamy, mixed, hyperthermic in cuesta summits unit; b) *Typic Torriorthents*, coarse loamy, mixed, hyperthermic in cuesta fronts units; c) *Typic Torripsammets*, siliceous, hyperthermic in aeolian plain unit and d) *Lithic Torripsammets*, siliceous, hyperthermic in dissected slopes unit.

The supreme and subsequent prior potential suitability of sixteen specific crops, *i.e.*, field crops (wheat, barley, maize, cotton and onion), oil crops (sesame and sunflower), fodder crops (alfalfa and sorghum), vegetables (tomato) and fruit trees (banana, citrus, guava, mango, oil palm and olive) to be cultivated in the studied physiographic-soil units was carried out by matching between the parametric approach of land evaluation classes and their crop-physiography adaptations. The obtained data show that the potential suitability classes differed according to the

Fayoum J. Agric. Res. & Dev., Vol.23, No.2, July, 2009

satisfaction conditions between different properties of soils developed on the studied physiographic-soil units and plant requirements. These adaptations can be promising for rather higher output as the major land improvements that are considered for the land qualities of drainage, salinity and sodicity to be achieved when the land considering free of those limitations.

Key words: Physiographic-soil units, land evaluation, soil taxa, El Fayoum region.

INTRODUCTION:

The progressive increase of human pressure on limited cultivated areas in El-Fayoum Governorate requires to pay an attention towards two main aspects, *i.e.*, conserving the productivity status of cultivated soils and the horizontal expansion for the desert outskirts of El Fayoum depression. The later soils have less productive desert siliceous or calcareous in nature. The desert soils under study at both north and east rims of Oarun lake are considered ones of the promising areas for agricultural utilization and developing the economical activities of El Favoum Governorate. According to **High Dam Soil Survey Project (1963)**, the studied area are identified into four soil units for the northern-east rim of Oarun lake, *i.e.*, saline loose sand soils, foot slopes, partly shale and soft sand stone. **Shendi (1984)** classified the soils adjacent to Oarun lake from the eastern and southern rims as Typic Calcigypsid, Typic Haplogypsid and Typic Torripsammments.

The scientific benefits of the current work should be created to update and support the local knowledge, concerning the best use of land whether be under demand for agriculture use or be planned for later on use. The objectives were to identify the physiographic-soil units of a unique area in El Favoum depression by mapping them to be a digital model in a harmony of physiography and soil data set, serving the extrapolation approach when other areas will be under study.

It is also to find the best adaptation between certain land units with specific crop to give the maximum output. For this purpose, the harmony of descriptive and processing systems, established by **Svs (1991) and Svs et al. (1993)** were considered, being highly required in this study. The collective findings of this study create and document copy data sets, using visual interpretation of Landsat image, basic and topographic maps to obtain a physiographic-soil map, soil taxa and land suitability classes. The result is a comprehensive land evaluation database for a certain area in Egypt. These data can be matched with the other products, produced as the same global standard.

MATERIAL AND METHODS:

Image interpretation:

Landsat images interpretation performed using the physiographic analysis as proposed by **Burnigh (1960) and Gossen (1967)**. Generally, Landsat image composite of Enhanced Thematic Mapper (ETM7) with bands 2,3 and 4 was used to add an extra landscape assessment to the photo-interpretation map. The image was helpful for getting a collective overall view of the studied area as well as using the spectral signatures of the used bands in detecting the cultivated areas and drainage conditions.

Visual analysis of Landsat TM5:

Images of Landsat 5 Thematic Mapper (TM) were used for the purpose of visual analysis. The overall view for delineating the promising areas in East desert soils of Menya characterized by the spectral signatures of an

Orthorectified Landsat Thematic Mapper (TM 5) Mosaic. It is a composite of the bands 4, 3 and 2. The pixel size is a mixture of 28.5 and 30.0 meters. The composite output was of benefit, especially when focusing on the infrared bands that permit the detection and discrimination of broad combinations of different vegetation cover types and identification of water bodies, active drainage, drainage conditions, cultivated areas, and rock types.

The image of Landsat 5 was used for the detailed physiographic analysis for modeling the study areas. This Landsat 5 was acquired during the year 2004 (path 175 rows 42, resolution 28.5 to 30 m). The Thematic Mapper of this Landsat is operating in eight spectral bands. The images are considered as a source of recent information that can be aimed at transferring the recent or modified infrastructures to the maps during the phase of cartography.

Field work:

The physiographic units of space images were checked in the field to confirm the boundaries of the physiographic units or to revise what were shifted. Soil profiles representing the predominant characteristics of the identified physiographic units of the studied area were taken, however, eleven soil profiles were dug to a depth of 150 cm or lithic contact, their locations are shown in Fig. (1).


Soil profiles were described, using the nomenclature of the **Soil Survey Division Staff Manual (USDA, 2003)**. The described soil characteristics included land surface configuration, soil slope, vegetation, parent material, soil depth, master horizons, texture, rock fragments, soil matrix color, soil structure, consistence, roots, pores, secondary formation of calcium carbonate and gypsum and boundaries of horizons and layers. Soil samples were air dried, crushed, with wooden hammer, sieved through a 2 mm sieve to obtain the fine earth used for physical and chemical analysis. The elements of soil color description, *i.e.*, the colour name and notations were determined using the **Munsell Soil Colour Chart (1975)**.

Laboratory analyses:

Particle size distribution was determined using the International Pipette method (**Piper, 1950**) and sodium hexametaphosphate as dispersing agent (**Baruah and Barthakur, 1997**). Calcium carbonate content was measured using the Collin's Calcimeter method (**Wright, 1939**). Gypsum was determined by the acetone method (**Bower and Huss, 1948**). Saturation soil paste extract analysis, and soil pH in the soil water suspension of 1:2.5 were determined according to the methods describe by **Jackson (1973)**. Cation exchange capacity and the exchangeable sodium were determined according to the methods describe by **Richards (1954)**.

Soil classification and evaluation:

Soils under study were classified into taxonomic units starting from the level of soil order down to the soil family level according to Soil Taxonomy System (**USDA, 1975**) and using the keys to Soil Taxonomy (**USDA, 2006**). Soils under investigation were evaluated using the parametric system for land evaluation undertaken by **Svs and Verheve (1978)**. Land suitability classification for specific crops was done according to **Svs et al. (1991)** and **Svs et al. (1993)**. Matching the land characteristics with the crop requirements was undertaken by limitation approach.



RESULTS AND DISCUSSION:

A general view on the physiographic-soil units and their taxonomic classes:

Physiographic-soil legend has been set up as shown in Fig. (1), associated with the morphological description of the representative soil profiles in Table (1). The identified physiographic-soil units were Landsat imagery delineated in two main formations, *i.e.*, 1) The Nile deposits (flat or almost flat Nile alluvial terraces) and 2) The desert formations (cuesta

SOIL TAXONOMY AND EVALUATION MODEL FOR SPECIFIC... 69

summits, cuesta fronts, bajada, wadis, pediplain, aeolian plain, dissected slopes, desert alluvial terraces). Soil taxa after soil physical and chemical analyses are presented in Tables (2, 3 and 4).

Table 1: Morphological description of the studied soil profiles.

Physiographic unit	Profile No.	Slope gradient	Horizon	Depth (cm)	Soil colour	Modified texture class	Soil structure	Soil consistency	Boundary	Pedological features
The Nile alluvial terraces	1	Almost flat	A	0-10	10YR 3/4	C	Medium sub-angular blocky	Hard	-- Diffuse	--
			By	10-30	10YR 3/3			Very hard		sfg
			Byz	30-80	10YR 3/3			Very hard		sfg
			By	80-150	10YR 3/2			Very hard		sfg
The desert alluvial terraces	2	Gently undulating	A	0-25	10YR 7/6	Sl	Massive Sub-angular blocky	Slightly hard	-- Diffuse	sfcg
			Bky	25-70	10YR 6/4					sfCa
			By	70-150	10YR 7/6					sfg
Dissected slopes	3	Gently slopping	A	0-10	10YR 7/6	LS	Massive	Slightly hard	--	--
			Lithic contact							
Aeolian plain	4	Almost flat	A	0-30	10YR 7/5	S	Single grain	Loose	-- Diffuse	--
			C	30-75	10YR 6/6					
			C ₂	75-150	10YR 6/6					
Pediplain	5	Gently undulating	A	0-25	10YR 7/4	LS	Single grain	Slightly hard	-- Clear	--
			C	25-80	10YR 6/3					sfCa
			C ₂	80-150	10YR 6/5					--
Bajada	6	Gently slopping	A	0-30	10YR 7/5	SCL	Massive Sub-blocky Platy	Slightly hard	-- Clear Clear	sfCa
			By	30-100	10YR 8/6			Hard		sfg
			C	100-150	10YR 6/4			Hard		--
Cuesta fronts	7	Very gently sloping	A	0-30	10YR 7/6	LS	Single grain	Slightly hard	-- Diffuse	--
			C	30-60	10YR 7/5					
			C ₂	60-150	10YR 6/6					
Cuesta summits	8	Gently undulating	A	0-25	10YR 7/5	GSL	Massive	Loose	-- Clear	--
			C	25-40	10YR 6/6	SCL		Hard		
Wadis	9	Almost flat	A	0-25	10YR 6/7	SCL	Massive Platy Platy	Slightly hard	-- Clear Diffuse	sfCa
			C	25-80	10YR 5/5			Hard		sfg
			C ₂	80-150	10YR 5/6			Hard		--

Soil texture: S=Sand, LS=Loamy sand, SL=Sandy loam, GSL=Gravelly sandy loam, SCL=Sandy clay loam, C=Clay.

Pedological features: sfg= secondary formations of gypsum, sfCa=secondary formations of lime

A brief note about each of the identified physiographic-soil units, which are belonging the previous two formations in the studied area, was carried out as follows:

1. The Nile alluvial deposits:

The Nile alluvium occurred under a specific depositional action of the ancient lake of Fayoum depression "Lake Moeris" with the Nile fresh-water in the Pliocene period resulting in different physiographic-soil units, particularly the Nile alluvial low terraces adjacent to Birkat Qarun. Thus, an injection was occurred between fluvial deposits and local lacustrine ones, which were richer in secondary formations of CaCO₃ and gypsum. **Said (1990)** stated that, in the case of Lake Moeris, where the Fayoum depression fell completely with the fresh Nile water, that reaches inundate its floodbasin. Due

to the depositional processes rather than erosional processes, hence the Nile suspended matter was deposited in different physiographic-soil units from the seasonal and periodic flooding. Discharge that is fully confined to El Fayoum basin maintains high competence, and when discharge exceeds basin capacity, there is a dramatic increase in cross-sectional area associated with expansion into the floodbasin.

Table 2: Particles size distribution %, calcium carbonate %, gypsum %, CEC and ESP of the studied profiles.

Physiographic units	Profile No.	Depth (cm)	Gravel %	Particle size distribution %				Modified texture class	CaCO ₃ %	CaSO ₄ ·2H ₂ O %	CEC (C mol _c kg ⁻¹)	ESP
				C. sand	F. sand	Silt	Clay					
The Nile alluvial terraces	1	0-10	--	15.0	7.9	33.8	43.3	C	9.10	1.50	38.55	7.58
		10-30	--	14.3	8.2	33.3	44.2		10.50	9.30	33.67	6.58
		30-80	--	15.7	6.8	34.6	42.9		8.10	11.30	34.08	5.65
		80-150	--	16.8	9.7	29.9	43.6		11.30	10.10	35.78	9.67
The desert alluvial terraces	2	0-30	--	56.4	20.0	8.9	14.7	SL	3.50	11.30	11.87	6.15
		30-70	--	60.1	16.1	10.2	13.6		17.30	11.70	10.43	7.84
		70-150	--	65.2	14.4	8.3	12.1		5.10	13.10	8.68	9.76
Dissected slopes	3	0-10	--	69.4	15.3	7.1	8.2	LS	2.44	1.12	5.35	11.47
		Lithic contact										
Aeolian plain	4	0-30	--	79.3	15.0	3.5	2.5	S	2.48	1.37	2.76	4.95
		360-75	--	77.5	17.4	2.8	2.3		1.97	0.84	1.89	5.84
		75-150	--	76.5	16.7	3.3	3.5		2.06	0.75	2.12	7.03
Pediplain	5	0-25	--	65.4	19.1	8.2	7.3	LS	14.97	3.75	5.96	9.74
		25-80	--	59.5	18.7	7.3	14.5	SL	27.50	4.02	6.87	12.35
		80-150	--	67.7	17.5	5.3	9.5	LS	21.34	3.44	8.39	13.89
Bajada	6	0-30	--	47.5	15.0	12.5	25.0	SCL	9.15	2.50	16.11	8.36
		30-100	--	50.2	15.5	11.3	23.0		3.10	13.30	14.97	9.15
		100-150	--	35.0	14.4	22.4	28.2		2.30	3.70	18.53	11.74
Cuesta fronts	7	0-30	--	63.7	20.0	6.8	9.5	LS	2.54	3.94	7.47	12.87
		30-60	--	67.3	16.0	6.0	10.7		2.88	2.58	8.95	14.35
		60-150	--	62.9	21.7	7.0	8.4		3.66	1.24	7.15	14.86
Cuesta summits	8	0-20	41.3	55.8	21.4	8.3	14.5	G SL	2.10	1.24	11.01	10.60
		20-40	5.1	34.4	26.2	15.3	24.1	SCL	7.30	1.97	19.54	13.42
Wadis	9	0-30	--	43.5	13.3	13.7	23.5	SCL	10.95	3.12	17.85	7.99
		30-80	--	45.4	16.6	12.8	25.2		4.55	11.84	18.40	8.87
		80-150	--	37.9	15.3	19.7	27.1		2.79	2.50	21.61	9.45

Soil texture: S=Sand, LS=Loamy sand, SL=Sandy loam, GSL=Gravelly sandy loam, SCL=Sandy clay loam, C=Clay.

The velocity and depth of water flowing inside of the floodbasin declines rapidly with distance away from the water courses. The coarsest sediment (usually fine sand and silt) undergoes rapid deposition immediately adjacent to the entrance of the Nile water at Al-Lahun gap, while the finest sediment was deposited away from the gap at the lowest level of Birket Qarun, resulting in low-lying flat to almost flat (slightly depressed) Nile alluvial terraces with somewhat well drained soils of heavy-textured soil material. These soils are subjected to the swelling and shrinkage process fitting the main requirement to be *Vertisols*. Also, these soils have a control section is fully characterized by

SOIL TAXONOMY AND EVALUATION MODEL FOR SPECIFIC... 71

clayey layers, and are represented by soil profile No. 1. The soils are more developed due to the occurrence of gypsum horizon "By", and are classified as *Chromic Gypsite*, *fine clayey, semectitic, hyperthermic*.

Table 3: Chemical analysis of soil paste extract of the studied soil profiles.

Physiographic units	Profile No.	Depth (cm)	pH	EC (dS/m)	Cations (meq / L)				Anions (meq / L)		
					Ca ⁺⁺	Mg ⁺⁺	Na ⁺	K ⁺	HCO ₃ ⁻	Cl ⁻	SO ₄ ⁻
The Nile alluvial terraces	1	0-10	7.50	46.15	150.50	16.28	356.07	0.95	2.28	51.00	470.52
		10-30	7.10	41.14	247.30	75.72	163.60	0.89	2.23	380.00	105.28
		30-80	7.20	84.99	344.01	62.46	632.87	0.95	2.54	620.00	417.74
		80-150	7.30	20.12	53.70	19.28	145.26	0.67	2.11	110.00	106.80
The desert alluvial terraces	2	0-30	7.50	5.54	38.71	6.72	10.46	0.56	1.91	42.00	12.54
		30-70	7.70	13.58	40.99	11.28	90.60	0.67	2.41	95.00	46.13
		70-150	8.00	13.25	47.37	14.55	79.97	0.74	2.47	80.00	60.16
Dissected slopes	3	0-10	7.80	11.13	29.10	10.19	74.95	0.84	1.98	55.00	58.10
		Lithic contact									
Aeolian plain	4	0-30	7.25	17.10	21.69	11.94	143.12	0.24	1.81	75.60	99.58
		360-75	7.24	55.50	87.25	55.68	514.97	0.89	2.26	391.00	265.53
		75-150	7.18	21.37	28.62	15.54	187.90	0.45	1.89	108.00	122.62
Pediplain	5	0-25	7.90	13.16	19.15	7.36	108.30	0.47	1.89	69.00	64.39
		25-80	7.80	75.10	35.12	19.35	788.90	0.75	1.86	541.00	301.26
		80-150	7.70	15.30	17.33	7.93	132.46	0.64	1.58	83.00	73.78
Bajada	6	0-30	7.60	13.82	43.01	14.37	86.22	0.54	2.36	115.00	26.78
		30-100	7.70	12.60	37.64	9.36	84.31	0.49	2.27	105.00	24.53
		100-150	7.30	5.58	32.38	4.15	20.17	0.44	1.96	40.00	15.18
Cuesta fronts	7	0-30	7.90	14.77	18.25	9.40	125.77	0.48	2.18	65.00	86.72
		30-60	7.70	94.19	40.78	21.73	1065.81	0.93	2.51	672.00	454.83
		60-150	7.60	12.25	32.39	16.27	77.17	0.59	2.06	65.00	59.36
Cuesta summits	8	0-20	7.70	13.26	40.94	6.78	94.68	0.68	2.31	85.00	55.77
		20-40	7.30	7.18	12.38	12.69	49.47	0.71	1.99	45.00	28.26
Wadis	9	0-30	7.68	15.13	39.93	13.18	106.05	0.61	2.11	67.60	90.06
		30-80	7.24	17.34	48.99	16.83	118.61	0.59	2.00	101.50	81.52
		80-150	7.19	9.78	34.87	7.94	59.23	0.55	2.29	59.80	40.50

2. The desert formations:

2.1. Alluvial terraces:

These terraces are beginning from the northern shoreline of Oarun lake and extended towards the north direction. According to **Said (1990)**, they are remnants of formerly erosional processes. On this area, consequent streams, that follows the initial slope of the land (southwards) were rejuvenated, resuming down-cutting, thereby forming terraces, resulting in gullied surfaces of concave convex complex slopes (gently undulating topography). The soils are classified as *Typic Calcigypsis*, *fine loamy, mixed, hyperthermic*. This soil family is characterized by sandy loam in texture, and it is represented by soil profile No. 2.

2.2. Dissected slopes:

This physiographic unit represents the remnants of a structural plateau, subjected to severe dissection resulting in a rocky structure that be located on rocky slopes covered by talus and pediments with a complex pattern of steep and rolling concave convex surfaces. The soils are very shallow depth (10 cm depth) underlain lithic contact bed-rock, and they are classified as *Lithic Torripsamments, siliceous, hyperthermic*. This soil family is characterized by sand in texture, and it is represented by soil profile No. 3.

Table (4): Soil taxonomic units of the studied soil profiles.

Order	Sub-order	Great group	Sub-group	Family	Representative soil profiles and physiographic units
Vertisols	Torrerts	Gypsi-torrerts	Chromic Gypsi-torrerts	Chromic Gypsi-torrerts, fine clayey, smectitic, hyperthermic	1 (The Nile alluvial terraces)
Aridisols	Gypsisds	Calci-gypsisds	Typic Calci-gypsisds	Typic Calcigypsisds, fine loamy, mixed, hyperthermic	2 (The desert alluvial terraces)
		Haplo-gypsisds	Typic Haplo-gypsisds	Typic Haplogypsisds, coarse loamy, mixed, hyperthermic	6 (Bajada)
	Calcids	Haplo-calcids	Typic Haplo-calcids	Typic Haplocalcids, coarse loamy, mixed, hyperthermic	5 (Pediplain)
				Typic Haplogypsisds, fine loamy, mixed, hyperthermic	9 (Wadis)
Entisols	Psamments	Torri-psamments	Lithic Torri-psamments	Lithic Torripsamments, siliceous, hyperthermic	3 (Dissected slopes)
			Typic Torri-psamments	Typic Torripsamments, siliceous, hyperthermic	4 (Aeolian plain)
	Orthents	Torri-orthents	Typic Torri-orthents	Typic Torriorthents, coarse loamy, mixed, hyperthermic	7 (Cuesta fronts)
			Lithic Torri-orthents	Lithic Torriorthents, fine loamy, mixed, hyperthermic	8 (Cuesta summits)

2.3. Aeolian plain:

This unit is found in areas of the Earth where erosion and deposition by wind are the dominant geomorphic forces, particularly under the dry climates that are classified as arid deserts and semi-arid steppe. Different processes are responsible for the transport of sediment by wind, like this rolling motion,

which called traction and suspension, this type of transport is called also saltation. This plain is located on the margins of the plateau remnant, immediately bordering the dissected slopes in the southern side. It is most probably that, these aeolian deposits are covering an old bed-rock that was naturally excavated. The plain is locally occupied by sinuous clustered low-dunes, which are currently retreated as subjected to the natural vegetation. This plain includes excessively well-drained coarse textured soils, that are classified as *Typic Torripsamment, siliceous, hyperthermic*, and its soils are represented by profile No. 4.

2.4. Pediplain:

This physiographic unit was formed under the prevailing aridic conditions through an action of physical weathering processes on the limestone parent rock. This unit has very gently slopping to gently undulating and including somewhat well drained soils. Its polygons are the remnants of weathered limestone rock, including residual parent material over limestone lithic contact at the depth of 90-150 from soil surface. This parent material developed to *Aridisols*, being with diagnostic calcic horizon (BK). The representative soils are with control sections dominated by loam sandy texture grade. The soils were classified as *Typic Haplocalcids, coarse loamy, mixed, hyperthermic*; and these soils are represented by soil profile No. 5.

2.5. Bajada:

This unit is a depositional belt in the studied area along the elevated structures of cuesta fronts when the fans coalesce laterally to form that bajada. It is somewhat relatively broad and gently inclined, alluvial piedmont slope extending from the base of cuesta range out into relatively low basin southern-eastwards. Bajada surface is gullied and gently slopping. The soils are well drained, classified as *Typic Haplogypsis, coarse loamy, mixed, hyperthermic*. These soils are more developed than those the pediplain, being with gypsic "By" horizon, and they are represented by soil profile No. 6.

2.6. Cuesta:

This physiographic unit represents the remnants of a structural plateau, subjected to severe dissection resulting in a rocky structure, that be divided into two sub units as:

a. Cuesta fronts:

These fronts are located on gently slopping area covered by talus deposits and with a complex pattern of relatively rolling concave convex surfaces. These soils are classified as *Typic Torriorthents, coarse loamy, mixed, hyperthermic*. Also, this soil family is characterized by loamy sand in texture, and it is represented by soil profile No. 7.

b. Cuesta summits:

These summits represent the original elevation of the limestone body before the dissection processes. The soils are very shallow depth (40 cm depth) underlain lithic contact bed-rock. This unit has gently undulating to gently sloping, gravely and stony surfaces including somewhat well drained soils, and they are classified as *Lithic Torriorthents, fine loamy, mixed, hyperthermic*. This soil family is characterized by sandy loam to sandy clay loam in texture, and it is represented by soil profile No. 8.

2.7. Wadis:

The surface of this physiographic unit is almost flat to gently undulating, partly natural vegetated on well drained soils. Also, this unit is the resultant of dissection action of the surrounding landscape as the interaction of erosional

and depositional processes in the fluvial period. They appear as dry wadis, that seasonally receive flush flooding, running from west to east or southeast, draining into the relatively low areas. The soils of these wadis occur in a complex pattern and dominated by one taxonomic unit of *Typic Haplogypsis*, *fine loamy, mixed, hyperthermic*. These soils have a relatively fine texture class within the control section (sandy clay loam) as compared to the other studied desertic formations. This soil family is represented by soil profile No. 9.

It is noteworthy to mention that, based on the detected soil morphological features and physico-chemical properties, the soils under investigation could be classified up to the family level into nine taxonomic classes according to **Keys to Soil Taxonomy (USDA, 2006)**, as shown in Table (4). It is most probably due to the studied soils are developed on different parent materials differ in their mineralogical compositions, which are more related to the physiographic positions. Also, the investigated area lies within the climatic conditions characterized by a long hot rainless summer and short mild winter with a rare rainfall. The majority of the studied soil profiles are rich in secondary CaCO_3 and gypsum accumulations, which satisfy the requirement of calcic and gypsic horizons in some soil profile layers such as Nos. 1, 2, 5, 6 and 9.

Soil limitations for productivity and land evaluation:

The physical parametric land evaluation system undertaken by **Sys and Verheye (1978)**, which is considered a favourable system under the conditions prevailing in the soils of Egypt (**Moussa, 1991**), was applied to determine the soil limitations and their intensities as well as land suitability classes according to the current suitability ratings. Since it is valid for irrigation purposes in arid and semi arid regions. By this approach, the classification was processed according to the **FAO Framework (1976)**, at the level of sub-classes.

The obtained data in Table (5) reveal that all the studied soils have no limitations for their topography (except profile Nos. 1 and 3), wetness (w), the effective soil depth (except profile Nos. 3 and 8) and gypsum content (except profile No. 2). On the other hand, most of the representative soil profiles are suffering from soil texture (s_1), CaCO_3 (s_3) content and salinity/alkalinity (n) as limitations for soil productivity, which are put into variable intensity degrees of (40-95, very severe-slight), (95, slight) and (50-85, severe-moderate), respectively.

According to the same parametric system and the estimated data of soil criteria, the suitability indices (Ci) for the studied nine profiles for current and potential suitability classes are also assessed and recorded in Table (5). The obtained results show that the estimated current ratings of the studied soil profiles ranged between 11.99 and 76.61, indicate that the soils of the studied areas could be categorized into four classes, as follows.

a. Not suitable soils (N):

The rating of this class is < 25 , and represented by soil profile Nos. 3 and 4.

b. Marginally suitable soils (S3):

The rating of this class is $25-<50$, and represented by profile Nos. 1, 2, 5, 7 and 8.

c. Moderately suitable soils (S2):

The rating of this class is $50-<75$, and represented by soil profile No. 9.

d. Highly suitable soils (S1):

The rating of this class is 75 or more, and represented by soil profile No. 6.

Table (5): Soil limitations and rating indices for the evaluation of the studied soil profiles.

Physiographic units	Profile No.	Suitability condition	Topography (t)	Wetness (w)	s				Soil salinity/Alkalinity (n)	Rating (Ci)	Suitability class	Suitability subclass
					Soil texture (s1)	Soil depth (s2)	CaCO ₃ (s3)	Gypsum (s4)				
Nile alluvial terraces	1	Current	90	100	85	100	95	100	50	36.33	S3	S3n
		Potential	100	100	85	100	95	100	100	80.75	S1	S1
Desert alluvial terraces	2	Current	100	100	75	100	95	80	85	48.45	S3	S3s1s4
		Potential	100	100	75	100	95	80	100	57.00	S2	S1s1s4
Dissected slopes	3	Current	90	100	55	30	95	100	85	11.99	N2	N2s1s2
		Potential	100	100	55	30	95	100	100	15.67	N2	N2s1s2
Aeolian plain	4	Current	100	100	40	100	95	100	58	22.04	N1	N1s1n
		Potential	100	100	40	100	95	100	100	38.00	S3	S3s1
Pediplain	5	Current	100	100	55	100	100	100	58	31.90	S3	S2s1n
		Potential	100	100	55	100	100	100	100	55.00	S2	S2s1
Bajada	6	Current	100	100	95	100	95	100	85	76.71	S1	S1
		Potential	100	100	95	100	95	100	100	90.25	S1	S1
Cuesta fronts	7	Current	100	100	55	100	95	100	58	30.30	S3	S2s1n
		Potential	100	100	55	100	95	100	100	52.25	S2	S2s1
Cuesta summits	8	Current	100	100	85	55	95	100	85	40.34	S3	S3s2
		Potential	100	100	85	55	95	100	100	44.41	S3	S3s2
Wadis	9	Current	100	100	95	100	95	100	80	72.20	S2	S2n
		Potential	100	100	95	100	95	100	100	90.25	S1	S1

For ameliorating the suitability of these soils, major improvement practices should be carried out such as removing the excess of soluble salts and sodicity through applying the gypsum and leaching requirements under an efficient drainage ditches, in addition to organic fertilization and other soil and water managements. Such agro-management practices will correct the ratings of soil potential suitability classes to be ranged 15.67-90.25, and potential soil suitability becomes as follows.

a. *Not suitable soils (N):*

The rating of this class is < 25, and represented by soil profile No. 3.

b. *Marginally suitable soils (S3):*

The rating of this class is 25-<50, and represented by soil profile Nos. 4 and 8.

c. *Moderately suitable soils (S2):*

The rating of this class is 50-<75, and represented by soil profile Nos. 2, 5 and 7.

b. *Highly suitable soils (S1):*

The rating of this class is 75<, and represented by soil profile Nos. 1, 6, and 9.

Land suitability for certain crops:

The physiographic-soil map was used as a base for presenting land suitability classes. The simple approach that proposed by **Sys (1991)** was selected for land suitability evaluation of the studied area, since it is valid for irrigation purposes in arid and semi arid regions. By this approach, the classification was processed according to the framework of **FAO (1976)**, at

the level of subclasses. Ratings, attributed to land qualities, were matched with each crop requirements. The land qualities are drainage (d), soil texture (x), stoniness (g, gravel %), soil depth (p), calcium carbonate % (c), salinity (s, ECe), sodicity (n, ESP) and fertility (f). Fertility ratings attributed to soil reaction (pH), cation exchange capacity (CEC) and sum of basic cations (exchangeable Ca, Mg and K). Suitability subclasses in Tables (6 and 7) reflect the kind of limitations as indicated in symbols, using lower-case letters synonymous with those limitations when any of them is moderate.

Land suitability for agricultural irrigated soils is the appraisal of specific areas of land from a general point of view without mentioning the specific kind of use. So, some soils may be suitable for a specific crop and unsuitable for another. The ideal approach for land evaluation is based on evaluating the land for utilization types which used as guides for the most beneficial use for a specific productivity by replacing a less adapted land utilization type by another promising one, and was applied in this study according to **Sys (1991)**. The evaluation indices of land characteristics are done by rating them and specifying their limitations for certain crops by matching the calculated rating with the crop requirements in different suitability levels as proposed by **Sys et al. (1993)**.

a. Current land suitability classification (Cs):

In the studied area, without major land improvements, the crop requirements were matched with the present land qualities for processing the current land suitability of the different land units. This approach enables management of different alternatives for specific utilizations that are adapted to the existing limitations to give maximum output. The current land suitability classification of different physiographic units for the different specific utilizations is shown in Tables (6 and 7).

b. Potential land suitability classification (Ps):

As for this purpose, the land utilization is applicable after executing specified major land improvements as proposed in the current study according to their necessity. Potential land suitability classification can be established if the main improvements for the studied area are considered regarding land qualities of drainage, salinity and sodicity. The potential land suitability classification of different physiographic units for the different specific utilizations is shown in Tables (6 and 7) for the studied area. The obtained potential land suitability subclasses were sorted in two productive levels. These two levels were designed to be guide charts for the best land utilization alternatives giving a possible maximum output. The two potential land suitability levels are as follows:

1. Supreme potential suitability for specific utilizations:

Matching charts of the supreme potential suitability for specific utilizations with the different physiographic-soil units of the studied area shown in Tables (6 and 7).

Tables 6, 7

The resultant adaptations are as follows:

*** Highly suitable (S1) adaptations:**

- a. Some soil sites of the desert alluvial terraces unit are suitable for sesame, alfalfa and olives.
- b. Some soil sites of bajada unit are suitable for maize, sesame, sunflower and olives.
- c. Some soil sites of cuesta fronts unit are suitable for olive.
- d. Some soil sites of wadis unit are suitable for sesame, sorghum and olives.

*** Moderately suitable (S2) adaptations:**

- a. Some soil sites of the Nile alluvial terraces unit are suitable for wheat, barley, maize, cotton, sesame, sunflower, alfalfa, sorghum, guava and olives.
- b. Some soil sites of pediplain unit are suitable for maize, sesame, sunflower and alfalfa.
- c. Some soil sites of cuesta summits unit are suitable for barley, sorghum and guava.

*** Marginally suitable (S3) adaptations:**

- a. Some soil sites of dissected slopes unit are suitable for wheat, barley, sorghum and guava.
- b. Some soil sites of aeolian plain unit are suitable for cotton, sesame, sunflower, alfalfa, sorghum, onion, tomato, citrus, guava, mango and olives.

2. Subsequent prior potential suitability for specific utilizations:

*** Moderately suitable (S2) adaptations:**

- a. Some soil sites of the desert alluvial terraces unit are suitable for sorghum and guava.
- b. Some soil sites of bajada unit are suitable for wheat, barley, cotton, alfalfa and guava.
- c. Some soil sites of cuesta fronts unit are suitable for maize, sunflower, alfalfa, onion, tomato, citrus, and mango.
- d. Some soil sites of wadis unit are suitable for wheat, barley, maize, cotton, sunflower, alfalfa and guava.

*** Marginally suitable (S3) adaptations:**

- a. Some soil sites of the Nile alluvial terraces unit are suitable for banana.
- b. Some soil sites of pediplain unit are suitable for wheat, barley, cotton, sorghum, onion, tomato, citrus, guava and mango.
- c. Some soil sites of cuesta summits unit are suitable for wheat, maize, cotton, sesame, alfalfa, banana and oil palm.

From the aforementioned discussion, it could be concluded that the results of the current work are created a local knowledge should be supported the future projects of agricultural utilization in El Fayoum area. Also, the obtained can be use as a guide to explain and correct the problems facing the future agricultural utilization projects in the area under investigation, and in turn the best use of land whether be under demand for agriculture use or be planned for later on use. That means the obtained results represent the best adaptation between certain land units with specific land use in agricultural purposes as well as to give the maximum outputs from the agricultural utilization projects.

Moreover, identifying the physiographic-soil features of a unique area in the adjacent desert outskirts of El Fayoum depression by mapping them to be a

model is in a harmony of physiographic and soil data set, serving the extrapolation approach when other areas will be under study. In addition, this work was carried out by using physiographic-soil units map obtained from the visual analysis of Enhanced Images of Landsat Thematic Mapper 7 of the studied area. These soil criteria represent a base for making a proper agricultural utilization and could be considered as promising items in soil potentiality and its sustainable agriculture on the long-term.

REFERENCES:

- Baruah, T.C. and Barthakur, H.P. (1997).** A Text Book of Soil Analysis. Vikas Publishing Housing, PVT LTD, New Delhi, pp. 35-67.
- Bower, C.A. and Huss, R.B. (1948).** A rapid conductometric method for estimating gypsum in soils. *Soil Sci.*, 66: 199-240.
- Burnigh, P. (1960).** The applications of aerial photographs in soil surveys. *Man. of Photogr. Interpr.*, Washington, D.C., USA.
- FAO (1976).** A framework for land evaluation. *Soil Bull. No.32*, FAD, Rome, Italy.
- Goosen, A.A.I. (1967).** Aerial photo-interpretation in soil survey. *FAO Soil Bulletin No. 6*, Rome, Italy.
- High Dam Soil Survey Project (1963).** Soil Survey Report. Soil Water and Environment Institute, Giza, Egypt.
- Jackson, M.L. (1973).** Soil Chemical Analysis. Prentice India Private LTD, New Delhi, Indian.
- Moussa, M.A. (1991).** Land suitability evaluation of El Saff area, Eastern Desert, Egypt for agriculture utilization. Ph. D. Thesis, Fac. of Agric. at Moshtohor, Zagazig Univ., Egypt.
- Munsell Soil Colour Charts (1975).** Edition Munsell Colour, Macbeth Division of Kollmorgen Corp., 2441 North Calvert Street, Baltimore, Maryland, USA.
- Piper, C.S. (1950).** Soils and Plant Analysis. Inter Science Publishers, Inc., New York, pp. 59-75.
- Richards, L.A. (1954).** Diagnosis and Improvement of Saline and Alkali Soils. U.S. Dep. of Agric., Hand Book No. 60, pp. 102.
- Said, R. 1990.** The Geology of Egypt. Published for the Egyptian Central Petroleum Corporation. Conoco Hurghada Inc, by Balkema, A.A., Rotterdam, The Netherlands.
- Shendi, M.M. (1984).** Pedological studies on soils adjacent to Qarun lake, Fayoum depression, Egypt. M. Sc. Thesis, Fac. of Agric., Cairo Univ., Egypt.
- Sys, C. (1991).** Land evaluation. Parts I, II and III, Lecture Notes, Ghent Univ., Ghent, Belgium.
- Sys, C. and Verheye, W. (1978).** An attempt the evaluation of physical land characteristics for irrigation according to the FAO framework for land evaluation. Int. Train Course for Post Grad. Soil, Sci. Univ., Ghent, Belgium.
- Sys, C.; Van Ranst, F.; Debaveye, J. and Bernaert, F. (1993).** Land evaluation. Part III., Crop Requirements Agricultural Publication No.7, General Administration for Development Cooperation, Ghent, Belgium.
- USDA (2003).** Soil Survey Manual. United State, Department of Agriculture, Handbook 18, U.S. Gov. Print Off., Washington, DC.,USA.
- U.S.D.A. (2006).** Keys to Soil Taxonomy. 10th (Ed), United States Department of Agriculture, Nature Resources Conservation Service.

نموذج تصنيفي وتقييمي لأراضي الوحدات الفيزيوجرافية لاستخدام زراعي معين في النطاق الصحراوي

الواقع شمال-شرق حافة بحيرة قارون - منخفض الفيوم - مصر

طلبه صالح عبد العال* ، ياسر ربيع أمين سليمان ، أحمد السيد أحمد حساتين

*قسم الأراضي والمياه - كلية الزراعة - جامعة الفيوم - مصر

معهد بحوث الأراضي والمياه والبيئة - مركز البحوث الزراعية - جيزة - مصر

تمثل أراضي النطاق الصحراوي الواقع شمال-شرق حافة بحيرة قارون أحد المناطق الواعدة في الظهير الصحراوي لمحافظة الفيوم تحت الطلب للإستخدام في الأغراض الزراعية أو في التخطيط المستقبلي لمشروع التنمية الزراعية. هذا حقيقي، لأنها تعتبر نموذجاً ممثلاً لمعظم ملامح الوحدات الارضية في منخفض الفيوم، قد اختيرت هذه المنطقة لتعريفها من خلال المفهوم الفيزيوجرافي وتصنيف التربة وتقييمها بغرض استخدامات زراعية معينة. وتضم هذه المنطقة الترسبات القارية لنهر النيل وأيضا الترسبات الصحراوية المتاخمة لها والتي اشتقت من صخور الأم المحلية. ولقد تم تعريف الوحدات الفيزيوجرافية عن طريق التحليل المرئي لصور القمر الاستطاعي (ETM 7) مع تطبيق المنهج الفيزيوجرافي. ومن خلال هذا التحليل وجد ان منطقة الدراسة تتكون من المصاطب النهرية الرسوبية The Nile alluvial terraces، المصاطب الصحراوية الرسوبية Desert alluvial terraces، سلسلة من الميول المتقطعة Dissected slopes، سهل الترسبات الهوائية Aeolian plain، سهل تجوية والذي يحتوي على مادة اصل متبقية Pediplain، الباجادا Bajada، من كويستا وبروزات Cuesta summit and fronts، الوديان Wadis. والملاحم الأرضية تضم أجزاء مستوية وأخرى ذات ميول بسيطة. والأجزاء ذات المناسيب المرتفعة تضم تتواجد في شكل بقايا مادة اصل هضبية الشكل Plateau تعرضت لعمليات تجوية شديدة حتى أصبحت في شكل كويستا وبروزات. ولقد حددت خمسة وأربعون موقعا لحفر نقاط ملاحظة أرضية صغيرة من اجل تصميم مصطلحات الخريطة حيث تم تمثيل تبايناتها بدراسة تفصيلية لتسعة قطاعات لأرضية اخذت منها عينات للتربة للتحليلات المعملية.

وقد تم تصنيفت خواص التربة حسب دليل نظام التصنيف الأمريكي لعام (٢٠٠٦) حتى مستوي عائلات

التربة، حيث وجد ان الأراضي المتكونة تتبع ثلاثة رتب تشتمل على العائلات التالية:

- i) The Aridisols, soil families are a) Typic Haplocalcids, coarse loamy, mixed, hyperthermic in pediplain unit b) Typic Haplogypsiids, coarse loamy, mixed, hyperthermic in bajada unit; c) Soils of wadis unit are found in a complex pattern of Typic Haplogypsiids, fine loamy, mixed, hyperthermic and d) Typic Calcigypsiids, fine loamy, mixed, hyperthermic in the desert alluvial terraces unit.
- ii) The Vertisols include a) Chromic Gypsitegrets, fine clayey, semectitic, hyperthermic in the Nile alluvial terraces unit.
- iii) Entisols include a) Lithic Torriorthents, fine loamy, mixed, hyperthermic in cuesta summits unit; b) Typic Torriorthents, coarse loamy, mixed, hyperthermic in cuesta fronts units; c) Typic Torripsamments, siliceous, hyperthermic in aeolian plain unit and d) Lithic Torripsamments, siliceous, hyperthermic in dissected slopes unit.

وقد تم تقييم أراضي الوحدات الفيزيوجرافية - سواء بالنسبة لصفات التربة الحالية أو بعد تحسينها - بغرض تحديد الأفضلية الأعلى لحاصلات زراعية معينة بلغت في جملتها ستة عشر محصول ممثلة في حاصلات الحقل (القمح، الشعير، الأذرة، القطن، البصل)، حاصلات زيتية (السمسم، عباد الشمس)، حاصلات عاف (البرسيم، السورجم)، خضر (الطماطم)، أشجار فاكهة (الموز، الموالح، الجوافة، المانجو، نخيل الزيت، نخيل البلح) والتي يمكن زراعتها في أراضي الوحدات الفيزيوجرافية تحت الدراسة، وذلك من خلال عمل توافق بين التقييم الكمي للتربة وإحتياجات تلك الحاصلات. وتشير النتائج المتحصل عليها إلى أن القدرة الكامنة لمدى صلاحية الأراضي تختلف تبعاً لمدى توفر خصائص التربة الملائمة المتكونة على كل من الوحدات الفيزيوجرافية تحت الدراسة. وأن هذه هذه الموائمات بين أراضي الوحدات الفيزيوجرافية والمحاصيل المختلفة تتوقف على مدى التأقلم المستقبلي ما بين خصائص التربة وإحتياجات الحاصلات المختارة. وهذا التأقلم المستقبلي واعدا بلا شك لتعظيم العائد من خلال تحسين حالة التربة خاصة فيما يتعلق بالصرف، مستويات الملوحة والصودية لتعزيز القدرة الكامنة للأراضي بعد خلوها من تلك العوامل المعوقة للإنتاجية.