

EFFECT OF POTASSIUM NITRATE, GARLIC AND ONION EXTRACTS ON BUD BREAK, GROWTH, YIELD AND SOME CHEMICAL CONSTITUENTS OF APPLE (*MALUS SYLVESTRIS*, MILL) TREES.

Morsi, M.E.* and Mohamed A. Seif El-Yazal**

* Dept. of Horticulture, Fac. of Agric., Fayoum Univ., Egypt.

** Dept. of Agric. Botany, Fac. of Agric., Fayoum Univ., Egypt.

ABSTRACT

This investigation was carried out during the two successive seasons of 2006 and 2007 to investigate the effect of potassium nitrate, garlic extract and onion extract on bud break, growth, yield and some chemical constituents of "Anna" apple (*Malus sylvestris*, Mill) variety. The trees were grown in loamy sand soil, and sprayed with six treatments (potassium nitrate 10%), garlic extract (20%), onion extract (20%), potassium nitrate (5%) mixed with garlic extract (10%), potassium nitrate (5%) mixed with onion extract (10%) and control. Generally, it was found that all studied growth parameters, date of flower bud break, percentage of bud break, fruit-setting, fruit weight, fruit size, fruit number/tree, yield/tree (kg) and some chemical constituents of leaves (total chlorophyll, total carbohydrates, total protein, nitrogen, phosphorous and potassium contents) and some chemical constituents of fruits (total soluble solids (T.S.S.), T.S.S./ acid ratio, vitamin C, water content %, total free amino acids, total carbohydrates, total sugars and reducing sugars) were increased with the application of the different treatments. The best results were obtained from the treatments of potassium nitrate at 5% mixed with onion extract at 10%, potassium nitrate at 5% mixed with garlic extract at 10% and potassium nitrate at 10%. On the contrary, the same treatments decreased total acidity and total phenols in fruits as compared to the control. It could be recommended to use potassium nitrate at 5% in combination with onion extract or garlic extract at 10% for improving bud break, growth, yield and chemical constituents of apple trees or fruits.

Key Words: apple (*Malus sylvestris*, Mill), potassium nitrate, garlic extract, onion extract, bud break, growth, yield, chemical constituents.

INTRODUCTION

Dormancy in deciduous fruit trees is a physiological syndrome occurring annually to enable plants to survive cold winters. Entrance into dormant state occurs in late summer and fall. Deciduous trees require low winter temperatures to overcome dormancy and grow in the following spring. Induction and release of buds from dormancy are linked to structural and metabolic modification occurring on the bud level (Ramina *et al.*, 1995).

Several chemicals can be used to induce bud break of deciduous fruit trees in areas lacking sufficient chilling units. Nitrogenous compounds such as potassium nitrate have a synergistic effect on bud break, growth, yield and some chemical constituents of many fruit trees (Kuden *et al.*, 1997 on apricot trees; Jackson *et al.*, 1997 and El-Shewy *et al.*, 1999 on apple trees).

Garlic and onion extract have been widely used for stimulating bud break in various fruit species. In this respect, Botelho and Muller (2007a) study the effect of garlic extract at different concentration on bud break of "Royal Gala"

Fayoum J. Agric. Res. & Dev., Vol.22, No.1, January, 2008

EFFECT OF POTASSIUM NITRATE, GARLIC AND ONION..... 83

apple trees compared with the effects of the conventionally used hydrogen cyanamide (H₂CN₂) immediately after winter pruning, different treatments were sprayed at dormant bud stage using a hand sprayer. All treatment showed similar effects compared with hydrogen cyanamide (H₂CN₂) alone, achieving 80% bud sprouting, whereas the control attained 18% bud sprouting. The 10% garlic extract + 2% mineral oil treatment was superior to the other treatments, reaching 95% bud sprouting at 50 days after treatment and advancing bloom.

The beneficial effect of garlic and onion extract on bud break, growth, yield and some chemical constituents of different fruit species were studied by several workers (Kubota *et al.*, 1999; Kim *et al.*, 2000a & b; Kubota *et al.*, (2000) and Botelho *et al.*, (2007) on grape trees; Botelho and Muller (2007a & b) on apple trees).

MATERIALS AND METHODS

This study was carried out during the two successive seasons of 2006 and 2007 in the orchard of the Horticulture Farm of Faculty of Agriculture, Fayoum University in an attempt to break dormancy of "Anna" apple variety (*Malus sylvestris*, Mill) grafted on Malling-Merton 106 (MM 106) root stock. The trees were 10 years old when experiment started and grown in loamy sand soil. Trees were selected in November, 2005 a uniform as possible for spray treatments.

The experiment involved the following treatments

- 1- Control (spraying with tap water)
- 2 Spraying with potassium nitrate at rate 10%
- 3- Spraying with garlic extract at rate 20%
- 4 Spraying with onion extract at rate 20%
- 5- Spraying with potassium nitrate at rate 5%+ garlic extract at rate 10%
- 6- Spraying with potassium nitrate at rate 5%+ onion extract at rate 10%

The physical and chemical characters of the orchard soil was determined according to Wilde *et al.*, (1985) and the results are shown in Table (1).

Table (1): Chemical and physical analysis of the soil

Depth	Physical characteristics									
	Particle size distribution				Texture	Bulk density g/cm ³	Organic mater %	Soil moisture constant %		
	Coarse sand%	Fine sand %	Silt %	Clay %				F. C	W.P	A.W
0-30	57.24	28.99	7.07	6.60	Loamy	1.39	0.81	17.5	7.16	10.75
30-60	52.87	28.98	8.86	8.37	Sand	1.41	0.74	20.1	8.74	11.77
chemical characteristics										
	Soluble cations (meq /L)				pH	ECe (dS/m)	Soluble anions (meq/l)			
	Ca ⁺⁺	Mg ⁺⁺	Na ⁺	K ⁺			Cl ⁻	Hco ₃ ⁻	So ₄ ⁻	
0-30	17.65	15.88	12.23	1.31	7.21	4.49	7.67	2.82	33.44	
30-60	10.20	11.81	5.01	0.62	7.45	2.63	3.65	2.95	22.46	

In all experiments, Phosphorous as calcium super phosphate (15.5% P₂O₅) at the rate of 200 kg/fed., was added in the orchard in the second week of February. Nitrogen as ammonium nitrate (33.5% N) at the rate of 250 kg /fed. was added in two doses for the orchard (first dose 150 kg/fed. in the second week of February and second dose 100 kg./fed. before top flowering (first

week of April) and potassium sulphate (48% K₂O) at the rate of 50 kg/fed., was given in two equal doses in alternative with nitrogen fertilizer. The first dose of fertilizer was added in (March) and the second dose given after 30 days from the first one. The other cultural practices were followed as normal. The control trees were sprayed with tap water, however, potassium nitrate (containing 13% nitrogen and 44% potassium), garlic extract and onion extract were sprayed at two equal doses, the first was sprayed before the end of dormancy (nearly 30th of December), and the second was applied two weeks later with a volume of 4 L/tree for each one. Triton B as a wetting agent at 0.1% was added to the spraying solutions.

Preparation of garlic and onion extracts :

Garlic or onion samples were ground using mortar and pestle and the active ingredients were extracted by ethyl alcohol (95%). The garlic or onion ethanol mixture was filtered and the alcohol was removed by evaporating under vacuum (30C°) using rotary evaporator, Buchi model 011. The extract was kept cool in refrigerator (4C°) until use. Garlic or onion extract was diluted by water to give the final concentration required (10 and 20%) directly before use.

Data recorded

A. Morphological characteristics

In both the two successive seasons of the study, each tree the dates of flower and vegetative bud opening were recorded. Number of vegetative and flower buds was counted when all buds were opened and the percentages were estimated. The dormant buds were also counted and were expressed as percentage from the total number of buds. The dates at which flowering reached 25, 50, 75 and 100 percent of the total flowers were estimated in each treatment. Flowers whose calyx began to extend were tagged in order to determine the percent of fruit set. The yield of fruits in kg/tree as well as the number of mature fruits/tree were recorded when fruits reached the commercial colour to be picked.

In order to determine fruit quality, 20 fruits were taken at random from each tree as a sample. Samples were transferred immediately to the laboratory. Each fruit was weighed to get the average fruit weight. Average fruit size was determined by emerging the fruit in a jar containing water and receiving the excess water in a graduated cylinder.

Chemical analysis

Fresh and dried leaves as well as fruits (May15th for chemical constituents & 30th July for mineral elements in leaves and at harvesting, for fruits) were taken to determine the following constituents: total chlorophyll was extracted from fresh leaves by acetone (80%) and its concentration was determined as mg/100g fresh weight according to **Welburn and Lichtenthaler (1984)**, total carbohydrates mg/g dry weight were determined colorimetrically by using phenol-sulphuric acid reagent according to the method described by **Herbert et al. (1971)**. Total free amino acids in fresh fruits were determined as mg/g fresh weight colorimetrically using ninhydrin reagent according to the method described by **Jayarman (1981)**. (Total and reducing sugars were determined as mg/g fresh weight using phosphomolybdic acid reagent., total phenols in fresh fruits were determined as mg/g fresh weight using Folin-Denis reagent. Water content in fruits was determined, total soluble solids (T.S.S.) in fruits were estimated using handle

EFFECT OF POTASSIUM NITRATE, GARLIC AND ONION..... 85

Refractometer model PZONr. 19877, total acidity was estimated in fruits as malic acids using sodium hydroxide for a known normality and phenolphthaline as an indicator. Total soluble solids/ acid ratio were calculated and vitamin C content in fruits, nitrogen %, crude protein percentage and phosphorus % in dry leaves were determined according to **A.O.A.C., 1995**). Potassium was determined by Flame Photometer, Parkin–Elmer model 52 according to the method described by **Page et al. (1982)**.

Statistical analysis:

The experiment was in a complete randomized block design with 6 treatments and 3 replicates for each treatment. One tree was used as a replicate. Results were statistically analyzed using the L.S.D. at probability level of 5% for comparisons according to **(Gomez and Gomez, 1983)**.

RESULTS:

A- Date of flower bud break:

Data presented in Table (2) clearly indicated that spraying apple trees with all the tested substances hastened the beginning of flower bud break as compared to the control. This earliness reached about 31 and 29 days for potassium nitrate at 10%, 25 and 22 days for garlic extract at 20%., 27 and 28 days for onion extract at 20%, 28 and 25 days for potassium nitrate at 5% mixed with onion extract at 10%. and 38 and 38 days for potassium nitrate at 5% mixed with onion extract at 10% over the control in both seasons, respectively.

As regards to the effect of the tested substances on 50% bud break, the present results clearly show that all treatments hastened 50 % bud break as compared to the control. This earliness reached about 35 and 36 days for potassium nitrate at 10%, 31 and 31days for garlic extract at 20%, 30 and 31 days for onion extract at 20%., 34 and 37 days for potassium nitrate at 5% mixed with onion extract at 10%. and 40 and 38 days for potassium nitrate at 5% mixed with onion extract at 10% over the control in both seasons, respectively.

Table (2): Effect of potassium nitrate, garlic and onion extract treatments on time of flower bud opening in "Anna" apple trees.

Treatments	Date of flower bud opening											
	Beginning		25%		50%		75%		End		Flowering period (No. of days)	
	2006	2007	2006	2007	2006	2007	2006	2007	2006	2007	2006	2007
Control	15/3	17/3	19/3	20/3	1/4	3/4	7/4	7/4	10/4	9/4	26	25
Potassium nitrate 10%	12/2	16/2	16/2	18/2	24/2	26/2	1/3	2/3	8/3	12/3	24	24
Garlic extract 20%	18/2	23/2	20/2	24/2	1/3	3/3	10/3	12/3	13/3	19/3	23	24
Onion extract 20%	16/2	17/2	20/2	20/2	1/3	2/3	5/3	6/3	9/3	12/3	21	23
KNO ₃ 5%+Garlic 10%	15/2	20/2	18/2	21/2	26/2	25/2	3/3	6/3	10/3	14/3	23	22
KNO ₃ 5%+Onion 10%	5/2	7/2	8/2	10/2	20/2	24/2	22/2	26/2	25/2	1/3	20	22

B-Percentage of bud break:

Data presented in Table (3) clearly show that all treatments gave a high percentage of flower bud break compared with the control. The maximum increases were recorded with potassium nitrate at 5% mixed with onion or garlic extract at 10% which recorded 12.56 and 11.33% in both seasons over

the control for potassium nitrate plus onion extract and 9.32 and 10.94% for potassium nitrate plus garlic extract, respectively.

C- Yields and its components:

Data in Table (3) indicated that all the tested substances increased apple yield and its components (fruit-setting, fruit weight, fruit size and fruit number) as compared to the control trees. Such trend was true during the two studied seasons. The maximum increases were recorded with potassium nitrate at 5% mixed with onion extract at 10% and potassium nitrate at 5% mixed with garlic extract at 10% which recorded 6.97 and 5.91 for fruit-setting, 33.86 and 33.28% for fruit weight, 26.52 and 25.47% for fruit size, 6.72 and 5.23% for fruit number and 42.82 and 40.22% for apple yield/tree in the first season, respectively over the control trees. Moreover, in the second season such increases were 7.13 and 6.42 for fruit-setting, 32.05 and 30.91% for fruit weight, 25.26 and 25.05% for fruit size, 5.56 and 5.30% for fruit number and 39.32 and 37.81% for yield/tree, over the control level, respectively.

Table (3): Effect of potassium nitrate, garlic and onion extract treatments on the percentage of bud break, fruit setting, Fruit weight (gm), Fruit size(CC³), total number of fruits/tree and yield/tree in "Anna" apple trees.

Treatments	Bud break %		Fruit set%		Fruit weight (gm)		Fruit size (CC ³)		Total number of fruits / tree		Yield per tree (Kg)	
	2006	2007	2006	2007	2006	2007	2006	2007	2006	2007	2006	2007
Control	79.6	78.33	13.13	13.18	84.63	86.28	95.00	95.64	154.79	153.56	13.10	13.25
Potassium nitrate 10%	86.30	86.70	18.31	19.13	111.62	112.70	118.10	119.10	161.98	160.57	18.08	18.09
Garlic extract 20%	84.60	84.36	16.13	16.58	111.15	112.16	117.60	118.00	157.61	156.91	17.51	17.59
Onion extract 20%	86.60	86.30	18.15	18.25	107.14	108.43	116.00	117.11	158.20	158.01	16.95	17.13
KNO ₃ 5%+Garlic 10%	86.95	86.90	19.04	19.60	112.80	112.95	119.20	119.60	162.90	161.70	18.37	18.26
KNO ₃ 5%+Onion 10%	89.60	87.21	20.10	20.31	113.29	113.94	120.20	119.80	165.20	162.10	18.71	18.46
L.S.D at 5%	2.24	2.66	1.39	1.42	1.12	1.13	1.56	1.63	2.89	2.93	1.04	1.05

C- Chemical constituents of leaves:

1- Total chlorophyll, total carbohydrates and total protein

Data presented in Table (4) clearly showed that, during the two successive seasons of the study, all treatments increased the concentrations of leaf constituents (total chlorophyll, total carbohydrates, total protein) as compared to the control. The best results were observed when apple trees were sprayed with potassium nitrate at 5% mixed with onion extract at 10% and potassium nitrate at 5% mixed with garlic extract at 10% which recorded 3.19 and 3.19% for total chlorophyll, 16.81 and 16.77% for total carbohydrates and 1.44 and 1.44 for total protein in the first seasons and 7.73 and 7.73% for total chlorophyll, 17.08 and 17.00% for total carbohydrates and 1.18 and 1.12 for total protein in the second seasons over the control plants, respectively.

2- Nitrogen, phosphorus and potassium concentrations in leaves:

Data presented in Table (4) revealed that, leaves of apple trees contained higher concentrations of nitrogen, phosphorus and potassium under foliar

EFFECT OF POTASSIUM NITRATE, GARLIC AND ONION..... 87

spray with any of the treatments than the control. The maximum increases were obtained when potassium nitrate at 5% mixed with onion extract at 10% as well as potassium nitrate at 5% mixed with garlic extract at 10% which recorded 0.23 and 0.23 for nitrogen, 0.05 and 0.04 for phosphorous and 0.03 and 0.06 for potassium in the first seasons and 0.18 and 0.18 for nitrogen, 0.05 and 0.04 for phosphorous and 0.06 and 0.06 for potassium in the second seasons respectively as over the control trees.

Table (4): Effect of potassium nitrate, garlic and onion extract treatments on chemical contents of leaves in "Anna" variety.

Treatments	Total chlorophyll mg/g.f.w		Total carbohy drate mg/g. d.w		Total protein %		N %		P %		K %	
	2006	2007	2006	2007	2006	2007	2006	2007	2006	2007	2006	2007
Control	1.88	1.81	76.20	76.15	10.56	11.00	1.69	1.76	0.20	0.21	1.52	1.50
Potassium nitrate 10%	1.93	1.94	88.10	89.01	11.94	12.06	1.91	1.92	0.24	0.24	1.53	1.55
Garlic extract 20%	1.93	1.92	87.09	86.60	11.68	11.56	1.87	1.85	0.23	0.22	1.52	1.51
Onion extract 20%	1.93	1.94	88.06	87.15	11.93	12.06	1.91	1.92	0.24	0.29	1.53	1.54
KNO ₃ 5%+Garlic 10%	1.94	1.95	88.98	89.10	12.00	12.12	1.92	1.94	0.24	0.25	1.54	1.56
KNO ₃ 5%+Onion 10%	1.94	1.95	89.10	89.16	12.00	12.18	1.92	1.94	0.25	0.26	1.55	1.56
L.S.D at 5%	0.04	0.08	1.11	1.20	0.23	0.22	0.04	0.04	0.01	0.01	N.S	N.S

D- Chemical constituents of fruits:

Data of Tables (5 and 6) clearly showed that spraying apple trees with any of the tested substances significantly improved the chemical constituents of fruits (total soluble solids (T.S.S.), total acidity, T.S.S/ acid ratio, vitamin C, water content %, total free amino acids, total carbohydrates, total sugars, reducing sugars, total phenols) as compared to the control trees. Such trend was true during the two seasons of the study. The maximum increases were recorded with potassium nitrate at 5% mixed with either onion or garlic extract at 10% which recorded 1.5 and 1.5 for total soluble solids, 5.24 and 4.84 for T.S.S/ acid ratio, 34.51 and 32.74% for vitamin C, 3.30 and 3.01 for water content, 16.94 and 15.96% for total carbohydrates, 20.66 and 19.84% for total sugars, 30.27 and 28.19% for reducing sugars and 27.07 and 24.86% for total free amino acids in the first seasons respectively, as compared to the control trees, and 1.38 and 1.26 for total soluble solids, 4.92 and 4.58 for T.S.S/ acid ratio, 31.30 and 29.56% for vitamin C, 4.9 and 4.51 for water content, 17.09 and 14.74% for total carbohydrates, 20.09 and 18.59% for total sugars, 30.30 and 28.78% for reducing sugars and 31.31 and 30.76% for total free amino acids in the second seasons, respectively, as compared to the control trees. On the other hand, the data in Table (5) also showed a marked decrease in total acidity and total phenols concentrations in fruits when trees were treated with any of the tested substances comparing with the control trees.

Table (5): Effect of potassium nitrate, garlic and onion extract treatments on chemical fruit quality of "Anna" apple fruits.

Treatments	TSS %		Acidity %		TSS/Acidity ratio		Vitamin C mg/100 ml juice	
	2006	2007	2006	2007	2006	2007	2006	2007
Control	11.30	11.40	1.05	1.05	10.76	10.85	1.13	1.15
Potassium nitrate 10%	13.10	12.60	0.88	0.87	14.26	14.36	1.48	1.48
Garlic extract 20%	12.55	12.60	0.92	0.91	13.58	13.84	1.45	1.48
Onion extract 20%	12.75	12.60	0.90	0.89	14.16	14.15	1.49	1.46
KNO ₃ 5%+Garlic 10%	12.80	12.66	0.82	0.82	15.60	15.43	1.50	1.49
KNO ₃ 5%+Onion 10%	12.80	12.78	0.80	0.81	16.00	15.77	1.52	1.51
L.S.D at 5%	0.05	0.07	0.04	0.05	0.89	0.89	N.S	N.S

Table (6): Effect of potassium nitrate, garlic and onion extract treatments on some chemical composition of "Anna" apple fruits.

Treatments	Water content %		Total carbohydrates mg/F.W		Total sugars mg/g F.W		Reducing sugars mg/g F.W		Total free amino acids mg/g F.W		Total phenols mg/g F.W	
	2006	2007	2006	2007	2006	2007	2006	2007	2006	2007	2006	2007
Control	77.60	76.50	141.60	145.10	85.61	87.10	55.50	56.10	1.81	1.82	0.45	0.47
Potassium nitrate 10%	79.88	80.26	156.6	159.60	100.60	102.60	68.60	69.10	2.20	2.30	0.31	0.28
Garlic extract 20%	79.60	80.20	150.10	152.50	90.10	91.20	60.61	63.15	1.99	1.98	0.35	0.36
Onion extract 20%	79.20	79.90	151.10	152.30	91.60	92.30	61.20	62.20	1.97	1.98	0.36	0.37
KNO ₃ 5%+Garlic 10%	80.61	81.01	164.20	166.50	102.60	103.30	71.15	72.25	2.26	2.38	0.30	0.28
KNO ₃ 5%+Onion	80.90	81.40	165.60	169.90	103.30	104.60	72.30	73.10	2.30	2.39	0.28	0.27
L.S.D at 5%	1.15	1.26	2.95	2.99	1.01	1.05	1.11	1.09	0.06	0.08	0.3	0.04

DISCUSSION

Spraying apple trees with any of the tested treatments (potassium nitrate at 10%, garlic extract at 20%, onion extract at 20%, potassium nitrate at 5% mixed with either garlic or onion extract at 10%) resulted in vigorous plant (tree) growth as well as high productivity with good fruit quality. Treatments increased the measured growth characters. This was due to the fact that these treatments resulted in more availability of macronutrients (N, K and S) to plants. Enhancement of growth parameters with N application would be expected since nitrogen is of extreme importance to plants. It is a constituent of many important substances within plant cells such as protoplasm, in addition to amino acids, nucleic acids, protein and chlorophylls (Salisbury and Ross, 1992). The high levels of endogenous auxin and gibberellins were found in those plants sprayed with high N fertilizer (Rajagopal and Rao, 1974), which encourage cell division and cell elongation, increases leaf number and produce a sufficient assimilation area for maximum rate of photosynthesis (Greenwood and Hunt, 1986). Moreover, Mengel and Kirkby (1987) reported that, the role of K in metabolism, growth and yield formation can be characterized by two major function: as an activator of enzymes and as K⁻ ions are very mobile within the plant as well as within a

EFFECT OF POTASSIUM NITRATE, GARLIC AND ONION..... 89

cell are transported through biological membranes with high rate and specificity. More than 60 enzymes are known to require K^+ as an activator. The high mobility of K^+ on photosynthesis phloem loading and phloem transport ...etc. Such important physiological roles enable potassium to perform its functions, which lead to an increase in various vegetative growth and yield. Also, **Kubota *et al.*, (1999)** investigated the active substances in garlic which is responsible for breaking bud dormancy in grapevines. The compounds were identified as diallyl mono-, di-, tri-, and tetra-sulfides, but only trace amounts of dimethyl mono- and di-sulfides were present. Exposure to volatiles of diallyl di- and tri-sulfides was the most effective treatment in promoting bud break, irrespective of the concentration and the duration of exposure. However, the effects of dimethyl sulfide and diallyl sulfide on budbreak varied among the concentrations and the duration of exposure. These results indicate that the active substances in garlic, responsible for breaking bud dormancy in grapevines, are sulfur- containing compounds with an allyl group (CH_2CHCH_2), particularly diallyl disulfide. Moreover, garlic and onion extract used in this study, have an improving effect on vegetative growth parameters. This may be attributed to the essential role of these substances in the synthesis of some amino acid and consequently, formation of growth regulators especially auxin, and ethylene.

Also, the favorable effect of the used substances on date of flower bud opening may be due to their stimulation effect of natural gibberellin. In this connection **Luna *et al.* (1993) and Subha-Drabandhu (1995)**, concluded that the induction of flowering could be correlated with a natural rise in gibberellin which promote flower formation in plants by either facilitating the formation of flowering hormone in the leaves or expressing it in the growing buds. Gibberellins also may be a primarily responsible for bolting which may be essential for the formation of the floral stimulus in leaves. Moreover, **Subha-drabandhu (1995) and Nashaat (1996)** reported that some different spray treatments may break dormancy by decreasing ABA content in buds.

The improving effect of potassium nitrate, garlic extract and onion extract on yield and its components was mainly attributed to its positive action on enhancing growth parameters (Table 2) and photosynthetic pigments of plant leaves (Table 4). In this respect, **Skene (1969)** reported that when a bud opens and attains the shape of a shoot, its tip acts as a strong sink for metabolites and thus being interception center for photosynthates and nutrients results in earlier start of the bloom.

The promotive effect of potassium nitrate, garlic extract and onion extract on chlorophyll formation might be attributed to their enhancing effect on the nutritional status of apple trees. Also the increase of total chlorophyll by spraying with N and K may be due to that N and K play an important role for stimulating chlorophyll synthesis enzymes which can be reflected on the formation of chlorophyll molecule. Moreover, the stimulating effect of potassium nitrate, garlic extract and onion extract as foliar spray on total carbohydrates concentrations in leaves of sprayed trees may be directly or indirectly due to certain enzymes which activate the anabolic processes leading to the accumulation of these substances. The increase of all mentioned constituents by foliar N application may be due to that certain enzymes may be activated as a result of these treatments leading to the accumulation of such substances. The increase of macronutrients (N, P, and K) and protein content were supported by the results of **El-Shewy *et al.*, (1999)** on apple trees. In this

connection **Tromp (1970)** found that there was a decrease in the nitrogen concentration of the woody tissues in the spring, particularly in the bark tissues of shoots. This might be attributed to the movement of nitrogenous compounds from the bark and wood to the developing flower buds and growing points. Moreover, the stimulating effect of potassium nitrate, garlic extract and onion extract on physical characters (fruit weight and size) and chemical fruit characters (T.S.S, total acidity, vitamin C, total carbohydrates, total sugars, reducing sugars, total free amino acids and total phenols) was mainly attributed to its positive action on enhancing growth parameters (Table 2) and photosynthetic pigments of plant leaves (Table 4). Concerning the effect on T.S.S. and acidity in fruits, the results showed that all treatments increased T.S.S. significantly and decreased the total acidity. This increase in T.S.S. may be due to the increase in synthesis of carbohydrates and its accumulation in the developing fruits of the treated trees. In this connection, **Boghdadi (1964)** mentioned that sugars represented about 70% of the total soluble solids (T.S.S) in apple fruits and the increase in sugars lead to increase in T.S.S. He also added that the increase in cellular sap lead to decrease in acidity as a result of dilution of the organic acids. Moreover **Dame et al. (1956)** and **Mann and Singh (1990)** on pear, found that the increase in T.S.S. may be due to rapid conversion of starch, and the decrease in total acids content with advancement of ripening period may be due to that the acids are converted into soluble solids. The increase in vitamin C may be due to that fruits synthesize ascorbic acid from hexose sugars and hence the adequate supply of these precursors would greatly depend on the photosynthetic activity (**Mapson, 1970**). In this connection **George et al. (1990)** suggested that water and nutrients may also be mobilized to the growing points at the expense of the developing fruits. Also, **Ahmed (1995)** found that large "Anna" apple fruits had significantly higher reducing and total sugars as well as lower starch and non-reducing sugars than small sized fruits. Moreover, **Dame et al. (1956)** found that the increase in accumulation of T.S.S. and sugars during maturation has been related to accumulation of glucose, sucrose and higher levels of fructose in "Bartlett" pear. On the other hand, **Mann and Singh (1990)** found that the total phenols content (as tannic acid) decrease during ripening period. The reduction in phenolic content during ripening process may be attributed to its hydrolysis to different components such as sugars, acids and other compounds (**Gangwar, 1972**).

Finally, from the results of the present investigation, it could be concluded that the application of potassium nitrate, garlic extract or onion extract greatly increased growth and apple yield as well as improved apple quality and its chemical constituents. The constituents of these substances participate in the different metabolic processes which increased syntheses of chlorophyll, carbohydrates, total free amino acids, and absorption of essential nutrients, so that the use of potassium nitrate, garlic extract and onion extract could increase apple productivity with high fruit quality.

REFERENCES

- Ahmed, E. Z. (1995):** Effect of Gibberellin, Cycocel, Calcium and Boron, fruit size and position within tree canopy on quality and mineral content of Anna apple fruits during storage. Ph. D. Thesis, Alexandria Univ.
- A.O.A.C. (1995):** Official Methods of Analysis of the Association of Official Agricultural Chemists. 16th ed., Washington D.C., USA.
- Boghdadi, H. A. (1964):** Principles of fruit production . Dar El-Maerif, Cairo, Egypt, 3rd ed., pp. 663-667.
- Botelho, R. V. and Muller, M. M. L. (2007a):** Evaluation of garlic extract on bud dormancy release of "Royal Gala" apple trees. Australian Journal of Experimental Agriculture 47 (6) 738-741
- Botelho, R. V. and Muller, M. M. L. (2007b):** Garlic extract as alternative on bud dormant break of apple trees cv. Fuji Kiku. Rev. Bras. Frutic., Vol. 29, No. 1, p. 37-41.
- Botelho, R. V.; Pavanello, A. P.; Pires, E. J. P.; Terra, M. M. and Mullr, M.M.L. (2007):** Effect of chilling and garlic extract on bud dormancy release in Cabernet Sauvignon grapevine cuttings. Am. J. Enol. Vitic. 58; 402-404.
- Dame, C.D.; Lonard, S. J.; Luh, B. S. and G. L. Mansh (1956):** The influence of ripeness on the organic acids, sugars and pectin of canned Bartlett pears. Fd. Techn. Champing 10: 23-33.
- El-Shewy, A.A.; Ibrahim, A.A.; Zeid, F.A. and El-Yazal, M.A.S. (1999):** Effect of some dormancy breaking compounds on chemical composition of leaves and buds of some apple cultivars. II-Chemical composition vegetative and generative buds. Annals of Agric., Sci., Moshtohor 87 (4): 2279-2306.
- Gangwar, B.M. (1972):** Biochemical studies on growth and ripening of guava. Indian Food Packer, 26: 13-15.
- Gomez, K.A. and Gomez, A.A. (1983):** Statistical Analysis Procedure of Agricultural Research. John Wiley and Sons, New York, pp. 25-30.
- George, A.P.; R.J. Nissen; Lioyed, J. and K. Richens (1990):** Factors affecting fruit quality of low chill stone fruit in subtropical Australia. Acta. Hort., 279: 559-564.
- Greenwood, D. J. and J. Hunt (1986):** Effect of nitrogen fertilizer on the nitrate contents of field vegetables grown in Britain. J. Sci. Food Agric., 37: 373 – 383 .
- Herbert, D.; Phipps, P.J. and Strange, R.F. (1971):** Determination of total carbohydrates. Methods in Microbian, 5 (B): 209-244.
- Jayarman, J. (1981):** Laboratory Manual in Biochemistry, Wiley Eastern Limited New York, pp. 61-73.
- Jackson J.E.; Barritt, B.H, and Kappel, F. (1997):** Branch induction using hydrogen cyanamide and Promalin. Acta-Horticulturae, No. 451, 679-681.
- Kim, S.; Kim, S.H.; Kim, S.K. and Kim, S.H. (2000a):** Effects of Allium, Merit Blue, and soluble P-K formulae on budbreak in differentially chilled "Campbell Early" grapevines. Journal of the Korean Society for Horticultural Science. 2000, 41: 3, 265-268.

- Kim, S.K.; Kim, S.H.; Lee-Young, C.; Kim, S.K.; Kim, S.H. and Lee, Y.C. (2000b):** Effects of anaerobiosis and subsequent treatments with Merit Blue and Allium formulae on budbreak in 'Campbell Early' and 'Muscat Bailey A' grapevine cuttings. *Journal of the Korean Society for Horticultural Science*. 41: 3, 269-272.
- Kubota, N. and Miyamuki, M. (1992):** Breaking bud dormancy in grapevines with garlic paste. *Journal of the American Society for Horticultural Science*. 117: 6, 898-901.
- Kubota, N.; Yamane, Y.; Toriu, K.; Kawazu, K.; Higuchi, T. and Nishimura, S.(1999):** Identification of active substances in garlic responsible for breaking bud dormancy in grapevines. *Journal of the Japanese Society for Horticultural Science*. Nov., 68 (6): 1111-1117.
- Kubota, N.; Matthews, M.A.; Takahagi, T. and Kliewer, W.M. (2000):** Budbreak with garlic preparations: effects of garlic preparations and of calcium and hydrogen cyanamides on budbreak of grapevines grown in greenhouses. *American Journal of Enology and Viticulture*. 2000, 51: 4, 409-414.
- Kuden, A.B.; Son-L; Kuden, A. and Dennis-FG (1997):** Dormancy breaking experiments on apricot . *Acta. Horticulture*. No. 441, 153-157.
- Luna, V.; Soriano, M.D.; Bottini, R.; Sheng, C.; and R.P. Pharis (1993):** Levels of endogenous gibberellins, abscisic acid, Indole 3 acetic acid and naringenin during dormancy of peach flower buds. *Acta Hort.*, 329: 165 -267.
- Mann, S. S. and B. Singh (1990):** Some aspects on development physiology of patharankh pear. *Acta. Hort.*, 279: 155-158.
- Mapson, L.W. (1970):** Vitamins in fruits. In *The Biochemistry of fruits and their production*. Vol. 1, Ac Hulme. Ed . Academic press. New York.
- Mengel, K. and Kirkby, E.A. (1987):** *Principal of plant nutrition* .4th Ed. International potash Institute. Pern, Switzerland ,pp.687.
- Nashaat, E.M.A. (1996):** Bud break, yield, fruit quality and some endogenous compounds of flame seedless grape vines and sultani fig trees in relation of dormex spray. M. Sc. Thesis, Univ. of Alexandria.
- Page, A. I.; Miller, R. H. and Keeny, D. R. (1982):** "Methods of Soil Analysis". Part II. Chemical and Microbiological Methods. 2nd Ed. Amer. Soc. Agron., Madison, Wisconsin, USA.
- Rajagopal, V. and I. M. Rao (1974):** Changes in the endogenous level of auxin and gibberellin like substances in the shoot apices of N- deficient tomato plant. *Aust. J. Bot.*, 22: 429 – 435.
- Ramina, A.; Colauzzi, M.; Masia, A.; Pitacco, A.; Caruso, T.; Messina, R. and G. Scalabrelli (1995):** Hormonal and climatological aspects of dormancy in peach buds. *Acta Hort*. 395: 35-45.
- Salisbury, F. B. and C.W. Ross (1992):** The photosynthesis, transpiration compromise. In: *Plant Physiology*, pp. 66-92. Wadsworth Pub. Comp. California USA.
- Skene, K. G. M. (1969):** A comparison of the effects of cycocell and tipping on fruit set *Vitis vinefra* L. *Aust. J. Bio. Sci.* , 22: 1305 – 1311.
- Subha-drabandhu, S. (1995):** Induction of bud break in apple trees that received insufficient chilling by hydrogen cyanamide. *Acta Hort.*, 409: 171-178.

- Tromp, J. (1970):** Storage and mobilization of nitrogenous compounds in apple trees with special reference to arginine. In physiology of trees crops., Eds. L.C. Luckwill and C.V. cutting, 14: 3-59. Academic Press, Londone and New York.
- Welburn, A.R. and Lichtenthaler, H. (1984):** Formula and program to determine total carotenoids and chlorophyll a and b of leaf extracts different solvents. In Advances in photosynthesis Research (Sybesma C.Ed.) Vol., II pp. 9-12, Mortinus Njihoff Dr. W. Junk publishers, the Hague
- Wilde, S.A.; Corey, R.B.; Lyer, J.J. and Voigt, G.K. (1985):** Soil and Plant Analysis For Tree Culture, 3rd Ed. Oxford IBLT Publishing Co., New Delhi: 9-100

تأثير نترات البوتاسيوم ومستخلص كل من الثوم والبصل على تفتح البراعم والنمو والمحصول والمحتوى الكيماوي لأشجار التفاح

محمد السيد مرسى* ومحمد احمد سيف اليزل**

* قسم البساتين - كلية الزراعة - جامعة الفيوم - مصر

** قسم النبات الزراعى - كلية الزراعة - جامعة الفيوم - مصر

أجرى هذا البحث خلال موسمين متتاليين ٢٠٠٦ و ٢٠٠٧م فى تربة رملية طميية بمزرعة التجارب بكلية الزراعة بالفيوم لدراسة تأثير رش النباتات بنترات البوتاسيوم ومستخلص الثوم والبصل على تفتح البراعم والنمو والمحصول والمحتوى الكيماوي لأشجار التفاح. تم رش الأشجار بستة معاملات وهى (نترات البوتاسيوم ١٠% ومستخلص الثوم ٢٠% ومستخلص البصل ٢٠% و نترات البوتاسيوم ٥%+ مستخلص الثوم ١٠% و نترات البوتاسيوم ٥%+ مستخلص البصل ١٠%) بالإضافة الى الكونترول (الرش بالماء فقط). أوضحت النتائج المتحصل عليها بصفة عامة أن كل صفات النمو (ميعاد تفتح البراعم الزهرية ونسبة تفتح البراعم) وايضا المحصول/شجرة و صفات المحصول (نسبة العقد ووزن الثمرة وحجم الثمرة وعدد الثمار) والمكونات الكيماوية للأوراق (الكلوروفيلات الكلية والكربوهيدرات الكلية والبروتينات) وبعض العناصر (نتروجين - فسفور - بوتاسيوم) وأيضا محتوى الثمار من المواد الصلبة الذاتية الكلية ونسبة المواد الصلبة الذاتية الكلية/ الحموضة وفيتامين ج والكربوهيدرات الكلية والسكريات الكلية والمختزلة والأحماض الأمينية الحرة الكلية قد زادت نتيجة الرش بالمعاملات المختلفة، وتم الحصول على أعلى القيم للقياسات السابقة عند استخدام نترات البوتاسيوم ٥% + مستخلص البصل ١٠% أو نترات البوتاسيوم ٥% + مستخلص الثوم ١٠% أو نترات البوتاسيوم ١٠% وعلى العكس من ذلك أدى الرش باستخدام نفس المعاملات إلى تناقص محتوى الثمار من الحموضة الكلية والفينولات الكلية مقارنة بالكنترول. وهكذا نوصى باستخدام نترات البوتاسيوم او مستخلص البصل او مستخلص الثوم سواء بصورة منفردة او متحدة مع نترات البوتاسيوم رشا على الاشجار لتحسين تفتح البراعم والنمو الخضري والمحصول والمكونات الكيماوية لأشجار التفاح.