

Characterization and Pharmacological Activities of an Isolated Compound from Alcoholic Extract of *Desmostachya bipinnata* Underground Parts

Anupama Singh^{1,2,*}, Vikas Anand Saharan² and Anil Bhandari³

¹ Faculty of Pharmaceutical Sciences, Jodhpur National University, Boranada, Jodhpur, Rajasthan, India.

² Sardar BhagwanSingh Postgraduate Institute of Biomedical Sciences & Research, Balawala, Dehradun, Uttarakhand, India.

³ Goenka College of Pharmacy, Lachhmangarh, Sikar, Rajasthan, India.

Received: 5 Jan. 2016, Revised: 12 Feb. 2016, Accepted: 15 Feb. 2016.

Published online: 1 May 2016.

Abstract: *Desmostachya bipinnata* (L.) Stapf [Gramineae (Poaceae)] is used in Indian traditional medicines system for treatment of various diseases such as asthma, kidney stone, diarrhea, wound healing, etc. Some of these traditional claims require authentication by performing animal testing. In present study, *D. bipinnata* was extracted in alcohol. An alkaloid was isolated from the prepared extract, characterized and evaluated for antidiarrheal and antiasthmatic activity. Isolated compound was characterized by melting point, chemical tests, FT-IR, LC-MS and NMR. Antidiarrheal activity of the isolated compound was evaluated by castor oil and magnesium sulphate induced diarrhea in Wistar rats. Castor oil and magnesium sulphate induced diarrhea in Wistar rats were treated with prepared *D. bipinnata* alcoholic extract, isolated compound and loperamide. Antiasthmatic activity of the isolated compound was evaluated by isolated guinea pig ileum and histamine induced lethality test. The isolated Compound I was 4-(2-(dimethylamino) ethyl) phenol. Isolated Compound I was 70.16% (P<0.01) effective in reducing feces in castor oil induced diarrheal rats and 70.94% (P<0.01) feces reduction in magnesium sulphate induced diarrhea model. Loperamide, a standard antidiarrheal drug, was effective in reducing number of feces by 70.94% in castor oil induced diarrheal rats and 71.77% reduction in feces in magnesium sulphate induced diarrheal rats. Isolated compound I showed 97.67% contraction, at 0.4 ml dose, where as chlorpheniramine showed 100% contraction, at 0.8 ml dose in isolated guinea pig ileum model. Isolated compound I and cetirizine, both, showed 100% protection against histamine induced lethality with no grasping and difficulty in breathing. The isolated compound was pharmacologically active as antidiarrheal and antiasthmatic in tested animals.

Keywords: *Desmostachya bipinnata*, diarrhea, asthma, castor oil, magnesium sulphate, rats.

1 Introduction

Desmostachya bipinnata (L.) Stapf [Gramineae (Poaceae)] (Syn: *Eragrostis cynosuroides*) is commonly known as sacrificial grass, *kusha* [1], *drabh* [2] and *dab* [3]. *D. bipinnata* contains flavonoids like kaempferol, quercetin, quercetin-3-glucoside, trycin, trycin-7-glucoside; coumarins like scopoletin and umbelliferone; sugars; amino acids; and carbohydrates [4]. Camphene, β -eudesmol, eseroline, and calarene are the main components of the *D. bipinnata* oil. Other phytoconstituents like diphenyliodonium bromide, limenone, 2-cyclohexene-1-one, and 8-nitro-12-tridecanolide have also been isolated in small amounts from *D. bipinnata* [5]. Leaf paste of *D. bipinnata* is used to cure cuts and wounds [6]. Roots are used in treatment of asthma, rheumatism [7], carbuncles, piles, cholera, dysuria [2], diuretic, galactagogue, astringent [1], dysentery, leucorrhoea and wounds [1].

Herbal treatment for diarrhea and asthma in natural and traditional medicinal practices includes use of plants or plant extracts such as *Semicarpus anacardium*, *Achyranthus aspera*, *Rhus semialata* [8] *D. bipinnata* [7], *Elytraria acaulis* [9, 10]. Antidiarrheal activity of *D. bipinnata* has been investigated for alcoholic and aqueous extracts of *D. bipinnata* [11] by castor oil induced diarrhea model in Wistar rats and charcoal meal stimulated gastrointestinal transit in albino mice. Literature review revealed that antiasthmatic activity of *D. bipinnata* has not yet been investigated. Furthermore, active constituents responsible for antiasthmatic and antidiarrheal activity have not been isolated from *D. bipinnata*.

The present study was carried out to identify active constituents responsible for traditional claims for antidiarrheal and antiasthmatic activity of *D. bipinnata*. Extract of *D. bipinnata* were prepared in alcohol. An isolated compound and prepared extract were evaluated for antidiarrheal activity by castor oil induced diarrhea and

magnesium sulphate induced diarrhea models in Wistar rats. Isolated guinea pig ileum and histamine induced lethality tests in guinea pig were conducted to evaluate antiasthmatic activity.

2 Materials and methods

2.1 Plant material: collection, authentication

The whole plants of *D. bipinnata* were collected from Chirawa and Jhunjhunu districts of Rajasthan, India. The collected plant was authenticated by Dr. R. P. Pandey from Botanical Survey of India, Jodhpur, India. A voucher specimen, JNU/PH/2010/D_b D₂, was deposited in the herbarium of Jodhpur National University, Jodhpur, India.

2.2 Extraction and isolation

Underground parts of *D. bipinnata* were grinded in electric mixer-grinder and screened using BSS standard sieves. Powder which passed through sieve no. 22 (average aperture size 710 μm) and retained on sieve no. 44 (average aperture size 355 μm) was selected and used for extraction. The powdered crude drug was packed in a paper cylinder made from a filter paper and placed in the body of soxhlet extractor. The solvent was poured in soxhlet extractor and allowed to run for 3-4 cycles. After that, the apparatus was fitted in appropriate manner and drug was extracted. The obtained extracts were filtered through Whatman filter paper, concentrated to syrupy mass, acidified with dilute sulphuric acid and filtered. Filtrate was decolorized with charcoal, made slightly alkaline by adding sodium carbonate and extracted with petroleum ether in separating funnel. On evaporation of ethereal extract, needle shaped crystals were obtained. These crystals of the isolated compound were further purified by recrystallization in boiling water.

2.3. Characterization of isolated compound

2.3.1. Melting point determination

Melting point of isolated compound was determined by melting in a glass capillary. Isolated compound was filled in a glass capillary tube fused at one end and heated in a melting point apparatus. The melting point was determined as the temperature when the isolated compound melted in the capillary tube.

2.3.2. Xanthoproteic test

Isolated compound was treated with concentrated nitric acid, a white precipitate was observed. It was boiled and cooled. Further to it, sodium hydroxide (20%) or ammonia was added. Orange color developed in solution indicated the presence of aromatic amino acids [12].

2.3.3 Million's test

Isolated compound was treated with Millon's reagent. The formation of white precipitate, which turned to red upon heating, indicated the presence of proteins and amino acids [12].

2.3.4 Ferric chloride test

Isolated compound was treated with ferric chloride. Reddish blue coloration was observed [12].

2.3.5. Thin layer chromatography

Thin layer chromatographic plates of isolated compound were prepared using solvent system, n-butanol: acetic acid: water (60:15:25). Iodine and Dragendroff's reagent were used as detecting reagents.

2.3.6 IR spectroscopy

KBr disk of Compound I was prepared and IR spectrum of Compound I was recorded using Perkin Elmer 1600 FTIR (USA) spectrophotometer.

2.3.7 LC-MS spectroscopy

The mass spectrum of Compound I was recorded using TOF MS ES micromass spectrometer.

2.3.8. NMR spectroscopy

The ^1H and ^{13}C NMR spectrum of Compound I was recorded using Bruker Advance II 400 NMR spectrometer in deuterium substituted water (D_2O) using tetramethylsilane (TMS) as internal standard (chemical shifts in δ ppm).

2.4. Pharmacological evaluation of isolated compound

2.4.1 Antidiarrheal activity

Magnesium sulphate induced diarrhea and castor oil induced diarrhea models in rats models were used for evaluation of antidiarrheal activity.

2.4.1.1 Animals

The study was conducted on healthy female Wistar rats of either sex weighing 150-200 g. The animals were randomly distributed to 7 different groups, each consisting of 5 animals. Prior approval by Institutional Animal Ethical Committee (IAEC), registration number 1258/ac/09/CPCSEA, was obtained for conduct of animal experiments. The animals were kept in colony cages at standard husbandry conditions. All animals had free access

to feed and water *ad libitum*.

2.4.1.2. Experimental procedure

Healthy Wistar rats were marked and randomly distributed to 4 groups, each consisting of 5 animals. The groups were given treatments as follows:

Group I: Normal control (1% CMC; 10 ml/kg, body weight)

Group II: Standard drug (loperamide 3 mg/kg body weight)

Group III: Diarrheal rats, treated with alcoholic extract of *D. bipinnata*

Group IV: Diarrheal rats, treated with isolated Compound I

All animals were initially screened for induction of diarrhea by administering 1 ml of castor oil.

Only animals which developed diarrhea were selected for antidiarrheal studies.

2.4.1.3. Castor oil induced and magnesium sulphate induced diarrhea in rats

Wistar rats weighing 150-200 g were selected and kept for overnight fasting. Loperamide, *D. bipinnata* extract (500 mg/kg body weight) and isolated Compound I were administered orally by gavage.

For castor oil induced diarrhea, 1 ml of castor oil was administered orally to each animal after one hour after administration of drug/extract. For magnesium sulphate induced diarrhea, magnesium sulphate was administered, at dose 2 g/kg orally to each animal, thirty minutes after administration of drug/extract. All animals were placed in cages, where floor was lined with non-wetting paper sheets of uniform weight. Non-wetting paper sheets were changed per hour up to 6 hour. Characteristic diarrheal droppings per hour up to 6th hour were recorded after draining the urine by gravity.

A numerical score based on stool consistency was assigned. Normal stools were assigned a score 1, semi solid stools as 2 and watery stools as 3. Mean of diarrheal droppings passed by treated animals in various groups were compared to control group [13-15].

2.4.1.4. Statistical analysis

The data obtained in the studies was analyzed by one way analysis of variance (ANOVA) followed by Dunnett's 't' test using GraphPad Prism Version 5.01 (Graph Pad Software Inc., U.S.A) to compare different groups with control. P-value <0.01 was considered significant and results were expressed as mean \pm SD.

2.4.2. Antiasthmatic activity

2.4.2.1. Isolated guinea pig ileum

Isolated guinea pig ileum was used for evaluating contractile responses of test compound and extract. Concentration response curves of various drugs have been prepared and compared to evaluate antiasthmatic activity [15, 16]. Overnight fasted guinea pigs were sacrificed, dissected to cut and remove ileum. Ileum was mounted in an organ bath containing tyrode solution continuously aerated at 37 \pm 0.5°C. Concentration response curves for histamine, chlorpheniramine with histamine, drug extract with histamine and isolated compound with histamine in Tyrode solution were plotted.

2.4.2.2. Histamine induced lethality in guinea pig

2.4.2.2.1. Animals

The study was conducted on healthy guinea pigs of either sex weighing 200-300 g. Prior approval by IAEC (registration number 1258/ac/09/CPCSEA) was obtained for conduct of experiments. The animals were kept in colony cages at standard husbandry conditions and had free access to feed and water *ad libitum*.

2.4.2.2.2. Experimental procedure

Healthy guinea pigs were marked and randomly distributed to 4 groups, each consisting of 5 animals. These animal groups were given treatments as follows:

Group I: Normal control

Group II: Standard drug (Cetirizine 0.5 mg/kg)

Group III: Asthmatic guinea pigs, treated with alcoholic extract of *D. bipinnata*.

Group IV: Asthmatic guinea pigs, treated with isolated Compound I

2.4.2.2.3. Induction of asthma and its treatment

Asthma was induced in guinea pigs as per standard procedure [17]. The plant extracts were dissolved in distilled water and administered by oral feeding needle. After one hour of treatment, fur was removed from either left or right leg of guinea pigs with help of blade and area was cleaned with warm cotton swab to dilate pedal vein. After that histamine hydrochloride solution (1.25 mg/kg body weight) was injected intravenously to each animal. Animals in each group were observed for convulsions and lethality up to one hour after histamine injection.

3 Results and discussion

Our previous studies reported the presence of alkaloids, flavonoids, steroids, glycosides, and coumarins in alcoholic extract of *D. bipinnata* [18].

3.1. Extraction and isolation

D. bipinnata underground parts (480 g) were extracted with alcohol from which 115 mg of alkaloid was obtained after isolation.

3.2. Characterization of isolated compound

Melting point of the isolated Compound I was 117-118°C as determined by melting a glass capillary tube. It gave positive xanthoproteic test, Millon's test and a reddish blue color with ferric chloride. TLC of isolated Compound I in solvent system n-butanol: acetic acid: water (60:15:25) gave R_f value 0.44 (Figure 1). IR spectrum is presented in Figure 2. IR spectrum had shown characteristic peaks at 1313 cm^{-1} (C-N stretch); 2797 and 2856 cm^{-1} (C-H stretch); 1464, 1514, 1589, and 1612 cm^{-1} (C=C stretch); 818 and 841 cm^{-1} (C-H out of plane bending, p-di-substituted benzene).



Figure 1. TLC of alcoholic extract of *D. bipinnata* and isolated Compound I




Figure 2. IR spectrum of isolated Compound I

The mass spectrum of compound I was recorded on TOF

MS ES micromass spectrometer. The m/z for compound I was found to be 165 ($\text{C}_{10}\text{H}_{14}\text{ON}^+$), 121 ($\text{C}_8\text{H}_9\text{O}^+$), 103 (C_8H_7^+), 93 ($\text{C}_6\text{H}_5\text{O}^+$), and 77 (C_6H_5^+). The mass spectrum is shown in Figure 3(a) and 3(b).




Figure 3(a). Mass spectrum of isolated Compound I




Figure 3(b). Mass spectrum of isolated Compound I

^1H NMR and ^{13}C NMR spectrum of compound I was recorded on Bruker Avance II 400 NMR spectrometer in deuterium substituted water (D_2O) using TMS as internal standard (Chemical shifts in δ ppm).

The observed NMR bands for different functional groups in the Compound I in ^1H NMR and ^{13}C NMR spectra are shown in Figure 4 and Figure 5, respectively. ^1H NMR bands for different functional groups in the isolated compound I were δ 4.70 (s, 1H, -OH group), δ 7.01- 6.99 (d, J 8.4, 2H, -Ar-H group), δ 6.64- 6.62 (d, J 8.8, 2H, -Ar-H group), δ 2.91- 2.71 (m, 4H, -N- CH_2 group), δ 2.5 (s, 6H, - CH_3 group). ^{13}C NMR bands for different functional groups in the isolated compound I were 158.26 δ (s, -Ar-H group), 132.26 δ (s, -Ar-H group), 129.97 – 126.77 δ (m, -Ar-H group), 116.88 δ (s, -Ar-H group), 59.35 δ (s, - CH_2 -N group), 42.99 δ (s, N- CH_3 group), 30.18 δ (s, -C- CH_2

group).




Figure 4. ^1H NMR spectrum of isolated Compound I




Figure 5. ^{13}C NMR spectrum of isolated Compound I

Compound I showed positive Xanthoproteic test and Millon's test. Also, water soluble phenols were indicated by positive ferric chloride test. IR, ^1H and ^{13}C NMR, mass interpretation results confirmed that isolated Compound I was 4-(2-(dimethylamino) ethyl phenol having molecular weight 165.23 and molecular formula $\text{C}_{10}\text{H}_{15}\text{NO}$.

3.3. Pharmacological evaluation of isolated compound

3.3.1. Antidiarrheal activity

Alcoholic extract successfully inhibited diarrhea in both models, *i.e.*, the castor oil and magnesium sulphate induced diarrhea. The antidiarrheal effect was evident from the

reduction of total number of feces in the test groups, which may be due to flavonoids, glycosides, tannins and alkaloids present in alcoholic extract [18]

3.3.1.1. Castor oil induced diarrhea in rats

In castor oil induced diarrhea model, the *D. bipinnata* extract in alcohol and isolated Compound I showed antidiarrheal effect in Wistar rats. Loperamide, being standard anti-diarrheal drug, was most effective in reducing number of feces by 70.94%, while among studied extracts alcoholic extract was found most effective which reduced the number of feces by 61.54%. Isolated Compound I was found effective as antidiarrheal agent and reduced the number of feces by 69.23%. All the tested extracts significantly ($P < 0.01$) reduced the total number of feces when compared to control group by ANOVA followed to Dunnett's 't' test (Table 1).

Table 1. Effect of *D. bipinnata* underground parts alcoholic extract and isolated Compound I on castor oil induced diarrhea in rats (n=5)

Treatment	Dose (mg/kg, p.o.)	Mean no. of feces in 6 h	Feces Reduction (%)
Alcoholic extract	500	9.0±1.00**	61.54
Compound I	3	7.2±0.84**	69.23
Loperamide	3	6.8±0.84**	70.94
Control	10 [#]	23.4±2.07	

** Significant difference at $P < 0.01$ vs. control and $P < 0.001$ vs. control; one-way ANOVA followed by Dunnett's 't' test

*Significant difference at $P < 0.01$ vs control; No significant difference at $P < 0.001$ vs. control; one-way ANOVA followed by Dunnett's 't' test

In ml/kg

Castor oil hydrolysis produces ricinoleic acid which induces diarrhea as hypersecretory response due to changes in the transport of water and electrolytes [19, 20]. Ricinoleic acid causes irritation and inflammation of gastric mucosa resulting in release of prostaglandins causing stimulation of secretion [21, 22]. Furthermore, ricinoleic acid also sensitizes intramural neurons of the gut. Several other mechanisms have also been reported to explain the diarrheal effect of castor oil which are adenylate cyclase activation or cAMP mediated active secretion [23] and inhibition of Na^+ , K^+ -ATPase activity [24].

3.3.1.2. Magnesium sulphate induced diarrhea model

In magnesium sulphate induced diarrhea model, the extract of *D. bipinnata* and isolated compound I showed antidiarrheal effect in Wistar rats (Table 2). Alcoholic extract of *D. bipinnata* showed 64.52% reduction in feces, which outperform slightly to the effect of standard antidiarrheal drug loperamide with 71.77% reduction in

feces. Isolated Compound I was effective in preventing diarrhea and reduced the number of feces by 70.16%. All extracts showed significant ($P < 0.01$) antidiarrheal effect [18] in reducing the total number of feces, when compared to control using by ANOVA and followed to Dunnett 't' test.

Table 2. Effect of *D. bipinnata* underground parts alcoholic extract and isolated Compound I on magnesium sulphate induced diarrhea in rats (n=5)

Treatment	Dose (mg/kg, p.o.)	Mean no. of feces in 6 h	Feces Reduction (%)
Alcohol	500	8.8±1.10**	64.52
Compound I	3	7.4 ±0.55**	70.16
Loperamide	3	7.0±0.71**	71.77
Control	10 [#]	24.8±1.92	

** Significant difference at $P < 0.01$ vs. control and $P < 0.001$ vs. control; one-way ANOVA followed by Dunnett's 't' test

*Significant difference at $P < 0.01$ vs control; No significant difference at $P < 0.001$ vs. control; one-way ANOVA followed by Dunnett's 't' test

In ml/kg

Diarrhea in rats is also induced by administration of oral magnesium sulphate which increases the accumulation of fluid in the intestinal lumen and enhances flow from the proximal to distal intestine. This mechanism also involves release of NO, probably through stimulation of the constitutive form of NO synthase [25]. Magnesium sulphate has also been reported to liberate cholecystokinin from duodenal mucosa resulting in increase of small intestine secretions and motility and thus preventing the reabsorption of water and sodium chloride [26, 27].

Alkaloids, flavonoids, and steroids, in combination, exists in various plant extracts which possess anti-diarrhoeal activity, viz. root extract of *Asparagus racemosus* (Liliaceae), leaf extract of *Clerodendrum phlomidis* (Verbenaceae), stem bark extract of *Cylicodiscus gabunensis* (Mimosaceae), leaf extract of *Emilia coccinea* (Asteraceae) and fruit extract of *Momordica cymbalaria* (Cucurbitaceae) [28]. However, it was also observed that root extracts of *Guiera senegalensis* (Combretaceae) containing alkaloid, flavonoid and glycosides are effective in controlling diarrhea [28]. Stem bark extracts of *Butea monosperma* (Fabaceae) contains glycosides, flavonoids and steroids, which are effective in curing diarrhea [28]. It was also observed that plants containing alkaloids, flavanoids, glycosides, and steroids, viz. stem bark extract of *Annona senegalensis* (Annonaceae), root extract of *Combretum sericeum* (Combretaceae) and leaf extract of *Dalbergia sissoo* (Fabaceae), are also effective in diarrheal

conditions [28].

Isolated Compound I, 4-(2-(dimethylamino) ethyl) phenol, is an alkaloid which showed antidiarrheal effect comparable to standard loperamide in both castor oil induced diarrhea and magnesium induced diarrhea model. Flavonoids and alkaloids have been reported to inhibit prostaglandins and autacoids release resulting in reduction of motility and secretion [28, 29]. Therefore, it is proposed that isolated Compound I acts by inhibition of prostaglandins and autacoids release resulting in reduction of motility and secretion.

3.3.2 Antiasthmatic activity

Standard drug chlorpheniramine showed 100% and alcoholic extract showed 97.72% contraction, at 0.8 ml dose (Figure 6 and Table 3). Preliminary phytochemical screening revealed the presence of alkaloids, flavanoids, steroids, glycosides, and coumarins in alcoholic extract of *D. bipinnata* [18]. Chlorpheniramine and alcoholic extract showed similar contractile responses. But, isolated compound I showed 97.67% contraction, at 0.4 ml dose (Figure 7), whereas chlorpheniramine showed 100% contraction, at 0.8 ml dose.




Figure 6. Concentration response curve of histamine and histamine with chlorpheniramine.




Figure 7. Concentration response curve of histamine and histamine with isolated Compound I.

Cetirizine (Standard drug), alcoholic extract and isolated

Compound I, showed 100% protection against histamine induced lethality with no grasping (Table 4). Therefore, it may be concluded that isolated Compound I, isolated from alcoholic extract, possess anti-histaminic activity and thus isolated Compound I may be pharmacologically active and potentially useful in humans for prevention of asthma.

Table 3. Concentration response curve of *D. bipinnata* underground parts alcoholic extract and isolated Compound I in isolated guinea pig ileum.

Histamine dose	Contraction (%)		
	Chlorpheniramine	Alcoholic extract	Isolated Compound I
0.05 ml	31.7	-	-
0.1 ml	63.41	-	30.23
0.2 ml	75.6	-	55.81
0.3 ml	-	-	83.72
0.4 ml	95.12	27.27	97.67
0.6 ml	97.56	-	-
0.8 ml	100	97.72	-

Table 4. *D. bipinnata* underground parts alcoholic extract and isolated Compound I effect in histamine induced lethality in guinea pigs (n=5).

Treatment	Body weight g (Avg. \pm Std. dev.)	Dose (mg/kg body weight)	Recovery/Death	Lethality (%)	Grasping/Asthma
Standard (Cetirizine)	314.86 \pm 47.07	0.5	Recovered	0	No grasping observed
Control	270.90 \pm 34.90		Death	100	Grasping observed
Alcoholic	496.33 \pm 10.87	500	Recovered	0	No grasping observed
Isolated Compound I	378.21 \pm 18.15	0.5	Recovered	0	No grasping observed

Histamine is an important chemical indicator involved in many allergic reactions. Histamine causes contraction of airway smooth muscle. Consequently, the muscles surrounding the airways constrict causing bronchospasm and shortness of breath, a life threatening condition [17]. Guinea pigs are highly sensitive to histamine induced bronchospasm. Present study deals with screening of anti-asthmatic activity of *D. bipinnata* underground parts extract and isolated Compound I. Different agonists like acetyl choline, histamine, 5-hydroxytryptamine and bradykinin are responsible for contractile responses [30].

Bronchial asthma is characterized by increased airway reactivity to spasmogens. An initial event in asthma appears to be the release of inflammatory mediators (e.g. histamine, tryptase, leukotrienes and prostaglandins). Some of these mediators directly cause acute bronchoconstriction, airway hyper responsiveness and bronchial airway

inflammation. Spasmolytic drugs like β adrenergic agonists, xanthine derivatives and anticholinergics relax the airway smooth muscles and are thus used as quick relief medications in acute asthmatic attacks. β adrenergic agonists promote bronchodilation by direct stimulation of β adrenergic receptors in the airway smooth muscle, that lead to relaxation of bronchial smooth muscle by rapid decrease in airway resistance *in vivo*. [31, 32]

β adrenergic agonists activates adenylate cyclase resulting an increases in concentration of intracellular cyclic adenosine 3', 5'-monophosphate (cAMP) leading to activation of specific cAMP-dependent protein kinases that cause relaxation [33, 34]. Relaxation may also be due to inhibition of myosin phosphorylation. β adrenergic agonists reverse bronchoconstriction irrespective of the contractile agent. β adrenergic agonists prevent release of mediators from a number of inflammatory cells *in vitro* [33, 34].

Leaf extract of *Adhatoda vasica* (Acanthaceae), *Fagopyrum esculaentum* (Polygonaceae), seed oil of *Hordeum vulgare* (Poaceae), and whole plant extract of *Passiflora incarnate* (Passifloraceae), *Passiflora foetida* (Passifloraceae), *Lepidium sativum*, *Solanum brevistigma* (Solanaceae), *Solanum xanthocarpum* (Solanaceae), *Solanum trilobatum* (Solanaceae), *Tylophora indica* (Asclepiadaceae) are different plant extracts in which active constituents such as alkaloids have been proven for their antiasthmatic action and mechanism of action in these drugs is bronchodilation [8, 35, 36].

Isolated compound, an alkaloid, is pharmacologically active as antiasthmatic agent. The proposed mechanism for isolated Compound I appear similar to β agonist as it antagonizes histamine contractile responses of airway smooth muscle, resulting in relaxation in smooth muscles.

4 Conclusion

An isolated Compound I was isolated from alcoholic extract of *D. bipinnata*. The isolated Compound I was proteinaceous and aromatic in nature. The isolated compound I was confirmed by IR, ^1H and ^{13}C NMR, and mass as 4-(2-(dimethylamino) ethyl phenol with molecular weight 165.23 and molecular formula $\text{C}_{10}\text{H}_{15}\text{NO}$. Animal studies with isolated compound had proved the antiasthmatic and antidiarrheal activity of *D. bipinnata*.

Conflicts of Interest

Authors declare no conflicts of interest.

Acknowledgements

The authors express their thanks to Dr. R. P. Pandey from

Botanical Survey of India, Jodhpur, India, for authentication of the plant.

References

- [1] C. E. Khare. Indian Medicinal Plants An Illustrated Dictionary. Berlin/Heidelberg: Springer-Verlag; 2007.
- [2] R. Qureshi, G. Bhatt and R. Memon. Ethnomedicinal uses of herbs from northern part of Nara desert, Pakistan. Pak J Bot 2010; **42**: 839-851.
- [3] Praveen, B. Upadhyay, S. Roy and A. Kumar. Traditional uses of medicinal plants among the rural communities of churu district in the Thar Desert, India. J Ethnopharmacol 2007; **113**: 387-399.
- [4] A. Awaad, N. Mohamed, D. Maitland and G. Soliman. Anti-ulcerogenic activity of extract and some isolated flavonoids from *Desmostachya bipinnata* (L.) Stapf. Rec Nat Prod 2008; **2**: 76-82.
- [5] K. Kumar, Sharvane, J. Patel and R. Choudhary. Chemical composition and antimicrobial activity of the essential oil of *Desmostachya bipinnata* Linn. Int J Phytomed 2010; **2**: 436-439.
- [6] S. Katewa and A. Jain. Traditional fold herbal medicines. Udaipur, Jaipur: Apex Publishing House; 2006.
- [7] F. Ahmad, M. Khan, M. Ahmad, M. Zafar, T. Mahmood, A. Jabeen and S. Marwat. Ethnomedicinal uses of grasses in salt range region of northern Pakistan. J Med Plant Res 2010; **4**: 362-369.
- [8] A. Alexander, Ajazuddin, A. Singh and Swarna. Herbal drugs used for the treatment of asthma: an overview. Int J Cur Biomed Phar Res 2011; **1**: 67-79.
- [9] A. Jain, S. Katewa, P. Galav and P. Sharma. Medicinal plant diversity of Sitamata wildlife sanctuary, Rajasthan, India. J Ethnopharmacol 2005; **102**: 143-147.
- [10] S. Katewa and P. Galav. Additions to the traditional folk herbal medicines of shekhawati region of rajasthan. Ind J Traditional Medicine 2006; **5**: 494-500.
- [11] M. Hegde, K. Lakshman, K. Giriya, B. Kumar and V. Lakshmiprasanna. Assessment of antidiarrhoeal activity of *Desmostachya bipinnata* L. (Poaceae) root extracts. BolLatinoam Caribe Plant Med Aromat 2010; **9**: 312-318.
- [12] K. Khandelwal. Practical pharmacognosy. 13th Edn ed. Pune: Nirali prakashan; 2005.
- [13] J. Shilpi, Md. T-U-Rahman, S. Uddin, MD.S Alam, S. Sadhu and V. Seidel. Preliminary pharmacological screening of *Bixa orellana* L. leaves. Journal of Ethnopharmacology 2006; **108**: 264-271.
- [14] R. Akter, S. Hasan, M. M. Hossain, M. Jamila, M. E. Hoque and M. Rahman. In Vitro Antioxidant and In Vivo Antidiarrhoeal Activity of Hydromethanolic Extract of *Xanthium Indicum* Koenig. Leaves. European Journal of Scientific Research 2009; **33**: 305-312.
- [15] G. Vogel and W. Vogel. Drug discovery and evaluation. Berlin: Springer Verlag; 1998.
- [16] P. Pandit, A. Singh, A. Bafna, P. Kadam and M. Patil. Evaluation of antiasthmatic activity of *Curculigo orchoides* Gaertn. Rhizomes. Ind J Pharm Sci 2008; **Jul-Aug**: 440-444.
- [17] S. Kulkarni. Practical pharmacology and clinical pharmacy. 1st ed. Delhi: Vallabh Prakashan; 2008.
- [18] A. Singh, V. Saharan and A. Bhandari. Pharmacognostic standardization with various plant parts of *Desmostachya bipinnata* (L.) Stapf. Pharm Biol 2014, **52**, 298-307.
- [19] J. Fioramonti, C. J. E. Niemegeers and F. Awouters. Diarrhoea and antidiarrhoeal drugs. In: Y. Ruckebusch, P.-L. Toutain and G. D.Koritz, editors. Veterinary Pharmacology and Toxicology, Springer Netherlands; 1983; pp. 307-320 LA - English.
- [20] C. J. E. Niemegeers, F. Awouters and P. A. J. Janssen. The castor oil test in rats: An in vivo method to evaluate antipropulsive and antisecretory activity of antidiarrheals? Drug Dev. Res. 1984; **4**: 223-227.
- [21] T. S. Gaginella, J. J. Stewart, W. A. Olsen and P. Bass. Actions of ricinoleic acid and structurally related fatty acids on the gastrointestinal tract. II. Effects on water and electrolyte absorption in vitro. Journal of Pharmacology and Experimental Therapeutics 1975 **195** 355-361
- [22] J. Galvez, A. Zarzuelo, M. E. Crespo, M. D. Lorente, M. A. Ocete and J. Jimenez. Antidiarrhoeic Activity of Euphorbia hirta Extract and Isolation of an Active Flavonoid Constituent. Planta Med 1993; **59**: 333-336.
- [23] F. Capasso, N. Mascolo, A. A. Izzo and T. S. Gaginella. Dissociation of castor oil-induced diarrhoea and intestinal mucosal injury in rat: effect of NG-nitro-L-arginine methyl ester. British Journal of Pharmacology 1994; **113**: 1127-1130.
- [24] T. S. Gaginella and P. Bass. Laxatives: An update on mechanism of action. Life Sciences 1978; **23**: 1001-1009.
- [25] A. A. Izzo, T. S. Gaginella, N. Mascolo and F. Capasso. Nitric oxide as a mediator of the laxative action of magnesium sulphate. British Journal of Pharmacology 1994; **113**: 228-232.
- [26] J. Galvez, M. E. Crespo, J. Jimenez, A. Suarez and A. Zarzuelo (1993). Antidiarrhoeic activity of quercitrin in mice and rats. Journal of Pharmacy and Pharmacology 1993; **45**: 157-159.
- [27] M. A. Zavala, S. PÃ©rez, C. PÃ©rez, R. Vargas and R. M. Parez. Antidiarrhoeal activity of *Waltheria americana*, *Commelina coelestis* and *Alternanthera repens*. Journal of Ethnopharmacology 1998; **61**: 41-47.
- [28] R. Kumar, R. Sharma, K. Bairwa, R. Roy and A. Kumar. Pharmacological review on natural antidiarrhoeal agents. Der Pharma Chemica 2010; **2**: 66-93.
- [29] V. F. Veiga, L. Zunino, J. o. B. Calixto, M. L. Patitucci and Ã-C. Pinto. Phytochemical and antioedematogenic studies of commercial copaiba oils available in Brazil. Phytotherapy Research 2001; **15**: 476-480.

- [30] M. Ghosh. Fundamental of experimental pharmacology. 2nd ed. Calcutta: Scientific Book Agency; 1984.
- [31] D. Cockcroft. Mechanism of perennial allergic asthma. *The Lancet* 1983; **2**: 253-256.
- [32] A. Mehta and B. Agrawal. Investigation into the mechanism of action of *Moringa oleifera* for its anti-asthmatic activity. *Oriental Pharmacy and Experimental Medicine* 2008; **8**: 24-31.
- [33] R. G. Mali and A. S. Dhake. A review on herbal antiasthmatics. *Oriental Pharmacy and Experimental Medicine* 2011; **11**: 77-90.
- [34] M. Church and J. Hiroi. Inhibition of IgE-dependant histamine release from human dispersed lung mast cells by anti-allergic drugs and salbutamol. *British Journal of Pharmacology* 1987; **90**: 421-429.
- [35] M. Yaheya and M. Ismail. Antiasthmatic herbal drugs review. *International journal of pharmacy and pharmaceutical sciences* 2010; **2**: 28-29.
- [36] A. Huntley and E. Ernst. Herbal medicines for asthma: a systematic review. *Thorax* 2000; **55**: 925-929.
-