

IMPACT OF BICARBONATE CONTENT OF IRRIGATION WATER, FARMYARD MANURE AND SULPHUR ON WHEAT PRODUCTION AND AVAILABLE MICRONUTRIENTS

Alaga, M. D.* , R. S. Abdel-Aal **; A. H. Abd El-Hamid** and A. A. Mousa**

*Soil Conservation Dept., Desert Res. Centre Cairo, Egypt

**Soil Sci. Dept., Faculty of Agric., Moshtohor, Benha Univ., Egypt

ABSTRACT

This study was carried out on a calcareous sandy loam soil at Sudr Research Station, South Sinai for conserving soil from the bicarbonate hazard in saline irrigation water. Four bicarbonate levels; 2.5, 5, 10 and 15 meqL⁻¹ in saline irrigation water were applied. Also, three farmyard manure rates (0, 10, 20 m³ fed⁻¹) and sulphure rates (0, 250 and 500 kg fed⁻¹) were applied individual and/or in combined. The organic manure and sulphure had positively significant effects on yield. The results also show that the highest value of grain and straw yields were obtained from (HCO₃)₁OM₂S₂ but the best treatment from economic point of view was (HCO₃)₂OM₁S₂. The highest values of available Fe⁺⁺, Mn⁺⁺ and Zn⁺⁺ were found with OM₂S₂ under all bicarbonate concentrations in saline irrigation water.

Keywords: Calcareous sandy loam soil, saline irrigation water, FYM, sulphur, wheat production and available micronutrients.

INTRODUCTION

In Egypt, calcareous soils represent about 25% of the new reclaimed area. Because of the limiting water supply, the need of expanding agricultural lands had to use saline water in irrigation which caused soil deterioration in some areas.

Mengel *et al.*, (1984) interpreted the reduction in the yield of plant due to HCO₃ to one or to more of the following reasons (a) the adverse effect of HCO₃ on the plant metabolic processes, (b) the disturbance of the plant nutrient balance, (c) the excess of HCO₃ induced the dominance of CO₃⁻, ions which react with free soluble Zn in soil solution.

El-Shall *et al.*, (1987) found that sulphur was referred to its influence on reducing soil pH, improving soil structure and increasing the availability of certain nutrients.

Aich *et al.*, (1997) stated that grain yield of wheat was decreased with increasing salinity of irrigation water in all three soils. Application of decomposed cow dung decomposed straw, gypsum or lime increased grain yield.

Clarck *et al.*, (1999) mentioned that sorghum aerial plant and root production decreased with increasing salinity. Plant Ca, Sr, Mn and Cd levels increased with increasing salinity. In contrast, sorghum K, P, and S levels declined with increasing salinity.

The aim of the present work is to understand the interactions between bicarbonate, farmyard manure and sulphur.

MATERIALS AND METHODS

A field experiment was carried out at Sudr Research Station, South Sinai, in split-split design. The field was divided into plots (1/400 fed), 3 X 3.5 m. All treatments received constant rates, from NPK (80, 30 and 48 kg fed⁻¹) as N, P₂O₅ and K₂O, respectively as form ammonium nitrate, super phosphate and potassium sulphate respectively. Farm yard manure and sulphur were applied at rates of (0, 10 and 20 m³ fed⁻¹) from FYM and 0, 250 and 500 kg S fed⁻¹. The wheat seeds (Sakha 8) were cultivated through two seasons 1999/2000 and 2000/2001. Grain and straw yields were evaluated at harvesting. After harvesting of wheat plants, representative soil samples were taken from 0-15, 15 -30 and 30 - 45 cm depths of each replicate to determine the soil chemical and physical properties under investigation. O.M%, Soil pH, EC, soluble cations and anions also ESP were determined according to Jackson (1967) and Chapman and Pratt (1961).

Table 1: Some physical properties of the experimental soils at Ras Sudr.

Soil depth (cm)	Particle size distribution (%)				Texture class	Bulk Density (g cm ⁻³)	Field capacity (%)	Wilting point (%)	Av. water (%)	CaCO ₃ (%)
	Coarse Sand	Fine Sand	Silt	Clay						
0-15	39.45	39.69	8.46	12.40	Sandy loam	1.44	18.90	9.52	50.90	50.90
15-30	37.20	43.31	12.22	7.27	Sandy loam	1.53	17.80	10.48	61.56	61.56
30-45	37.40	42.22	13.10	7.28	Sandy loam	1.52	17.30	10.63	51.32	51.32

Table 2: Some chemical properties of the experimental soils at Ras Sudr.

Soil depth (cm)	pH	EC (dSm ⁻¹)	Soluble cations (meqL ⁻¹)			Soluble anions (meq L ⁻¹)				ESP
			Ca ⁺² + Mg ⁺²	Na ⁺	K ⁺	CO ₃ ⁼	HCO ₃ ⁻	Cl ⁻	SO ₄ ⁼	
0-15	7.4	14.3	97.71	79.80	0.01	--	3.54	133.47	50.90	13.60
15-30	7.4	13.60	84.00	82.54	0.01	--	6.80	1.70	61.56	14.92
30-45	7.4	11.70	84.88	58.75	1.80	--	5.16	74.97	51.32	10.75

Well water analyses were determined as previously mentioned in soil extracts and are given in Table 3.

Table 3: Analyses of the saline well water used for irrigation.

EC (dS m ⁻¹)	Soluble cations (meq L ⁻¹)			Soluble anions (meq L ⁻¹)				SAR
	Ca ⁺⁺ + Mg ⁺⁺	Na ⁺	K ⁺	CO ₃ ⁼	HCO ₃ ⁻	Cl ⁻	SO ₄ ⁼	
10.20	62.80	80.13	1.18	--	2.50	81.90	39.60	10.70

FYM analyses were determined according to Jackson (1967), Lindsay and Norvell (1978) and Frie *et al.*, (1964). The obtained data are showed in Table 4.

Table 4: Analyses of the applied farmyard manure.

EC (dSm ⁻¹)	pH	O. C. (%)	Water holding capacity (%)	Total N (%)	C/N (ratio)	Total P (%)	Total K (%)	Total sulphate (%)	Total micronutrients (mg kg ⁻¹)		
									Fe	Mn	Zn
34.9	7.3	20.3	180	1.52	13.4	0.92	1.48	1.24	212	127	98

Finally, the statistical analysis of the obtained data was done according to the method described by Gomez and Gomez (1984) using LSD to compare the mean values of treatments.

RESULTS AND DISCUSSION

Data in Tables 5 and 6 reveal that grain and straw yields of wheat (ton fed⁻¹) were affected by the interaction between bicarbonate, organic manure and sulphur application during the two seasons. Table 5 shows that the effect of organic manure and sulphur was the best in case of OM₂ S₂ through the concentration of 5 meqL⁻¹ bicarbonate, while the same effect was in case of 10 meqL⁻¹ HCO₃⁻ less than 5 meqL⁻¹ where, the bicarbonate increased the deteriorious effect, while the more pronounced effects of FYM and S applications were noticed in the 1st season.

Table 5: Grain yields of wheat (ton fed⁻¹) as affected by bicarbonate, organic manure and sulphur application during the two studied seasons.

Treatments	1 st season											
	Control			HCO ₃ ⁻ (5meq L ⁻¹)			HCO ₃ ⁻ (10meq L ⁻¹)			HCO ₃ ⁻ (15meq L ⁻¹)		
	OM ₀	OM ₁	OM ₂	OM ₀	OM ₁	OM ₂	OM ₀	OM ₁	OM ₂	OM ₀	OM ₁	OM ₂
S ₀	1.17	2.10	2.22	1.16	2.05	2.21	0.92	1.46	1.50	0.50	0.74	0.86
S ₁	1.51	2.20	2.36	1.49	2.17	2.35	0.98	1.47	1.62	0.55	0.88	0.93
S ₂	1.94	2.25	2.45	1.91	2.24	2.44	1.10	1.51	1.70	0.68	0.91	0.97
LSD at 0.05	0.28											
Treatments	2 nd season											
	OM ₀	OM ₁	OM ₂	OM ₀	OM ₁	OM ₂	OM ₀	OM ₁	OM ₂	OM ₀	OM ₁	OM ₂
	S ₀	0.80	0.84	0.85	0.51	0.62	0.71	0.50	0.52	0.71	0.38	0.39
S ₁	0.86	0.93	0.94	0.56	0.69	0.82	0.56	0.68	0.81	0.53	0.64	0.71
S ₂	0.94	1.04	1.26	0.64	0.75	1.11	0.63	0.72	0.86	0.60	0.70	0.77
LSD at 0.05	0.21											

Table 6: Straw yields of wheat (ton fed⁻¹) as affected by bicarbonate, organic manure and sulphur application during the two studied seasons.

Treatments	1 st season											
	Control			HCO ₃ ⁻ (5meq L ⁻¹)			HCO ₃ ⁻ (10meq L ⁻¹)			HCO ₃ ⁻ (15meq L ⁻¹)		
	OM ₀	OM ₁	OM ₂	OM ₀	OM ₁	OM ₂	OM ₀	OM ₁	OM ₂	OM ₀	OM ₁	OM ₂
S ₀	2.65	2.96	3.46	2.61	2.69	2.80	1.83	2.12	2.25	1.65	1.70	1.97
S ₁	3.63	4.27	4.62	2.74	3.46	4.15	2.12	2.24	2.28	1.68	2.06	2.10
S ₂	4.15	4.59	5.28	3.30	4.07	4.18	2.15	2.32	2.90	1.75	2.25	2.56
LSD at 0.05	0.78											
Treatments	2 nd season											
	OM ₀	OM ₁	OM ₂	OM ₀	OM ₁	OM ₂	OM ₀	OM ₁	OM ₂	OM ₀	OM ₁	OM ₂
	S ₀	1.76	1.90	2.79	1.52	1.78	1.91	1.35	1.51	1.61	1.10	1.15
S ₁	1.94	2.34	2.88	1.90	2.06	2.62	1.88	2.06	2.39	1.26	1.67	2.03
S ₂	2.40	2.71	2.98	2.27	2.59	2.78	2.18	2.54	2.64	1.56	2.24	2.34
LSD at 0.05	0.69											

The second season data (2000-2001) show that there is sharply decreasing trend was noticed in grains yield. This may be due to bicarbonate salt accumulation. The treatment of (HCO₃)₁ OM₂S₂ was the best treatment, but from economic point of view the best treatment was (HCO₃)₂OM₁S₂. The

bicarbonate concentration at 10 and 15 meqL⁻¹ had harmful negatively effect on the yield for both grains and straw.

The reduction in the yield of plant may be due to the following reasons, a) Adverse effect of HCO₃⁻ on plant metabolic processes; b) Disturbance of nutrients balance; and C) Excess of (HCO₃⁻) induce of CO₃⁻ ions, which react with free soluble Zn⁺⁺ in soil solution and induce chlorosis. Similar results were obtained by Mengle *et al.* (1984).

Sulphur was referred to its influence on reducing soil pH, improving soil structure and increasing the availability of certain plant nutrients (El-Shall *et al.*, 1987 and Aziz, 1998).

With respect to available micronutrients in soil, Figs 1 to 3 show that the highest values of available Fe, Mn and Zn were obtained in case of OM₂S₂ at all treatments of HCO₃⁻ in saline irrigation water. This may be due to the nutrients content of FYM and the different acids, which were produced from decomposition of FYM and may decrease soil pH. Sulphur application also may have a role in decrease soil pH and increasing micronutrients availability.




Fig. 1: Available Fe in (mg kg soil⁻¹) as affected by bicarbonate, organic manure and sulphur application.




Fig 2: Available Mn in (mg kg soil⁻¹) as affected by bicarbonate, organic manure and sulphur application.




Fig. 3: Available Zn in (mg kg soil⁻¹) as affected by bicarbonate, organic manure and sulphur application.

Conclusion

Grains and straw yield of wheat was the highest value in case of (HCO₃)₁ OM₂S₂ in the first and second seasons. The harmful negatively effect of HCO₃ ions was clear through the second season. This due to HCO₃ salt accumulation at 10 and 15 meqL⁻¹.

From the economic point of view the best treatment was (HCO₃)₂ OM₁S₂. Available micronutrients in soil raised with increasing FYM and S rates.

The highest values of available Fe, Mn and Zn were in case of OM₂S₂ at all treatments of HCO₃⁻.

REFERENCES

- Aich, A. C., A. H. M. Ahmed, and R. Mandal (1997). Impact of organic matter, lime and gypsum on grain yield of wheat in salt affected soil irrigated with different grades of brackish water. Journal of physiological research. 10: 1-2, 79-84
- Aziz, M. A., M. F. A. Sallam, A. M. El-Gendy and M. A. El-Moniem(1998). Effect of natural soil conditioners and irrigation conditions on some chemical properties of sandy soils of Inshas and cucumber yield. Egyptian Journal of Soil Science. 38:1-4, 377-411.
- Chapman, H. D and D. F. Pratt (1961). Methods of Analysis for Soil, Plants and Water . Univ. of California, Division of Agric. Science.
- Clarck, D. R., C. J. Green, V. G. Allen and C. P. Brown (1999) Effect of salinity in irrigation water on forage sorghum and soil chemical. Journal-of-plant-Nutrition. 22: (12), 1905-1920.
- El-Shall, A., M. Wassif., M. Hilal and I. El-Bagouri (1987). Response of barley to sulphur application in a calcareous soil under saline water. Desert-Institute-Bulletin, Egypt. 37, 1, 117-130.
- Frie, E; K. Peyer and E. shatg (1964). Determination of phosphorus by ascorbic acid. Schw landwirtschaft, Forschung Heft. 3.318:328
- Gomez, K. A. and A. A. Gomez (1984). "Statistical Procedures for Agricultural Research". 2nd Ed. John Wiley and Sons New York, pp. 680.Inc.
- Jackson, M. L. (1967). Soil Chemical Analysis Constable & Co., Ltd., London.
- Lindsay and Norvell, (1978). Development of a DTPA soil test for Fe, Mn, Zn and Cu. Soil Sci. Amer. Proc. 42:421-428.
- Mengel, K., M. Th. Breininger and W. Bubl (1984). Bicarbonate, the most important factor inducing iron chlorosis in Vine grapes on calcareous soil. Plant and Soil. 81: 333 – 344.

تأثير محتوى مياه الري من البيكربونات وسماد المزرعة والكبريت على انتاجية القمح وتيسر العناصر الصغرى

محمود ضياء الدين الأجه*, رافت سرور عبد العال**, على عبد الحميد** و عزمي عبد الله محمد موسى**

* قسم صيانة الأراضي - مركز بحوث الصحراء - القاهرة- مصر

** قسم الأراضي - كلية الزراعة بمشهر- جامعة بنها- مصر

أجريت تجربة حقلية في محطة بحوث رأس سدر جنوب سيناء حيث التربة جيرية رملية طميية وماء الري مالح خلال موسمي ٢٠٠٠/١٩٩٩ و ٢٠٠١/٢٠٠٠. المعاملات ٢,٥، ٥، ١٠، ١٥ و ١٥ ملليمكافى بيكربونات فى اللتر وسماد مزرعة FYM بالمعدلات صفر، ١٠ و ٢٠ متر مكعب للفدان. أيضا الكبريت المعدنى أضيف بالمعدلات ٢٥٠، ٥٠٠ و ٥٠٠ كجم للفدان وري التجربة بمعدل ٠,٥ متر مكعب للحوض (٤٠٠/١ فدان) والمحصول المنزرع قمح سخا ٨ واشتملت الدراسة على بعض التقديرات.

تتلخص فى الآتى :

١. إنتاجية الحبوب لمحصول القمح كانت أفضل معاملة OM_2S_2 (HCO_3)₀ لكن الأفضل من الناحية الاقتصادية كانت المعاملة OM_1S_2 (HCO_3)₁ حيث أن الفرق كان غير معنوي البيكربونات أكثر من المعاملة السابقة.
٢. إنتاجية القش : السماد العضوى والكبريت كان لهما تأثيرا معنويا وجيدا على إنتاجية القش حيث أفضل معاملة كانت OM_1S_2 (HCO_3)₀ لكن المعاملة الأفضل من الناحية الاقتصادية كانت OM_1S_2 (HCO_3)₁.
٣. المغذيات الصغرى الميسرة فى التربة (Fe, Mn and Zn) زادت مع زيادة السماد العضوى والكبريت وكانت أعلى قيم لهم فى معاملة OM_2S_2 تحت كل تركيزات البيكربونات فى ماء الري المالح .