

EFFECT OF SOME ORGANIC MANURES AND POTASSIUM ADDITION ON GROWTH, YIELD AND FRUIT QUALITY OF SULTANI FIG TREES

Gowda, A. M.

Olive and Semi – Arid Zone Fruits Res., Hort. Res. Instit., Agric. Res. Center, Giza, Egypt

ABSTRACT

An experiment was carried out in a private orchard situated at Nasser District, Beni-Sweif Governorate to investigate the effect of some organic manure sources (cattle, sheep and chicken manures) as soil and / or foliar application of potassium and their combinations on growth, yield and fruit quality of Sultani fig trees grown on clay loamy soil during two successive seasons of 2005 and 2006. Data revealed that, the highest records of trunk and canopy circumferences were obtained from the fertilized trees with sheep manure and cattle manure, while chicken manure application gave the tallest shoots. Moreover, the highest values of leaf area, fresh weight and number of leaves/ shoot were recorded from trees treated with sheep and chicken manures. Meantime, cattle manure gave the heaviest leaf dry weight. Trees received 750 gm K₂O in potassium sulphate form (48 % K₂O) plus foliar potassium (1%) gave the highest values in most of the studied growth parameters in both seasons.

The data also cleared that, Sultani fig trees fertilized with sheep and cattle manure gave the highest number of breba fruits/ tree, number of main fruits/ shoot and /tree, yield (kg/ tree), main fruit weight, diameter and height of fruit, TSS and total sugars. On the contrary, the same treatment gave the lowest values of total acidity and fruit moisture content. However, the application of 750 or 500 gm K₂O and foliar potassium 1% don't give significant differences on yield of Sultani fig trees .Whereas, the addition of 750 gm K₂O + foliar 1% potassium surpassed of the treatments for fruit quality characters (except for total acidity and fruit moisture content). Moreover, the results showed that sheep manure when supplemented with 750 gm K₂O and foliar 1% potassium gave the highest values for most of the studied characters in both seasons. As for Leaf nitrogen, phosphorus and potassium contents, sheep manure treatments increased leaf nitrogen and phosphorus contents, however chicken manure gave the highest content of potassium. The application of 500 gm K₂O + foliar 1% potassium gave the highest leaf nitrogen and phosphorus contents in both seasons.

It can be recommended that, sheep or cattle manure and potassium addition at 500 gm K₂O/ tree as soil application + foliar 1% potassium can give the highest yield with the best fruit quality under the same conditions of this study .

INTRODUCTION

Figs (*Ficus carica* Risso) are considered one of the most popular deciduous fruit trees in Egypt. Sultani fig cultivar is the most widely grown and considers the local standard cultivar. The total fig acreage in Egypt is estimated by 77.227 feddans and fig production attained 160.124 tons of fresh fruits (according to the statistics of the Ministry of Agricultural, 2004). A great attention has been focused on using organic fertilizers in fruit production in order to reduce plant and soil pollutions and its effect on human health. Organic manures improve the physical, chemical and biological

properties of nearly all soil types, adjusting soil Ph, increasing solubility and nutrients availability to plant consequently, influence the growth and production of the plants (Madison *et al.*, 1986; Mengel and Kirkby, 1987 and Zhou *et al.*, 2001). The using of animals manures will be monitored more closely, especially in terms of nutrient leaching on highly permeable soils (Ferguson,1994). The addition of organic manures to the soil encouraged proliferation of soil microorganisms, increased microbial population and activity of microbial enzymes i.e. dehydrogenase, urease and nitrogenase (Youssef *et al.*, 2001 and Abou-Hussein *et al.*, 2002). Some investigators studied the effect of organic manure application as compared with mineral fertilizers on different fruit crops (El-Adawy,1987; Mordogan, *et al.*, 2002; Caetano, *et al.*., 2006) on fig , (Abou- Taleb , 2004) on pecan, (Kassem & Marzouk, 2002) on grapevines (Foud *et al.*, 2002; Gamal & Ragab, 2003 ; Abd El-Naby, 2004 and Abd El- Naby *et al.*, 2004) on citrus and (Haggag, 1996) on olive trees. They reported that, under organic system soil biotic life increased as a result of the plant synthesis of more vitamins and amount of total sugars. Moreover, the addition of organic fertilizer is necessary for the best growth, greater yield and fruit quality when compared to mineral fertilizers. Potassium is one of the most important essential elements, when supplied to the soil . It plays a dominant role in the mineral nutrition of fruit crops. The need for potassium may be widespread because the fruits consume more potassium than any other nutrient element (Chapman, 1968 and Koo, 1985). Potassium availability in soils could also be influenced by moisture content, plant withdrawal rates, temperature and other factors (Koo, 1985). The objective of this study is to demonstrate the effect of different organic manure sources and potassium addition on tree growth, yield and fruit quality of Sultani fig trees.

MATERIALS AND METHODS

This study was conducted during 2005 and 2006 growing seasons on twenty years old Sultani fig trees grown in a private orchard situated at Nasser District, Beni-Sweif Governorate (15 kms North-East, Beni-Sweif city).The trees were planted in a square system of four meters apart and received the same horticultural management . They were vase trained to equal number of bearing units (20 per tree with 8 buds each) and grown on clay loamy soil under flood irrigation system. Physical and chemical properties of the surface soil (0.0 – 90 cm) were determined according to Wild *et al.*, (1985) and data are shown in Table (1).

A factorial experiment was carried out to investigate the response of Sultani fig trees to some organic manure sources and potassium fertilization addition during 2005 and 2006 seasons.

Organic manure treatments:

Cattle manure, sheep manure, chicken manure and control (without organic manure).The chemical analysis of the three used manures are shown in Table (2).

The three organic fertilizers were applied superficially and mixed into the root zone under the shedding of the tree canopy in mid–December of each season at the recommended basal dressing fertilizer for fig tree 300 gm N from mineral source as ammonium sulphate form (20.5 % N) and another 300 gm N from organic source, consequently, each tree received 600 gm N. The rates of application of organic manure sources and potassium fertilization addition are shown in Table (3).

Potassium fertilization treatments; Potassium treatments were applied as follows:

k1 : 500 gm K₂O / tree as soil application (1.042 kg potassium sulphate).

k2 : 750 gm K₂O /tree as soil application (1.563 kg potassium sulphate).

k3 : k1 + foliar spray with 1 % potassium.

k4 : k2 + foliar spray with 1 % potassium.

k5 : only foliar spray with 1 % potassium .

Table (1): Mechanical and chemical analysis of the experimental soil.

Parameters	Value
Mechanical analysis	
Sand %	22.22
Silt %	29.32
Clay %	48.46
Texture	Clay loam
Chemical analysis	
Organic matter	1.58
PH (1: 2.5 suspension)	7.78
E.C mmhos/cm 1: 2.5	1.04
Available nutrients	
N %	0.08
P ppm	15.20
K ppm	202.5
Fe ppm	33.0
Mn ppm	19.0
Zn ppm	6.0

Table (2): Analysis of the three selected manures.

Character	Cattle manure		Sheep manure		Chicken manure	
	1 st season	2 nd season	1 st season	2 nd season	1 st season	2 nd season
Weight of m ³ /kg	340	340	464	420	260	510
Humidity%	8.0	10.3	9.6	8.6	8.7	13.9
Organic matter%	39.47	71.51	72.32	74.95	36.76	52.3
Organic carbon%	22.89	41.48	41.55	43.47	21.32	24.51
N%	1.38	1.85	2.35	2.72	4.17	4.1
P%	0.68	0.94	1.02	1.09	0.73	1.02
K%	1.86	2.37	2.11	2.29	1.90	1.58
C:N ratio	19.4	22.4	18.8	16.5	5.1	8.4
Fe ppm	4174	4410	6645	6100	4296.6	4483.1
Mn ppm	327.8	299	242.3	259	212.5	257
Zn ppm	79.3	43	109	61	792.9	250

Table (3): Organic fertilizers and potassium application rates

Rates / Treatment	Cattle manure		Sheep manure		Chicken manure	
	1 st season	2 nd season	1 st season	2 nd season	1 st season	2 nd season
Organic manure rates as the basal of 300 gm N / tree	21.8	16.2	12.8	11.0	7.2	7.4
Potassium in the rate of organic source (gm K ₂ O /tree)	405.5	383.9	270.1	251.9	136.8	116.9
Supplementary of potassium treatments from potassium sulphate (48% K₂O)						
500 gm K ₂ O / tree	94.5	116.1	229.9	248.1	363.2	383.1
750 gm K ₂ O / tree	344.5	366.1	479.9	498.1	613.2	633.1

Potassium soil application was in two equal doses during May and August of each year, while potassium addition as foliar application was added at three times May, June and July from a compound contains 36 % potassium. The applied rates of potassium were calculated on the basal of potassium content in organic manure application rate as shown in Table (3).

Experimental layout:

Sixty homogenous Sultani fig trees were chosen and arranged in a factorial experiment split plot design of twenty treatments, three replicates /each and represented with one tree. The main is organic manure sources (control, cattle manure, sheep manure and chicken manure) and the sub plot is potassium treatments (500, 750 gm K₂O / tree and /or foliar potassium 1 %).

This study was evaluated through the following measurements:**I- Growth parameters.**

1-Tree characteristics: Trunk circumference for each tree was measured at the beginning and the end of the experiment in the two seasons of study, the net increase of trunk circumference (cm) was calculated. Canopy circumference (m) was measured. Shoot length (cm) was recorded at November of each season on ten shoots per tree which tagged at random for all tree directions at the beginning of each growing season.

2- Leaf characteristics: Leaf area (cm²) was measured by planimeter on 20 mature leaves from each tree in August , cleaned to remove the dust and washed to record fresh weight of leaf (gm). Leaf samples were dried at 70° c until constant weight to determine the dry weight of leaf (gm).

Number of leaves / shoot was recorded at the end of October of each season.

II -Yield and fruit quality.

1- Yield: Number of breba fruits / tree , number of main fruits per shoot and per tree and main yield (kg) / tree (number of main fruits / tree × average main fruit weight) .

2-Fruit quality: At the second half of August, 15 fruits per each selected tree were randomly harvested for measuring the following determination :

- Average main fruit weight.
- Fruit height and diameter (cm) were estimated using a Varnier Caliper.

- Fruit moisture percentage was determined by drying at 70° c till constant weight.
- Total soluble solids percentage (T.S.S. %) was estimated using hand refractometer.
- Total acidity was determined as citric acid by titrating diluted flesh against 0.1 N NaOH using phenolphthalein as an indicator.
- Total sugars was estimated according to the method of Lane and Eynon as outlined in A.O.A.C. (1975).

III- Leaf mineral composition. Dry leaf samples were grounded and digested according to (Chapman and Pratt, 1978) for the following determinations :

-Total nitrogen: was determined by micro-Kjeldahl method (Jackson, 1973).

-Phosphorus: was determined according to the method of Murphy and Riley (1962).

-Potassium: was determined by Flame Photometer (Jackson, 1973).

Statistical analysis :

All data obtained during the three experimental seasons were subjected to analysis of variances according to Snedecor and Cochran, (1980) and means were differentiated using Multiple Range Test (Duncan, 1955) at the 5 (%) level of probability.

RESULTS AND DISCUSSION

I-Growth parameters .

1-Tree characteristics:

Results in Table (4) clearly show that, cattle, sheep and chicken manure sources significantly improved trunk & canopy circumference and shoot length compared to the control. In this respect, sheep manure and cattle manure had the best values of net increase in trunk circumference and canopy circumference, while chicken manure gave the tallest shoot length in 2005 and 2006 seasons .

Meanwhile, the specific effect of potassium treatments was significant for all characteristics. It can be noticed that, there was a gradual increase in trunk & canopy circumference and shoot length values. For the same characters, the trees which treated with 750 gm K₂O plus foliar potassium 1% had the highest values.

Regarding the interaction between organic manure sources and potassium additions increased significantly shoot length in 2006 season . In this concern, the combination representing chicken manure + 750 gm K₂O plus foliar potassium 1% was the most effective treatment. These results are in agreement with El-Adawy,(1987) and Mordogan, *et al.*,(2002) on fig ; Abou-Taleb, (2004) on pecan and Abd El-Naby *et al.*, (2004) on citrus.

2- Leaf characteristics:

It is quite evident as shown from data in Tables (5 & 6) that, the four investigated leaf parameters (leaf area, leaf fresh & dry weight and number of leaves / shoot) were significantly different in response to organic and potassium treatments.

Gowda, A. M.

4

1278

Generally, organic manure sources additions significantly succeeded to improve the aforesaid leaf characters compared with the control treatment. Herein, chicken and sheep manures had the highest values of leaf area, leaf fresh weight and number of leaves /shoot during both seasons of study. Moreover, sheep manure fertilizer (in 1st season) and cattle and sheep manures (in 2nd season) gave the highest records of leaf dry weight.

Potassium treatments show significant values of leaf parameters. Whereas the trees received both soil and foliar potassium had the highest values compared with the applied trees with soil or foliar application in 2005 and 2006 seasons. In this respect, soil potassium addition at 750 gm K₂O + foliar potassium at 1% was superior in both seasons.

The interaction effect on leaf dry weight was significant in both seasons. However, its effect on leaf area, leaf fresh weight and number of leaves / shoot was not significant especially in the first season. The highest values of leaf area and leaf fresh weight were obtained when the trees treated with chicken, sheep and cattle manures with 750 gm K₂O + foliar 1% potassium. Sheep manure + 750 gm K₂O + foliar potassium 1% gave the heaviest leaf dry weight and the highest number of leaves / shoot.

The enhancement of different growth parameters due to organic fertilizers may be attributed to the positive effect of organic materials on increasing the availability of most nutrients and improving physical and chemical properties of soil which in turn increase nutritional status and growth of the tree (Nijjar, 1985). Moreover, organic manures increase the microorganisms activity in soils which produce growth promoting substances consequently increase the plant growth. This increase in plant growth may increase the photosynthetic rates leading to an increase of the assimilation rates. The present results of using organic fertilizers was supported by the results of El-Adawy, (1987) and Mordogan, *et al.*, (2002) on fig; Abou - Taleb, (2004) on pecan and Abd El-Naby *et al.*, (2004) on citrus. Moreover, the improvement of growth parameters reflects enhancing potassium for uptake of more nutrients via roots and accumulation in the leaves, that activated the growth processes of the shoots. Similar response was reported by Abdel-kader and Sabbah, (2002) on pomegranate.

II -Yield and fruit quality.

1- Yield:

Number of breba fruits / tree: number of breba fruits /tree as affected by different organic manure sources and potassium additions is presented in Table (6). The data indicated that, both organic treatments and potassium addition significantly increased the number of breba fruits /tree as compared to the control during the two seasons of study. Sheep and cattle manures were more effective than chicken manure.

Increasing the potassium addition from 500 to 750 gm K₂O/ tree with or without foliar potassium 1% increased number of breba fruits /tree in 2005 and 2006 seasons. The interaction was significant specially in the second season and the highest values were recorded from those trees treated with cattle manure and supplemented with 750 gm K₂O + foliar potassium 1% (K4).

Number of main fruits per shoot: It was also noticed from data in Table (6) that, the differences as a result of treatments were significant. Trees fertilized with sheep manure had the best significant values as compared to all other treatments in the first and second seasons.

In addition, soil potassium application with foliar spraying exhibited the highest records of number of main fruits /shoot in both studied seasons. The differences between 500 or 750 gm K₂O/ tree + foliar potassium 1% treatments were not significant in 2005 and 2006 seasons. The interaction was insignificant in both seasons.

Number of main fruits / tree: Data presented in Table (7) indicated that, Sultani fig trees produced higher number of main fruits when fertilized with sheep and cattle manure followed by chicken manure treatment than control in both seasons.

Regarding the effect of potassium addition, the trees fertilized with 750 gm K₂O/ tree + foliar potassium 1% (K4) had the greatest number of main fruits /tree. Moreover, the differences between the treatments (K4) and 500gm K₂O + foliar potassium 1%(K3) were not significant during both studied seasons. The interaction was not significant in 2005 and 2006 seasons.

Main yield (kg) / tree: Data in Table (7) shows that applying organic manures significantly increased the yield /tree compared with the control. The trees fertilized with sheep manure gave higher yield followed by cattle manure and chicken manures in both seasons.

The effect of potassium addition was significant in 2005 and 2006 seasons. Whereas, values of the treatment included addition of 750gm K₂O/ tree + foliar potassium 1% (K4) don't give significant values related to that treated with 500gm K₂O/ tree + foliar potassium 1% (K3) in the two seasons of study. The interaction was not significant in both seasons.

The enhancement of fruiting as a result of using organic fertilizer in general and potassium addition in particular may be due to the organic materials which improve soil physiochemical conditions and reserved the sufficient amounts of N and K. In addition, the release of much more less available elements (Fe, Zn and Mn) as well as they increase the soil content of IAA and cytokenins (Li *et al.*,1998). Vigorous vegetative growth produced a high photosynthetic efficiency which causes an increment of reproductive growth and yield (Maksoud, 2000). The results of tree yield are in line with that of El-Adawy, (1987) ; Mordogan, *et al.*, (2002) and Caetano, *et al.* ,(2006) on fig; Abou - Taleb (2004) on pecan and Foud, *et al.*, (2002) on citrus.

2-Fruit quality:

Average main fruit weight: Data concerning the effect of different organic sources and potassium addition on average main fruit weight during 2005 and 2006 seasons are presented in Table (7).

Regarding the specific organic manures, all the tested sources were significantly effective comparing with the control. The heaviest fruit weight was recorded when sheep manure was added, while the differences between values of trees fertilized with cattle and chicken manure were not significant compared with values of sheep manure in both seasons of study. Meanwhile, using 750 gm K₂O/ tree + foliar potassium 1% (K4) gave statistically higher records of main fruit weight in 2005 and 2006 seasons.

Gowda, A. M.

6

1282

Concerning the interaction effect, it was significant in the second season, whereas Sultani fig trees treated with sheep manure + 750 gm K₂O/ tree as soil addition and foliar potassium 1% increased significantly the fruit weight.

Fruit diameter, height and fruit moisture percentage: In this respect, the effect of organic sources and potassium addition are presented in Table (8).

As for the specific effect of different organic manure sources, data reveals obviously that, fruit diameter and fruit height significantly respond to organic manure source compared with the control. Whereas, sheep and cattle manure had the highest values followed by chicken manure in 2005 and 2006 seasons. On the other hand, the lowest values of fruit moisture percentage were recorded in fruits of the trees fertilized with sheep and cattle manures. Anyhow, the highest percentage of moisture in fruits were obtained from fruits of trees subjected to chicken manure and control and the differences were not significant in 2005 and 2006 seasons.

Referring to the specific effect of potassium addition, Table (8) also shows that, fruit diameter and height were significantly affected by the different potassium fertilization techniques with soil and / or foliar application in the two seasons of study. Meantime, Sultani fig trees fertilized with 750 gm K₂O/ tree (soil) + foliar potassium 1% (K4) produced the highest fruit diameter and height followed by 500gm K₂O/ tree + foliar potassium 1% (K3). The highest percentage of fruit moisture content was recorded from the treatment included foliar potassium 1% in both seasons (Table, 8).

With regard to the effect of the interaction on fruit diameter and height, it was significant, while the same treatment didn't affect fruit moisture content in 2005 and 2006 seasons. The highest fruit diameter and height were obtained from trees fertilized with sheep manure and supplemented by 750 gm K₂O/ tree + foliar potassium 1% (K4) during both seasons. These findings were in agreement with those obtained by Mordogan, *et al.*, (2002) and Caetano, *et al.*, (2006) on fig; Abou -Taleb, (2004) on pecan.

The improvement of fruit quality (fruit weight, diameter, height and fruit moisture content) as a result of applying organic matter to soil is mainly due to the improvement in structure, essential elements, soils ability to hold water and nutrients as well as resist compacting and crusting (Madison *et al.*, 1986). The positive effect of potassium fertilization may be due to its important role in promoting and enhancing the metabolic process during uptake, root activation, regulate water balance and translocation compounds which in turn increase the growth and reflects on yield and fruit quality (Najjar, 1985). Sen and Chaunan (1983) found that rising potassium rates increased the yield of pomegranate trees.

Total soluble solids (T.S.S.%), total acidity (%) and total sugars (%) :

As for the effect of the investigated organic fertilizer, data in Table (9) show that, the fruits of sheep manure fertilized trees had the highest values of total soluble solids (T.S.S.%) and total sugars. Moreover, cattle and sheep manure treatments gave the highest total soluble solids values compared to the control treatment in the 1st season. In the contrast, in the 2nd one, the differences of trees which fertilized with cattle, sheep and chicken manures were not significant compared with the control.

Gowda, A. M.

9

1286

Regarding fruit total sugars content , the trees fertilized with cattle, sheep and chicken manures had insignificant values related to the control in 2005 season .

While, in 2006 season, the differences between the values of trees fertilized with cattle and chicken manures were not significant .

Concerning the response to potassium addition, it was so clear that fertilization with 750 gm K₂O/ tree + foliar potassium 1% (K4) treatment was superior for raising fruit TSS and total sugars contents during both seasons. As for interaction effect, it was not significant in the first season. Sultani fig trees fertilized with sheep manure and (k4) exerted statistically the highest stimulate effect on fruit TSS and total sugars in the 2nd one .

The three sources of organic manures had a significant effect on total acidity values of Sultani fig tree fruits compared with the control. On the contrary, the same character decreased significantly during 2005 and 2006 seasons when sheep manure was applied in the first season and cattle manure in the second one.

Meanwhile, the addition of potassium improved the acidity content of fruits, hence, the lowest acidity content was recorded when the trees fertilized with 500 gm K₂O/ tree + foliar potassium 1% (K3) and 750 gm K₂O/ tree + foliar potassium 1% (K4) in the two seasons of study .

In addition, total acidity responded significantly to the interaction between organic manure sources and potassium addition in both seasons. Application of cattle and sheep manures with (k4) treatment gave the pronounced effect on decreasing fruit total acidity content in the two studied seasons. The results are in line with those of El-Adawy, (1987) ; Mordogan, *et al.*, (2002) and Caetano, *et al.* , (2006) on fig and Abd El-Naby *et al.*, (2004) on citrus

III- Leaf mineral composition .

Data concerning the effect of organic and potassium addition on leaf N, P and K contents during 2005 and 2006 seasons are presented in Table (10). Regarding the effect of organic manure sources, the highest nitrogen and phosphorus values were recorded in leaves of trees treated with sheep manure in both seasons. While, the differences between sheep and chicken manure on phosphorus content were not significant in both seasons. Leaf potassium content increased significantly due to the addition of chicken and sheep manure compared with the control.

As for potassium treatments effect, the highest level of nitrogen and phosphorus content was observed in leaves of the fertilized trees with 500 gm K₂O/ tree + foliar sprayed with potassium 1% (K3) followed by 750 gm K₂O/ tree + foliar sprayed with potassium 1% (K4) treatment in 2005 and 2006 seasons. Leaf potassium content was significantly increased by adding 750 gm K₂O/ tree + foliar potassium 1% (K4) in both seasons.

Concerning the interaction effect on N , P and K contents. Data reveals that Sultani fig trees received combination of chicken or sheep manure and 500 gm K₂O/ tree + foliar potassium 1%(K3) induced the highest level of nitrogen and phosphorus in the 1st season. While, sheep manure + (k3) gave the highest N content in the 2nd one .

Gowda, A. M.

10

1288

The combination which included chicken manure and 750 gm K₂O/ tree + foliar sprayed with potassium 1% (K4) gave the highest values of leaf potassium content in 2005 and 2006 seasons. These results are in conformity with El-Adawy,(1987); Irget *et al.*, (1999); Mordogan,*et al.*, (2002) and Caetano, *et al.*, (2006) on fig; Kassem and Marzouk, (2002) on grapevine; Abou -Taleb,(2004) on pecan and Abd El-Naby *et al.*, (2004) on citrus.

The promotion in leaf mineral content due to organic fertilizers which improving the structure of soil, aeration and drainage, amount of water available and favorable conditions of root growth and nutrient absorption. On the other hand, the organic nitrogen added to the soil in the form of plant and animal residues is largely proteinceous in nature. Similar results were reported by Cook, (1982). Moreover, the improvement of leaf nutrients content as a result of potassium addition may be due to its active role in enhancing the absorption, translocation and accumulation of mineral contents in leaves (Hikal, 2000).

As a conclusion, sheep or cattle manure when supplemented with 500 gm K₂O/ tree + foliar potassium at 1% can be recommended in potassium fertilization program to obtain higher yield and better fruit quality of fig trees (Sultani cv.) under the same conditions of this study.

REFERENCES

- A.O.A.C.(1975). Official methods of analysis. Association of Official Analysis Chemist. Published by A.O.A.C. Washington, D.C.(U.S.A.).
- Abdel- Kader.Hayat M. and S.M.Sabbah (2002). Effect of potassium fertilization on vegetative growth, yield, fruit quality and leaf mineral composition of some pomegranate cultivars. *Minufiya J. Agric. Res.* Vol. 27 (2): 337-353.
- Abd El-Naby ,S. K.M. (2004). Response of Washington Navel orange trees growing on sandy soil to mixture of some organic fertilizers. *Assiut J. of Agric. Sci.* Vol. 35 (2): 215-235.
- Abd El-Naby ,S. K.M.; Abd El- Moneim , Eman A.A. and A.S.E.Abd-Allah (2004). Effect of source and date of organic manure application on growth, yield, fruit quality and mineral content of Washington Navel orange trees grown in sandy soil. *Minufiya J. Agric. Res.* Vol. 2 (29): 515-540.
- Abou- Hussein S.D.; I.El- Oksha; T.El-Shorbagy and A.M.Gomaa (2002). Effect of cattle manure ,bio fertilizers and reducing mineral fertilizer on nutrient content and yield of potato plant. *Egypt J. Hort.* 29 (1): 99-115.
- Abou-Taleb, Safia, A (2004). Effect of cattle manure and reducing mineral fertilizer on growth, fruit quality and nutrient content of pecan trees. *Annals of Agric. Sc. Moshtohor* , Vol. 42 (3) : 1197- 1214.
- Caetano, L.C.S.; A.J. Cordeiro de Carvalho and J.M. Jasmim (2006). Preliminary report on yield productivity and mineral composition of the fig tree as a function of boron and cattle manure fertilization in Brazil. *Fruits*, 61: 341- 349.
- Chapman, H.D. (1968). The mineral nutrition of citrus. *Citrus Ind.* Vol.II, Chap. 3: 127- 289. Univ. Calif., USA.
- Chapman, H.D. and P.E.Pratt (1978).Methods of Analysis for Soils, Plants and Waters . Univ. of California, Div. Agric. Sci., priced pub 4034.

- Cook, G.W.(1982).Fertilizing for Maximum Yield . 3rd ed. Granada Publishing Limited, pp. 465.
- Duncan, D. B. (1955). Multiple range and multiple F. Tests biometrics, 11: 1-42.
- El- Adawy, Samia A.F.(1987). Response of irrigated and non-irrigated Sultany fig trees grown in the Egyptian North Western Coast to soil and foliar fertilization.Ph.D.Thesis. Fac. Agric.Alex.Univ., Egypt
- Ferguson, j,j (1994). Growth and yield of bearing and non- bearing citrus trees fertilized with fresh and processed chicken manure .Proc. of The Florida State Hort. Soc. 1994 , 107 : 29- 32.
- Foud, Amara A. ; L.L.Khalil, Fekrya ;E.M.Abdallah and E.A.Shaban (2002). Studies on the effect of organic nitrogen and / or mineral nitrogen, phosphorus and potassium fertilizers on the yield and fruit quality of Washington Navel orange trees . Proc. Minia 1st Conf. For Agric. & Environ. Sci. , Minia , Egypt , March 25- 28.
- Gamal,A.M. and M.A.Ragab (2003). Effect of organic manure source and its rate on growth, nutritional status of the trees and productivity of Balady Mandarin trees. Assiut J. of Agric. Sci. Vol. 34 (6): 253-264.
- Haggag, Laila, F. (1996). Response of Picual olive trees in sandy soil to various forms of N fertilization (organic and chemical). Annals Agric. Sci., Ain Shams Univ., Cairo .41 (1): 313- 319.
- Hikal, A.R.F.(2000). Physiological studies on nutrition of Washington Navel orange trees. Ph.D.Thesis. Fac. Agric.Mansoura Univ., Egypt.
- Irget, M.E.; S.Aydin; M.Oktay; M.Tutam; U.Aksoy and M.Nalbant (1999). Effects of foliar potassium nitrate and calcium nitrate application on nutrient content and fruit quality of fig . Improved crop quality by nutrient management .Kluwer Academic Publishers. Dordrecht , Boston- London. 81-86.
- Jackson, M.H. (1973). Soil Chemical Analysis. Prentice Hall . Inc. N.J. Privatle Limited and New Delhi.
- Kassem ,H.A. and H.A.Marzouk (2002). Effect of organic and / or mineral nitrogen fertilization on the nutritional states, yield and fruit quality of Flam Seedless Grapevines grown in calcareous soil. J. Adv. Agric. Res., 7 (1): 117- 125.
- Koo, R.C.J.(1985). Effect of nitrogen and potassium fertilization on winter injury of citrus trees . Pro. Fla. State Hort., 98 : 53-56.
- Li, X.J.; S. F.Dong and Y.S. Liu (1998) .Determination of IAA and cytokinins in the soil with different organic manure for pot cultured apple. Plant Physiology Communications, 34(3):183-185.
- Madison, F.;K. Kelling ; J. Peterson; I. Daniel; G.Jackson and L.Massie (1986).Managing manure and waste : Guidelines for applying manure to pasture and cropland in Wisconsin. Report A 3392. Madison: University of Wisconsin- Extension.
- Maksoud, M.A. (2000).Response of growth and flowering of Manzanillo olive trees to different sorts of nutrient . Egypt J. of Hort. 27(4): 513-583.
- Mengel, K. and A.Kirkby .(1987). "Principles of plant nutrition" , 4th Ed., International Potash Institute , Norblafen- Bern , Switzerland.
- Mordogan,N.; H.Hakerlerler; S.Ceylan; S.Aydin; B.Yagmur and U.Aksoy (2002).Effect of organic fertilization on fig leaf nutrients and fruit quality. International Conference On Sustainable Land Use and Management, 2002. Canakkale, Turkey.
- Murphy, J. and J.P.Riley (1962). A modified single solution for the determination of phosphate natural water . Anal. Chem. Acta., 27 : 31-36.

- Nijjar, G.S. (1985). Nutrition of fruit trees .Kaalyani Puplichers, New Delhi, India. pp. 10- 52.
- Sen,N.L. and K.S.Chauhan (1983).Effect of differential NPK fertilization on physico- chemical characters of pomegranate. Punjab – Hort. J.,23: 1/2 , 59 – 63.
- Snedecor, G.W. and W.G. Cochran (1980). Statistical methods . 7th ed. Iowa State Univ. Press, Ames , Iowa, U.S.A. pp . 507.
- Wilde, S.A.;R.B. Corey; J.G.Lyer and G.K.Voigt. (1985).Soil and Plant Analysis for Tree culture. Oxford and IBH Publishing Co., New Delhi India.
- Youssef,A.H.;A.H.El- Fouly ; M.S.Youssef and S.A.Mohamadien (2001). Effect of using organic and chemical fertilizers in fertigation system on yield and fruit quality of potato plants (*Solanum turosum* L.). Resent Technologies in Agriculture.Proc. Of the 2nd congress . Vol.(1) : 79-94 .
- Zhou, X.W. ;Z.Y.Li; B.Lu; X.N.Chen; L.J.Xu; Y.W.Yi; X.W.Zhou;Z.Y.Li; B.Lu; X.N.Chen; L.J.Xu and Y.W.Yi (2001).Study on the improvement of the soil of the newly established orchard on the reclaimed purple soil .J. of Fruit Sci. 18 : 1, 15-19.(C.F. CAB Abst. 1001 – 7364).

تأثير بعض الأسمدة العضوية و إضافة البوتاسيوم على النمو والمحصول وجوده

ثمار التين السلطاني

عادل محمد جوده

قسم بحوث الزيتون وفاكهة المناطق شبه الجافة- معهد بحوث البساتين- مركز البحوث الزراعية – الجيزة- مصر

أجريت هذه التجربة بمركز ناصر – محافظه بنسوف لدراسة تأثير بعض مصادر الأسمدة العضوية و أضافه البوتاسيوم على النمو، المحصول وصفات الجودة لثمار أشجار التين السلطاني النامية في أرض طميبه خلال موسمي ٢٠٠٥ ، ٢٠٠٦ . حيث استخدمت أسمده الماشية أو الغنم أو الدواجن مع أضافه البوتاسيوم أرضي أو رش أو الاثني معا .وقد أتضح الآتي:

بالنسبة لقياسات النمو الخضري: كانت أعلى القيم بالنسبة لصفاف الزيادة في محيط الجذع ومحيط نمو الشجرة عندما سمدت الأشجار بسمد الغنم والماشية وأطول الأفرع للأشجار المسمد ه بسمد الدواجن أما أعلى القيم لمساحة الورقة والوزن الطازج للورقة وعدد الأوراق / فرخ فقد سجلت بتسميد الأشجار بسمد الغنم و الدواجن و أعلى القيم للوزن الجاف للورقة فكان للأشجار التي تم تسميدها بسمد الماشية . أعطت المعاملة باضا فه ٧٥٠ جم أكسيد بوتاسيوم في صورته سلفات بوتاسيوم أضافه أرضيه مع الرش الورقي للبوتاسيوم بتركيز ١ % أعلى القيم لجميع خصائص النمو الخضري المدروسة.

أما المحصول وجوده الثمار: فقد أوضحت النتائج أن الأشجار المسمدة بسمد الغنم والماشية أعطت القيم الأعلى في عدد الثمار البوني / شجره ، عدد ثمار المحصول الرئيس لكل فرع و لكل شجره ، المحصول (كجم / شجره) ، متوسط وزن الثمرة في المحصول الرئيس وطول و قطر الثمرة ومحتوى الثمار من المواد الصلبه الكلية الذائبة والسكريات الكلية و أقل قيم لمحتوى الثمار من الرطوبة والحموضة .وكان تأثير أضافه البوتاسيوم غير معنوي بين الأشجار التي سمدت ب ٧٥٠ جم أكسيد بوتاسيوم/ شجره أو ٥٠٠ جم أكسيد بوتاسيوم / شجره مع الرش الورقي للبوتاسيوم بتركيز ١ % على محصول الأشجار . ولكن كانت المعاملة باضا فه ٧٥٠ جم أكسيد بوتاسيوم / شجره+ الرش الورقي للبوتاسيوم بتركيز ١% هي المتفوقة في صفات جودة الثمار ماعدا الحموضة ومحتوى الثمار من الرطوبة.كما تفوقت المعاملة باضا فه سمد الغنم مع ٧٥٠ جم أكسيد بوتاسيوم / شجره+ الرش الورقي للبوتاسيوم بتركيز ١% في معظم الصفات المدروسة.

و بالنسبة لمحتوى الأوراق من عناصر النتروجين والفسفور والبوتاسيوم: أدى التسميد بسمد الغنم إلى رفع محتوى الورقة من النتروجين والفسفور أما سمد الدواجن فأدى لرفع محتوى الورقة من البوتاسيوم.أدت المعاملة باضا فه ٥٠٠ جم أكسيد بوتاسيوم / شجره+ الرش الورقي للبوتاسيوم بتركيز ١% إلى زيادة محتوى الورقة من النتروجين والفسفور

وعلى ضوء هذه النتائج فإنه يمكن التوصية بتسميد أشجار التين السلطاني باستخدام أسمدة الغنم أو الماشية مع أضافه ٥٠٠ جم أكسيد بوتاسيوم / شجره+ الرش الورقي للبوتاسيوم بتركيز ١% وذلك للحصول على أعلى محصول و أفضل جودة ثمار تحت الظروف المماثلة لهذه الدراسة .

Table (4) : Effect of different organic manure sources and potassium addition on net increase in trunk circumference (cm), canopy circumference(m) and shoot length (cm) during 2005 and 2006 seasons.

Charact Treat.	Net increase in trunk circumference (m)					Canopy circumference (m)					Shoot length (cm)				
	Cattle manure	Sheep manure	Chicken manure	Control	Mean	Cattle manure	Sheep manure	Chicken manure	Control	Mean	Cattle manure	Sheep manure	Chicken manure	Control	Mean
2005 season															
K 1	6.04**	5.77	5.21	4.73	5.44 D	9.4	9.6	9.8	8.1	9.2 C	68.45	69.37	72.49	63.60	68.48 D
K 2	6.32	6.07	5.64	5.08	5.78 C	10.4	11.2	10.4	8.7	10.2 B	72.44	76.61	76.02	64.00	72.27 C
K 3	6.64	6.20	5.58	5.90	6.08 B	10.4	11.9	10.7	9.2	10.6 B	73.77	77.20	82.02	64.99	74.50 B
K 4	6.97	6.74	6.00	5.88	6.40 A	11.9	13.0	11.6	9.4	11.5 A	76.57	76.73	83.47	69.54	76.58 A
K 5	5.66	5.24	4.71	4.08	4.92E	9.0	9.5	9.1	7.6	8.8 C	65.74	68.98	70.93	56.38	65.51 E
Mean	6.33 A*	6.00 A	5.43 B	5.13 B		10.2 B	11.0 A	10.3 B	8.6 C		71.39 C	73.78 B	76.99 A	63.70 D	
2006 season															
K 1	6.32	6.24	5.72	4.79	5.77 C	13.2	14.2	12.0	11.2	12.7 C	72.48 gh	78.48 def	80.66 cde	67.58 i	74.80 C
K 2	6.70	6.64	6.28	5.28	6.23 B	15.9	16.2	13.3	12.7	14.5 B	78.13 def	80.24 de	84.79 bc	73.15 gh	79.08 B
K 3	7.16	6.88	5.94	5.82	6.45 AB	14.8	15.6	13.7	12.5	14.2 B	79.10 de	82.35 bcd	84.86 bc	71.00 hi	79.33 B
K 4	7.11	7.14	6.22	5.90	6.59 A	16.7	17.1	15.1	13.3	15.6 A	86.87 b	85.85 b	94.33 a	76.34 efg	85.85 A
K 5	5.79	5.71	5.70	4.54	5.44 D	11.4	13.5	11.4	10.1	11.6 D	73.99 fgh	72.93 gh	74.33 fgh	59.62 j	70.22 D
Mean	6.62 A	6.52 A	5.97 B	5.27 C		14.4 B	15.3 A	13.1 C	12.0 D		78.11 C	79.97 B	83.79 A	69.54 D	

Means having * the same letters or ** not having letters are not significantly differ at 5% level.

Table (5) : Effect of different organic manure sources and potassium addition on leaf area (cm)², leaf fresh and dry weight (gm) during 2005 and 2006 seasons.

Characteristics	Leaf area (cm) ²					Leaf fresh weight (gm)					Leaf dry weight (gm)					
	Treat.	Cattle manure	Sheep manure	Chicken manure	Control	Mean	Cattle manure	Sheep manure	Chicken manure	Control	Mean	Cattle manure	Sheep manure	Chicken manure	Control	Mean
2005 season																
K 1	354.9**	418.8	342.0	304.0	354.9 C	15.56	16.36	17.07	13.45	15.61 C	7.63 ghi	7.40 hi	6.97 hij	5.76 i	6.94 D	
K 2	379.9	420.9	365.9	322.3	372.2 BC	17.89	17.90	18.07	14.93	17.20 B	9.03 cde	8.36 efg	7.83 fgh	6.72 ilk	7.98 C	
K 3	415.3	441.1	385.6	322.4	391.1 AB	18.20	19.52	19.00	15.13	17.96 B	9.52 cd	11.21 b	8.68 def	6.41 jkl	8.96 B	
K 4	432.8	449.8	418.9	351.0	413.1 A	19.27	20.64	20.48	16.63	19.26 A	10.99 b	12.46 a	9.59 c	7.67 gh	10.18 A	
K 5	339.9	371.4	433.9	288.9	358.5 BC	14.31	15.02	15.50	12.25	14.27 D	6.89 hij	7.63 ghi	6.38 jkl	5.93 kl	6.71 D	
Mean	384.6 B*	420.4 A	389.3 B	317.7 C		17.05 B	17.89 AB	18.03 A	14.50 C		8.81 B	9.41 A	7.89 C	6.50 D		
2006 season																
K 1	334.7 de	321.24ef	322.5 ef	302.8 fg	320.3 D	16.07 fg	18.30 cde	19.40 bc	16.78 fg	17.64 D	8.56 ef	8.22 efg	7.73 fgh	6.91 h	7.85 C	
K 2	342.9 cde	327.8 ef	364.3 bc	316.7 ef	337.9 C	19.55 bc	20.24 b	20.26 b	18.38 cd	19.61 B	10.03 bcd	9.88 cd	8.61 ef	7.68 fgh	9.05 B	
K 3	371.3 ab	357.1 bcd	374.2 ab	304.8 fg	351.8 B	17.36 def	19.67 bc	20.51 b	16.55 fg	18.52 C	10.29 bc	10.88 b	9.22 de	7.28 gh	9.42 B	
K 4	394.1 a	375.9 ab	396.3 a	329.5 ef	374.0 A	22.92 a	21.89 a	22.25 a	19.51 bc	21.64 A	14.91 a	14.33 a	10.00 bcd	8.50 ef	11.94 A	
K 5	284.5 gh	306.3 fg	318.1 ef	273.0 h	295.4 E	15.70 g	16.42 fg	17.01 efg	14.07 h	15.80 E	8.44 ef	7.88 fgh	7.58 fgh	5.85 i	7.44 C	
Mean	345.5AB	337.7 B	355.1 A	305.4 C		18.32BB	19.30AA	19.89 A	17.06 C		10.45A	10.24A	8.63 B	7.25 C		

Means having * the same letters or ** not having letters are not significantly differ at 5% level.

Table (6) : Effect of different organic manure sources and potassium addition on number of leaves / shoot, number of breba fruits / tree and number of main fruits/ shoot during 2005 and 2006 seasons.

Characteristics	Number of leaves / shoot					Number of breba fruits/ tree					Number of main fruits / shoot				
	Treat.	Cattle manure	Sheep manure	Chicken manure	Control	Mean	Cattle manure	Sheep manure	Chicken manure	Control	Mean	Cattle manure	Sheep manure	Chicken manure	Control
2005 season															
K 1	11.72**	13.41	14.76	10.32	12.55 D	12.7	12.0	10.0	8.7	10.8 D	12.93	13.48	13.10	12.17	12.92 C
K 2	12.91	14.07	15.74	11.90	13.65 C	14.7	15.0	13.0	9.3	13.0 C	14.04	14.21	13.28	13.15	13.67 B
K 3	13.72	15.33	16.71	12.20	14.49 B	18.0	19.7	14.3	10.0	15.5 B	14.43	16.26	14.40	13.40	14.62 A
K 4	15.05	16.38	17.65	12.89	15.49 A	23.7	24.7	15.7	11.7	18.9 A	15.00	16.53	14.57	13.70	14.95 A
K 5	10.11	12.01	13.39	8.96	11.12 E	15.0	13.7	10.0	7.0	11.4 CD	12.09	13.24	13.07	11.80	12.55 C
Mean	12.70 C	14.24 B	15.65 A	11.25 D		16.8 A*	17.0 A	12.6 B	9.3 C		13.70 B	14.74 A	13.69 B	12.84 C	
2006 season															
K 1	13.47 gh	14.84 e	14.98 e	13.02 hi	14.08 D	15.3 gh	16.7 fg	16.0 gh	17.7 d-g	16.4CD	14.10	14.87	14.34	12.82	14.03 C
K 2	14.48 ef	15.89 d	15.96 d	14.28 ef	15.15 C	18.3 c-g	20.0 b-f	17.7 d-g	12.7 h	17.2 C	14.95	15.21	14.60	13.25	14.50 BC
K 3	16.63 cd	16.6 cd	17.07 c	16.09 d	16.60 B	20.7 a-e	21.7 abc	18.0 c-g	15.7 gh	19.0 B	16.39	16.44	15.61	13.47	15.48 A
K 4	19.09 a	18.18 b	18.23 b	16.54 cd	18.01 A	24.3 a	22.3 ab	21.3 a-d	20.3 b-f	22.1 A	16.51	16.30	16.29	14.29	15.85 A
K 5	12.34 i	13.80 fg	13.97 fg	11.0 j	12.78 E	17.3 efg	18.0 c-g	16.0 gh	8.7 i	15.0 D	13.62	14.53	13.95	11.70	13.45 D
Mean	15.20 B	15.86 AB	16.04 A	14.19 C		19.2 A	19.7 A	17.8 B	15.0 C		15.11AB	15.47 A	14.96 B	13.11 C	

Means having * the same letters or ** not having letters are not significantly differ at 5% level.

Table (7) : Effect of different organic manure sources and potassium addition on number of main fruits / tree, fruit weight (gm) and yield (kg / tree) during 2005 and 2006 seasons.

Characteristics	Number of main fruits/ tree					Fruit weight (gm)					Yield (kg / tree)					
	Treat.	Cattle manure	Sheep manure	Chicken manure	Control	Mean	Cattle manure	Sheep manure	Chicken manure	Control	Mean	Cattle manure	Sheep manure	Chicken manure	Control	Mean
2005 season																
K 1	778.3**	803.7	741.7	493.3	704.3 C	45.40	46.8	46.90	38.03	44.28 C	35.30	37.61	34.74	18.76	31.60C	
K 2	842.7	890.3	812.0	571.7	779.2 B	47.07	51.30	48.70	40.90	46.99 B	39.68	44.46	39.54	23.40	36.77B	
K 3	864.3	908.3	805.7	584.3	790.7 AB	48.27	52.57	47.63	41.37	47.46 B	41.72	47.75	38.38	24.17	38.01AB	
K 4	893.7	905.3	873.0	599.3	817.8 A	49.67	53.5	49.97	41.90	48.76 A	44.39	48.43	43.62	25.11	42.02A	
K 5	775.3	762.3	641.7	469.0	662.1 D	44.77	44.63	45.13	36.57	42.78 D	34.71	34.02	28.93	17.14	28.70D	
Mean	830.9 A*	854.0 A	774.8 B	543.5 C		47.03 B	49.76 A	47.67 B	39.75 C		39.16B	42.45A	37.04B	21.72C		
2006 season																
K 1	704.0	816.0	702.3	614.7	709.3 C	45.70 efg	47.27d-g	46.40d-g	40.67 j	45.01 C	32.18	38.51	32.61	24.98	32.07C	
K 2	780.3	877.7	764.3	677.0	774.8 B	50.53 defg	49.83 bc	49.33 cd	43.33 i	48.26 B	39.40	43.75	37.70	29.32	37.54B	
K 3	856.7	885.0	804.7	728.0	818.6 AB	55.03 de	57.57 b	51.83 def	45.90 hi	52.58 B	47.14	50.95	41.71	33.42	40.69A	
K 4	876.0	901.0	823.0	747.0	836.8 A	58.47 bc	58.50 a	54.40 b	51.50 gh	55.72 A	51.23	52.70	44.71	38.48	46.78A	
K 5	703.7	706.0	670.3	562.7	660.7 D	47.23 fg	46.13 fg	45.23 efg	37.07 j	43.92 D	33.21	32.60	30.33	20.86	29.25C	
Mean	784.2 B	837.1 A	752.9 C	665.9 D		51.39 B	51.86 A	49.44 B	43.69 C		40.63B	43.70A	37.41C	29.41D		

Means having * the same letters or ** not having letters are not significantly differ at 5% level.

Gowda, A. M.

Table (8) : Effect of different organic manure sources and potassium addition on fruit diameter (cm), fruit height (cm) and fruit water content (%)

Characteristics	Fruit diameter (cm)					Fruit height (cm)					Fruit water content %				
	Treat.	Cattle manure	Sheep manure	Chicken manure	Control	Mean	Cattle manure	Sheep manure	Chicken manure	Control	Mean	Cattle manure	Sheep manure	Chicken manure	Control
	2005 season														
K 1	4.25 d-g	4.23 efg	3.99 ij	3.87 jk	4.08 C	5.36 gh	6.31 cde	5.49 fgh	5.04 hi	5.55 C	81.70 **	80.30	82.16	82.56	81.68 B
K 2	4.36 b-e	4.44 bc	4.12 ghi	4.01 hij	4.23 B	5.63 fg	6.50 cd	6.10 de	5.25 gh	5.87 B	81.42	79.12	81.62	81.67	80.96 C
K 3	4.34 b-e	4.41 bcd	4.12 ghi	4.14 ghi	4.25 B	6.67 bc	6.42 cd	5.59 fg	5.52 fg	6.05 B	80.54	79.20	81.63	81.54	80.73 C
K 4	4.41 bcd	4.63 a	4.44 b	4.32 b-f	4.45 A	6.95 ab	7.20 a	6.31 cde	5.55 fg	6.50 A	79.53	77.89	80.91	81.53	79.97 D
K 5	4.16 fgh	4.27	3.83 k	3.80 k	4.02 C	5.61 fg	5.94 ef	5.34 gh	4.75 i	5.41 C	82.40	80.84	82.56	83.80	82.48 A

during 2005 and 2006 seasons.

	c-g														
Mean	4.31 B	4.40 A	4.10 C	4.03 C		6.04 B	6.48 A	5.77 B	5.22 C		81.12 B	79.47 C	81.84 A*	82.22 A	
2006 season															
K 1	4.48 def	4.49 def	4.33 f-i	4.19 ijk	4.37 C	5.62 gh	6.14 e	5.93 efg	5.49 h	5.79 D	76.42	77.67	82.70	83.30	80.02 B
K 2	4.63 cd	4.60 de	4.37 f-i	4.25 hij	4.46 B	6.10 ef	6.24 de	6.11 ef	5.73 fgh	6.05 C	75.48	76.50	81.50	81.82	78.83 C
K 3	4.44 e-h	4.79 bc	4.46 d-g	4.1 jk	4.45 BC	6.25 de	6.95 b	6.53 cd	5.42 hi	6.29 B	74.38	75.67	79.22	80.82	77.52 D
K 4	4.81 b	5.03 a	4.62 cde	4.40 fgh	4.72 A	6.56 cd	7.52 a	6.88 bc	5.99 efg	6.74 A	73.22	72.50	78.76	79.95	76.11 E
K 5	4.34 f-i	4.30 f-i	4.28 g-j	4.04 k	4.34 D	5.66 gh	6.56 cd	6.04 ef	5.11 i	5.84 D	78.56	78.67	83.78	83.88	81.22 A
Mean	4.54 A	4.64 A	4.41 B	4.20 C		6.04 C	6.68 A	6.30 B	5.55 D		75.61 B	76.20 B	81.19 A	81.96 A	

Means having * the same letters or ** not having letters are not significantly differ at 5% level.

Table (9) : Effect of different organic manure sources and potassium addition on TSS, acidity and total sugars during 2005 and 2006 seasons .

Characteristics	TSS %					Acidity %					Total sugars %				
	Treat.	Cattle manure	Sheep manure	Chicken manure	Control	Mean	Cattle manure	Sheep manure	Chicken manure	Control	Mean	Cattle manure	Sheep manure	Chicken manure	Control
2005 season															
K 1	14.87**	15.40	14.80	14.53	14.90 D	0.386 cd	0.321 ij	0.354 fg	0.417 b	0.370 B	16.16	16.50	16.86	14.12	15.91 B
K 2	15.33	15.47	15.00	15.20	15.25 C	0.381 cd	0.311 jk	0.337 ghi	0.390 cd	0.355 C	16.77	17.22	17.50	14.31	16.45 AB
K 3	16.20	16.53	15.67	15.27	15.92 B	0.372 de	0.307 jk	0.332 hi	0.334 hi	0.336 D	16.66	16.68	16.11	14.50	15.99 B
K 4	16.33	17.27	16.47	16.33	16.60 A	0.345 fgh	0.295 k	0.344 fgh	0.374 de	0.340 D	16.84	17.00	18.09	15.81	16.94 A
K 5	15.20	15.20	15.20	15.33	15.23 C	0.394 c	0.357 ef	0.383 cd	0.455 a	0.397 A	15.36	15.57	15.12	14.18	15.06 C
Mean	15.59 B*	15.97 A	15.43 B	15.33 B		0.376 B	0.318 D	0.350 C	0.394 A		16.36 A	16.60 A	16.74 A	14.59 B	
2006 season															
K 1	16.27 de	16.13 de	16.73 cde	15.73 de	16.22 CD	0.339 f	0.340 f	0.432 d	0.527 b	0.410 B	15.44 ef	16.15 cde	15.82 def	14.53 gh	15.48 C
K 2	16.53 de	17.33 bcd	17.20 b-e	16.07 de	16.78 BC	0.314 g	0.313 g	0.420 d	0.493 c	0.385 C	16.38 cd	17.89 b	15.73 def	15.23 fg	16.31 B
K 3	17.07 b-e	18.20 bc	17.00 b-e	16.00 de	17.07 B	0.268 i	0.302 gh	0.378 e	0.426 d	0.344 D	16.05 c-f	16.89 c	15.91 def	15.70 def	16.14 B
K 4	20.00 a	20.33 a	18.40 b	16.47 de	18.80 A	0.250 j	0.286 h	0.349 f	0.394 e	0.320 E	18.10 b	18.87 a	16.18 cde	15.86 def	17.25 A
K 5	15.93 de	16.47 de	15.73 de	15.53 e	15.92 D	0.345 f	0.350 f	0.433 d	0.578 a	0.426 A	15.21 fg	15.99 def	15.90 def	14.32 h	15.35 C
Mean	17.16 A	17.69 A	17.01 A	15.96 B		0.303 D	0.318 C	0.402 B	0.484 A		16.24 B	17.16 A	15.91 B	15.13 C	

Means having * the same letters or ** not having letters are not significantly differ at 5% level .

Characteristics	Nitrogen %					Phosphorus %					Potassium %					
	Treat.	Cattle manure	Sheep manure	Chicken manure	Control	Mean	Cattle manure	Sheep manure	Chicken manure	Control	Mean	Cattle manure	Sheep manure	Chicken manure	Control	Mean
2005 season																
K 1	1.90 fg	2.02 de	1.89 fg	1.83 gh	1.91 C	0.26 d-h	0.27 cde	0.27 d-g	0.22 kl	0.26 B	0.81 jk	0.90 ghi	1.09 d	0.86 ij	0.91 C	
K 2	1.70 ij	1.79 g-j	1.72 hij	1.73 hij	1.74 E	0.26 e-h	0.29 abc	0.28 bcd	0.23 kl	0.26 B	0.85 ij	0.97 ef	1.18 c	0.95 fg	0.99 B	
K 3	2.14 bc	2.37 a	2.21 b	1.98 ef	2.18 A	0.28 bcd	0.29 ab	0.30 a	0.25 hij	0.28 A	0.88 hi	0.93 fgh	1.27 b	0.92 fgh	1.00 B	
K 4	2.04 cde	2.17 b	2.13 bcd	1.86 fg	2.05 B	0.25 fgh	0.27 d-g	0.27 c-f	0.26 d-h	0.26 B	1.08 d	1.25 b	1.34 a	1.01 e	1.17 A	
K 5	1.86 g	1.84 gh	1.81 ghi	1.68 j	1.80 D	0.23 ijk	0.26 e-h	0.25 ghi	0.21 l	0.24 C	0.78 k	0.86 ij	0.96 ef	0.64 l	0.81 D	
Mean	1.93 B	2.04 A	1.95 B	1.82 C		0.26B	0.28A	0.27A	0.24C		0.88 C	0.98 B	1.17 A	0.88 C		
2006 season																
K 1	1.96 h	2.09 ef	1.93 hi	1.66 k	1.91 D	0.22 **	0.23	0.25	0.19	0.22 D	0.85 hi	0.94 efg	1.00 de	0.74 j	0.88 D	

Table (10): Effect of different organic manure sources and potassium addition on leaf nitrogen, phosphorus and potassium percentages during 2005 and 2006 seasons .

Gowda, A. M.

K 2	2.00 gh	2.00 gh	2.06 fg	1.81 j	1.97 C	0.24	0.25	0.25	0.20	0.24 C	0.95 efg	1.15 c	1.16 c	0.90 fgh	1.04 C
K 3	2.26 bc	2.45 a	2.15 de	1.88 ij	2.18 A	0.25	0.27	0.26	0.20	0.25 B	1.05 d	1.25 b	1.30 b	0.84 hi	1.11 B
K 4	2.21 cd	2.31 b	2.10 ef	1.85 ij	2.12 B	0.26	0.29	0.29	0.25	0.27 A	1.29 b	1.31 b	1.48 a	0.98 def	1.26 A
K 5	1.86 ij	1.86 ij	1.86 ij	1.60 k	1.80 E	0.20	0.20	0.23	0.15	0.20 E	0.77 ij	0.87 gh	0.95 efg	0.65 k	0.81 E
Mean	2.06 B	2.14 A	2.02 C	1.78 D		0.24 B	0.25 A*	0.26 A	0.20 C		0.98 C	1.11 B	1.18 A	0.82 D	

Means having * the same letters or ** not having letters are not significantly differ at 5% level.

Means having the same letters or not having letters are not significantly differ at 5% level.

Means having the same letters or not having letters are not significantly differ at 5% level.

Means having the same letters or not having letters are not significantly differ at 5% level.

Means having the same letters or not having letters are not significantly differ at 5% level.

Means having the same letters or not having letters are not significantly differ at 5% level.

Means having the same letters or not having letters are not significantly differ at 5% level.

Means having the same letters or not having letters are not significantly differ at 5% level.

Means having the same letters or not having letters are not significantly differ at 5% level.

Gowda, A. M.

Means having the same letters or not having letters are not significantly differ at 5% level.