

FERTILIZATION CAPACITY OF ROOSTER SPERMATOZOA UPON THE BIOCHEMICAL CHANGES OF SEMEN

(Received: 11. 12. 2012)

By

E. A. El-Gendy, M. M. Ahmed*, Sh. M. El-Tantawy and Sh. A. Ibrahim*

Department of Animal Production, Faculty of Agriculture, Cairo University, Giza - Egypt.

**Department of Animal Production, National Research Center, Dokki, Giza - Egypt.*

ABSTRACT

An experiment was carried out to study the effects of biochemical changes in rooster semen on the fertilization capacity of the spermatozoa. Chickens of two lines (CE2 and CE4) were used. Seven treatments of semen were designed and included the incubation of sperm with a bacterial plasmid (pUK18), a mixture of the plasmid and lipofectin at 5 or 2.5% concentration and the incubation of spermatozoa with lipofectin and a semen extender (BPSE). The progenies were obtained from the insemination of hens by the semen of different treatments. Sperm motility was greatly influenced by the treatments. Motility was significantly the highest in the control semen and averaged 92.42% and highly significantly declined to the renege of 52.08 and 58.75% in the semen samples treated with the plasmid, lipofectin at 2.5 or 5% concentration and diluted with BPSE. The percentage of live sperm was not affected by the addition of the plasmid. The addition of the plasmid and lipofectin or the diluent BPSE resulted in a significant reduction in the percentage of live sperms. The percentage of live sperms ranged 59-62% when the plasmid, lipofectin and BPSE were all together added to the semen samples. The percentages of dead and abnormally-shaped sperm reached 26.88 and 17.13% respectively, in the semen treated with plasmid, lipofectin 5% and BPSE. Fertility averaged 88.22% in the eggs of hens inseminated with the control semen, and significantly decreased to 42.14% when semen was incubated with the plasmid pUC18, and reached 58.98% when semen was treated with plasmid, lipofectin (5%) and BPSE.

Key words: *bacterial plasmid, lipofectin, rooster spermatozoa, semen dilution.*

1. INTRODUCTION

Semen characteristics can be affected by age of the rooster, lighting schedule, season, body weight, and diet (Sexton, 1986 & 1987), as well as the changes in the biochemical composition of the semen (El-Gendy *et al.*, 2007). The characteristics include the concentration, motility, viability and morphology of spermatozoa. Many of these parameters correlate with the fertilizing capacity of the spermatozoa.

Fowl spermatozoa show similar patterns of motility regulation with more than 60% of spermatozoa motile at 30°C and virtually no spermatozoa motile at 40°C (Wishart and Wilson, 1999). Florescent strains have successfully been used to determine live and dead sperm ratios (Lake and Stewart, 1978; Bilgili and Renden, 1984; and Bayyari *et al.*, 1990).

Fowl semen is low in volume, but is highly concentrated with spermatozoa. Therefore, the dilution of semen is common to increase its volume and to increase the number of hens that

can be inseminated. The proper dilution uniformly distributes spermatozoa, so that the proper dosage can be delivered during insemination. It also sustains and protects spermatozoa during short or long term storage. Semen dilution is based on the biochemical composition of chicken semen. Because glutamic acid is the most prominent anionic constituent of avian seminal plasma, so it is considered a standard component of most of the diluents (Lake and McIndoe, 1959). Sexton (1977) developed a phosphate buffered semen diluent known as Beltsville Poultry Semen Extender (BPSE). This diluent was successfully used to assess the insemination dose required for optimum fertility following short-term storage. Fertility value of more than 88% was reported for white leghorn hens weekly inseminated with 20 million spermatozoa suspended in BPSE.

Fertility levels of more than 90% were achieved when a dose of 100 million sperm cells suspended in BPSE was used. It was reported that a dilution rate of 1:4 and a weekly insemination

dose of 20 million sperms was determined to be the maximum extension rate of chicken semen diluted in the BPSE. The function of the extender BPSE as a storage medium was assessed by Sexton and Fewlass (1978), using various diluent components for fowl semen stored at 5°C. Spermatozoa maintained the capability of fertilization in diluents with osmolarities ranging 250-460 mosM/kg H₂O. In hypo-osmotic conditions, spermatozoa displayed increased incidence of bent necks. This defect was frequently found in diluted chicken and turkey semen, and was negatively correlated with fertility (Bajpai and Brown, 1964; and Clark *et al.*, 1984). Giesen and sexton (1982) observed a disappearance of turkey spermatozoa over an 18-hour storage period, suggesting that spermatozoa swell and burst *in vitro*. It was hypothesized that hypertonic diluents could improve sperm survival *in vitro* (El-Gendy *et al.*, 2007).

(1987) and Sato *et al.* (2003) indicated that lipofectin stabilizes exogenous DNA and keeps it intact. The fertility of cock spermatozoa was detracted when treated with lipofectin (Rottmann *et al.*, 1992; Squires and Drake, 1993). However, Trefil *et al.* (1996) observed continuous egg fertility for three weeks of hens inseminated with lipofectin-treated spermatozoa.

Because of their natural role in insemination, that spermatozoa are interestingly able to seek out the female pronucleous with such precision seems reasonable enough to think that they may be utilized to deliver exotic DNA to the target ovum and also at the same time resulting in the successful integration into the genome of the introduced gene (Khoo *et al.*, 1992; and Yin *et al.*, 2009).

The objective of this study was to evaluate the fertilization capacity of cock sperms under different conditions of dilution, loading with lipofectin and bacterial plasmid.

Table (1): The chemical composition of the Beltsville poultry semen extender (BPSE)

Constituent	grams/liter
Potassium diphosphaate.3H ₂ O	12.70
Sodium glutamate	8.67
Fructose (anhydrous)	5.00
Sodium acetate.3 H ₂ O	4.30
TES ¹	1.95
Potassium citrate	0.64
Potassium monophosphate	0.65
Magnesium chloride	0.34
d H ₂ O , was added to the mixture to reach 1000 ml	
pH	7.50
Osmotic pressure (m.Osm./kg.H ₂ O)	333

¹, N-tris (Hydroxymethyl) methyl-2-aminoethane sulfonic acid

Light microscopic autoradiograms showed that radioactive DNA was associated with a number of sperm cells but not all the sperm cells were labeled indicating that some sperm cells were unable to pick up DNA. Evidence from electron microscope examination of the autoradigraphs obtained after *in situ* hybridization showed that foreign DNA was present within the sperm nucleus. This clearly showed that DNA had been internalised into the sperm head. The fluorescence *in situ* hybridization used by Nakanishi and Iritani (1993) showed that 51.6% of the exogenous DNA-lipofectin-treated sperm retained the exogenous DNA. Fellgner *et al.*

2. MATERIALS AND METHODS

2.1. Genetic background of the chickens

Two lines of chickens (CE2 and CE4) were used in this experiment. Line CE2 is normally feathered and line CE4 is naked neck. Both lines have been originally derived from a heterozygous naked-neck local chicken population in Egypt (El-Gendy, 2009). Both lines are being maintained in small randombred flocks for 10 generations.

2.2. Experimental design

Ten males and 20 females were randomly assigned in each line. Semen samples were individually collected from the males of each line

and were immediately pooled to two samples, 5 males each. Seven different treatments were applied to the pooled semen samples. In treatment 1 (T₁), semen was used as the control treatment. In treatment 2 (T₂), semen was treated with the plasmid pUC18 (2.5 µg /100 µl semen), and then heat incubated. In treatments 3 and 4 (T₃ and T₄), semen was treated with a mixture of the plasmid pUC18 (2.5 µg /100 µl semen) and lipofectin (Life Technologies, Carlsbad, California, USA) at 5.0 µl /100 µl semen (T₃) or 2.5 µl/100 µl semen (T₄), and then heat incubated. In treatments 5 and 6 (T₅ and T₆), semen was diluted (1:4, v:v) with Beltsville poultry semen extender (BPSE, Table 1) proposed by Sexton (1977), treated with a mixture of the plasmid pUC18 (2.5 µg /100 µl semen) and lipofectin at 5.0 µl /100 µl semen (T₅) or 2.5 µl /100 µl semen (T₆), and then heat incubated. In treatment 7 (T₇), semen was diluted with BPSE (1:4, v:v), mixed with the plasmid pUC18 (2.5 µg /100 µl semen), and then heat incubated. Heat incubation was by the exposure of semen to 37°C for 30 minutes using water bath. Each treatment was repeated three times, forming three replicates. For all treatments, hens of each line were inseminated with the treated semen. Four hatches were obtained. Eggs were collected for 10-15 days for each hatch and were incubated. Upon hatch, chicks were wing banded, by line and dam. All chicks per hatch were housed in floor chambers in a conventional open house with routine management.

2.3. Sperm capacity

Sperm capacity was assessed in all treatments by the estimation of:

2.3.1. Sperm concentration: Sperm concentration was determined, where the semen of the given treatment was diluted at a rate of 1:200 using a physiological saline solution (1ml NaCl 3%, 2ml eosin 2%, and 50ml deionized water). A drop of the diluted semen was put on a standard hemacytometer slide which is equally divided into 25 squares, each is also equally sub-divided into 16 cells, each cell is 1/20 x 1/20 x 1/10 mm³. Sperms were counted as the sum of sperm counts in five squares on the slide, the four squares at the corners and the square at the center. Sperm concentration was then calculated according to Sorensen (1979):

Sperm concentration (sperm/ml) = [counted sperm x dilution rate x 4000 x 1000] ÷ (5x16) The dilution rate was 200, and the value of 4000 denotes to the product of the cell volume.

2.3.2. Sperm motility: Individual motility of the sperm was determined. A drop of the semen of

each treatment was mixed with 200 µl of BPSE and warmed to 37°C under light microscope. The individual motility was subjectively and numerically estimated on a rank from zero for no individual motility to 100% for maximum individual motility (Morisson *et al.*, 1997).

2.3.3. Sperm viability: The parameters of spermatozoa viability included percentages of live, dead and abnormally-shaped sperm. Forty microliters of semen samples of each treatment were added to 150 µl of the staining solution (eosin 16 g/l and nigrosin 60 g/l in BPSE) and put onto slide. Two minutes later, smears were performed for each sample and spermatozoa were observed, using microscope with an oil immersion objective. Live spermatozoa were seen white in color because they were eosin- impermeable. However, dead spermatozoa were pink because they became eosin-permeable. Two hundred spermatozoa per sample were observed and the percentages of live, dead and abnormally-shaped spermatozoa were estimated (Sorensen, 1979).

2.3.4. Fertility and hatchability: Fertility and hatchability were calculated for each treatment and chicken line.

2.3.5. Statistical analysis

The data set of sperm characteristics was subjected to the analysis of variance using the statistical analysis system (SAS, 1999). The sources of variances included effects of line, treatment, replication and the interaction between line and treatment. Line and treatment significance were assessed by least squares means.

3. RESULTS AND DISCUSSION

3.1. Sperm characteristics

Table (2) presents the levels of significance of different effects on sperm capacity. The results indicate that semen characteristics were highly significantly influenced by treatment differences. Also, replication influenced only sperm viability measurements (percentages of live, dead and abnormally-shaped spermatozoa). Neither line differences nor line by treatment interaction influenced the sperm characteristics. These results reveal that sperm capacity influenced by the biochemical changes in semen composition. The effect of the genetic background on sperm characteristics was totally absent.

Semen characteristics under different treatments are shown in Table (3). Sperm concentration was significantly the highest (4.1*10⁹ sperm/ml³) in the control sample (T₁). The incubation of semen with the plasmid pUC18 (T₂) significantly reduced the sperm concentration

Table (2): Levels of significance of different sources of effects on sperm capacity

Source	Df	Concentration	Motility	Live	Dead	Abnormal
Model	15	< 0.0001	< 0.0001	< 0.0001	< 0.0002	< 0.0001
Line (L)	1	0.7735	0.3759	0.6862	0.1504	0.4703
Treatment (T)	6	< 0.0001	< 0.0001	< 0.001	< 0.0001	< 0.0001
Replication	2	0.2104	0.0566	0.0014	0.0106	< 0.0001
L * T	6	0.5278	0.7290	0.5183	0.4429	0.8753

by 18.14%. The incubation of semen with the plasmid in the presence of lipofectin (T₃ and T₄) reduced the sperm concentration by about 26-42%. Also, the dilution of semen with BPSE (T₇) resulted in a significant reduction in sperm concentration by 51.55%, and was not significantly different from that in the semen samples treated with plasmid, lipofectin and BPSE (T₅ and T₆). Therefore, it is obvious that sperm concentration was greatly changed when semen samples were subjected to different treatments. Hafez (1974) reported that fowl semen although is low in volume it is highly concentrated. A cock eject varies from 0.2 to 0.5 ml and contains an average density of about 3×10^9 sperm/ml.

Sperm motility was significantly the highest in the control semen (T₁) and was not significantly changed when incubated with the plasmid (T₂). A dramatic and significant decline by 32.83 and 26.96% in sperm motility accompanied the treatment of semen with lipofectin at 5 and 2.5% concentrations, respectively (T₃ and T₄). Motility significantly declined by 22.46% when semen was diluted with BPSE (T₇). The motility declined more when lipofectin was at 5% concentration than when was at 2.5% concentration, and with or without BPSE. This means that lipofectin primarily impedes the sperm motility, and this motility impedance positively correlated with lipofectin concentration. The depression in fowl spermatozoa motility in response to different environmental changes was also reported by Wishart and Wilson (1999), where more than 60% of spermatozoa motile at 30°C and virtually no spermatozoa motile at 40°C.

The percentage of live sperms was 84.83% in the control semen (T₁) and was not significantly affected by the incubation with the plasmid (T₂). The incubation of semen with the mixture of the plasmid and lipofectin (T₃ and T₄) or the dilution of semen by BPSE (T₇), resulted in a significant reduction by 15.12 - 21.22% in the percentage of live sperms and the reduction was even more and reached 26.28 and 30.35% when the plasmid, lipofectin (5 or 2.5% concentration) and BPSE were all together incubated with the semen (T₅

and T₆). Accordingly, the percentage of dead sperms significantly increased in all treatments compared to that observed in the control treatment (7.75%). The increase in percentage of dead sperms varied from about 2-folds in the semen treated with the plasmid (T₂) to about 2.5-folds in the semen treated with the plasmid, lipofectin at 2.5% concentration and BPSE (T₆). The treatment of semen with only BPSE (T₇) resulted in increasing the percentage of dead sperms by about 1.5-folds.

Similar to the increasing trend of the percentage of dead sperms due to the different treatments of semen, the percentage of abnormally-shaped sperms increased but with less magnitude. The abnormally-shaped sperms basically represented 7.42% in the control semen (T₁), and increased to as high as about 1.3-folds in the semen treated with plasmid, lipofectin at 2.5% concentration and BPSE (T₆). It seems that the percentages of live, dead and abnormally-shaped spermatozoa were significantly influenced by the addition of the plasmid, lipofectin and BPSE to the semen; however the addition of plasmid only to the semen did not alter the percentages of live, dead and abnormally-shaped sperms. These results indicate that many sperms were damaged by the changes in semen biochemical composition, resulting in significant increases in the sperm mortality and misshaping. Morisson *et al.* (1997) reported significant differences in semen characteristics between two lines of Rhode Island Red chickens. The percentages of live, dead and abnormal spermatozoa were 92, 3.6, and 4.3% in one line *versus* 89, 6.1 and 5.2% in the other line.

3.2. Fertility and hatchability

Because the concentration, motility, viability and morphology of spermatozoa correlate with the fertilizing potential of the spermatozoa, so the fertility and hatchability were obtained for the collected eggs of each treatment and line (Table 4). Fertility was 90.72 and 85.71% in lines CE2 and CE4 respectively, with an average of 88.22% in the eggs of hens inseminated with control semen (T₁). Fertility remarkably and significantly decreased in both lines among different treatments

Table (3): Sperm Characteristics (LSM±SE), overall lines, under different semen treatments.

Treatment	No.	Concentration		Motility		Live		Dead		Abnormally-shaped	
		* 10 ⁹	↓	%	↓	%	↓	%	↑	%	↑
T ₁	12	4.19 ± 0.26 ^a		92.42 ± 3.00 ^a		84.83 ± 2.95 ^a		7.75 ± 3.03 ^c		7.42 ± 1.45 ^d	
T ₂	12	3.43 ± 0.26 ^b	18.14	85.92 ± 3.00 ^a	7.03	80.88 ± 2.95 ^a	4.66	10.67 ± 3.03 ^{bc}	37.68	9.88 ± 1.45 ^{cd}	33.15
T ₃	12	3.08 ± 0.26 ^{bc}	26.49	62.08 ± 3.00 ^c	32.83	72.00 ± 2.95 ^b	15.12	18.87 ± 3.03 ^{ab}	143.48	8.00 ± 1.45 ^{cd}	7.82
T ₄	12	2.41 ± 0.26 ^{cd}	42.48	67.50 ± 3.00 ^{bc}	26.96	71.54 ± 2.95 ^b	15.67	21.13 ± 3.03 ^a	172.65	9.29 ± 1.45 ^{cd}	25.20
T ₅	12	2.55 ± 0.26 ^{cd}	39.14	52.08 ± 3.00 ^d	43.65	62.54 ± 2.95 ^c	26.28	26.13 ± 3.03 ^a	237.16	14.75 ± 1.45 ^{ab}	98.79
T ₆	12	2.31 ± 0.26 ^{cd}	44.87	58.75 ± 3.00 ^d	36.43	59.08 ± 2.95 ^c	30.35	26.88 ± 3.03 ^a	246.84	17.13 ± 1.45 ^a	130.86
T ₇	12	2.03 ± 0.26 ^d	51.55	71.66 ± 3.00 ^b	22.46	66.83 ± 2.95 ^{bc}	21.22	20.25 ± 3.03 ^a	161.29	11.83 ± 1.45 ^{bc}	59.43

^{a-d}, LS means of same trait with different superscripts are significantly different (P ≤ 0.05).

↓ or ↑, indicates the percentage of reduction or increase in the sperm character due to different treatments, compared to the control treatment (T₁).

T₁ = non-treated semen (control semen). T₂ = semen treated with the plasmid. T₃ = semen treated with the plasmid and lipofectin 5%.

T₄ = semen treated with the plasmid and lipofectin 2.5%. T₅ = semen treated with the plasmid and lipofectin 5% and BPSE.

T₆ = semen treated with the plasmid and lipofectin 2.5% and BPSE. T₇ = semen treated with the plasmid and BPSE.

Table (4): Effects of different semen treatments on fertility and hatchability.

Treatment	Fertility			Hatchability		
	Line CE2	Line CE4	X ± SE	Line CE2	Line CE4	X ± SE
T ₁	90.72	85.71	88.22 ± 2.51 ^a	75.00	91.67	83.34 ± 8.34 ^a
T ₂	35.62	48.65	42.14 ± 6.52 ^{bc}	69.23	55.56	62.40 ± 6.84 ^{ab}
T ₃	63.89	44.83	54.36 ± 9.53 ^b	41.30	46.15	43.73 ± 2.43 ^b
T ₄	24.64	22.54	23.59 ± 1.05 ^c	82.35	68.75	75.55 ± 6.80 ^a
T ₅	45.45	72.50	58.98 ± 13.53 ^b	40.00	37.93	38.97 ± 1.04 ^b
T ₆	44.57	63.33	53.95 ± 9.38 ^b	90.24	57.89	74.07 ± 16.18 ^a
T ₇	31.03	46.81	38.92 ± 7.89 ^c	55.56	68.18	61.87 ± 6.31 ^{ab}
X ± SE	47.99 ± 8.57	54.91 ± 7.84		64.81 ± 7.45	60.88 ± 6.62	

^{a-c}, trait means accompanied by different superscripts are significantly different (P ≤ 0.05).

T₁ = non-treated semen (control semen). T₂ = semen treated with the plasmid. T₃ = semen treated with the plasmid and lipofectin 5%.

T₄ = semen treated with the plasmid and lipofectin 2.5%. T₅ = semen treated with the plasmid and lipofectin 5% and BPSE.

T₆ = semen treated with the plasmid and lipofectin 2.5% and BPSE. T₇ = semen treated with the plasmid and BPSE.

of semen, with no obvious trend. The incubation of spermatozoa with the plasmid pUC18 (T₂) significantly decreased the average fertility to 42.14%. Fertility was 23.59% when semen was incubated with the plasmid and lipofectin 2.5% (T₄), and reached 53.95% when semen was treated with plasmid, lipofectin 2.5% and BPSE (T₆). In three of four lipofectin treatments of semen, fertility was not reduced as much as the reduction in other semen treatments in which lipofectin was not used. No significant difference in fertility was obtained between both lines.

Hatchability averaged 83.34% in both lines for the eggs of hens inseminated with untreated control semen (T₁), and was significantly decreased to 62.40% upon the treatment of semen with the plasmid (T₂). The reduction in hatchability was obvious and reached 43.73 and 38.97% when the semen was treated with the plasmid and lipofectin 5% (T₃), and the diluents BPSE as well (T₅). But the treatment of semen with the plasmid, lipofectin 2.5% without the diluents BPSE (T₄) and with the diluents BPSE (T₆) resulted in hatchability of 75.55 and 74.07%, respectively. It might be of meaning that lipofectin in general sustained and protected the plasmid and also diminished the deleterious effect of incubation of semen with the plasmid on the fertilizing potential and hatching.

According to Rottmann *et al.* (1992) and Squires and Drake (1993), the fertility of cock spermatozoa was detracted when treated with lipofectin. Trefil *et al.* (1996) observed continuous egg fertility for three weeks of hens inseminated with lipofectin-treated spermatozoa, and the fertility reached 52.3% during the third week. The fluorescence *in situ* hybridization used by Nakanishi and Iritani (1993) showed that 51.6% of the exogenous DNA-lipofectin treated sperm retained the exogenous DNA. The fertility of eggs was 47% for the DNA incubated with sperm, 23% for the electroporated sperm and 67% for the lipofectin-treated sperm.

4. REFERENCES

- Bajpai P.K. and Brown K.E. (1964). The effect of different temperatures on the metabolic activity, morphology, and fertilizing capacity of turkey semen. *Poult. Sci.*, 43:1501-1508.
- Bayyari G.R., Cook J.R., Harris G.C., Macy L.B., Slavik M.F., and Skeeles J.K. (1990). The evaluation of chicken spermatozoa using fluorescent staining in a 96-well format. *Poult. Sci.*, 69:1602-1605.
- Bilgili S.F. and Renden J.A. (1984). Fluorometric determination of avian spermatozoa viability and concentration. *Poult. Sci.* 63: 2275-2277.
- Clark R.N., Bakst M.R. and Ottinger M.A. (1984). Morphological changes in chicken and turkey spermatozoa incubated under various conditions. *Poult. Sci.* 63:801-805.
- El-Gendy E.A. (2009). A model for the genetic employment of chickens local to warm climate. 1. Crossing with a fast growing strain and growth patterns of the crossbreds. *International Journal of Poultry Science*, 8:299-306.
- El-Gendy E. A., Gad A.Y. and Mostageer A. (2007). Sperm-mediated gene transfer in poultry 1. The relationship with cock sperm viability. *Arab. J. Biotech.*, 10 (1):1-12.
- Fellgner P.L., Gadek T.R., Holm M., Roman R., Chan H.W., Wenz M., Nortro J.P., Ringold G.M. and danielsen M. (1987). Lipofection: A highly efficient, lipid-mediated DNA-transfection procedure. *Proc. Natl. Acad. Sci., USA* , 84: 7413-7417.
- Giesen A.F. and Sexton T. J. (1982). Boltsville Poultry Semen Extender:7.Comparison Of Commercial Diluents For Holding Turkey Semen Six Hour 15°C .*Poult. Sci.*, 62:379-381.
- Hafez E.S.E. (1974). *Reproduction in Farm Animals 3rd ed.*, LEA & FEBIGER, Philadelphia, USA, pp. 312-317.
- Khoo HW., Ang LH., Lim H.B. and Wong KY. (1992). Sperm cells as vectors for introducing foreign DNA into zebrafish. *Aquaculture*, 107:1-19.
- Lake P.E. and McIndoe W.M. (1959). The glutamic acid and creatine content of cock seminal plasma. *Biochem. J.*, 71:303-306.
- Lake P.E. and Stewart J.M. (1978). Preservation of fowl semen in liquid nitrogen –an improved method. *Poult. Sci.*, 19:187-194.
- Morisson M., Bordas A., Petit J.M., Jayat-Vignoles C., Julien R. and Minvielle F. (1997). Associated effects of divergent selection for residual feed consumption on reproduction, sperm characteristics, and mitochondria of spermatozoa. *Poult. Sci.*, 76: 425-431.
- Nakanishi A. and Iritani A. (1993). Gene transfer in the chicken by spermmediated methods. *Mol. Reprod. Dev.*, 36:258-61.
- Rottmann O.J., Antes R., Höfer P. and Maierhofer G. (1992). Liposomes mediated gene transfer

- via spermatozoa into avian eggs. J. Anim. Breed. Genet., 109: 64-70.
- SAS (1999). SAS/STAT User's Guide: Statistics. SAS Institute Inc., Cary, NC, USA.
- Sato F. Soh T., Hattori M. and Fujihara N. (2003). Evaluation of deoxyribonuclease activity in seminal plasma of ejaculated chicken semen. Asian J. Androl., 5: 213-216.
- Sexton T.J. (1986). Effects of dietary protein and season on fertility of turkey semen stored 18 hours at 58°C. Poult. Sci., 65:604-606.
- Sexton T.J. (1987). Effect of semen treatments and age of tom on fertility of unstored semen and semen held 18 hours. Poult. Sci.,66:1721-1726.
- Sexton T.J. (1977). A new poultry semen extender 1. Effect of extension on the fertility of chicken semen. Poult. Sci.,56: 1443-1446.
- Sexton T.J. and Fewlass T.A. (1978). A new poultry semen extender . Effeect of the diluent components on the fertility of chicken semen stored at 5°C. Poult. Sci.,57:277-284.
- Sorensen A. M. (1979). Animal Reproduction: Principles and Practices. McGraw-Hill, New York, USA.
- Squires J.E. and Drake D. (1993). Liposome-mediated DNA transfer to chicken sperm cells. Anim. Biotechn., 4:71-88.
- Trefil P., Thoraval P., Mika, J., Coudert F. and Dambrine G.(1996). Intramaginal insemination of hens can eliminate negative influence of lipofectin on fertilizing ability of spermatozoa. Br. Poult. Sci., 37: 661-664.
- Wishart G.J. and Wilson Y.I.(1999). Temperature-dependent inhibition of motility in spermatozoa from different avian species. Anim. Reprod. Sci., 57:229-235.
- Yin J., Zhangb J., Shic G., Xied S., Wange X. and Wange H. (2009). Sperm mediated human coagulation factor VIII gene transfer and expression in transgenic Mice. Swiss Med Wkly, 139(25 – 26): 364 – 372.

كفاءة الإخصاب للحيوانات المنوية للديوك عند تغيير التركيب الكيميائي الحيوي للسائل المنوي

عصام عباس الجندي - مروه مصطفى محمد* - شكرى محمد الطنطاوى - شوقى أحمد إبراهيم*

قسم الإنتاج الحيوانى - كلية الزراعة - جامعة القاهرة - * قسم الإنتاج الحيوانى - المركز القومى للبحوث - الدقى - الجيزة - مصر

ملخص

استهدفت هذه التجربة دراسة تأثير التغيرات البيوكيميائية فى السائل المنوي لذكور الدجاج على كفاءة الإخصاب للحيوانات المنوية. استخدم فى هذه الدراسة سلالتين من الدجاج CE4 ، CE2 ، وتم تصميم سبع معاملات للسائل المنوي تشتمل على تحضين السائل المنوي مع بلازميد بكتيرى، تحضين السائل المنوي مع خليط من البلازميد واللييوفكتين بتركيز ٥ أو ٢.٥% بالإضافة إلى مخفف للسائل المنوي (BPSE). تم تلقيح الإناث بالسائل المنوي المعامل بالمعاملات المختلفة. وجد أن حركة الحيوانات المنوية تأثرت معنوياً بالمعاملات المختلفة وكانت الأعلى (٩٢.٤٢%) فى المعاملة الكونترول وإنخفضت معنوياً فى مدى تتراوح بين ٥٢.٠٨ ، ٥٨.٧٥% وبترتيب تنازلى فى السائل المنوي المعامل بالبلازميد والمخفف واللييوفكتين بتركيز ٥ ، ٢.٥% على التوالى. لم تتأثر نسبة الحيوانات المنوية الحية عند إضافة البلازميد إلى السائل المنوي، بينما إنخفضت معنوياً فى مدى ٥٩ - ٦٢% فى العينات المعاملة بالبلازميد واللييوفكتين بتركيزه والمخفف. كانت نسبتي الحيوانات المنوية الميتة والشاذة ٢٦.٨٨ ، ١٧.١٣% على التوالى فى السائل المنوي المعامل بالبلازميد واللييوفكتين بتركيز ٥% والمخفف. كانت نسبة الخصوبة فى بيض الدجاج الملقح بالسائل المنوي الكونترول ٨٨.٢٢% وإنخفضت معنوياً إلى ٤٢.١٤% فى بيض الدجاج الملقح بالسائل المنوي المعامل بالبلازميد ووصلت إلى ٥٨.٩٨% فى بيض الدجاج الملقح بالسائل المنوي المعامل بالبلازميد واللييوفكتين بتركيز ٥% والمخفف.

المجلة العلمية لكلية الزراعة - جامعة القاهرة - المجلد (٦٣) العدد الرابع (أكتوبر ٢٠١٢): ٤٠٦-٤١٢.