

USING YEASTS IN LAYING HEN DIETS

Abd-Elsamee, M. O.; A. S. Shalaby and A. M. Ali.

Depart. of Animal Production, Faculty of Agric. Cairo University.

ABSTRACT

The current study was conducted to evaluate the effect of adding inactive or active dried yeast (*Saccharomyces Cerevisiae*) into laying hen diets on their performance and egg quality. For this purpose a total number of 900 white Hy line hens 24 weeks old were divided into nine treated groups of 100 laying hens each. For 16 weeks experimental period the groups were fed on basal diet supplemented with 5 graded levels of active or inactive dried yeast as 0.0% (control), 2%, 4%, 6% and 8%. These experimental diets were formulated to be iso-caloric. The obtained results showed that:

- 1- Adding yeast either inactive or active to layer diets at most tested levels increased both egg production and egg weight compared to the control group.
- 2- Adding yeast culture to layer diets at most tested levels significantly reduced the average feed intake and significantly ($P \leq 0.05$) improved feed conversion ratio compared to the control group.
- 3- Values of most egg quality traits especially shell thickness, yolk index and shape index were not affected significantly by supplying layer diets with either inactive or active yeast at most tested levels in comparison to the control group.
- 4- Supplying layer diets with inactive or active yeast at each tested level improved both crude protein and ether extract digestibilities and nitrogen balance % compared to the control group.

Conclusively, it could be recommended to incorporate 2% inactive yeast or 6% active yeast to laying hen diets for improving their performance, egg quality and economic efficiency.

Keywords: (yeasts, laying hens, performance, egg quality.)

INTRODUCTION

Probiotics have been defined as live microbial feed supplements that beneficially affect host animals by improving their intestinal microbial balance (Fuller, 1977; Bardly *et al.*, 1994 and Abd El-Azeem, 2002). They found that addition of yeast to diets improve growth performance and feed conversion ratio in layers (Scott *et al.*, 1982 and Santin *et al.*, 2003) and egg production (Abdulrahim *et al.*, 1996; Maia *et al.*, 2001 and Osman, 2003) feed consumption (Kornegay *et al.*, 1995 and Park *et al.*, 2001) egg weight (Tortuero and Fernandez, 1995 and Jin *et al.*, 1997) as well as egg mass and egg size in layers (Park *et al.*, 2002 and Santin *et al.*, 2003) and egg specific gravity (Mohan *et al.*, 1995 and Yildirm and Parlat, 2003).

Recently, active live yeast and non active live dry yeast have been successfully examined as satisfactory alternative to antibiotics feed additives due to its antagonistic effect against harmful pathogenic bacteria (Wakwak *et al.*, 2003). Furthermore, live yeast and dry yeast are effective in counteracting aflatoxin or redestine (Stanley *et al.*, 1993).

The objective of the present study to determine the effect of both live yeast and inactive live dry yeast on laying hen performance and egg quality.

MATERIALS AND METHODS

The experimental work of the present study was carried out at the poultry farm of the El-Shark El-Aousat Company (MELARKGYPT). The chemical analytical part was performed at the Laboratories of Animal Production Department (Animal Nutrition), Faculty of Agriculture, Cairo University.

The objective of the study was to investigate the effect of using different sources and levels of yeast as a partial replacing for soy bean meal on laying hen performance, egg quality, nutrients digestibility and economic efficiency.

Experimental design:

Two sources of yeast and four levels of replacement were studied in this work. Yeast sources were inactive and active yeast. The different levels of replacement were 2, 4, 6 and 8% of soybean meal. Accordingly, 8 (2x4) experimental diets were formulated, in addition, the control diet containing corn-soy without either inactive or active yeast was used. Therefore, a total of 9 experimental treatments were performed.

Experimental diets:

Composition of the 9 basic experimental diets used in this study and their calculated analysis are presented in table (1).

Table (1): Composition and chemical analysis of the experimental diets:

Ingredients	Basal Diets (%)								
	Control	Yeast ₁ (Inactive)				Yeast ₂ (Active)			
		2%	4%	6%	8%	2%	4%	6%	8%
Yellow Corn	57.7	57.6	57.5	57.5	57.4	57.6	57.5	57.5	57.4
Soy bean meal (44% CP)	24.5	22.5	20.5	18.5	16.5	22.5	20.5	18.5	16.5
Gluten (48 % CP)	3	3	3	3	3	3	3	3	3
Yeast	0	2	4	6	8	2	4	6	8
Oil	2.5	2.6	2.7	2.7	2.8	2.6	2.7	2.7	2.8
Bone meal	2.8	2.8	2.8	2.8	2.8	2.8	2.8	2.8	2.8
Lime stone	8.5	8.5	8.5	8.5	8.5	8.5	8.5	8.5	8.5
Salt	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4
Premix *	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
Methionine	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Lysine	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Calculated Values **									
Crude protein%	18	18	18	18	18	18	18	18	18
ME K cal/kg	2850	2850	2850	2850	2850	2850	2850	2850	2850
Crude fiber %	2.58	2.58	2.58	2.58	2.58	2.58	2.58	2.58	2.58
Ether extract %	3.04	3.04	3.04	3.04	3.04	3.04	3.04	3.04	3.04
Methionin %	0.52	0.52	0.51	0.51	0.51	0.52	0.51	0.51	0.51
Lysine %	0.98	0.96	0.95	0.94	0.92	0.96	0.95	0.94	0.92
Ca %	4	4	4	4	4	4	4	4	4
Av. Phosphorus%***	0.47	0.48	0.52	0.54	0.56	0.48	0.52	0.54	0.56
Linoleic acid%	1.37	1.37	1.37	1.37	1.37	1.37	1.37	1.37	1.37

* Vitamin and mineral premix at 0.3% of the diet supplies the following per kg of the diet: Vit. A 10000 IU, Vit. D₃ 3000 IU, Vit. E 20 mg, Vit. K₃ 3 mg, Vit. B₁ 2 mg, Vit. B₂ 6 mg, Pantothenic acid 10 mg, Folic acid 1mg, Biotin 5mcg, Choline Chloride 500 mg, Niacin 66mg, Vit. B₆ 5 mg, Vit. B₁₂ 20 mcg, Mn 100 mg, Fe 100 mg, Zn 75mg, Cu 8 mg, I₂ 45 mcg and Se 10mcg.

** According to NRC (1994).

*** Av = Available.

The control diet (No. 1) was formulated to meet the requirement of crude protein without using the tested yeast sources. The different sources of yeast used in this study were *Saccharomyces Cervisiae* in inactive or active form. Two yeast protein products were used in this study. The first one was inactive dried yeast, grown on molasses, product of the Egyptian Sugar and Distillation Company, Hawamdia. While, the second source was active dried yeast, product of the Egyptian Strach Products and Yeast Company, Alexandria. Each one of the tested yeast sources was replaced for soybean meal at levels of 2, 4, 6 and 8%. In all experimental diets, amino acids, vitamins and minerals were adjusted according to the strain recommended catalog. The experimental diets were formulated to meet the nutrient requirements according to the recommended allowances of the Hy-lin breed, which 18% CP and 2850 kcal ME/kg.

Experimental birds and management:

A total number of 900 white Hy-lin laying hens, 24 weeks of age were used. Hens were randomly divided into 9 groups of 100 hens in four replicates of 25 hens each. Birds were kept in cleaned and fumigated cages of wire floored batteries in closed system house. Feed and water were offered ad-libitum all over the experimental period (16 weeks) from 24 to 40 week of age, under a total of 16 hours light per day regimen.

Measurements and methods of interperetating results Laying hen performance:

The daily eggs laid were recorded on a hen day (HD) basis. The average daily egg production was calculated per hen every four weeks intervals. The average weight was recorded per hen every four weeks intervals to the nearest 0.10g. Also, the daily feed consumed per hen was calculated every four weeks intervals during the experimental periods. Records of egg production, egg weight and feed consumption were used to calculate the amount of feed (kg) required to produce one kg of eggs/hen. Therefore, feed conversion was calculated as follow:

$$\text{Feed conversion} = (\text{kg. feed}) / (\text{kg. Eggs}).$$

Egg quality:

A total of 20 eggs were taken from each treatment (5eggs from each replicate) every four weeks for testing their quality. Shape index was determined according to *Romanoff and Romanoff (1949)*. Shape index % = egg width (cm)/egg length (cm) × 100. Dry shell was weighed to the nearest 0.10g. Shell thickness was measured by using Ames shell thickness Gauge. Yolk index was calculated according to *Funk (1948)*.
Yolk index = Yolk height / diameter × 100.

The height was measured using tripod micrometer reading to the nearest 0.01cc, while the yolk diameter was measured by vernier caliber to the nearest mm. Albumen index was calculated as:
height / diameter mean × 100.

Digestion trials:

Digestion trials were conducted at the end of the experimental period to estimate the digestion coefficients and nitrogen retention of the experimental diets. A total number of 72 laying hens were used in digestion trials (8 hens from each treatments, 2 hens from each replicate). The analysis

of both diets and excreta were performed according to the Association of official Agricultural Chemists. (A.O.A.C., 1990). Faecal nitrogen was determined according to *Jakobsen et al.* (1960) and urinary organic matter was calculated according to *Abo-Raya and Galal* (1971).

Economic efficiency:

To determine the economic efficiency of egg production, the amount of feed consumed during the experimental period and the total eggs produced per treatment were obtained. The price of experimental diets was calculated according to the price of local market as well as the price of the yeast and prices of the ingredients at the time of the experiment.

Economic efficiency = the net revenue / total cost.

Statistical analysis:

Data collected in this study were statistically analyzed using the general linear models (GLM) of (SAS, 1996). One-way model was used to study the main effect of treatments on different parameters, the used model was:

$$Y_{ij} = \mu + T_i + E_{ij}$$

Where:

Y_{ij} = the observation of the parameter measured.

μ = overall mean.

T_i = effect of treatments, i (1 to 9).

E_{ij} = experimental error.

Significant differences among means were achieved using the Duncan's Multiple Range test (*Duncan*, 1955). Significant level was set at 5%.

RESULTS AND DISCUSSION

Laying hen performance:

Egg Production and egg weight:

Effect of the experimental diets on egg production and egg weight at the defferent period is presented in Table (2).

During the first period (24- 28 weeks of age). It could be noticed that layer groups fed 2% yeast₁ or 6% yeast₂ diets produced higher ($P \leq 0.05$) eggs and had higher ($P \leq 0.05$) egg weight than the control group. Nevertheless, the opposite trend was noticed when birds group fed yeast₂ 8% in comparison to the control group. While, at the second period (28-32 weeks of age), results revealed that layer groups fed yeast₁ (2, 4, 6 and 8 %) or yeast₂ (2, 4 and 6%) diets had higher ($P \leq 0.05$) egg production and egg weight compared to the control group. However, layers group fed yeast₂ 8% diet recorded less ($P \leq 0.05$) egg production and a lower insignificant egg weight compared to the control group. At the third period (32-36 weeks of age), layer groups fed yeast₁ 2% or 6% and yeast₂ 2% or 6% had higher ($P \leq 0.05$) egg production compared to the control group. Besides, egg weight values were higher ($P \leq 0.05$) with layer groups fed 2% yeast₁ or 6% yeast₂ than the control group. At the forth period (36- 40 weeks of age). layer group fed 2% yeast₁ diet produced egg higher ($P \leq 0.05$) than the control group,

while the production was significantly ($P \leq 0.05$) lower with that fed 8% yeast₂ in comparison to the control group. Also, there were insignificant differences in egg weight values among the different experimental groups except 8% yeast₂ for egg weight whereas the values were significantly ($P \leq 0.05$) lower than the control group.

Table (2): Effect of the experimental diets on egg production and egg weight.

Treatments	The first period (24-28 wk)		The second period (28-32 wk)		The third period (32-36 wk)		The fourth period (36- 40 wk)		Overall period (24-40 wk)	
	EP%	EW(g)	EP%	EW(g)	EP%	EW(g)	EP%	EW(g)	EP%	EW(g)
Control	70.08 ^b	51.33 ^b	72.67 ^c	52.91 ^c	74.88 ^b	60.50 ^b	82.33 ^b	63.34 ^a	74.99 ^b	58.02 ^{ab}
2% - Y ₁	81.43 ^a	60.00 ^a	85.02 ^a	61.55 ^a	84.91 ^a	64.36 ^a	88.88 ^a	64.89 ^a	87.56 ^a	62.70 ^a
4% - Y ₁	74.81 ^{ab}	50.66 ^{ab}	80.90 ^{ab}	57.26 ^{ab}	81.16 ^{ab}	62.19 ^{ab}	83.81 ^b	62.45 ^{ab}	80.17 ^{ab}	58.14 ^{ab}
6% - Y ₁	77.90 ^{ab}	50.00 ^b	81.95 ^{ab}	57.88 ^{ab}	82.79 ^a	62.69 ^{ab}	85.00 ^{ab}	63.03 ^a	81.91 ^{ab}	58.14 ^{ab}
8% - Y ₁	74.81 ^{ab}	52.33 ^{ab}	79.16 ^b	57.91 ^{ab}	79.70 ^{ab}	62.76 ^{ab}	83.77 ^b	63.40 ^a	79.36 ^b	59.10 ^{ab}
2% - Y ₂	78.36 ^a	47.33 ^c	80.04 ^{ab}	56.33 ^{ab}	82.86 ^a	60.73 ^b	85.14 ^{ab}	61.41 ^{ab}	81.60 ^{ab}	56.45 ^b
4% - Y ₂	74.52 ^{ab}	50.00 ^b	74.85 ^b	57.26 ^{ab}	79.31 ^{ab}	62.40 ^{ab}	83.12 ^b	62.52 ^{ab}	77.95 ^b	58.04 ^{ab}
6% - Y ₂	81.00 ^a	56.66 ^a	84.55 ^a	59.64 ^a	85.21 ^a	64.08 ^a	87.72 ^{ab}	64.34 ^a	84.62 ^a	61.18 ^a
8% - Y ₂	67.16 ^c	57.33 ^a	67.16 ^d	59.65 ^a	72.16 ^b	60.48 ^b	75.76 ^c	60.62 ^b	70.56 ^c	56.02 ^b

a, b, c, d . . . Mean each column bearing the same superscripts don't differ significantly ($P \leq 0.05$)
 Y₁: Inactive yeast 2 %, 4%, 6% & 8%. Y₂: Active yeast 2 %, 4%, 6% & 8%.

Generally, feeding layer hens on diets supplemented with 2% yeast₁ or 6% yeast₂ significantly ($P \leq 0.05$) improved egg production in comparison to the control group. Nevertheless, the values were ($P \leq 0.05$) lower than the control group when layer hens were fed diets supplemented with 8% yeast₂. The differences among the other experimental groups in comparison to the control group were insignificant as shown in Table (2).

It is worthy to note that addition yeast culture to layer diets at most tested levels resulted in an increase in both egg production and egg weight in comparison to the control group. These results are in agreement with those reported by *Tortuero and Fernandez (1995)*; *Osman et al. (2003)*; *Siam et al. (2004)*; *Mohdavi et al. (2005)*; *Abu-Taleb et al. (2005)* and *Chin et al. (2006)* who found that average egg production significantly increased due to adding active dried yeast by 1% or 1.5% into diets of laying quail for eight weeks experimental periods. The improvement in the productive performance may be attributed to the large amount of metabolites which with some viable yeast cell can be used as a probiotic as reported by *Miles and Bootwell (1991)*.

Feed intake and feed conversion ratio:

Effect of the experimental diets on feed intake (FI) and feed conversion ratio (FCR) at the different periods is listed in Table (3). The

differences in (FI) values among the experimental groups during the different experimental periods were nearly in significant. Nevertheless the values of (FCR) were significantly ($P \leq 0.05$) improved with adding yeast₁ or yeast₂ to layer diets during most experimental periods in comparison to the control group.

Table (3): Effect of the experimental diets on Feed Intake (FI) and Feed Conversion Ratio (FCR).

Treatments	The first period (24-28 wk)		The second period (28-32 wk)		The third period (32-36 wk)		The fourth period (36- 40 wk)		Overall period (24 – 40wk)	
	FI	FCR	FI	FCR	FI	FCR	FI	FCR	FI	FCR
Control	87.00 ^a	2.42 ^a	94.25 ^a	2.45 ^a	101.50 ^a	2.24 ^a	101.60 ^a	1.95 ^b	96.09 ^a	2.21 ^a
2% - Y ₁	85.00 ^{ab}	1.74 ^d	86.74 ^b	1.66 ^f	95.50 ^b	1.75 ^c	95.60 ^b	1.66 ^d	90.71 ^b	1.65 ^e
4% - Y ₁	87.09 ^a	2.29 ^b	92.40 ^{ab}	1.99 ^d	98.80 ^{ab}	1.96 ^b	98.91 ^{ab}	1.89 ^{bc}	94.3 ^a	2.02 ^b
6% - Y ₁	85.79 ^{ab}	2.20 ^b	93.90 ^a	1.98 ^d	98.60 ^{ab}	1.89 ^b	99.67 ^a	1.86 ^{bc}	94.49 ^a	1.98 ^c
8% - Y ₁	85.70 ^{ab}	2.19 ^b	92.40 ^{ab}	2.02 ^{cd}	98.80 ^{ab}	1.97 ^b	99.93 ^a	1.88 ^{bc}	94.21 ^a	2.01 ^{bc}
2% - Y ₂	85.00 ^{ab}	2.29 ^b	92.40 ^{ab}	2.05 ^{cd}	96.50 ^{ab}	1.92 ^b	95.50 ^{ab}	1.83 ^c	92.44 ^{ab}	2.01 ^{bc}
4% - Y ₂	85.00 ^{ab}	2.28 ^b	92.25 ^{ab}	2.15 ^c	95.56 ^b	1.93 ^b	96.64 ^{ab}	1.86 ^{bc}	92.36 ^{ab}	2.041 ^{bc}
6% - Y ₂	86.80 ^a	1.89 ^c	92.25 ^{ab}	1.83 ^e	95.50 ^b	1.75 ^c	95.63 ^b	1.69 ^d	92.55 ^{ab}	1.79 ^d
8% - Y ₂	86.80 ^a	2.25 ^b	92.51 ^{ab}	2.13 ^b	98.80 ^{ab}	2.26 ^a	99.94 ^a	2.18 ^a	94.51 ^a	2.25 ^a

a, b, c, d Mean in each column bearing the same superscripts don't differ significantly ($P \leq 0.05$).

Y₁: Inactive yeast 2 %, 4%, 6% & 8%. Y₂: Active yeast 2 %, 4%, 6% & 8%.

Generally, addition yeast culture inactive or active to layer diets at the most tested levels insignificantly reduced FI and Significantly ($P \leq 0.05$) improved FCR in comparison to the control group.

Its worthy to note that, the decline of feed intake resulted from addition yeast culture to layer diets at most tested levels was supported by several researches reported by *Koudela and Nyirenda* . (1995); *Balev et al.* (2001) and *Soliman* (2002) who reported that feed intake was reduced as active dried yeast added at 4g/kg into Bovans white laying hens diets at 25 weeks old for 90 days.

Egg quality:

Results of egg quality as indicated by shell weight (SW), shell thickness (ST), yolk index (YI), albumin index (AI) and shape index (SI) are presented in Table (4). Results showed that supplementation layer diets with yeast₁ at levels of 2, 4, 6 and 8% did not affect significantly on ST and YI compared to the control group. Also, the same trend was noticed with ST and SI when layer diets were supplemented with yeast₂ at levels of 2, 4, 6 and 8%. However, both AI and SI values were significantly ($P \leq 0.05$) improved with feeding layers on 4 or 6% yeast₁ diets in comparison to the control

group. The AI value was only improved ($P \leq 0.05$) when layers fed on 8% yeast₂ diets compared to the control group (74.81 vs 78.53).

Table (4): Effect of experimental diets on egg quality.

Parameteres	SW	ST	YI	AI	SI
Treatment					
Control	9.68 ^{ab}	36.69 ^{abc}	44.03 ^a	74.81 ^b	69.02 ^{bc}
2% - Y ₁	10.67 ^{ab}	39.59 ^b	42.85 ^{ab}	78.16 ^{ab}	71.38 ^{ab}
4% - Y ₁	10.03 ^{ab}	37.60 ^b	48.13 ^a	79.26 ^a	75.90 ^a
6% - Y ₁	8.53 ^b	35.41 ^{abc}	42.4 ^{ab}	79.56 ^a	76.23 ^a
8% - Y ₁	12.47 ^a	34.06 ^{bc}	42.1 ^{ab}	76.10 ^{ab}	72.48 ^{ab}
2% - Y ₂	6.95 ^{bc}	37.39 ^{abc}	40.3 ^{ab}	76.30 ^{ab}	69.58 ^b
4% - Y ₂	11.24 ^a	32.70 ^c	39.9 ^b	76.47 ^{ab}	72.90 ^{ab}
6% - Y ₂	8.15 ^{bc}	40.34 ^a	40.5 ^{ab}	76.88 ^{ab}	73.95 ^{ab}
8% - Y ₂	7.55 ^{bc}	33.38 ^{bc}	40.6 ^{ab}	78.53 ^a	67.15 ^c

a, b, c, d . . . Mean each column bearing the same superscripts don't differ significantly ($P \leq 0.05$).

SW: shell weight.

ST: shell thickness.

YI: yolk

index.AI: albumin index.

SI : Shape index.

Generally, values of most egg quality traits especially ST, YI and SI were not affected significantly by supplying layer diets with dried yeast₁ or yeast₂ at most tested levels in comparison to the control group. These observations were supported with those reported by *Piva et al. (2003)* and *Nursoy et al. (2004)* who found that egg breaking strength, egg shell thickness, egg yolk index, egg albumin index and yolk colour scores were similar among leghorn hens groups fed diets supplemented with different levels of yeast culture (0, 0.2 and 0.3% of the diet).

Digestion coefficient and nitrogen balance.

Data in Table (5) showed that supplying layer diets with 6 or 8% yeast₁ improved significantly ($P \leq 0.05$) CP and EE digestibility and nitrogen balance (%) compared to the control group. The obtained values were 94.22, 96.63 and 96.94% for CP.; 74.06, 79.13 and 79.72% for EE and 78.60, 80.43 and 80.39% for nitrogen balance %, respectively. Besides, there was a significant ($P \leq 0.05$) improvement in CP digestibility due to fed laying hens diet containing 8% yeast₂ compared to the control group (96.74 vs 94.22%). However, the differences in NFE and OM digestibility among most experimental groups were insignificant.

In general, supplementation yeast 1 or 2 in layer diets at levels of 2, 4, 6 and 8% resulted in an improvement in both CP and EE digestibilities and nitrogen balance % in comparison to the control group. This improvement may be due to the benefit effect of yeast in altering metabolism

by increasing digestive enzyme activity and decreasing bacterial enzyme activity and ammonia production and improving digestion neutralizing enterotoxins (Guillot, 2000 and Nir and Senkoylu, 2000).

Table (5): Effect of the experimental diets on digestion coefficient and nitrogen balance:

Parameters ----- Treatment	CP(%)	EE(%)	CF(%)	NFE(%)	OM(%)	NB(%)
	Control	94.22 ^b	74.06 ^b	26.04 ^b	81.10 ^{abc}	79.38 ^{ab}
2% - Y ₁	97.85 ^a	75.58 ^{ab}	26.85 ^a	79.61 ^c	75.38 ^C	78.68 ^b
4% - Y ₁	95.91 ^{ab}	76.19 ^{ab}	24.72 ^d	80.47 ^{bc}	77.02 ^{abc}	79.19 ^{ab}
6% - Y ₁	96.63 ^a	79.13 ^a	25.38 ^c	81.02 ^{abc}	76.66 ^{bc}	80.43 ^a
8% - Y ₁	96.94 ^a	79.72 ^a	26.00 ^b	81.57 ^{abc}	76.45 ^{bc}	80.39 ^a
2% - Y ₂	96.18 ^{ab}	78.55 ^{ab}	26.22 ^{ab}	81.52 ^{abc}	77.99 ^{abc}	80.13 ^{ab}
4% - Y ₂	96.20 ^{ab}	77.61 ^{ab}	26.82 ^a	80.93 ^{abc}	77.54 ^{abc}	79.77 ^{ab}
6% - Y ₂	96.25 ^a	77.19 ^{ab}	25.38 ^{ab}	83.85 ^a	76.22 ^{bc}	79.45 ^{ab}
8% - Y ₂	96.74 ^a	78.15 ^{ab}	26.05 ^b	83.00 ^{ab}	80.26 ^a	79.34 ^{ab}

a, b, c, d . . . Mean each column bearing the same superscripts don't differ significantly (P≤0.05) .

CP: Crude Protein. EE: Ether Extract. CF: Crude Fiber.
 NFE: Nitrogen Free Extract
 OM: Organic Matter. NB: Nitrogen Balance.

Economic efficiency:

Effect of the experimental diets on economic efficiency is shown in Table (6).

Table (6) : Effect of experimental diets on economic efficiency .

(1) L.E = 1 pound Egyptian Currency = 100 piasters.

Items	Control	Experimental treatments				Percentage of MF substitution			
		Yeast (1)				Yeast (2)			
	0 %	2 %	4 %	6 %	8 %	2 %	4 %	6 %	8 %
Price / kg feed (L.E.) ⁽¹⁾	1.45	1.52	1.49	1.47	1.45	1.53	1.52	1.52	1.51
Total Feed Intake Kg/hen	11.36	11.99	11.42	11.47	11.44	11.22	11.15	11.20	11.41
Total Feed cost/hen (L.E)	16.47	18.25	17.02	16.86	16.59	17.17	16.95	17.02	17.23
Total number of eggs / hen	92	106	98	100	98	100	95	103	86
Price of total egg production /hen (L.E).	26.84	30.92	28.59	29.17	28.59	29.17	27.16	30.05	25.09
Net Revenue / hen (L.E) ⁽²⁾	10.37	12.67	11.57	12.31	12.2	12.00	10.21	13.03	7.86
Economic Efficiency (E :E) ⁽³⁾	0.63	0.85	0.68	0.73	0.74	0.70	0.60	0.77	0.46
Relative Economic Efficiency ⁽⁴⁾	100	135	108	116	148	118	95	122	73

(2) Net Revenue / hen (L.E) = Price of total egg production /hen (L.E) – total feed cost /hen (L.E).

(3) Economic Efficiency (E.E) = Net revenue / price of total feed intake.

(4) Relative Economic Efficiency = assuming that the relative economic efficiency (E.E) of the control = 100.

Data showed that addition of yeast₁ or yeast₂ except 4% and 8% yeast₂ improved the relative economic efficiency as compared to the control group. This improvement could be due to reducing the amount of the feed required to produce egg or improving the feed conversion values. These results coincided with those reported by *Abd-Elwahed et al.* (2003) who reported that the economic efficiency values of the layers groups fed 0, 2, 4 and 6% dry yeast substituted for soya bean meal were 0.03, 0.38, 0.351 and 0.265, respectively.

Conclusively, addition of 2% inactive yeast or 6% active yeast to laying hen diets can be recommended for the performance, egg quality and economic efficiency.

REFERENCES

- Abd El-Azeem, F.(2002). Digestion, neomycin and yeast supplementation in broiler diets under Egyptian summer conditions. *Egypt. Poult. Sci.*, 22: 235-257.
- Abdel Wahed, H.M.; E.A. El-Full; A.M.R. Osman; and N.A. Hataba (2003). Effect of replacing soybean meal with graded levels of dried yeast on growth of Dandarawi and Golden Montaza Chicks. *Egypt. Poult. Sci.*, 23: 507-522.
- Abdulrahim, S. M; M. S. Y. Haddadin; E.A.R. Hashlamoun and R.K. Robinson (1996). The influence of lactobacillus acidophilus and bacitracin on layer performance of chickens and cholesterol content of plasma and egg yolk. *Bri. Poult. Sci.*, 37:341-346.
- Abo-Raya, A.k; and A.G.H. Galal (1971). Evaluation of poultry feeds in digestion traits with reference to some factors involved. *Egypt. J. Anim. Prod.* 11:207
- Abu-Taleb, A.M; Sh.F. El Afifi and M.M. Wakwak (2005). Effect of supplementing active dried yeast into laying Japanese quail on performance hatching traits and immune response. *Egyptian, J. Nutrition and Feed*, 9:33-42.
- Association of Official Analytical Chemists A.O.A.C. (1990). *Official Method of Analysis 14th ed.* Published by the A.O.A.C. Washington, D.C., USA.
- Balev, T; U.S. Ucan; B. Coskun; V. Kurtoglu; and I.S. Cetnglu(2001). Effect of dietary probiotic on performance and humoral immune response in layer hens. *Bri. Poult. Sci.*, 42:456-461.
- Bardly, G. L; T.F. Savage; and T.I. Karen(1994). Effect of supplementing diets with *Saccharomyces cerevisiae*, var. *boulardii* on male poult performance and ileal morphology. *Poult. Sci.*, 73:1766-1770.
- Chin, H.J; T.F. Shen; H.P. SU; and S.T. Ding(2006). *Schizochytrium limacinum* SR-21 as a source of docosahexaenoic acid: optimal growth and use as a dietary supplement for laying hens. *Australian. Journal. Of Agricultural. Research.*, 57:13-20.
- Duncan, D.B. (1955). Multiple range and multiple F –test, *Biometrics*, 11: 1-42.
- Fuller, R.(1977). The importance of lactobacilli in maintaining normal microbial balance in the crop. *Bri. Poult. Sci.*, 18:85-94.
- Funk, E.M. (1948). The relationship of the yolk index determined in natural position to the yolk as determined after separating the yolk from the albumen. *Poult. Sci.*, 27:367-378.
- Guillot. J.F. (2000). The pros and cons of probiotics. Make probiotics work for poultry. *World's Poult. Sci.*, 16:18-21.

- Jakobson, P.E.; S.G. Kirston and H. Nelson(1960). Digestibility trials with poultry. 322 bertning fraforsgs laboratoriet, udgivet of stants. Husdyrbugsud Valy-Kaben havn. Poult. Sci.73:1260-1266
- Jin, L. Z; Y.W. Ho; N. Abdullah and S. Jalaludin(1997). Probiotics in poultry : modes of action. World's Poult. Sci. J. 53: 351-368.
- Kornegay, E. T; D.D. Rhein-Welker; M.D. Lindermann and C.M. Wood(1995). Performance and nutrient digestability in weanling pigs as influenced by yeast culture additions to starter diets containing dry whey or one of two fiber sources. J. Anim. Sci., 73: 1381-1389.
- Koudela, K and C.C.S. Nyirenda (1995). Influences of probiotics lactiferm on methaemoglobin biosynthesis in Gallus donesticus. Scientia-Agriculturae – Bohemica. 26:209 – 218.
- Maia, G.A.R; J.B. Fonseca; R.T.R.N. Soares; and G.L.M. Souza(2001). Performance of commercial brown egg layers fed dried yeast of sugar cane. Revista-Brasileira-de-Sciencia-Avicola 3:163-171.
- Miles, R. D; and S.M. Bootwella(1991). Direct-fed microbials in animal production in direct-fed microbial in animal production. A review, National Food Ingredient Association, West Des Moines, Iowa, USA., PP:117-132.
- Mohan, B.R; B.M Kadirvel; and A.O.N. Natoraj(1995). Effect of probiotic supplementation on serum/yolk cholesterol and on egg shell thickness in layers. Bri. Poult. Sci., 36:799-803.
- Mohdavi, A-H; H.R. Rahmani; Y. Pourreza; and M.A. Edriss(2005). Effect of probiotic inclusion in different levels of barley substitution for corn diets on egg quality and laying hen's. Pakistan. Journal. Of - Bio. Sci. 8:1521-1528.
- Nir, I and N. Senkoylu(2000). Kanatlılar için sindiri mi destekleyen yem Katkı Maddeleri (Tekidag). Food Additives and contantnats Vol 21:817-823.
- NRC, (1994). Nutrient requirements of poultry. National Research Council, 9th revised ed. Nat. Acad. Press, Washington, D.C.
- Nursoy, H; O. Kaplan; M. N. Oguz; and O. Yilmaz(2004). Effect of varying levels of live yeast culture on yield and some parameters in laying hen diets. Indian – Veterinary – Journal., 81: 59-62.
- Osman, A.M.R; H.M.A. El-Wahed; E.A. El-Full; and N.A. Hataba(2003). Effect of replacing soybean meal with graded levels of dried yeast on Dandarawi and Golden Montazah layer performance. Egypt. Poult. Sci., 23:469-484.
- Osman, M (2003). The influence of probiotic inclusion on the productive performance of commercial layers. Egypt. Poult. Sci., 23: 283-297.
- Park, D.Y; H. Namkung; and I.K. Paik(2001). Effect of supplementary yeast culture on the performance of laying hens. J. Animal Sci. & Technology 43: 639-646.
- Park, J.H; G.H. Park and K.S. Ryu(2002). Effect of feeding organic acid mixture and yeast culture on performance and egg quality of laying hens. Korea. J. Poult. Sci., 29:109-115.
- Piva, A.M.E; P.P. Gatta; G. Biagi, Castellani, A.L. Gimordenti; J.B. Luchansky; S. Silva, and A. Moddenti(2003). The effect of dietary supplementation with trivalent chromium on production performance of laying hens and the yolk. Animal – Feed – Science – and – Technology. 106:149–163.
- Romanoff, A.L. and A.L. Romanoff (1949). The avian egg. Johan Wiley Ssons, Inc., New York.
- Santin, E; A.C. Paulillo; A. Maiorka; A.V.F. Silva and A.C. Aless(2003). Evaluation of the efficiency of Saccharomyces cerevsiae cell wall to ameliorate the toxic effects of aflatoxin in broilers. International J. Poult. Sci., 2:341-344.

- SAS Institute (1996). SAS / STAT user's guide: statistics, 8 edu. SAS Institute Inc., Cary NC.
- Scott, M.L.; M.C. Nesheim and R.J. Young, (1982). Nutrition of the Chicken. Published by M.L. Scott & Associates, Ithaca, New York.
- Siam, S.S; S.A. Riad; F.R. Mohamed and A.K.A. Eldein(2004). Influence of using two different levels of probiotics on egg performance, blood and yolk cholesterol of laying hens. Egypt. Poult. Sci., 24:267-269.
- Soliman, A.Z.M. (2002). Bacitracin and active yeast supplementation in layer diets varying in energy content. Egypt. Poult. Sci., 23:37-51.
- Stanley; V.G.R.O; S. Woldesenet; D.H. Hutchinson and L.F. Kubena(1993). The use of saccharomyces cerevisia to suppress the effects of aflatoxicosis in broiler chicks. Poult. Sci., 27:1867-1872.
- Tortuero, F and E. Fernandez(1995). Effects of inclusion of microbial cultures in barley –based diets fed to laying hens. Animal – Feed – Science – and – Technology., 53:255-265.
- Wakwak, M.M; Sh.F. El-Affifi; K. Nagla; Soliman and M. Attia(2003). Effect of adding active dried yeast into Japanese quail diets on performance, some immunity and microbiological aspects. J. Agric. Sci. Mansoura Univ.,28:2621-2632.
- Yildirm, I. and S.S. Parlat(2003). Effects of dietary addition of live yeast on hatching traits, testis and Ovary Weights of Japanese quail in aflatoxicosis. Archiv. Fur Gefluerkund. 67:208-211.

استخدام الخمائر في علائق الدجاج البياض.

ممدوح عمر عبد السميع، عادل صلاح الدين شلبي ، علي متولي علي.
قسم الإنتاج الحيواني – كلية الزراعة – جامعة القاهرة – الجيزة – مصر.

أجريت هذه الدراسة لتقييم تأثير إضافة الخميرة الجافة النشطة أو غير النشطة إلى علائق الدجاج البياض على الأداء الإنتاجي وصفات جودة البيضة والكفاءة الاقتصادية. ولهذا الغرض تم تقسيم ٩٠٠ دجاجة بياضة من سلالة هاي لاين الأبيض عمر ٢٤ أسبوعاً إلى ٩ مجاميع بكل مجموعة ١٠٠ دجاجة موزعة على ٤ مكررات بكل منها ٢٥ دجاجة. وتم تغذية هذه الطيور لمدة ١٦ أسبوعاً على عليقة أساسية ممددة بخمسة مستويات متدرجة من الخميرة الجافة النشطة أو غير النشطة (صفر للكنترول ، ٢% ، ٤% ، ٦% ، ٨%) وكانت جميع العلائق متماثلة في البروتين الخام والطاقة الممتلئة. ولقد أوضحت النتائج المتحصل عليها ما يلي: -

- 1) إضافة الخميرة سواء نشطة أو غير نشطة إلى علائق الدجاج البياض عند معظم المستويات المختبرة حسن كلاً من النسبة المئوية لإنتاج البيض ووزن البيضة مقارنة بمجموعة الكنترول.
- 2) إضافة الخميرة إلى علائق الدجاج البياض عند معظم المستويات المختبرة قلل معنوياً متوسط الغذاء المأكل وحسن معنوياً الكفاءة التحويلية للغذاء مقارنة بمجموعة الكنترول.
- 3) لم تتأثر معنوياً معظم صفات جودة البيضة خاصة سمك القشرة ، دليل الصفار ، دليل الشكل بإمداد علائق الدجاج البياض بالخميرة النشطة أو غير النشطة عند معظم المستويات المختبرة مقارنة بمجموعة الكنترول.
- 4) إمداد علائق الدجاج البياض بالخميرة غير النشطة أو النشطة عند كل مستوى مختبر حسن من معاملات هضم البروتين الخام ومستخلص الإثير وميزان الأزوت (كنسبة مئوية) مقارنة بمجموعة الكنترول. كما استنتج يمكن أن يوصى باستخدام ٢% خميرة غير نشطة أو ٦% خميرة نشطة في علائق الدجاج البياض من أجل الحصول على أفضل أداء إنتاجي وتحسين صفات جودة البيضة وزيادة العائد الاقتصادي.

2434 2435 2436 3437 2438 2439 2440 2441 2442 2443

2434 2435 2436 3437 2438 2439 2440 2441 2442 2443