

GENETIC BEHAVIOR FOR SOME IMPORTANT ECONOMICAL TRAITS AND THE NATURE OF RESISTANCE FOR POWDERY MILDEW IN SQUASH.

Abd El-Hadi, A. H.*; G.A.R. El-Sherbeny** and A.A.El-Magrabi***

* Dept. of Genetics Fac. of Agric. Mansoura Univ. Egypt

** Dept. of Agric. Botany (Genetics), Sohag Fac. of Agric., South Valley Univ. Sohag, Egypt

*** Plant Dis. Inst. Agric. Res. Center. Egypt.

ABSTRACT

In this investigation six parental varieties of squash obtained from different sources were used. These varieties were : Eskandrani (P_1), Giado (P_2), Zucchini mezza lung bianco (P_3), Zucchini 544-005 (P_4), White bush scallop (P_5) and Zucchini nano verda di Milano (P_6). These parental varieties were crossed among them to obtain 15 F_1 hybrids through diallel crosses mating design excluding reciprocals. All genotypes which included the six parental varieties and their 15 F_1 hybrids were evaluated. Different yield and yield component traits as well as the nature of resistance for powdery mildew were studied.

The results indicated the presence of highly significance differences among genotypes. This finding was expected, where the parental varieties were obtained from different sources. The results also revealed that no parental variety exceeded the other parents for mean performances for all studied traits. The parent P_3 was the earliest and gave the highest yield. In the same time, the P_5 and P_6 were the highest for some traits.

The F_1 hybrid $P_3 \times P_5$ was the earliest hybrid, while the F_1 hybrid $P_4 \times P_6$ was the latest. $P_5 \times P_6$ was the lowest hybrid for F.Y./P. and No. F./P. The highest F_1 hybrid was $P_2 \times P_4$ for No. F./P. $P_5 \times P_6$ showed the high level of resistance for powdery mildew.

Heterosis values were observed for most hybrids. The best heterosis ($H_{M,P}\%$) values were: -19.72($P_3 \times P_5$); -26.65($P_1 \times P_2$); -9.55($P_1 \times P_6$); 111.43($P_4 \times P_5$); 20.62($P_4 \times P_6$); 76.05($P_1 \times P_5$); 20.63($P_4 \times P_5$); 34.28($P_5 \times P_6$) and 800.3($P_1 \times P_6$) for D.^{1st} F.F., No.1st F.F.N., 1st P.D., F.Y./P kg., W.F./P.g., No.F./P., F.L.cm., F.D.cm. and D.S., estimated from the mid-parents. respectively.

The results also cleared that both additive and non-additive genetic variances including dominance could not be neglected in the inheritance of all studied traits of squash.

INTRODUCTION

In squash and other cucurbits, heterosis was investigated aiming to increase the productivity. In addition, the improvement of fruits quality is necessary. In this respect, El-Gazar (1981) and Abd El-Maksoud (1986) cleared that heterosis values in squash were significant and/or highly significant for fruits length, total number of fruits and total weight of fruits. In this respect, Dogra *et al.*, (1997) evaluated heterosis in cucumber. They mentioned that the highest values of heterosis was noticed in the cross K 75 X Gynt (57-35%). Similarly, Abd El-Hadi *et al.*, (2001), Sadek (2003), Abd El-Hadi and El-Gendi (2004) and Abd El-Hadi *et al.*, (2005a,b) evaluated heterosis in different F_1 hybrids of squash. They revealed that all studied F_1 hybrids exhibited significant or highly significant values of heterosis estimated

from the mid-parents or against the better parent for most studied traits. They also added that the highest values of heterosis were 135.4 and 194.6% for fruits yield per plot estimated versus the M.P. and the B.p, respectively.

Concerning the nature of gene action, Farid (1990), El-Adl *et al.*, (1996), Abd El-Hadi *et al.*, (2004) and Abd El-Hadi and El-Gendi (2004) evaluated many F₁ hybrids of squash and other cucurbits. They cleared that both general and specific combining abilities were important indicating the importance of additive and non-additive genetic variances including dominance in the inheritance of all studied traits. On the other hand, Abd El-Hadi *et al.*, (2001) and Sadek (2003) illustrated that the non-additive genetic variances including dominance were the most important source of genetic variance for all studied traits. In another study by Abd El-Hadi *et al.*, (2005a). They cleared that additive gene action was more important in the inheritance of vegetative and earliness traits of squash.

The nature of resistance of powdery mildew in squash was evaluated by many authors. Arora *et al.*, (1992) claimed that the resistance of powdery mildew in squash was controlled by additive and dominant genes, although the additive component being predominant. In the same time, Michelle *et al.*, (1999) cleared that the varieties PX47592 and PX50592 showed high resistance for powdery mildew. In this respect, James and Stevenson (2001) indicated that the highest resistance for powdery mildew was observed in the yellow crookneck squash cultivar sunglo (PMT) and the Zucchini line HMX 0710.

MATERIALS AND METHODS

In this investigation, 15 F₁ hybrids of squash were obtained among six varieties through diallel crosses mating design excluding reciprocals. These varieties were : Eskandrani (P₁), Giado (P₂), Zucchini mezza lung bianco (P₃), Zucchini 544-005 (P₄), White bush scallop (P₅) and Zucchini nano verda di Milano (P₆). The seeds of varieties were obtained from different countries i.e., F₁ from Egypt, P₂, P₄ and P₆ from Italy, P₃ from Germany and P₅ from United States of America (U.S.A.). All these parental varieties represented a wide range of variability for all studied traits. In the summer growing season of 2003, the parental varieties were crossed to produce 15 F₁ hybrids. In addition, the six parental varieties were selfed to obtain enough seeds. All 21 genotypes (six parental varieties and their 15 F₁ hybrids) were evaluated in the growing season of 2004 in a field trial experiment. This investigation was conducted at the experimental station, Faculty of Agriculture, Mansoura University. The experimental design was a randomized complete blocks design with three replications. Each block was 21 plots. Plot was one ridge 5.0 m. long and 1.0 m wide. The distance between hills was 0.5m apart. Data were recorded on the following traits.

- Date of the first female flower (D.1st F.F),
- First picking date (1st P. D.),
- Number of the first female flowering node (No. 1st F.F.N),
- Fruits yield per plant (F.Y. /P. Kg),

- Weight of fruits per plant (W.F./P.g),
- Number of fruits per plant (No. F./P),
- Fruit length in centimeters (F. L. cm),
- Fruit diameter in centimeters (F. D. cm) and
- Disease severity (D. S.).

All agricultural practices were carried out as recommended for squash. Different analyses of variance were made to estimate different genetic parameters. The form of the analysis of variances and the expectations of the mean squares were made as outlined by Steel and Torrie (1960). The amounts of heterosis were calculated as the deviation of F_1 hybrids from the mid-parents ($H_{M.P.}\%$) and the better parent ($H_{B.P.}\%$). The analysis of variances of diallel crosses mating design were made according to Geriffing (1956) method-2 model-1. Genetic parameters were obtained according to Singh and Chaudhary (1985).

Under natural condition, disease severity percentage (powdery mildew) was recorded by counting disease plants in 10 plant from each plot. The number of spots (disease severity) (D.S.) appearing on the leaves of diseased plants were counted. This trait was estimated according to Sherwood and Hagedorn (1958) as follows:

$D.S.1\% = (\text{Disease class})(\text{No. of diseased plants in that class}) \times 100 / \text{Total No. of plants} \times 3\%$

0 = No. symptoms,

1 = 1-10 lesion/plant,

2 = 11-20 lesion/plant,

3 = 21-30 lesion/plant and

4 = More than 30 lesions/plant.

RESULTS AND DISCUSSION

Squash, (*Cucurbita pepo*, L) consider as one of the important vegetable crops in Egypt. Thus, the improvement of quality and increasing the productivity were very important. In this respect, this investigation was conducted as an attempt to obtain new squash hybrids. In addition, the nature of resistance for powdery mildew disease in squash was also studied. In this investigation, six varieties of squash were crossed among them to obtain 15 F_1 hybrids through diallel crosses mating design excluding reciprocals. Then, the 21 genotypes which included the six parental varieties and their 15 F_1 hybrids were evaluated to study the genetic behavior of some important economical traits and disease tolerance specially for powdery mildew.

The analyses of variance were made for all studied traits. The obtained results of the analysis of variances and the mean squares are presented in Table 1.

Table 1: The results of the analysis of variances for all studied traits.

S.V.	d.f.	D.1 st F.F	No.1 st F.F.N.	1 st P.D.	F.Y./P.kg	W.F./P.g	No. F./P.	F.L. cm	F.D. cm	D.S.
Replications	2	4.854	0.067	7.323	0.019	31.39	4.18	0.034	0.011	13.349
Genotypes	20	50.05**	0.993**	52.35**	0.659**	229.9**	36.61**	29.55**	4.139**	775.91**
Error	40	3.842	0.071	3.624	0.004	18.98	1.794	0.036	0.005	2.299

*,** significant differences at 0.05 and 0.01 levels of probability, respectively.

The results revealed that the mean squares of genotypes showed highly significance for all studied traits. This finding indicated the presence of genetic variability among all the evaluated genotypes. This variability mainly due to the different sources of the parental varieties and small values of error mean squares. On the other hand, the mean squares of replications were insignificant (desirable) for all studied traits. The genetic variability would be estimated through the analysis of variance of diallel crosses mating design.

The means of the parental varieties and their 15 F₁ hybrids were calculated for all studied traits and the results are presented in Table 2.

The results illustrated that no parental variety showed high performance means for all studied traits. It could be also noticed that the parental variety P₃ was the earliest for date of the first female flower (D1st F. F) (43.7days) and first picking date (1st P.D.) (46.13 days) and gave the highest fruit yield per plant (F.Y./P) (1.96 kg) and number of fruits per plant (No. F./P) (16.9). In the same time, the variety P₁ was the best for of fruit per plant (W.F./P) (118.2g) and fruit length (F. L.) (13.8 cm.), while the parent P₅ showed the highest value for weight of fruit per plant (W.F./Pg) (123.8) and fruit diameter (F.D.) (5.62 cm.). Concerning disease resistance, one trait was studied: disease severity which indicate the percentage of infected plants. The parental variety (P₆) showed the highly resistance for powdery mildew with the mean of (O: no symptoms) (resistance) for this trait. On the other hand, the variety P₅ showed highest values with the mean of 50 for disease severity.

The results also revealed that there were no F₁ hybrid exceeded all the F₁ hybrids for all studied traits. It could be also noticed that the earliest F₁ hybrid was P₃xP₅, where showed the lowest number of days to give the first female flower (D₁st F.F.), (desirable), first picking date (1st P. D.) and short distance to show the first female flowering node with the mean values of 40.5 days, 42.63 days and 3.50, respectively. On the other hand, the F₁ hybrid P₄xP₆ was the latest hybrid with the mean values of 50.2, 51.7 days and 5.45 for the same traits, respectively. The results also cleared that the F₁ hybrid P₅xP₆ was the lowest hybrid for F. Y./P.kg and No. F./P.kg with the means of 2.09 kg and 16.10, respectively. While the highest F₁ hybrid for these traits was P₂xP₄ with the means of 2.86 kg and 23.9, respectively. In this respect, the F₁ hybrid P₄xP₅ gave the heaviest fruit per plant (W.F./P.g) with the mean of 140.4g, while the F₁ hybrid P₂xP₄ showed the lowest weight of fruits/plant with the mean of 114.1 g. Concerning disease resistance, the results indicated that the F₁ hybrid P₅xP₆ showed high resistance for powdery mildew, where showed (O) disease severity.

Table 2: The mean performances of the parental varieties and their 15 F₁ hybrids for all studied traits.

Genotypes	D.1 st F.F	No.1 st F.F.N	1 st P.D.	F.Y./P.kg	W.F./P.g	No.F./P.	F.L.cm	F.D.cm	D.S.
P ₁	50.73	5.35	52.96	1.65	118.2	13.53	13.8	3.17	9.33
P ₂	52.60	4.33	55.53	1.59	108.9	15.06	12.96	2.93	19.33
P ₃	43.70	3.95	46.13	1.96	112.1	16.90	13.8	3.10	10.33
P ₄	45.40	3.78	47.53	1.54	109.3	14.40	13.06	3.11	29.33
P ₅	57.20	4.12	60.53	0.91	123.8	7.10	2.74	5.62	50
P ₆	44.70	3.81	46.76	1.54	108.9	13.9	12.33	2.84	0.0
P ₁ xP ₂	41.96	3.55	44.8	2.25	118.4	18.8	13.76	3.09	46.33
xP ₃	44.88	4.07	47.30	2.26	118.2	18.9	14.53	3.16	38.0
xP ₄	42.43	3.52	45.13	2.32	114.8	19.16	13.46	3.00	60.0
xP ₅	46.73	4.45	49.03	2.49	134.1	18.16	8.23	5.73	30.0
xP ₆	42.63	3.78	45.10	2.34	114.2	19.5	13.73	2.98	42.0
P ₂ xP ₃	43.20	4.11	45.60	2.50	115.8	21.3	14.23	3.12	40.66
xP ₄	41.93	3.40	44.66	2.86	114.1	23.9	13.73	3.20	17.33
xP ₅	44.53	4.061	46.86	2.59	125.8	18.83	8.33	5.24	20.0
xP ₆	46.70	4.85	48.96	2.24	115.1	19.06	14.2	3.13	20.0
P ₃ xP ₄	42.30	4.27	44.90	2.43	120.2	20.26	14.2	3.06	50.33
xP ₅	40.50	3.50	42.63	2.56	127.9	19.70	8.86	5.80	20.33
xP ₆	46.23	4.50	49.03	2.28	124.3	19.20	14.66	3.27	22.0
P ₄ xP ₅	43.53	3.91	46.03	2.59	140.4	19.40	9.53	5.65	16.33
xP ₆	50.20	5.45	51.70	2.15	131.6	17.50	14.93	4.34	20.0
P ₅ xP ₆	46.73	4.73	48.96	2.09	129.3	16.10	8.66	5.68	0.0
L.S.D.o.05	3.23	0.439	3.14	0.104	7.19	2.21	0.312	0.116	2.502
L.S.D.o.01	4.32	0.587	4.20	0.139	9.60	2.95	0.418	0.155	3.347

*,** significant differences at 0.05 and 0.01 levels of probability , respectively .

On the other hand, the F_1 hybrid $P_3 \times P_4$ showed highly sensitivity for powdery mildew disease with the means of (50.33) for the same trait. These results indicated that the parental variety P_6 showed high resistance or played an important role for the appearance of resistance for powdery mildew. This parent could be use in breeding programs to obtain resistance genotypes.

The deviation of F_1 hybrids against their parents as heterosis values were estimated from the mid-parents ($H_{M,P}\%$) and the better parent ($H_{B,P}\%$) for all studied traits and the results are presented in Tables 3 and 4, respectively.

The results cleared that all F_1 hybrids significantly exceeded the mid of their parents for all studied traits with few exceptions. Consequently, significant heterosis values versus the mid-parents ($H_{M,P}\%$) were observed for all studied traits with few exceptions. The best recorded heterosis values (desirable) were -19.72% ($P_3 \times P_5$), -26.65% ($P_1 \times P_2$), -20.06% ($P_3 \times P_5$), 111.43% ($P_4 \times P_5$); 20.62% ($P_4 \times P_6$); 80.47% ($P_4 \times P_5$); 20.63% ($P_4 \times P_5$); 34.28% ($P_5 \times P_6$) and -325.94% ($P_3 \times P_6$) for D. 1^{st} F. F.; No. 1^{st} F.F. N; 1^{st} P. D.; F. Y./P. kg; W.F. g.; No. F./P.; F.L. cm; F.D. cm and D.S., respectively. It could be also noticed that the hybrids which contained the parents P_4 and/or P_5 and P_6 showed desirable values of heterosis versus the mid-parents. This finding cleared the importance of general combining ability effect (g_i) for those parental varieties. Similar results were obtained by many authors among them: El-Gazar (1981), Abd-El-Maksoud (1986), Dogra *et al.*, (1997), Abd El-Hadi and El-Gendy (2004) and Abd El-Hadi *et al* (2005 A and B).

The results in Table 4 indicated that most F_1 hybrids showed heterosis values against the better parent ($H_{B,P}\%$) for most studied traits. Concerning the earliness traits most F_1 hybrids exhibited negative (desirable) and significant heterosis values for that traits. In the same time, most F_1 hybrids exhibited significant and highly significant heterosis values for yield traits. The values of heterosis for yield traits ranged from 16.33% ($P_3 \times P_6$) to 79.87% ($P_2 \times P_4$); 0 ($P_1 \times P_3$) to 20.40% ($P_4 \times P_6$) and 11.83 ($P_1 \times P_3$) to 58.70% ($P_2 \times P_4$) for F. Y./P. kg, W. F. g. and No. F./P., respectively. It could be also regarded that the F_1 hybrids $P_1 \times P_5$, $P_2 \times P_6$, $P_3 \times P_6$, $P_4 \times P_6$ and $P_5 \times P_6$ showed high levels of resistance for powdery mildew.

The analysis of variances of diallel crosses mating design were made and the results are shown in Table 5. The results indicated that the mean squares of general combining ability (GCA) and specific combining ability (SCA) showed highly significance values for all studied traits. This finding indicated that both GCA and SCA play an important role in the expression of these traits.

These results were in agreement with those of Farid (1990), El Adl *et al.* (1996), and Abd El-Hadi and El-Gendi (2004).

The results also illustrated that the magnitudes of general combining ability (GCA) mean squares were larger than those of the mean squares for specific combining ability (SCA) for most yield traits (W. F./P g), (No. F./P.), (F. L. cm) and (F. D. cm). The results also showed that the earliness traits: D. 1^{st} F. F., N. 1^{st} F. F. N and 1^{st} P. D. and disease severity were non additively controlled.

Table3: The amounts of heterosis versus the mid-parents ($H_{M,P}\%$) for all studied traits.

Crosses	D.1 st F.F	No.1 st F.F.N.	1 st P.D.	F.Y./P.kg	W.F./P.g	No.F./P.	F.L.cm	F.D.cm	D.S.
P ₁ xP ₂	-18.78**	-26.65**	-17.71**	38.89**	4.27**	31.51**	2.84**	1.31*	223.31**
XP ₃	-4.95**	-12.47**	-4.53**	25.21**	2.65*	24.22**	5.29**	0.80	286.57**
XP ₄	-11.72**	-22.89**	-10.18**	45.45**	0.92	37.20**	0.22	-4.46**	210.40**
XP ₅	-13.41**	-6.02**	-13.60**	94.53**	10.83**	76.05**	-0.48**	30.38**	1.13
XP ₆	-10.66**	-17.47**	-9.55**	46.71**	0.57	42.91**	5.09**	-0.83	800.30**
P ₂ x P ₃	-10.28**	-0.72	-10.29**	40.85**	4.80**	33.29**	6.35**	3.48**	174.17**
XP ₄	-14.43**	-16.15**	-13.33**	82.75**	4.58**	62.25**	5.53**	5.96**	-28.77**
XP ₅	-18.89**	9.11**	-19.25**	107.20**	8.12**	69.95**	6.11**	22.57**	-42.30**
XP ₆	-4.01	19.16**	-4.27**	43.13**	5.69**	31.63**	12.30**	8.49**	106.93**
P ₃ x P ₄	-5.05**	10.48**	-4.12**	38.86**	8.58**	29.46**	5.73**	-1.45**	153.81**
XP ₅	-19.72**	-13.26**	-20.06**	78.40**	8.44**	64.17**	7.13**	33.03**	-32.60**
XP ₆	4.54**	15.98**	5.57**	30.29**	12.49**	24.68**	12.21**	10.10**	-325.94**
P ₄ x P ₅	-15.15**	-1.01	-14.81**	111.43**	20.46**	80.47**	20.63**	29.44**	-58.83**
XP ₆	11.43**	37.97**	9.66**	39.61**	20.62**	23.67**	17.61**	12.27**	36.38**
P ₅ x P ₆	-8.28**	19.29**	-8.73**	70.61**	11.13**	53.33**	14.93**	34.28**	-100.00**
L.S.D _{0.05}	1.104	0.150	1.072	0.035	2.455	0.754	0.106	0.039	0.854
L.S.D _{0.01}	1.475	0.200	1.433	0.047	3.279	1.00	0.142	0.053	1.141

**, * significant differences at 0.05 and 0.01 levels of probability, respectively.

Table 4: The amounts of heterosis versus the better parent ($H_{B.P.}\%$) for all studied traits.

Crosses	D.1 st F.F	No.1 st F.F.N.	1 st P.D.	F.Y./P.kg	W.F./P.g	No.F./P.	F.L.cm	F.D.cm	D.S.
P ₁ xP ₂	-17.29**	-18.01**	-15.41**	36.36**	0.17	24.83**	-1.43**	-2.52	396.57**
XP ₃	2.7	11.51**	2.54	15.31**	0	11.83**	5.29**	-0.32	307.29**
XP ₄	-6.54**	-6.88**	-5.05**	40.61**	-2.88	33.06**	-2.46	-5.36**	543.09**
XP ₅	-7.88**	8.01	-7.42**	50.91**	8.32**	34.22**	-40.36	1.96**	221.54
XP ₆	-4.63**	-0.79**	-3.55**	41.82**	-3.38	41.01**	-0.51**	-5.99	0
P ₂ x P ₃	-1.14**	4.05	-1.15**	27.55**	3.30*	26.04**	3.12**	0.65*	293.61**
XP ₄	-7.64**	-10.05**	-6.04**	79.87**	4.39	58.70**	5.13**	2.89**	-10.35**
XP ₅	-15.34**	11.89*	-15.61**	62.89**	1.62**	25.03**	-35.73**	-6.76**	3.47**
XP ₆	4.47	27.30**	4.70	40.88**	5.69*	26.56**	9.57**	6.83**	0
P ₃ x P ₄	-3.20	12.96*	-2.67	23.98**	7.23**	19.88**	2.90**	-1.61**	387.22**
XP ₅	-7.32**	-7.41**	-7.59**	30.61**	3.31**	16.57**	-35.80**	3.20**	96.81**
XP ₆	5.79	18.11**	6.29*	16.33**	10.88**	13.61**	6.23**	5.48**	0
P ₄ x P ₅	-4.12**	3.44	-3.16**	68.18**	13.41**	34.72**	-27.03**	0.53**	-44.32**
XP ₆	12.30**	44.18**	10.56**	39.61**	20.40**	21.53**	14.32**	7.40**	0
P ₅ x P ₅	4.54**	24.15**	4.70**	35.71**	4.44**	15.83**	-29.7**	1.07**	0
L.S.D _{0.05}	2.362	0.321	2.294	0.076	5.250	1.614	0.228	0.085	1.827
L.S.D _{0.01}	3.155	0.428	3.064	0.101	7.01	2.156	0.305	0.113	2.441

**, * significant differences at 0.05 and 0.01 levels of probability , respectively .

Table 5: The analysis of combining ability and the mean squares for all studied traits.

S. V.	D. f.	D.1 st F.F	No.1 st F.F.N.	1 st P.D.	F.Y./P.kg	W.F./P.g	No.F./P.	F.L.cm	F.D.cm	D.S.
GCA	5	45.22**	0.414**	49.31**	0.216**	409.2**	38.28**	114.02**	14.31**	672.277**
SCA	15	51.67**	1.187**	53.36**	0.806**	170.07**	36.06**	1.389**	0.750**	810.45**
SCA/ GCA	-	0.875	0.348	0.924	0.267	2.406	1.061	82.08	19.07	2.299

**, * significant differences at 0.05 and 0.01 levels of probability , respectively

Moreover, on the calculated ratio of GCA/SCA variances, the results showed that it more than the unity in W.F./Pg, No. F./P, F.L.cm, F.D.cm and D.S. This result cleared that the GCA variance or the additive genetic variance was more important than of the SCA or non-additive genetic variance including dominance in the inheritance of these traits. The reverse was true in the cases of D.1st F.F., No. 1st F.F.N, 1st P.D. and F.Y./P. kg. Which GCA/SCA ratios were less than unity. This result indicated that SCA variance was more important in the inheritance of these traits. These results were in common agreement with similar results obtained by Abd El-Hadi *et al.* (2001) and Sadek (2003).

These findings revealed that additive genetic variances played the important role in the inheritance of studied yield traits, while non-additive genetic variances including dominance controlled the inheritances of the earliness and disease severity trait. It could be also mentioned that the obtained values of heterosis for that traits which were obtained and described earlier mainly due to dominance effect (earliness and disease severity). Whereas, the values of heterosis which were obtained for studied yield traits mainly due to epistatic effects (add. x add.), (add. x dom) and (dom. x dom). Finally, the additive and non-additive gene action were more important in the inheritance of all studied traits and each of them could not be neglected.

In general, the choice of parents to start breeding program was very important. It could be also concluded that the possibility of producing high yield of squash through hybridization among some promising parental varieties to obtain heterosis. These F₁ hybrids could be used to select in their segregating generations to obtain promising genotypes with great emphasis to powdery mildew resistance.

Similar results were obtained by many authors among them: Farid (1990), El-Adl *et al.*, (1996), Abd El-Hadi *et al.*, (2004) and Abd El-Hadi and El-Gendi (2004).

General combining ability effects (g_i) for the six parental varieties were estimated for all studied traits and the results are presented in Table 6.

The results indicated that the parental varieties P₂ showed desirable positive highly significant values of g_i for all studied yield traits but P₃ for F.Y./P. kg, No. F./P. and F.L. cm. Mean while, the GCA effects were found to be highly significant and positive (desirable) for P₄ (F.L. and No. F./P) and P₅ (W.F./P.g). On the other hand, the best and desirable negative g_i for earliness traits were noticed for the parental varieties P₃ and P₄. On the same time P₂ and P₅ showed desirable negative and highly significant values for D.S. In general, there were no parental variety showed the highest values of g_i for all studied traits.

The estimates of specific combining ability effects (S_{ij}) for all studied traits were obtained and the results are shown in Table 7.

The results illustrated that negative significant values (desirable) for earliness traits in 7 and 5 F₁ hybrids for D.1st F.F. and 1st P.D., respectively.

Table 6: General combining ability effects (g_i) of the parental varieties for all studied traits.

Parents	D.1 st F.F	No.1 st F.F.N.	1 st P.D.	F.Y./P.kg	W.F./P.g	No.F./P.	F.L.cm	F.D.cm	D.S.
P ₁	0.061	0.089	0.071	-0.012	-0.713	-0.238	0.841**	-0.306**	5.556**
P ₂	0.490	-0.019	0.654**	0.073**	4.338**	1.054**	0.700**	-0.387**	-0.944**
P ₃	-1.889**	-0.124*	-875**	0.113**	-1.404	1.196**	1.19**	-0.266**	0.181
P ₄	-1.051**	-0.154**	-1.154**	-0.048	0.258	0.679**	0.925**	-0.283**	4.014**
P ₅	2.103**	0.012	2.233**	-0.113**	7.904**	-2.15**	-4.41**	1.574**	0.764**
P ₆	0.286	0.196**	0.071	-0.109**	-1.192	-0.542*	0.783**	-0.332**	-9.569**
LSD _{0.05}	0.737	0.101	0.716	0.020	1.64	0.504	0.072	0.028	0.571
LSD _{0.01}	0.986	0.135	0.958	0.027	2.19	0.673	0.097	0.038	0.763

** significant differences at 0.05 and 0.01 levels of probability, respectively.

Table7: Specific combining ability effects (s_{ij}) for the 15 F₁ hybrids for all studied traits.

Hybrids	D.1 st F.F	No.1 st F.F.N.	1 st P.D.	F.Y./P.kg	W.F./P.g	No.F./P.	F.L.cm	F.D.cm	D.S.
P ₁ xP ₂	-4.243**	-0.712	-4.03**	0.042	3.231	0.326	0.140	-0.040	14.5**
XP ₃	1.036*	-0.091	0.999	0.012	0.098	0.285	0.419**	-0.089	5.042**
XP ₄	-2.235*	-0.107	-1.888	0.137**	-4.448	1.068	-0.385**	-0.234**	23.208**
XP ₅	-1.089	0.153	-1.376	0.461**	6.623**	2.897**	-0.258*	0.639**	-3.542**
XP ₆	-3.373**	-0.701	-3.146**	0.308**	-4.182	2.722**	0.023	-0.205**	18.792**
P ₂ xP ₃	-1.060	0.064	-1.284	0.164**	1.269	1.393*	0.260*	-0.044	14.208**
XP ₄	-3.164**	-0.615	-2.938**	0.593**	-1.557	4.51**	0.023	0.054	-12.958**
XP ₅	-3.718**	0.428	-4.126**	0.480**	2.014	2.27**	-0.015	0.234**	-7.042**
XP ₆	0.265	0.477	0.137	0.130**	0.377	0.897	0.631**	0.026	3.292**
P ₃ xP ₄	-0.418	0.36	-0.176	0.119**	1.610	0.735	0.002	-0.203**	18.917**
xP ₆	-5.373**	-0.583	-5.83**	0.413**	1.114	2.997**	0.030	0.669**	-7.833**
XP ₆	2.177*	0.239	2.733	0.123**	6.643	0.889	0.610**	0.048	4.167**
P ₄ xP ₅	-3.177**	-0.143	-3.151	0.508**	12.47**	3.214**	0.959**	0.540**	-15.667**
XP ₆	5.307	1.213**	4.679	0.062	12.80**	0.295	1.14**	0.136**	-1.667**
P ₅ xP ₆	-1.314	0.330*	-1.442	0.166**	2.37	1.135	0.234*	0.619**	-8.417**
LSD _{0.05}	1.67	0.274	1.97	0.063	4.504	1.384	0.196	0.078	1.568
LSD _{0.01}	2.24	0.367	2.63	0.085	6.020	1.850	0.263	0.104	2.098

*** significant differences at 0.05 and 0.01 levels of probability, respectively.

While, all the F_1 hybrids had insignificant estimated values for No.1st F.F.N. Concerning yield traits, there were 12,3 and 7 F_1 hybrids showed positive significant values (desirable) for F.Y./P. kg, W.F. g and No. F./P., respectively.

On the other hand, the results indicated that the studied traits of resistance for powdery mildew showed negative significant (desirable) in 7 F_1 hybrids for D.S. trait. The results also cleared that positive values of SCA effects for these traits were found in some hybrids. It also noticed that the F_1 hybrid $P_3 \times P_5$ had the desirable negative highly significant for earliness traits such as D.1st F.F. and 1st P.D. On the other hand the F_1 hybrid $P_2 \times P_4$ had the desirable positive highly significant values for studied traits such as F.Y./P. kg and No. F./P. While, $P_4 \times P_5$ had desirable negative highly significant estimates for D.S. In general, it could be concluded that the best combinations for earliness and powdery mildew resistance was the hybrids which had desirable negative significant SCA effects for these trait. On the other hand, the best combinations for studied traits were the hybrids which had desirable positive significant SCA effects for yield traits.

REFERENCES

- Abd El-Hadi, A.H.; A.M. El-Adl; M.S. Hamada and M.A. Abdein (2005-A). Manifestation of heterosis and genetic parameters associated with it for some vegetative and earliness traits in squash. *J. Agric. Sci. Mansoura Univ.*, 30(3): 1363-1379.
- Abd El-Hadi, A.H.; A.M. El-Adl; M.S. Hamada and M.A. Abdein (2005-B). Heterosis and genetic behavior of some quantitative traits of squash at different environmental conditions. *Alex. J.Agric. Res.* 50 (2) 107 – 120.
- Abd El-Hadi, A.H.; A.M. El-Adl; M.S. Hamada and M.A. Abdein (2005-B). Manifestation of heterosis and genetic parameters associated with it for some vegetative and earliness traits in squash. *J. Agric. Sci. Mansoura Univ.*, 30(3): 1363-1379.
- Abd El-Hadi, A.H. and Soher E.A. El-Gendy (2004). Effect of genotypes by locations interaction on economical traits of squash. *J. Agric. Sci., Mansoura Univ.*, 29(10), 5667 -5687.
- Abd El-Hadi, A.H.; M.M. Zaghoul and A.H. Gaber (2004). Nature of gene action, heterosis and inbreeding depression of yield and yield components traits in squash (*Cucurbita pepo*, L.). *Zagazig J. Agric. Res.*, 31(6): 2707-2725.
- Abd El-Hadi, A.H.; Z.A. Kosba; Z.M. El-Diasty; El-S.H. Askar and G.M. Shamloul (2001). Evaluation of F_1 hybrids among new selected inbred lines of sweet melon (*Cucumis melo* var. *aegyptiacus*, L.). *J. Agric. Sci., Mansoura Univ.*, 26(5): 2831-2845.
- Abd El-Maksoud M.M. (1986). Nature of gene action of economic traits in squash (*Cucurbita Pepo*, L.) M.Sc. Thesis, Fac. of Agric., Mansoura Univ.

- Arora, S.K.; B. Singh; K.C Dhanju., and T.R Ghai,. (1992). Genetics of resistance in summer squash to powdery mildew and cucumber mosaic virus. *Plant disease Research* 7:2, 215-220.
- Dogra, B.S.; K.B. Raslogi; U.K. Kahli and H.K. Sharma (1997). Studies on combining ability in cucumber (*Cucumis sativus*, L.). *annals Agric. Res.*, 18(4): 559-560 (C.F. Pl.Br. Abst., 68(7): 7112).
- El-Adl, A.M.; Z.A. Kosba; Z.M. El-Diasty and A.H. Abd El-Hadi (1996). Types of gene action associated with the performances of hybrids among newly developed inbred lines of aggor, (*Cucumis melo var. chat*, L.). *J. Agric. Sc., Mansoura Univ.*, 21(8): 2821-2835.
- El-Gazar, T.M. (1981). Combining ability and manifestation of heterosis in squash (*Cucurbita pepo*, L.) Ph.D. Thesis, Fac. of Agric. Mansoura Univ.
- Farid, S.D.M. (1990). Efficiency of diallel crosses and factorial mating design in estimating genetic parameters of quantitative traits in cucumber, Ph.D. Thesis, Fac. of Agric., Mansoura Univ.
- Griffing, B. (1956). Concept of general and specific combining ability in relation to diallel crosses system. *Aust. J. Biol. Sci.*, 9: 463-493.
- James, R.V. and W.R. Stevenson (2001). Evaluation of pumpkin and squash varieties for resistance to powdery mildew. *Plant path. Wise.edu/wivegdis* 201.
- Michelle, L.I; S.A. Johaston and S.A. Garrison (1999). Disease tolerance of Zucchini and yellow squash varieties to choose for late season production. Uconn IPM. Integrated pest management, vegetables cucurbits diseases.
- Sadek, Mariam S.S. (2003). Inheritance of some economical traits in squash (*Cucurbita pepo*, L.). M.Sc. Thesis, Fac. of Agric., Mansoura Univ.
- Singh, R.K. and B. Chaudhary (1985). *Biometrical Methods In Quantitative Genetic Analysis*. Katyani Publishers, New Delhi, Revised Ed.
- Steel, G.D. and H. Torrie (1960). *Principles and procedures of statistics*. Mc. Raw. Hill Book Company, INC, New York, PP. 431.
- Sherwood, R.J. and D.G. Hagedorm (1958). Determined common root rot potential of bea fields. *Wis. Agric. Expt. Stn. Bull.*, 531-43.

السلوك الوراثى لبعض الصفات الاقتصادية الهامة وطبيعة المقاومة لمرض البياض الدقيقي فى قرع الكوسة.

أشرف حسين عبد الهادى*، جلال أحمد رزق الشربيني**، أمين علي المغربي***
*قسم الوراثة - كلية الزراعة - جامعة المنصورة - مصر.
**قسم النبات الزراعى (وراثة) - كلية الزراعة بسوهاج - جامعة جنوب الوادى - مصر.
***معهد بحوث أمراض النبات - مركز البحوث الزراعية - مصر.

- استخدم فى هذا البحث ستة من أصناف قرع الكوسة والتي تم الحصول عليها من مصادر مختلفة. وهذه الأصناف هى: Eskandrani (P_1), Giado (P_2), Zucchini mezza lung bianco (P_3), Zucchini 544-005 (P_4), White bush scallop (P_5) and Zucchini nano verda di Milano (P_6). تم التهجين بين هذه الأصناف وذلك للحصول على ١٥ هجيناً للجيل الأول وذلك من خلال نظام التهجينات الدورية مع استبعاد الهجن العكسية. وقد تم تقييم الأصناف الستة وهجينا الخسر عشرة وتم دراسة العديد من الصفات الخضرية و صفات المحصول وكذا المقاومة لمرض البياض الدقيقي.
- أوضحت النتائج وجود فروق عالية المعنوية بين الأصناف وبين الهجن ، وقد كانت هذه النتيجة متوقعة وذلك لأن الأصناف المستخدمة فى هذه الدراسة قد تم الحصول عليها من مصادر مختلفة.
- ولوحظ أيضا أنه لا يوجد أحد الأصناف قد تفوق أو كان هو الأفضل لكل الصفات التى درست وأن كان P_3 هو الأكثر تبيكيرا وكذلك أعطى أعلى محصولا. وفى نفس الوقت كان P_5 و P_6 الأفضل لبعض الصفات.
- وفيما يخص بالهجن فقد كان الهجين $P_3 \times P_5$ هو أكثر هجين تبيكيرا فى النضج بينما كان $P_4 \times P_6$ أكثر تأخرا فى النضج. وكان $P_5 \times P_6$ أقل الهجن فى محصول الثمار/النبات وعدد الثمار فى النبات. وكان أعلى الهجن هو $P_2 \times P_4$ وذلك لصفة عدد الثمار/النبات. ومن ناحية أخرى كان الهجين $P_5 \times P_6$ هو الأكثر مقاومة لمرض البياض الدقيقي.
- أظهرت الدراسة وجود قوة الهجين لغالبية الهجن: حيث أن أعلى قيمة لقوة الهجين تم الحصول عليها هى: $-(P_3 \times P_5) \% 19.72$ ، $-(P_1 \times P_2) \% 26.16$ ، $-(P_1 \times P_6) \% 9.55$ ، $-(P_4 \times P_5) \% 11.43$ ، $-(P_4 \times P_6) \% 20.62$ ، $-(P_1 \times P_5) \% 76.05$ ، $-(P_4 \times P_5) \% 20.62$ ، $-(P_5 \times P_6) \% 34.28$ ، $-(P_1 \times P_6) \% 8.03$.
- عدد العقد حتى أول زهرة مؤنثة (No.1st F.F.N.) ، تاريخ أول جمعه (1st P.D.) ، محصول الثمار/نبات (F.Y./Pkg) ، وزن الثمار/نبات بالجرام (W.F./P.g) ، عدد الثمار/نبات (No.) ، طول الثمار (F.L.cm) ، قطر الثمار (F.D. cm) ، شدة المرض (D.S.) على الترتيب. كما أوضحت النتائج أيضا أهمية كل من التباين الوراثى الإضافى وغير الإضافى شاملا السيادة فى وراثة الصفات محل الدراسة فى قرع الكوسة.

