

DIALLEL ANALYSIS OF NINE WHITE MAIZE INBRED LINES FOR DIFFERENT CHARACTERS UNDER DIFFERENT LOCATIONS

Mosa, H.E.

Maize Research Section, FCRI, Sakha ARS, ARC, Egypt

ABSTRACT

A Number of 36 single crosses obtained directly from crossing of nine inbred lines in a half diallel system and two checks were evaluated in a randomized complete block design, with four replications, in three locations; i.e., Sakhá, Gemmeiza and Sids Res. Stations in 2005 summer season. The objectives of this study were to estimate heterosis and combining ability. The combined analysis of data across environments showed highly significant for locations (Loc), general combining ability (GCA), specific combining ability (SCA) and GCA x Loc. interaction for all studied characters except GCA x Loc interaction for ear height. Additive genetic effects exhibited to have played an important role than the non-additive genetic effects in the expression of all studied characters, except for grain yield under and over the three locations. Also, the additive genetic effects were much more influenced by change of location than the non-additive genetic effects for all studied characters. The highest desirable values of GCA effects were obtained from Sk5170 and Sk8170 inbreds for silking date, plant and ear height, Sk6017, Sk8014 and Sk9195 inbreds for grain yield and ear length and Gm152 inbred for number of rows/ear. The best crosses having desirable values for both SCA and heterotic effects relative to the best check were Sk8012 x Sk8170 for silking date; Sk63 x Sk8012, Sk5170 x Sk8170 and Sk8014 x Sk9195 for plant and ear height; Sd63 x Sk6017, Sk6017 x Sk8014 and Sk6018 x Sk9195 for grain yield, Sd63 x Sk6017 for ear length and Gm152 x Sk6017 and Gm152 x Sk9195 for number of rows/ear. It is worth noting that a cross having good SCA effects, may come from two parents when at least one of them possesses high GCA effects.

INTRODUCTION

Combining ability of inbred lines is the ultimate factor determining future usefulness of the lines for hybrids. Combining ability initially was a general concept considered collectively for classifying an inbred line relative to its cross performance. Sprague and Tatum (1942) refined the concept of combining ability; general combining ability (GCA) and specific combining ability (SCA) have a significant impact on inbred line evaluation and population improvement in maize breeding. Dhillon *et al.* (1978), Maggiore *et al.* (1979), Nevado and Cross (1990), Melchinger *et al.* (1990), Lima *et al.* (1995), Tulu and Ramachandrapa (1998), Choukan (1999), Rameeh *et al.* (2000), Zelleke (2000), Desai and Singh (2001), Mahto and Ganguly (2001), San Vicente *et al.* (2001) and Fan Xing Ming *et al.* (2001) found that both GCA and SCA were significant for silking date, plant and ear heights, grain yield, ear length and number of rows/ear. However Debnath *et al.* (1988), Nawar *et al.* (1979), Crossa *et al.* (1990), Pal and Prodhan (1994), Lima *et al.* (1995), Dehghanpour *et al.* (1996) and Geetha and Jayaraman (2000) reported that the SCA (or non additive gene effects) was more important than GCA (or additive gene effects) in the inheritance of grain yield. While, Nawar *et al.* (1979), Cross *et al.* (1990), Tulu and Ramachandrapa (1998), Choukan (1999), and Nigussie and Zelleke (2001) found that the additive gene effects

were the primary type of gene actions operative in the crosses for silking date, plant and ear heights, ear length and number of rows/ear. This study aimed to estimate of type and relative amount of genetic variance components and their interaction with locations and heterotic effects relative to check variety.

MATERIALS AND METHODS

The materials used in this study were nine white maize inbred lines obtained by the maize research program i.e., Sd63, Gm152, Sk5170, Sk6017, Sk6018, Sk8012, Sk8014, Sk8170, Sk9195. In 2004, summer growing season, the nine inbred lines were hand crossed in half diallel fashion to obtain hybrid seeds of 36 genotypes i.e., 36 F_{1S} . At the 2005 growing season, the obtained hybrid seeds of 36 genotypes plus the two commercial hybrids i.e., SC10 and SC129 were grown at three locations i.e., Sakha, Gemmeiza and Sids Experimental Stations. The maize genotypes were grown in four replications of a randomized complete block design. Each plot was a single row, 6m long, 80cm apart and 25cm hill spacing. After 21 days from planting date, plants were thinned to one plant per hill. The recommended cultural practices were applied at the proper time. Data were recorded on grain yield ard/fed (1 ardab = 140 Kg and 1 feddan = 4200 m²), adjusted on 15.5 basis grain moisture content and shelling percentage, ear length (cm), number of rows/ ear, plant height (cm), ear height (cm) and number of days from planting to 50% silking emergence. An ordinary analysis of variance for the data was performed for each location then combined over the three locations according to Steel and Torrei (1980). The effect of hybrids was considered fixed effect while that of locations was considered to be random effect. Variations among the 36 hybrids was partitioned into GCA and SCA and their interactions with locations according to Griffing (1956) for Method-4, Model-1.

RESULTS AND DISCUSSION

Mean squares of diallel analysis of 9x9-inbred lines combined over three locations for six characters are presented in Table (1). Locations mean squares were highly significant for all characters with mean values for Sakha Res. Station (North Egypt) being higher than those for Gemmeiza and Sids Res. Stations (Middle and South Egypt, respectively) for all characters. Frey and Maldonado (1967) defined the stress environment as the one in which mean performance for certain attribute is low. Therefore, Sakha Station seemed to be the non-stress environment. The mean squares of GCA and SCA were highly significant for all studied characters. Thus both additive and non-additive gene actions were found to be important in controlling all studied characters. These results are in agreement with those obtained by Nawar *et al.* (1979) for silking date, El-Hosary (1988) for plant height, El-Shamarka (1995) for ear height, Abdel-Sattar *et al.* (1999) for ear length and number of rows/ear, Mosa (2005) for grain yield. However, the additive genetic effects (K^2 GCA) exhibited to have played an important role than the non-additive genetic effects (K^2 SCA) in the expression of all studied characters, except for grain yield under three locations and their combined Table (2). These results support the findings of Nawar *et al.* (1988) for plant and ear heights, Mosa (2005) for silking date, Mosa (2003) for grain yield, Motawei (2005) for ear

length and El-Shenawy (2005) for number of rows/ear. Meanwhile the mean squares of interaction between GCA and locations were significant or highly significant for all studied characters, except for ear height, The magnitude of interactions for GCA x locations was generally higher than that for SCA ones for all characters (Table 2), revealing that additive genetic effects were much more influenced by change of locations than non-additive genetic effects in all studied characters. These results are in agreement with those previously attained by Matzinger *et al.* (1959) El-Rouby *et al* (1973), Debnath and Sarkar (1987), El-Hosary (1988), Mahmoud (1996) and Mosa (2003 and 2005) suggested that the additive gene effects were more influenced by interaction with environments than the non-additive gene effects.

Table (1): Mean squares of diallel analysis of 9 x 9 inbred lines combined over three locations for six characters.

S.O.V	d.f	Silking date (days)	Plant height (cm)	Ear height (cm)	Grain yield (ard/fed)	Ear length (cm)	No.of rows/ear
Locations (L)	2	1004.22**	25069.02**	7856.81**	9422.70**	347.08**	21.79**
Rep/Loc	9	22.11	1876.47	1099.99	30.43	5.16	0.60
GCA	8	302.64**	12246.09**	9928.08**	438.69**	117.08**	31.56**
SCA	27	11.74**	1059.9**	548.28**	208.56**	17.56**	1.66**
GCA x L	16	2.85*	676.17**	115.97	75.07**	11.64**	3.66**
SCA x L	54	1.61	207.44	95.67	12.38	2.85	0.68
Error	333	1.42	134.63	86.42	12.12	2.15	0.51

*,** significant at the 0.05 and 0.01 levels of probability, respectively
+ Including checks

Table (2): Estimates of genetic and genetic x environment parameters for six characters under three locations.

Loca	Silking date (days)		Plant height (cm)		Ear height (cm)		Grain yield (ard/fed)		Ear length (cm)		No.of rows/ear	
	K ^o GCA/K ^o SCA	σ ^o GCAxL/σ ^o SCAxL	K ^o GCA/K ^o SCA	σ ^o GCAxL/σ ^o SCAxL	K ^o GCA/K ^o SCA	σ ^o GCAxL/σ ^o SCAxL	K ^o GCA/K ^o SCA	σ ^o GCAxL/σ ^o SCAxL	K ^o GCA/K ^o SCA	σ ^o GCAxL/σ ^o SCAxL	K ^o GCA/K ^o SCA	σ ^o GCAxL/σ ^o SCAxL
Sakha	2.58	----	3.28	----	5.45	----	0.35	----	1.30	----	4.01	----
Gemm	6.84	----	0.36	----	1.31	----	0.20	----	0.80	----	1.60	----
Sids	4.52	----	4.80	----	6.20	----	0.81	----	1.05	----	7.5	----
mb.	4.22	1.25	1.93	1.06	3.09	3.8	0.26	37.3	1.02	1.9	4.06	2.75

Estimates of GCA effects for the nine parental inbred lines in the combined analysis are presented in Table (3). The parental inbred lines Sk5170, Sk8012, Sk8170 and Sk9195 showed desirable values of GCA effects for silking date and plant and ear height, indicating that the four inbred lines could be considered as good combiners for developing early and short stalk genotypes. The inbred lines Sk6017, Sk8014 and Sk9195 had significant positive GCA effects for grain yield and ear length, while inbred line Gm 152 exhibited positive GCA effects for grain yield and number of rows/ear. Inbred lines Sd63 and Sk6018 for ear length and inbred line Sk9195 for number of rows/ear. These results suggested the importance of these lines to be involved for improving maize.

Table (3): Estimates of general combining ability effects of nine inbred lines for six characters over three locations.

Line	Silking date (days)	Plant height (cm)	Ear height (cm)	Grain yield ard/fed	Ear length (cm)	Number of rows/ear
Sd 63	2.46*	2.26	4.37*	-1.14*	0.35*	-0.41*
Gm 152	0.81*	-4.75*	3.93*	1.32*	-0.53*	1.45*
Sk5170	-2.78*	-15.96*	-17.54*	-1.91*	-1.20*	0.07
Sk6017	1.46*	24.17*	20.44*	3.23*	1.79*	-0.12
Sk6018	1.86*	10.71*	7.25*	-0.21	0.58*	-0.44*
Sk8012	-0.60*	-2.96*	-2.74*	-3.12*	-1.20*	-0.19*
Sk8014	0.42*	1.36	-0.49	0.90*	0.98*	-0.50*
Sk8170	-2.69*	-13.06*	-9.69*	-2.12*	-1.60*	-0.23*
Sk9195	-0.94*	-1.77	-5.54*	3.06*	0.84*	0.37*
LSD g. 0.05	0.24	2.33	1.87	0.70	0.29	0.14

* significant from zero at 0.05 level of probability.

Mean performance (M), % heterosis relative to the best check (H) and specific combining ability effects for six characters over three locations are shown in Table (4).

Table (4): Mean performance (M); % heterosis relative to the best check (H) and specific combining ability effects (SCA) for six characters over three locations.

Cross	Silking date (days)			Plant height (cm)			Ear height (cm)		
	M	% H	SCA	M	% H	SCA	M	% H	SCA
Sd 63 x Gm152	65.58	3.56*	0.06	275.83	-4.33*	3.02	166.00	3.75	3.43
x Sk-5170	61.33	-3.15*	-0.59*	256.58	-7.89*	3.99	144.58	-9.63*	3.49
x Sk-6017	65.83	3.94*	-0.34	309.25	7.25*	7.51*	186.16	16.35*	7.09*
x Sk-6018	65.66	3.67*	-0.89*	284.66	-1.27	-3.61	162.25	1.40	-3.64
x Sk-8012	66.83	6.52*	2.73*	247.08	-14.30*	-27.51*	137.08	-14.32*	-18.80*
x Sk-8014	65.00	2.63*	-0.13	282.25	-2.10	3.32	158.00	-1.25	-0.14
x Sk-8170	61.00	-3.87*	-1.01*	267.50	-7.22*	3.00	147.83	-7.60*	-1.10
x Sk-9195	63.91	0.91	0.16	286.08	-0.78	10.29*	162.75	1.71	9.66*
Gm152x Sk-8170	60.25	-4.86*	-0.02	255.25	-11.47*	0.66	138.25	-13.59*	-2.40
x Sk-6017	64.66	2.10*	0.15	300.00	4.04*	6.27	179.33	12.08*	0.70
x Sk-6018	64.75	2.24*	-0.16	276.33	-4.16*	-4.93	160.33	0.20	-5.11*
x Sk-8012	61.41	-3.03*	-1.03*	264.91	-8.12*	-2.67	153.50	-4.06	-1.95
x Sk-8014	63.41	0.12	-0.05	271.58	-5.80*	-0.34	159.50	-0.31	1.8
x Sk-8170	60.50	-4.46*	0.15	262.91	-8.81*	5.43	153.16	-4.27	4.67*
x Sk-9195	63.00	-0.52	0.89*	262.33	-9.01*	-6.44*	151.50	-5.31*	-1.15
Sk-5170xSk-6017	60.42	-4.59*	-0.51	282.66	-1.96	-0.86	158.66	-0.83	1.51
x Sk-6018	61.08	-3.55*	-0.23	271.66	-5.78*	1.62	137.75	-13.9*	-6.22*
x Sk-8012	58.08	-8.28*	-0.76*	267.83	-7.10*	11.46*	141.50	-11.56*	7.53*
x Sk-8014	59.58	-5.92*	-0.29	262.41	-8.98*	1.71	141.58	-11.56*	5.36*
x Sk-8170	59.33	-6.31*	2.87*	226.50	-21.44*	-19.78*	116.83	-26.98*	-10.18*
x Sk-9195	58.33	-7.89*	-0.17	258.75	-10.25*	1.19	132.08	-17.45*	0.91
Sk-6017xSk-6018	65.08	2.76*	-0.48	295.66	2.54	-14.53*	182.66	14.16*	0.71
x Sk-8012	64.41	1.70*	1.32*	300.25	4.13*	3.74	171.58	7.23*	-0.38
x Sk-8014	64.16	1.31	0.04	303.66	5.31*	2.82	175.75	9.84*	1.54
x Sk-8170	60.75	-4.07*	-0.26	293.50	1.79	7.08*	167.33	4.58	2.33
x Sk-9195	62.83	-0.78	0.08	286.66	-0.57	-11.04*	155.66	-2.71	-13.49*
Sk-6018xSk-8012	63.25	-0.12	-0.24	287.25	-0.37	4.20	166.33	3.95	7.57*
x Sk-8014	66.08	4.34*	1.57*	294.33	2.08	6.95*	164.25	2.65	3.23
x Sk-8170	62.00	-2.10*	0.60*	273.91	-6.00*	0.96	150.5	-5.93*	-1.32
x Sk-9195	63.00	-0.52	-0.15	293.58	1.82	9.34*	160.75	0.46	4.78*
Sk-8012x x Sk-8014	61.50	-2.88*	-0.55	270.75	-8.09*	-2.96	146.41	-8.49*	-4.60*
x Sk-8170	57.83	-8.68*	-1.10*	269.00	-6.70*	9.72*	147.33	-7.91*	5.52*
x Sk-9195	60.33	-4.73*	-0.35	274.58	-4.76*	4.02	151.08	-5.67*	5.11*
Sk-8014 x Sk-8170	59.41	-6.18*	-0.54	258.33	-10.40*	-6.28	143.41	-10.36*	-0.65
x Sk-9195	61.66	-2.63*	-0.04	268.66	-6.82*	-6.23*	141.66	-11.46*	-6.55*
Sk-8170 x Sk-9195	58.16	-8.16*	-0.42	259.33	-10.05*	-1.13	139.75	-12.65*	0.73
Check SC10	65.33	---	---	301.50	---	---	172.0	---	---
Check SC129	63.33	---	---	288.33	---	---	160.0	---	---

* significant from zero at the 0.05 level of probability.

Cont. (4):

Cross	Grain yield (ard/fed)			Ear length (cm)			No. of rows/ear		
	M	% H	SCA	M	% H	SCA	M	% H	SCA
Sd 63 x Gm152	29.04	-14.83*	0.23	20.78	-7.84*	0.34	14.81	-0.13	-0.28
x Sk-5170	30.86	-9.50*	4.96*	19.80	-12.19*	0.19	14.4	-2.89	0.59*
x Sk-6017	35.42	3.87	4.56*	24.31	7.80*	1.59*	13.96	-5.86*	0.38*
x Sk-6018	25.65	-24.78*	-1.82*	21.4	-5.09	-0.10	13.03	-12.13*	-0.21
x Sk-8012	12.72	-62.69*	-11.99*	15.65	-30.59*	-4.06*	12.90	-13.01*	-0.62*
x Sk-8014	30.01	-11.99*	1.48	21.9	-2.88	-0.01	13.16	-11.26*	-0.06
x Sk-8170	27.90	-18.18*	2.26*	20.33	-9.84*	1.08*	14.0	-6.59*	0.57*
x Sk-9195	31.11	-8.76*	0.32	22.91	1.59	0.96*	13.66	-7.88*	-0.36*
Gm152x Sk-8170	27.32	-19.88*	-0.75	18.33	-18.71*	-0.51	15.83	6.74*	0.22
x Sk-6017	34.70	1.81	1.26	20.93	-7.18*	-0.92*	15.80	6.54*	0.35*
x Sk-6018	31.15	-8.65*	1.30	19.91	-11.70	-0.62	15.10	1.82	0.01
x Sk-8012	27.71	-18.73*	0.63	19.55	-13.30	0.65	15.53	4.72*	0.08
x Sk-8014	30.33	-11.05*	-0.74	21.48	-4.74	0.38	14.66	-1.14	-0.43*
x Sk-8170	28.47	-16.51*	0.38	19.3	-14.41*	0.89*	15.10	1.82	-0.29
x Sk-9195	30.91	-9.35*	-2.31*	20.58	-8.82*	-0.22	16.2	9.23*	0.35*
Sk-5170xSk-6017	31.95	-6.30	1.75*	22.60	0.22	1.32*	14.13	-4.72*	-0.02
x Sk-6018	26.97	-20.9*	0.20	20.10	-10.86*	0.12	13.66	-7.88*	-0.03
x Sk-8012	26.50	-22.28*	2.70*	16.71	-17.02*	0.49	14.23	-4.04*	0.05
x Sk-8014	27.67	-18.85*	-0.25	20.60	-8.84*	0.30	13.26	-10.55*	-0.30
x Sk-8170	13.85	-59.38*	-10.08*	15.48	-31.35*	-2.27*	13.03	-12.13*	-0.83*
x Sk-9195	32.39	-5.01	2.26*	20.53	-9.95*	0.36	14.86	0.20	0.31
Sk-6017xSk-6018	29.57	-13.28*	-2.37*	22.48	-0.31	-0.46	13.16	-11.26*	-0.33
x Sk-8012	26.01	-17.85*	-0.95	21.40	-5.09	0.15	13.73	-7.41*	-0.08
x Sk-8014	35.21	3.25	2.35*	22.88	1.46	-0.44	13.76	-7.21*	0.22
x Sk-8170	30.68	-10.02	0.63	21.13	-6.29*	0.39	13.63	-8.09*	0.03
x Sk-9195	27.85	-18.32*	-7.23*	21.61	-4.16	-1.64*	13.73	-7.41*	-0.56*
Sk-6018xSk-6012	25.74	-24.51*	0.33	20.80	-7.76*	0.87*	13.50	-8.96*	0.07
x Sk-8014	27.06	-20.64*	-2.45*	22.76	0.93	0.68	13.40	-9.64*	0.13
x Sk-8170	28.66	-15.95*	2.17*	18.91	-16.14*	-0.64	13.60	-8.29*	0.19
x Sk-9195	34.70	0.41	2.84*	22.18	1.64	0.15	14.16	-4.51*	0.16
Sk-8012x x Sk-8014	27.61	-19.03*	0.88	20.33	-9.84*	-0.11	13.66	-7.88*	0.13
x Sk-8170	27.15	-20.38*	3.50*	19.31	-14.36*	1.39*	13.86	-6.54*	0.11
x Sk-9195	33.69	-1.20	4.89*	20.78	-7.84*	0.61	14.53	-2.02	0.25
Sk-8014 x Sk-8170	28.32	-16.95*	0.63	19.35	-14.19*	-0.71*	13.66	-7.88*	0.33
x Sk-9195	30.86	-9.50*	-1.89*	22.33	-0.97	-0.08	13.86	-6.54*	-0.02
Sk-8170 x Sk-9195	31.11	-8.76	1.31	19.65	-12.86*	-0.15	14.16	-4.51*	-0.12
Check SC10	34.10	—	—	22.65	—	—	13.4	—	—
Check SC129	30.60	—	—	21.18	—	—	14.83	—	—

* significant from zero at 0.05 level of probability.

Six crosses showed significantly negative SCA effects for silking date, where five of them significantly had heterotic effects for earliness than the best check SC129 (63.3 days). The cross Sk8012 x Sk8170 had the best desirable values for both SCA and heterotic effects. Six crosses showed significantly negative SCA effects for plant height, where three of them significantly had shorter stalks than the best check SC129 (288.33 cm). Meanwhile seven crosses exhibited significantly negative SCA effects for ear height, five of them significantly were shorter than SC129 (160 cm). Concerning plant and ear height, the crosses Sd 63 x Sk8012 and Sk5170 xSk8170 and Sk8014 x Sk9195 had best values for both SCA and heterotic effects among tested genotypes and could be of value for breeding for lodging resistance. Eleven crosses showed significantly positive SCA effects for grain yield, where three of them; i.e. Sd63 x Sk6017, Sk6017 x Sk8014 and Sk6018 x Sk9195 outyielded the best check SC10 (34.1 ard/fed). Hence it could be concluded that these crosses offer good possibility for improving grain yield of maize.

Seven crosses showed significantly positive SCA effects for ear length; the cross Sd63 x Sk6017 had the highest values for both SCA and heterotic effects. Five crosses showed desirable values for SCA effects for number of rows/ ear, out of them i.e. Gm152 x Sk6017 and Gm152 x Sk9195 significantly had more rows/ear than the best check SC129. It is worth noting that a cross having good SCA effects may come from two parents possessing good GCA or from one with good GCA and the other with poor GCA effects , similar findings were obtained by Nawar *et al.*, (1979), El-Hosary (1988), Mahmoud (1996) and Mosa (2003).

REFERENCES

- Abdel-Sattar, A.A.; A.A. El-Hosary and M.H. Motawea (1999). Genetic analysis of maize grain yield and its components by diallel crossing. *Minufiya J. Agric. Res.*, Vol.24 No.1: 43-63.
- Choukan, R. (1999). General and specific combining ability of ten maize inbred lines for different traits in diallel crosses. *Seed and plant*, Vol. 15 No.3: 280-295.
- Crossa, J; S.K. Vasal and D.L. Beck (1990). Combining ability estimates of CIMMYT's tropical late yellow maize germplasm. *Maydica.*, Vol.35 No.3: 273-278.
- Debanth, S.C. and K.R. Sarkar (1987). Genetic analysis of grain yield and other characters in maize (*Zea mays* L.). *Egypt J. Genet. Cytol.*, 17: 49-61.
- Debnath, S.C.; K.R. Sarkar and Daljit Singh (1988). Combining ability estimates in maize (*Zea mays* L.). *Annals of Agric. Res.*, Vol.9 No.1: 37-42.
- Dehghanpour, Z.; B. Ehdai and M. Moghaddam (1996). Diallel analysis of agronomic characters in white endosperm corn. *Journal of Genetic & Breeding.*, Vol.50 No.4: 337-365.
- Desai, S.A. and R.D. Singh (2001). Combining ability studies for some morphophysiological and biochemical traits related to drought tolerance in maize (*Zea mays* L.). *Indian J. Genetic and plant Breed.*, Vol. 61 No.1: 34-36.
- Dhillon, B.S.; Joginder Singh; A.S. Seth and N.N. Singh (1978). Combining ability in maize under varying plant densities. *Indian J. Genetics and plant Breed.*, Vol.38 No.3 304-312.
- El-Hosary, A.A. (1988). Heterosis and combining ability of ten maize inbred lines as determined by diallel crossing over two planting dates. *Egypt.J.Agron.*, Vol.13 No.1,2:13-25.
- El-Rouby, M.M.; Y.S. Koraim and A.A. Nawar (1973). Estimation of genetic variance and its components in maize under stress and non-stress environment. I. Planting date. *Egypt J. Genet. and Cytol.*, 2: 10-19.
- El-Shamarka, Sh.A. (1995). Estimation of heterotic and combining ability effects for some quantitative characters in maize under two nitrogen levels. *Minufiya J Agric. Res.*, Vol.20 No.2: 441-462.

- El-Shenawy, A.A. (2005). Combining ability of prolific and non-prolific maize inbred lines in their diallel crosses for yield and other traits. *J. Agric. Res. Tanta Univ.*, 31(1):16-31.
- Fan Xing Ming; Tan Jing; Huang Bihue and Liu Feng (2001). Analysis of combining ability and heterotic groups of yellow grain quality protein maize inbred. *Hereditas (Beijing)*, Vol.23 No.6: 547-552.
- Frey, K.J. and Maldonado,U. (1967). Relative productivity of homogenous and heterogenous oat cultivars in optimum and suboptimum environments. *Crop Sci.*, 7: 532-535.
- Geetha, K. and N. Jayaraman (2000). Genetic analysis of yield in maize (*Zea mays* L.). *Madras Agric. J.*,Vol.87 No.10/12: 638-640.
- Griffing, B. (1956). Concept of general and specific combining ability in relation to diallel crossing systems. *Aust. J. Biological Sci.*, 9:463-493.
- Lima, M.; J.B.de Miranda Filho and P.R. Furlani (1995). Diallel cross among inbred lines of maize differing in aluminum tolerance. *Brazil. J. Genetics*, Vol.18 No.4: 579-584.
- Maggiore, T.; C. Lorenzoni; M. Motto; M. Perenzin and F. Salamini (1979). Total dry matter and grain yield in late maize hybrids. *Rivista di Agronomia.*, Vol. 13 No.4: 431-435.
- Mahmoud, A.A. (1996). Evaluation of combining ability of newly developed inbred lines of maize. Ph. D. Thesis, Fac. Agric., Cairo Univ. Egypt.
- Mahto, R.N. and D.K. Ganguly (2001). Heterosis and combining ability studies in maize (*Zea mays* L.). *J. Res., Birsa Agric. Univ.*, Vol.13 No.2: 197-199.
- Matzinger, D.F.; G.F. Sprague and C.C. Cockerham (1959). Diallel crosses of experiments repeated over locations and years. *Agron. J.*, 51:346-349.
- Melchinger, A.E.; M. Lee; K.P. Lamkey; A.R. Hallauer and W.L. Woodman (1990). Genetic diversity for restriction fragment length polymorphisms and heterosis for two diallel sets of maize inbreds. *Theor. Appl. Genet.*, 80: 488-496.
- Mosa, H.E. (2003). Heterosis and combining ability in maize (*Zea mays* L.). *Minufiya J. Agric Res.*, 28: 1375-1386.
- Mosa, H.E. (2005). Combining ability of eight yellow maize (*Zea mays* L.) inbred lines for different characters in diallel crosses. *J Agric. Res. Tanta Univ.*, 31: 604-615.
- Motawei, A.A. (2005). Combining ability and heterotic effect of nine maize inbred lines via diallel cross analysis. *Minufiya J. Agric. Res.*, 30: 197-214.
- Nawar, A.A.; M.E. Gomaa and M.S. Rady (1979). Heterosis and combining ability in maize. *Egypt J. Genetics. Cytol.*, 9: 255-269.
- Nawar, A.A.; M.M. El-Sayed; H.A. Dawwan and M.A. Shehata (1988). Nitrogen fertilizer level effects on heterosis and combining ability in maize. *Minufiya J. Agric. Res.*, 13: 919-935.
- Nevado, M.E. and H.Z. Cross (1990). Diallel analysis of relative growth rates in maize synthetics *Crop Sci.*, 30: 549-552.
- Nigussie, M. and H. Zelleke (2001). Heterosis and combining ability in a diallel among eight elite maize populations. *African Crop sci. J.*, Vol.9 No.3: 471-479.

- Pal, A.K. and H.S. Prodhon (1994). Combining ability analysis of grain yield and oil content along with some other attributes in maize (*Zea mays* L.). Indian J. Genetics and Plant Breed., Vol.54 No.4: 376-380.
- Rameeh, V.; A. Rezai and A. Arzani (2000). Estimates of genetic parameters for yield and yield components in corn inbred lines using diallel crosses. J. Sci. and Technol. Agric. and Natural Resources, Vol.4 No.2: 95-104.
- San Vicente, F.; A. Bejarano; J. Crossa and C. Marian (2001). Heterosis and combining ability of tropical yellow endosperm maize populations. Agronomia Tropical (Maracay), Vol.51 No.3: 301-318.
- Sprague, G.F. and L.A. Tatum (1942). General vs. specific combining ability in single crosses of corn. J. Amer.Sco. Agron., 34: 923-932.
- Steel, R.G. and J.H. Torrie (1980). Principals and procedures of statistics. A Biometrical Approach. 2nd Ed. Me. Graw Hill, N. Y. USA.
- Tulu, L. and B.K. Ramchandrapa (1998). Combining ability of some traits in a seven parent diallel cross of selected maize (*Zea mays* L.) populations. Crop Research (Hisar), Vol.15 No.2/3:232-237.
- Zelleke, H. (2000). Combining ability for grain yield and other agronomic characters in inbred lines of maize (*Zea mays* L.). Indian J. Genetics and Plant Breed., Vol.60 No.1: 63-70.

تحليل التهجينات الدائرية لتسع سلالات من الذرة الشامية الأبيض لصفات مختلفة تحت مواقع مختلفة

حاتم الحمادى موسى

مركز البحوث الزراعية- معهد المحاصيل الحقلية- محطة بحوث سخا- قسم بحوث الذرة الشامية

تم تقييم ٣٦ هجيناً فردياً ناتجة من التهجين بين ٩ سلالات من الذرة الشامية البيضاء بنظام التزاوج الدائري مع استبعاد التزاوجات العكسية بالإضافة الي هجينين تجاريين في تصميم القطاعات الكاملة العشوائية في أربع تكرارات في ثلاث مواقع بحثية سخا و الجميزة و سنس في موسم صيف ٢٠٠٥. وذلك لدراسة القدرة علي الانتلاف وقوة الهجين. أظهر التحليل المشترك معنوية عالية بين المواقع والفترة العامة والخاصة علي الانتلاف والتفاعل بين الفترة العامة علي الانتلاف والمواقع لكل الصفات تحت الدراسة ما عدا تفاعل الفترة العامة علي الانتلاف مع المواقع لصفة ارتفاع الكوز. أظهرت النتائج أن الفعل الوراثي المضيف للجين كان أكثر أهمية من الفعل الوراثي غير المضيف في وراثته جميع الصفات تحت الدراسة ما عدا صفة المحصول. كذلك كان الفعل الوراثي المضيف للجين أكثر تلقراً بالبيئة من الفعل الوراثي غير المضيف لكل الصفات تحت الدراسة. كانت أعلى قيم مرغوبة تحصل عليها لتأثيرات الفترة العامة علي الانتلاف في السلالات سخا ٥١٧٠ وسخا ٨١٧٠ لصفات تاريخ ظهور ٥٠% من الحرائر للنورات المؤنثة وارتفاع كل من النبات والكوز، وكذلك السلالات سخا ٦٠١٧ وسخا ٨٠١٤ وسخا ٩١٩٥ لصفات محصول الحبوب وطول الكوز والسلالة جميزة ١٥٢ لصفة عدد الصفوف في الكوز. كذلك كانت أعلى قيم مرغوبة تم التحصل عليها لكلا من تأثيرات الفترة الخاصة علي الانتلاف وقوة الهجين نسبة لافضل هجن المقارنة في الهجن سخا ٨٠١٢ X سخا ٨١٧٠ لصفة تاريخ ظهور ٥٠% من حرائر النورات المؤنثة و الهجين سنس ٦٢ X سخا ٨٠١٢، وسخا ٥١٧٠ X سخا ٨١٧٠ وسخا ٨٠١٤ X سخا ٩١٩٥ لصفات ارتفاع النبات والكوز والهجن سنس ٦٢ X سخا ٦٠١٧ وسخا ٦٠١٧ X سخا ٨٠١٤ وسخا ٦٠١٨ X سخا ٩١٩٥ لصفات محصول الحبوب والهجين سنس ٦٢ X سخا ٦٠١٧ لطول الكوز والهجين جميزة ١٥٢ X سخا ٦٠١٧ وجميزة ١٥٢ X سخا ٩١٩٥ لصفة عدد الصفوف في الكوز. وتفيد نتائج الدراسة انه يمكن الحصول على هجين يتميز بفترة خاصة مرتفعة علي الانتلاف من أبوين يتميز أحدهما على الأقل بفترة عامة عالية على الانتلاف