

## STUDY ON THE PRODUCTION OF LOW CHOLESTEROL CREAM-LIKE.

Elhadidi, A.A.; M.A.M. Omar and M. A. Ali

Dairy Dept., Fac. of Agriculture, Al-Azhar University Cairo, Egypt.

### ABSTRACT

The mixed cream with coconut and palm seed oils may affects their properties, as fat constants and fatty acids composition. Among the vegetable oils, these fats proved to be the most suitable source of fatty products and to provide nutritional value. Therefore, these study was planned to investigate the possibility and the effect of adding these oils to the natural cream, From the obtained results, one can conclude the following: The addition of 25% and 50% of coconut oil to the cream resulted to a good quality cream, and decreased cholesterol from 210 to 162 and 115 mg/100g, respectively, with no effect on fat constant or the fatty acids composition. The addition of coconut and date palm oils to cream in the level of 75%, failed to produce cream of good quality. The addition of coconut and date palm oils to cream at the levels 25 and 50% gained the highest score points among the examined treatments. The addition 25% and 50% of date palm oil to the cream resulted in a good quality cream and decreased cholesterol from 210 to 157 and 97 mg/100g respectively, with no effect on fat constant or the fatty acids composition

### INTRODUCTION

There is more than one source for the world consumption of fats and oils which are animal and plant sources, the preference to consume one of them than the other depends on many factors. The healthy and economical factors are the most important ones, the milk fats is decreased in production, besides the high price due to the increment of the production cost. So, the consumers aimed to use the vegetable oils as a substitute of milk fats. Since the fats and oils are the concentrate source of energy compared with protein or starch substances, and one gram of protein or starch produces only four calories, while one gram of fats or oils produces nine calories, the fats and oils are even in their calories produced. Besides, the lockage of proteins sources in societies suffering from nutrition shortage, so, it is recommended to use the oils as an alternative source of energy. Consuming the animal fats comprising milk fat may cause a healthy problems due to the cholesterol content of these fats, while the vegetable oils contains the phytosterols (Bindal *et al.*, 1973). On the other hand, the triglycerides of vegetable oils containing unsaturated fatty acids, gave the oils its chemical character is ties characteristics concerning the oxidation. It was reported that the unsaturated fatty acids increase the high density cholesterol of blood, while decrease the low density cholesterol in blood, which causes the healthy problems. Mixing of vegetable oils with animal fats may partially led to reduce the costs of the produced admixture, besides the reducing of cholesterol content consequently may avoid the healthy problems occurred due to the high content of cholesterol in food (Katan *et al.*, 1984, Pantulu *et al.*, 1975 and Younes *et al.*, 1986).

Furthermore, the addition of 75% coconut oil to the cream, reduced the cholesterol content of the produced cream to be 67 mg/100 g (Katan *et al.*, 1984).

**Table (1): Fat constants and cholesterol content of buffaloe's Cream mixed with 25, 50 and 75% Coconut oil:**

Fatty acids	Cholesterol	RM	PN	IV	SN
Cream	210	27	3	37.9	230
25% Coconut oil	162	22	6.38	30.6	236.8
50% Coconut oil	115	17	9.75	23.45	243.5
75% Coconut oil	67	12	13.13	16.23	250.3

RM = Reichert-Meissl number  
IV = Iodine value

PN = Polenske number  
SN = Saponification number

Table (2), reports the results of adding palm seed oil to buffaloes' cream and its effect on some constants of produced cream. From these data, the use of palm seed oil as a milkfat substitute reduced the RM and IV, while such addition increased the values of SN and PN of the produced cream. These results may be attributed to the lower RM and IV and higher PN and SN in the substitute oil. However, the addition of 25% palm seed oil led to obtain a cream with normal values of RM, PN, IV and SN similar to those in the natural butter (Williams, 1950b and Bailey, 1964).

**Table (2): Fat constants and cholesterol content of buffaloe's Cream mixed with 25, 50 and 75% Palm seed oil:**

Fatty acids	Cholesterol	RM	PN	IV	SN
Cream	210	27	3	37.9	230
25% Palm oil	154	21.75	2.44	33.18	235
50% Palm oil	97	16.5	1.88	28.45	240
75% Palm oil	52	11.25	1.31	23.73	245

RM = Reichert-Meissl number,  
IV = Iodine value,

PN = Polenske number  
SN = Saponification number

On the other hand, it was also reported that the 50% and 75% added palm seed oil did not keep the RM and PN within the normal range.

From the same table it was reported that the addition of palm seed oil to the cream at the ratios 25%, 50% and 75% led to a decrease in the cholesterol content of the produced cream. The addition of 25% palm seed oil decreased the cholesterol content to 154 mg/100 g, 50% to 97 mg/100 g and 75% to 52 mg/100 g (Sebastian and Rao, 1974).

From the above results, it could be concluded that the effect of coconut oil addition to the cholesterol content was less than the palm seed oils, due to the unsaturated fatty acids content of the coconut oil which was higher than the unsaturated fatty acids content of the palm seed oil (Younes and Soliman, 1987).

It could also be concluded that the best treatment concerning the cream constants using coconut oil at the rate of 25% and 50% substitution, except the RM value with 50% substitution was failed, while the use of palm seed oil in the substitution of cream in the ratios 50% and 75%, failed in

keeping the RM and IV within the normal range of the natural cream SN value in spite of the decreasing of the cholesterol content significantly. Also, the addition of 75% resulted in lower RN value than the minimum limit of the natural cream.

**B) Effect of add coconut oil on the fatty acid composition of buffaloes' cream:**

Fatty acid composition of vegetable oils plays a great role in the fat characteristics is ties and its properties.

The vegetable oil characterized with the high content of short chain fatty acids, which the other fats are missing these components (Hilditch, 1949). The use of vegetable oil such as coconut oil as a milkfat substitute is reported in (Table 3). Coconut oil was mixed with buffaloes' cream. The ratios of added coconut oil were 25%, 50% and 75%. As shown in the table, coconut oil contains high short chain fatty acids C4-C12, while it is characterized with a low prportion of C16, C18 and C18:1. The latter acid was the predominant among the saturated and unsaturated fatty acids. The total unsaturated fatty acids of the coconut oil accounted 8.35% of the total fat. On the other hand, the fatty acid composition of the buffaloes' cream is characterized with the short chain acids which represented 12.32%, of total fat, and the unsaturated acids were 31.02% which means that coconut oil was softer than the milkfat (Gulyaev-Zaitsev and Tverdokhle, 1965).

However, the addition of coconut oil in the previous percentages to the buffaloes' cream led to an increase in the short chain fatty acids C4-C12 which was 24.71%, 37.67% and 50.75%, respectively, (Bailey, 1964), while the total short chain of the control cream was 12.32%.

Furthermore, the saturated fatty acid C14 increased from 9.60% in the control to 14.88% when 75% of coconut oil was added. The saturated fatty acids C16 and C18 slightly decreased. The total saturated fatty acids was gradually increased when 25%, 50% and 75% coconut oil were added. On the other hand, the unsaturated fatty acid C18:1, decreased as the added coconut oil increased (Echizen and Deki, 1975). The other unsaturated fatty acids C14:1, C14:2, C16:1 and C18:2 decreased in their values. Gnerally, according to the fatty acid content of the produced cream, specially the saturated acids, it was observed that the addition of coconut oil improved the quality of cream concerning the saturated/unsaturated fatty acids ratio (Younes and Soliman, 1986).

The treatment with palm seed oil was reported in Table (4). The buffaloe's cream was mixed with palm seed oil in the ratios of 25%, 50% and 75%. The effect of adding palm seed oil to the cream on the fatty acid composition of the cream admixture is illustrated in the same table. From these data, it was found that the fatty acid composition of palm seed oil greatly differs from the fatty acid composition of the buffaloe's cream. The short chain fatty acids of palm seed oil accounted 57.25% of the total fat (Inada, 1976). While the total unsaturated fatty acids was 16%. The addition of palm seed oil at the ratios 25%, 50% and 75% led to an increase in the total short chain fatty acids of the produced cream to 23.13%, 33.73% and 44.88%, respectively. These values of the short chain fatty acids was an

indication of the addition of substitute fat has a very high content of short chain, fatty acids, specially the latter values, (Carisano and Riva,1976).

**Table (3): Fatty acid composition of Cream and mixed cream with 25, 50% and 75% coconut oil:**

No. of C. atoms	Cream	C oil	25% C oil	50% C oil	75% C oil
4	3.5	-	2.45	1.52	0.88
6	1.5	0.5	1.15	0.9	0.75
8	1	7	2.23	4	5.5
10	3	8	4.25	5.5	6.75
12	3.5	48	14.63	25.75	36.88
14	11.5	16	12.63	13.75	14.88
16	27.5	9.5	23.38	18.35	14
18	11.5	2.4	9.23	6.95	4.68
20	2	0.2	1.8	1.3	0.80
10:01	-	-	-	-	0
12:01	-	-	-	-	0
14:01	2	-	1.7	1.2	0.6
16:01	3	0.5	2.38	1.9	1.13
18:01	27	6.5	21.88	16.55	11.43
18:02	2	1.25	1.81	1.63	1.44
18:03	1	0.1	0.78	0.55	0.33
Short chain	12.5	63.5	24.71	37.67	50.75
Total USA	35	8.35	28.34	21.68	15.01
Sat : Unsat	1.86	10.97	2.52	3.61	5.66

C oil = Coconut oil

Furthermore, the value of the saturated fatty acids C16 and C18 acids were greatly affected by increasing the palm seed oil ratio. They decreased gradually, while C14 acid was slightly affected. However, the total unsaturated acids decreased, being 30.25%, 25.50% and 20.75% regarding the 25%, 50% and 75% added palm seed oil (Soliman *et al.*, 1979). The results indicated that the adding of sunflower oil in the ratios 25% led to an improvement in the consistency of the produced cream, due to the high percentages of C12 when compared to the control cream, while the addition of 50% and 75% palm seed oil led to a production of cream expected to be softer than the control butter due to the high content of short chain fatty acids in spite of the improvement of the cream consistency, the fatty acid profile of the produced cream indicated a great evidence of the vegetable oil existence.

Furthermore the saturated:unsaturated acids of the produced cream by adding 25%, 50% and 75% palm seed oil was greatly changed than the saturated:unsaturated acids of the control cream, which emphasize the treatment with a vegetable oil (Hewela *et al.*, 1978). So, it must refer to such addition when such product deals with the market.

**Table (4): Fatty acid composition of cream and mixed cream with 25, 50 and 75% Palm seed oil:**

No. of C. atoms	Cream	P oil	25% P oil	50% P oil	75% P oil
4	3.5	-	2.33	1.71	0.85
6	1.5	0.75	1.24	1	0.93
8	1	3	1.41	2	2.5
10	3	4	3.25	3.21	3.75
12	3.5	49.5	14.9	25.8	38
14	11.5	16.5	12.6	14	15.25
16	27.5	8	23	18	12.68
18	11.5	3	9.58	7.35	5.13
20	2	-	1.3	1.1	0.7
10:01	-	-	-	-	-
12:01	-	-	-	-	-
14:01	2	-	1.7	0.80	0.5
16:01	3	-	2.05	1.5	0.75
18:01	27	14	23.75	20.5	17.25
18:02	2	2	1.95	2.1	1.9
18:03	1	-	0.8	0.6	0.35
Short chain	12.50	57.25	23.13	33.73	44.88
Total USA	35	16	30.25	25.5	20.75
Sat : Unsat	1.86	5.30	2.30	2.9	3.79

**C) Effect of addition of coconut and palm seed oils on the organoleptic properties of buffaloes' cream:**

For the completion of the present study organoleptic tests were undertaken to determine the real and final acceptability of the product by consumer. Results of most food researches lie in the human palate and from these the decision on success or failure is determined. The organoleptic tests were thought of almost importance to clear the final decision. Judging and scoring were carried out by a group of eminent dairy technology who examined with great care all the different organoleptic characteristics namely appearance, body & texture and Flavour of the resultant cream.

**Table (5): Organoleptic properties of buffalo's Cream mixed with Coconut and Palm seed oils:**

Properties	Control	25% C oil	25% P oil	50% C oil	50% P oil	75% C oil	75% P oil
Appearance (15)	14	13	12	12	10	9	7
Body & texture (30)	28	26	25	24	22	20	15
Flavour (55)	53	50	48	45	43	40	35
Total (100)	95	89	85	81	75	69	57

C oil = Coconut oil, P oil = Palm seed oil

The organoleptic properties of the examined cream were presented in Table (5). It could be observed that the addition of coconut oil to cream at the levels 25 and 50% (W/W) gained the highest score points (after the control score points) namely 89 and 81, respectively. The corresponding score points

of cream mad from cream mixed with palm seed oil were 85 and 75 at the same two the added levels respectively. When the addition coconut and palm seed oil to cream at the level 75% (W/W) were 69 and 57, respectively.

In conclusion, cream could be successfully produced by mixing coconut and palm seed oils with traditional cream at levels 25 and 50%.

## REFERENCES

- A.O.A.C. (1970). Official methods of analysis of the association of the official analytical chemists, 11th ed. Published by A.O.A.C. Washington, D.C.
- A.O.A.C. (1975). Official methods of analysis of the association of the official analytical chemists, 12th ed. Published by A.O.A.C., Washington, D.C.
- Bailey's (1964) Industrial oil and fat products 3rd. ed. Wiley, New York, London, Sidney, p. 168.
- Bergman, T. and Joost, K. (1953). The iodine value of milk fat during different parts of the year, and its influence on the quality of the butter. *Int. Dairy Congr.* 2: 89.
- Bindal, M.P. and Jain, M.K. (1973). Studies on the cholesterol content of cow and buffalo ghee. *Indian J. Anim. Sci.* 43 (10): 918-24.
- Bird, Emerson. W. (1964). Dairy chemistry. Iowa State University, Ames. Iowa.
- Braut, K. (1990). Butter quality in 1989. *Meieriposten* 79 (12) 352-354 .C.F. Dairy Sci. Abst. 53 (4) 3551 (1991).
- Carisano, A. and Riva, M. (1976). Chromatographic method for detecting adulteration of butter. *Rivista Italiana delle sestance grasse* 53 (11) 297-300 .C.F. Dairy Sci. Abst.; 39 (6): 3438 (1977).
- Echizen, A. and Deki, M. (1975). Determination of butterfat in mixed fats. III. Relation between butyric acid and butterfat content in mixed fats. Reports of the central customs laboratory No. 15, 111-114 .C.F. Dairy Sci. Abst. 41 (7) 4056 (1979).
- Gulyaev-Zaitsev, S. and Tverdokhleb, G. (1965). Chemical composition of butterfat and characteristics of its solidification and melting. *Moloch. Prom.* 26 (2) 7-11 C.F. Dairy Sci. Abst. 28 (3) 993 (1966).
- Helal, F.R.; Rifaat, I.D. and El-Sadek, G.M. (1976): Chemical structure of buffalo milk fat and hardness of butter. *Egyptian J. Dairy Sci.* 4 (2) 128-132.
- Hewella, Y.; Difrawi, E.A.; Shehata, A.A.Y.; Mohamed, M.S. and Ismail, A.A. (1978). Chemical and physical properties of some local and foreign butter oil substitutes. *Alexandria Journal of Agric. Research* 26: 355-361.
- Hilditch, T.B. (1949). The chemical constitution of natural fats. Capman and Hall Ltd., 37 Essex Street, W.C. 2, London Second Edition.
- Inada, T. (1976). Determination of butyric acid in mixed butterfats. Reports of the central customs laboratory No.16, 67-70. C.F. Dairy Sci. Abst. 41 (7) 4057 (1979).
- Katan. M.B.; Bovenkamp, P.V. and Breussaard, J.H. (1984). Fatty acid composition, trans-fatty acid and cholesterol contents of margarines and other edible oils. *Voading*, 45 (4): 27.
- Ling, E.R. (1963). A Text book of dairy chemistry, Vol. 2, Practical 3rd ed. Chapman and Hall, Limited London.

- Ortin, L. and Sanchez-Algaba, T., (1990). Mean composition of milk and products. 10 years of analytical control: 1977-1987. VI. Butter and cream. *Alimentaria*, 27 (217) 25-27.
- Pantulu, P.C.; M.B. Rao; S.D. Rao and C.P. Anantkrishman (1975). Application of liebermann-Burchard reaction to the unsaponifiable portion of milk lipids for the determination of cholesterol in milk and milk products. *Milchissenschaft* 30 (12): 735-738.
- Sebastian, J. and Rao, M.B. (1974). Detection of the presence of vegetable oils and fats in ghee butter fat. *Milchwissenschaft* 29 (9) 533-534. C.F. Dairy Sci. Abst. 37 (6) 3732 (1975).
- Shahin, Y. (1977). Contents of row milk free fatty acids in respect of production factories. Thesis, Justus Liebig. Univ. Giessen, W. Germany.
- Soliman, M.A.; Mohamed, A.A.; Hagrass, A.E. and El-Shabrawy, S.A. (1979). Fatty acid composition of buffaloe's milk fat. *Egyptian J. Dairy Sci.*, 7: 177.
- Williams, K.A. (1950a). Oils, fats and fatty foods. Their practical examination. J. and A. Churchill Ltd., 104, Gloucester place, W.I. London, Third Edition, p. 195.
- Williams, K.A. (1950b). Oils, fats and fatty foods. Their practical examination. J. and A. Churchill Ltd., 104, Gloucester place, W.I. London, Third Edition, p. 204.
- Younes, N.A. and Soliman, M.A. (1986). Adulteration of butterfat with other sources of fat. *Grasas Y Aceites*, 37 (4) 200-203.
- Younes, N.A. and Soliman, M.A. (1987). Adulteration of butterfat: sterol composition. *Grasas Y Aceites*, 38 (6) 372-374.

دراسة على إنتاج قشدة منخفضة الكولسترول  
علاء الدين أحمد الحديدي وممدوح أحمد محمد عمر، ومنصور عبده علي  
قسم الألبان - كلية الزراعة - جامعة الأزهر - مصر

تتميز الزيوت النباتية مثل زيت جوز الهند وزيت نوى النخيل بخلوها من الكولسترول وارتفاع محتواها من الأحماض الدهنية قصيرة السلسلة الكربونية مما يساعد عند خلطها بالقشدة الطبيعية الجاموسي في إنتاج قشدة منخفضة الكولسترول.

وقد اعتمدت هذه الدراسة على استبدال دهن القشدة بنسب مختلفة من نوعي الزيت ودراسة تأثيرها على ثوابت دهن اللبن وكذلك تركيب الأحماض الدهنية للقشدة ونسبة الكولسترول في القشدة الناتجة. حيث كانت نسب الاستبدال ٢٥، ٥٠، ٧٥% وزن لوزن وكانت النتائج كما يلي:

نجحت إضافة نوعي الزيت بنسبة ٢٥ و ٥٠% في خفض محتوى القشدة من الكولسترول من ٢١٠مجم/١٠٠جم إلى ١٦٢ و ١١٥مجم/١٠٠جم على التوالي بالنسبة للقشدة الناتجة من خلط زيت جوز الهند بينما انخفضت إلى ١٥٤ و ٩٧ لنسبتي خلط زيت نوى النخيل. دون التأثير على ثوابت دهن اللبن أو محتواه من الأحماض الدهنية.

أما عند الاستبدال بنسبة ٧٥% فقد أدى إلى خفض الكولسترول إلى ٦٧مجم/١٠٠جم عند إضافة زيت جوز الهند وإلى ٥٢مجم في حالة زيت نوى النخيل ولكن كان هناك تأثير واضح على ثوابت دهن اللبن والمحتوى من الأحماض الدهنية.

وأظهرت درجات التحكيم أن أقرب العينات إلى الكونترول كانت نسبة استبدال ٢٥% من زيت جوز الهند تليها نفس النسبة لزيت نوى النخيل ثم نسبة ٥٠% بنفس الترتيب لنوعي الزيت بينما انخفضت درجات التحكيم لنسبة استبدال ٧٥%.