

INTEGRATED CONTROL OF THE RENIFORM NEMATODE, *Rotylenchulus reniformis* ON SUNFLOWER.

Khalil, A.E.M.; S.M. Abdel-Momen and Hanaa S. Zawam

Plant Pathology Research Institute, Agricultural Research Centre, Giza,
Egypt.

ABSTRACT

A greenhouse experiment was conducted in order to determine the impact of *Datura stramonium*, *Artemisia herba*, *Morus nigra*, *Salix babylonica* and *Eucalyptus camaldulensis* powders singly or integrated with oxamyl on reproduction of *R. reniformis* on sunflower cv. Vidoc-5 and the resulting effect on plant growth. Results indicated that all treatments improved sunflower growth, exceeded that of the nematode alone. The application of *D. stramonium* integrated with oxamyl, accomplished the highest values regarding the percentage of increase of the whole sunflower plant fresh weight which was amounted to 62.24 %, followed by *D. stramonium* alone (45.37 %), *A. herba* plus oxamyl (44.68 %) and *E. camaldulensis* plus oxamyl (39.51 %). Nematode population density in soil and within sunflower roots infected with *R. reniformis* were significantly suppressed in all treatments. Great suppression in nematode population was recorded with oxamyl giving reduction of 89.72 % followed by *D. stramonium* integrated with oxamyl (88.54 %) and then *E. camaldulensis* plus oxamyl (86.96 %). Egg-mass numbers of *R. reniformis* on sunflower roots were significantly suppressed in all treatments compared to plants inoculated with nematode alone.

Keywords: sunflower, *Helianthus annuus*, *Rotylenchulus reniformis*, oxamyl, plant products, *Datura stramonium*, *Artemisia herba*, *Morus nigra*, *Salix babylonica* and *Eucalyptus camaldulensis*.

INTRODUCTION

Sunflower (*Helianthus annuus*, L.) is considered one of the most important economic cash crops in Egypt. It is used as source of edible oil. The reniform nematode, *Rotylenchulus reniformis* (Linford & Oliveira) is a serious pest on agricultural crops, mainly, sunflower in temperate and tropical regions results in yellowish leaves, stunting, poor plant growth and yield losses.

The chemical control of *R. reniformis* has successfully limited the detrimental effect of this nematode below damaging levels (Reddy & Khan, 1988; Lawrence *et al.*, 1990 and Sipes *et al.*, 1992). However, the environmental hazards, health risks and disturbance of the biological balance of nature are restrictions for the absolute use of nematicides. Also, the development of nematicide – resistant races of the target pest and the high cost are another restrictions for using nematicides. The previous restrictions enhanced scientists to find other alternative tactics to replace or minimize the use of nematicides.

Nowadays, integrated pest management using several control techniques with minimal use of nematicides received greater interest among nematologists, providing effective control against the target nematode and avoiding environmental pollution (Shahzad & Ghaffar, 1987; Reddy & Khan, 1988; Zaki & Bhatti, 1990; Vicente *et al.*, 1991; Mahmood & Siddiqui, 1993; Khan & Goswami, 1995; Oduor-Owino & Waudu, 1995; Khan & Saxena, 1997;

Nagesh *et al.*, 1997,1998; Rao *et al.*, 1997 a, b, 1998; Anjun-Ahmed *et al.*, 1998; Walia *et al.*, 1999; Almihanna *et al.*, 1999; Farahat *et al.*, 1999 and Saikia *et al.*, 1999).

On the other hand, integrated control of nematodes using combination of nematicide and organic amendments have been studied by many workers with respect to root-knot nematodes (Bhattacharya & Goswami, 1988; Jain & Bhatti, 1988; Oduor-Owino *et al.*, 1996 and Amin & Youssef, 1997), however, little attention has been given to the combined effect of nematicide and organic amendments on *R. reniformis* (Gaur & Mishra, 1989 and Prasad *et al.*, 1997). In Egypt, El-Sherif *et al.* (2001) reported that *Vinca rosea* powder integrated with oxamyl caused significant and better increment in sunflower growth, head diameter, seed weight, seed yield/plant and seed oil percentage as well as significant reduction in number of the reniform nematode in soil and roots under greenhouse and field microplots conditions.

The present investigation was conducted to study the impact of *Datura stramonium*, *Artemisia herba*, *Morus nigra*, *Salix babylonica* and *Eucalyptus camaldulensis* powders singly or combined with oxamyl on reproduction of *R. reniformis* infecting sunflower and the resulting effect on plant growth under greenhouse conditions.

MATERIALS AND METHODS

Fresh leaves of thorn apple, *Datura stramonium*; santonica, *Artemisia herba*; mulberry, *Morus nigra*; weeping willow, *Salix babylonica* and blue gum, *Eucalyptus camaldulensis* were sun-dried and powdered. Plastic pots 10- cm diam. were filled with 250 gm steam sterilized sandy : loam soil (1:1) and amended with particular dose of 0.50 gm powder of the previous mentioned plants, separately.

Oxamyl 10% G as nematicide was used alone at the recommended dose (0.03 gm) or in combination with powder at half recommended dose (0.015 gm). All components were added simultaneously, following the incorporation of powder alone or in combination with oxamyl into soil pots. Seeds of sunflower (*Helianthus annus*, L.) cv. Vidoc-5 were planted in pots following the addition of the six components (*Datura stramonium*, *Artemisia herba*, *Morus nigra*, *Salix babylonica*, *Eucalyptus camaldulensis* and oxamyl). Plants were thinned to one plant per pot after germination and inoculated with 2000 4th stage juveniles (immature females) of *R. reniformis*. Inocula were obtained from a pure culture of *R. reniformis* propagated on pigeon pea, *Cajanus* spp. in the greenhouse. Inoculated plants free of powders or oxamyl were served as control. Pots neither treated with nematode nor powders were also included.

Treatments were as follows:

- | | |
|---------------------------------------|--|
| 1- N + thorn apple, | 2- N + thorn apple + oxamyl (0.015 gm), |
| 3- N + santonica, | 4- N + santonica + oxamyl (0.015 gm), |
| 5- N + mulberry, | 6- N + mulberry + oxamyl (0.015 gm), |
| 7- N + weeping willow, | 8- N + weeping willow + oxamyl (0.015 gm), |
| 9- N + blue gum, | 10- N + blue gum + oxamyl (0.015 gm), |
| 11- N + oxamyl (0.03 gm), | 12- N alone (control), and |
| 13- Plant free of N or any treatment. | |

Each treatment was replicated four times. Pots were randomly arranged on a greenhouse bench at 28 ± 5 °C. Plants received water as needed. Forty-five days after inoculation, plants were removed. Data dealing with lengths and weights of fresh shoot and root were recorded. Shoot dry weights were also measured. Vermiform stages of *R. reniformis* were extracted from soil by sieving and modified Baerman techniques (Goodey, 1957). Roots were stained by acid fuchsin in acetic acid according to Byrd *et al.* (1983), and examined for number of nematode stages, and egg-masses.

Data were also subjected to analysis of variance (ANOVA) (Gomez and Gomez, 1984) and means were compared by Duncan's multiple-range test (Duncan, 1955). This work was undertaken in the greenhouse of Tag El-ez, Research Station, Tag El-ez, Temai El-Amdeed, Dakahlia Governorate.

RESULTS AND DISCUSSION

The impact of plant powders of *D. stramonium*, *A. herba*, *M. nigra*, *S. babylonica* and *E. camaldulensis* with or without oxamyl on plant growth response of sunflower cv. Vidoc-5 infected with *R. reniformis* are presented in Table (1). It was evident that all treatments improved sunflower growth, exceeded that of the nematode alone as well as nematode-free plants.

Regarding data on the fresh weight of the whole plant, *D. stramonium* mixed with oxamyl accomplished the highest percentage of increase which was amounted to 62.24 %, followed by *D. stramonium* alone (45.37 %), *A. herba* plus oxamyl (44.38 %) and *E. camaldulensis* plus oxamyl (39.51 %), whereas, the least values of this plant growth parameter were achieved by the application of *M. nigra* (20.88 %), followed by *S. babylonica* plus oxamyl (20.00 %) and then *S. babylonica* alone (19.32 %) (Table 1). Similar results were noticed with shoot dry weight. However, concomitant treatments using powders mixed with oxamyl showed better increase of the whole plant percentages than those of any component alone. Moreover, significant differences for plant growth parameters were also noticed among most treatments as compared with nematode alone (Table 1).

Concerning data in Table (2), it was evident that nematode population density in soil and within sunflower roots infected with *R. reniformis* were significantly suppressed in all treatments with rates of nematode build-up ranged from 0.167 to 0.278 as compared to nematode alone (1.624). Oxamyl gave great suppression in nematode population with reduction of 89.72 % followed by *D. stramonium* mixed with oxamyl (88.54 %) and then *E. camaldulensis* plus oxamyl (86.96 %) (Table 2). With respect to the egg-mass number of *R. reniformis* on sunflower roots, a significant suppression in all treatments was detected as compared to plants inoculated with nematode alone. However, no significant differences were observed among other treatments (Table 2).

Table (1): Plant growth response of sunflower cv. Vidoc - 5 infected with *R. reniformis* as influenced by some organic soil amendments singly or integrated with oxamyl under greenhouse conditions.

Treatments	Plant growth response*									
	Length (cm)		Fresh weight (gm)		Fresh Wt. of the whole plant (gm)	Increase %	Shoot dry* weight (gm)	Increase* %		
	Shoot	Root	Shoot	Root						
<i>D. stramonium</i>	74.75 b	12.20 ab	7.90 bc	7.00 a	14.90	45.37	2.59 ab	181.52		
<i>D. stramonium</i> + oxamyl	78.30 a	12.90 a	9.55 a	7.08 a	16.63	62.24	2.75 a	198.91		
<i>A. herba</i>	72.50 bcd	12.00 ab	6.96 cde	6.65 a	13.61	32.78	2.20 cd	139.13		
<i>A. herba</i> + oxamyl	73.25 bc	11.55 bc	8.10 b	6.73 a	14.83	44.68	2.48 abc	169.56		
<i>M. nigra</i>	69.22 de	10.55 cd	6.45 e	5.94 bc	12.39	20.88	1.99 d	116.30		
<i>M. nigra</i> + oxamyl	70.32 cde	10.73 cd	7.04 cde	5.78 cd	12.82	25.07	2.02 d	119.56		
<i>S. babylonica</i>	68.93 e	10.52 cd	6.45 e	5.78 cd	12.23	19.32	1.98 d	115.22		
<i>S. babylonica</i> + oxamyl	69.65 de	10.57 cd	7.00 cde	5.30 d	12.30	20.00	2.00 d	117.39		
<i>E. camaldulensis</i>	71.70 bcde	11.48 bc	6.61 de	6.90 a	13.51	31.80	2.40 bc	160.87		
<i>E. camaldulensis</i> + oxamyl	73.00 bc	11.57 bc	7.55 bcd	6.75 a	14.30	39.51	2.43 abc	164.13		
Oxamyl	71.25 cde	11.98 ab	6.98 cde	6.43 ab	13.41	30.83	2.06 d	123.91		
Nematoda alone	54.63 f	9.73 d	5.06 f	5.19 d	10.25		0.92 e			
Plant free of nematode	71.88 bcde	12.55 ab	7.04 cde	6.85 a	13.89	35.51	2.15 cd	133.70		

* Each value presented the mean of four replicates.

Means in each column followed by the same letter(s) did not differ at < 0.50 according to Duncan's multiple-range test.

Table (2): Rate of build-up and egg-mass numbers of *R. reniformis* infecting sunflower cv. Vidoc-5 as affected by some organic soil amendments singly or integrated with oxamyl under greenhouse conditions.

Treatments	Nematode population in *				Total	Rate of build-up	Reduction %	Egg-masses *
	Soil	Develo-pm ental stage	Root					
			Females					
<i>D. stramonium</i>	395.00 cd	19.30 bc	15.73 b		430.00 cd	0.215	86.76	8.73 b
<i>D. stramonium</i> + oxamyl	347.50 de	12.43 cd	12.35 b		372.30 de	0.186	88.54	7.08 b
<i>A. herba</i>	407.50 c	18.42 bc	16.90 b		442.80 c	0.221	86.37	10.02 b
<i>A. herba</i> + oxamyl	397.50 cd	11.70 cd	15.18 b		424.40 cd	0.212	86.93	8.45 b
<i>M. nigra</i>	495.00 b	17.81 bc	17.84 b		530.60 b	0.265	83.66	10.34 b
<i>M. nigra</i> + oxamyl	480.00 b	16.25 bcd	15.82 b		512.10 b	0.256	84.23	8.52 b
<i>S. babylonica</i>	512.50 b	22.02 b	20.45 b		555.00 b	0.278	82.91	10.23 b
<i>S. babylonica</i> + oxamyl	492.50 b	18.46 bc	17.39 b		528.30 b	0.264	83.73	8.05 b
<i>E. camaldulensis</i>	420.00 c	18.25 bc	15.57 b		453.80 c	0.227	86.03	8.58 b
<i>E. camaldulensis</i> + oxamyl	395.00 cd	15.27 bcd	13.20 b		423.50 cd	0.212	86.96	6.75 b
Oxamyl	315.00 e	9.55 d	11.06 b		333.90 e	0.167	89.72	6.43 b
Nematode alone	2975.00 a	123.7 a	149.6 a		3248.00 a	1.624	0.00	60.30 a

* Each value presented the mean of four replicates.

Means in each column followed by the same letter(s) did not differ at < 0.50 according to Duncan's multiple-range test.

From the previous findings it can be concluded that all treatments tested showed better results in reducing nematode population and improving plant growth of sunflower. *Datura. stramonium* integrated with oxamyl showed better performance in promoting plant growth as well as reducing nematode population. These findings are in accordance with those reported by Gaur & Mishra (1989) and Prasad *et al.*, (1997) who stated that combination of chopped leaves of *Calotropis procera* and seed treatment with nematicides was effective in reducing *R. reniformis*. Results of El-Sherif *et al.* (2001) in respect to *Vinca rosea* powder integrated with oxamyl on *R. reniformis* infecting sunflower support also our findings.

In conclusion, it may be noted that powders of certain plants singly or integrated with oxamyl could possess antihelminthic properties against plant parasitic nematodes on such crops and could promise to be used in crop management practices.

REFERENCES

- Almihanna, A.A.; M. Belal and A.A. Farahat (1999). The addition of chicken manure and eucalyptus dry leaves in relation to the management of *Heterodera avenae* and the incidence of soil-borne fungi. J. Agric. Sci. Mansoura Univ., 24(8): 4221-4228.
- Amin, A.W. and M.M.A. Youssef (1997). Comparative study on the efficiency of dry and green manuring of certain plant leaves against *Meloidogyne incognita* and *Rotylenchulus reniformis* infecting sunflower and their effect on plant growth. Abstract Book Sixth Arab Congress of Plant Protection, October 27-31, Beirut, Lebanon.
- Anjum-Ahmad; M.M. Alam and A. Ahmad (1998). Effect of organic amendments alone and in combination with *Paecilomyces lilacinus* on *Rotylenchulus reniformis* attacking some economically important crop plants. Archives of Phytopathology and Plant Protection, 31(5): 439-448.
- Bhattacharya, D. and B.K. Goswami (1988). Effect of oil-cakes used alone and in combination with aldicarb on *Meloidogyne* infecting tomato. Nematologia Mediterranean, 16(1): 139-141.
- Byrd, D.W.; T. Kirkpatrick and K.R. Barker (1983). An improved technique for clearing and staining plant tissues for detection of nematodes. J. Nematol., 15 (1): 142-143.
- Duncan, D.B. (1955). Multiple range and multiple, F-test Biometrics, 11: 1-42.
- El-Sherif, A.G.; F.A. Mostafa and A.A. Khalil (2001). Impact of *Vinca rosea* powder and oxamyl on controlling, *Rotylenchulus reniformis* infected sunflower. (Abstract) in IV International Nematology Symposium, Moscow, Russia.
- Farahat, A.A.; A.A. Almihanna and M.M. Belal (1999). Effect of adding chicken manure and eucalyptus dry leaves powder on root galling and growth of sunflower. J. Agric. Sci., Mansoura Univ., 24(8): 4229-4236.
- Gaur, H.S. and S.D. Mishra (1989). Integrated control of nematodes in lentil with aldicarb, neem cake and seed treatment with rhimet and its residual effect on the subsequent mung crop. Indian J. Entomol., 51(3): 283-287.

- Gomez, K.A. and A.A. Gomez (1984). Statistical procedures for agricultural research. 2nd Ed., John Wiley & Sons. Inc. New York.
- Goodey, J.B. (1957). Laboratory methods for work with plant and soil nematodes. Tech. Bull. No. 2 Min. Agric. Fish Ed. London, pp. 47.
- Jain, R.K. and D.S. Bhatti (1988). Integrated control of root-knot nematode, *Meloidogyne javanica* infecting tomato. *Indian J. Nematol.*, 18(1): 30-34.
- Khan, M.R. and B.K. Goswami (1995). Nematicidal efficacy of decomposed neem cake and fungal filtrates of *Paecilomyces lilacinus* and *Aspergillus flavus* against *Meloidogyne incognita*. Proceeding national symposium on sustainable agriculture in sub humid zone, March, 3-5, 1995.
- Khan, T.A. and S.K. Saxena (1997). Integrated management of root knot nematode *Meloidogyne incognita* infecting tomato using organic materials and *Paecilomyces lilacinus*. *Bioresource Technology*, 61(3): 247-250.
- Lawrence, G.W.; K.S. McLean; W.E. Batson; D. Miller and J.C. Borbon (1990). Response of *Rotylenchulus reniformis* to nematicide applications on cotton. *Suppl. J. Nematol.*, 22: 707-711.
- Mahmood, I. and Z.A. Siddiqui (1993). Integrated management of *Rotylenchulus reniformis* by green manuring and *Paecilomyces lilacinus*. *Nematologia Mediterranean*, 21(2): 285-289.
- Nagesh, M.; P.P. Reddy; M.V.V. Kumar; B.M. Nagaraju; P.P. Reddy; N.K.K. Kumar and A. Verghese (1998). Management of *Meloidogyne incognita* on *Polianthes tuberosa* using parasitic fungus, *Paecilomyces lilacinus* in combination with oil cakes. Proceeding of the first National Symposium on Pest Management in Horticultural Crops: environmental implications and thrusts, Bangalore, India, 15-17 October 1998, 339-344.
- Nagesh, M.; P.P. Reddy and M.S. Rao (1997). Integrated management of *Meloidogyne incognita* on tuberose using *Paecilomyces lilacinus* in combination with plant extracts. *Nematologia Mediterranean*, 25(1): 3-7.
- Oduor-Owino, P.; R.A. Sikora; S.W. Waudu and R.P. Schuster (1996). Effects of aldicarb and mixed cropping with *Datura stramonium*, *Ricinus communis* and *Tagetes minuta* on the biological control and integrated management of *Paecilomyces lilacinus*. *Nematologica*, 42(1): 127-130.
- Oduor-Owino, P. and S.W. Waudu (1995). Effects of antagonistic plants and chicken manure on the biological control and fungal parasitism of root-knot nematode eggs in naturally infested field soil. *Pakistan J. Nematol.*, 13(2): 109-117.
- Prasad, D.; K.N. Dubey and A. Mittal (1997). An integrated approach for management of plant parasitic nematodes in groundnut (*Arachis hypogaea*, L.). *Ann. Plant Prot. Sci.*, 5(2): 217-219.
- Rao, M.S.; P.P. Reddy and M. Nagesh (1997 a). Integration of *Paecilomyces lilacinus* with neem leaf suspension for the management of root-knot nematodes on egg plant. *Nematologia Mediterranean*, 25(2): 249-252.
- Rao, M.S.; P.P. Reddy and M. Nagesh (1997 b). Management of root-knot nematode, *Meloidogyne incognita* on tomato by integration of *Trichoderma harzianum* with neem cake. *Zeitschrift fur Pflanzenkrankheiten and Pflanzenschutz*, 104(4): 423-425.

- Rao, M.S.; P.P. Reddy; M. Nagesh; P.P. Reddy; N.K.K. Kumar and A. Verghese (1998). Integrated management of *Meloidogyne incognita* on tomato using oil cakes and a bioagent, *verticillium chlamyosporium*. Proceedings of the First National Symposium on Pest Management in Horticultural Crops: environmental implications and thrusts, Bangalore, India, 15-17 October 1998, 345-348.
- Reddy, P.P. and R.M. Khan (1988). Evaluation of *Paecilomyces lilacinus* for the biological control of *Rotylenchulus reniformis* infecting tomato, compared with carbofuran. *Nematologia Mediterranean*, 16(1): 113-115.
- Saikia, M.K.; K.C. Puzari and U.N. Saikia (1990). An integrated approach to control root-knot nematode, *Meloidogyne incognita* on vegetables. *Annals of Biology Ludhiana*, 15(1): 81-84.
- Shahzad, S. and A. Ghaffar (1987). Field application of *Paecilomyces lilacinus* and Furadan for the control of root-knot disease of okra and mung. *Int. Nematol. Network Newsl.*, 4(1): 33-34.
- Sipes, B.S.; D.P. Schmitt and C. Oda (1992). Multiple nematicide applications for the control of *Rotylenchulus reniformis* on pineapple, *Ananas comosus*. *J. Nematol.*, 24(4): p. 618.
- Vicente, N.E.; L.A. Sanchez and N. Acosta (1991). Effect of granular nematicides and the fungus *Paecilomyces lilacinus* in nematode control in watermelon. *J. Agric. Puerto Rico Univ.*, 75(3): 307-309.
- Walia, R.K.; S.N. Nandal and D.S. Bhatti (1999). Nematicidal efficacy of plant leaves and *Paecilomyces lilacinus*, alone or in combination, in controlling *Meloidogyne incognita* on okra and tomato. *Nematologia Mediterranean*, 27(1): 3-8.
- Zaki, F.A. and D.S. Bhatti (1990). Effect of castor (*Ricinus communis*) and the biocontrol fungus *Paecilomyces lilacinus* on *Meloidogyne javanica*. *Nematologica*, 36(1): 114-122.

المكافحة المتكاملة للنيماتودا الكلوية "روتيلنكيولس رينوفورمس" على عباد الشمس
أشرف السعيد محمد خليل – صلاح محمد عبدالمؤمن – هناء سيدهم زوام
معهد بحوث أمراض النباتات – مركز البحوث الزراعية – الجيزة

تم دراسة تأثير إضافة مسحوق الأوراق الجافة لنباتات الداتورة والشيح والتوت الأسود والصفصاف المتهدل والكافور مع أوبدون مبيد الأوكساميل على تعداد النيماتودا الكلوية ونمو نباتات عباد الشمس صنف فيدوك - ٥ تحت ظروف الصوبة ولوضحت النتائج ما يلي:-
- سجلت جميع المعاملات المختبرة تحسنا ملحوظا في النمو الكلي لنباتات عباد الشمس مقارنة بالكنترول (النيماتودا فقط).
- حققت معاملة الداتورة سواء كانت مع أو بدون المبيد عند نصف الجرعة (٠,٠١٥ جرام/نبات) أعلى نسبة زيادة في الوزن الكلي للنبات مقارنة بالكنترول يليها معاملة الشيح والمبيد معا ثم معاملة الكافور مع المبيد وذلك بنسبة زيادة تصل إلى ٦٢,٢٤% ، ٤٥,٣٧% ، ٤٤,٦٨% ، ٣٩,٥١% على التوالي.
- كان لجميع المعاملات المختبرة تأثير واضح في خفض أعداد النيماتودا معنوياً مقارنة بالكنترول.
- سجلت معاملة المبيد فقط أعلى نسبة انخفاض في تعداد النيماتودا (٨٩,٧٢%) يليها معاملة الداتورة والمبيد معا (٨٨,٥٤%) ثم معاملة الكافور والمبيد معا (٨٦,٩٦%).
- حققت جميع المعاملات المختبرة إنخفاضا معنوياً في تعداد كتل بيض النيماتودا الكلوية مقارنة بالكنترول.