

EFFECT OF SOME AMELIORATION PROCESSES ON NUTRIENTS AVAILABILITY AND UPTAKE IN SALT AFFECTED SOILS

Gazia, E.A.E.

Soil, Water and Environment Research Institute - ARC

ABSTRACT

Salt affected soils in Egypt are located, mostly, at North Delta. Reclamation processes of such soils are commonly using chemical amendments as gypsum (G) or phosphogypsum (PG) and/or organic amendments as farmyard manure (FYM), in addition to deep ploughing using sub-soiler. The current study was carried out for two seasons (2000 and 2000/2001) to study the effect of some amelioration processes on macro and micronutrients availability, elemental uptake and yield productivity of salt affected soil at North Delta. Amelioration treatments (FYM and/or PG application) have a significant effect on increasing the available N, P, and K contents in the sequent layers of soil profile especially in the surface layers. Also, amelioration treatments resulted in a significant increase in the available Fe, Mn, Zn and Cu contents in the surface layers.

Moreover, amelioration processes have a highly significant effect on sorghum and barley yields. PG has a highly significant effect on the yield of both crops. These results may be attributed to that PG affected the soil properties such as porosity, ESP, pH and nutrients availability, which enhance plant growth. Regarding subsoiling and FYM, it could be noticed that subsoiling and FYM as well have a highly significant effect on the yield of both crops.


In addition, the uptake of the determined macronutrients (N, P, K) and micronutrients (Fe, Mn, Zn and Cu) were increased, while Na was decreased under the different amelioration processes.

INTRODUCTION

Salt affected soils are scattered all over the world, especially in arid and semi-arid regions like Egypt. It is located, mostly, in Egypt at North Delta. Soil salinity and alkalinity affect soil properties and availability of plant nutrients and hence soil productivity. Reclamation processes of such soils are commonly using chemicals (as gypsum or phosphogypsum) and/or organic amendments (as farmyard manure), in addition to deep ploughing using sub-soiler.

Gypsum ($\text{CaSO}_4 \cdot x\text{H}_2\text{O}$) is used in the agriculture either as sources of Ca and S for crops or as soil conditioners to improve certain physical properties of problem soils. They are available as mined gypsum or as industrial by-products; such as: phosphogypsum (PG). They, likewise, can be used as source of Ca and S for crops, soil ameliorants for sodic dispersed soils, soil conditioners for hard setting clay soils and hardpans, bulk carriers for micronutrients (Alcorido, 1993) and/or in modifying Ca ratios in soil (Alva and Gascho, 1991) and in reducing $\text{NH}_3\text{-H}$ losses from urea fertilizers and farm manure (Da Gloria et al. 1991, C.F. Rechcigl 1995). PG is produced

during phosphoric acid production from rock phosphate according to the following reaction:



Many countries produce PG, e. g. USA, Russia, Canada, etc. (Novikov *et al.* 1990 and Collings, 1980). PG is highly acidic with pH in water ranging from 2.0 to 5.0 while mined gypsum ranged from 6.7 to 7.4. PG solubility is 2.6 g/l in water while that of mined gypsum is 2.41 g/l in water (Weast, 1981).

The availability of macro and micro-nutrients is affected by amendments application so that the highest nitrogen content of the soil was achieved with gypsum application combined with sandy mole, while the highest phosphorus content was detected with gypsum application combined with sub-soiling. (Shams El-Din *et al.*, 2000). The availability of micro-nutrients (e.g. Zn, Mn, Fe, Cu, ...etc.) is increased with increasing the amounts of applied gypsum (Abd El-Fattah *et al.*, 1987; Mahmoud *et al.*, 1996; and Youssef, 1992). Macro and micro-nutrients uptake, such as N, P, K, Ca, Mg, Zn, Mn and Fe as well as Na is increased by increasing rates of gypsum application, (El-Fakharani, 1997; Sonbol *et al.*, 2001; Gazia, 2001; Genaidy *et al.*, 1989 and Ismail *et al.*, 1991). Moreover, it is stated that gypsum addition leads to an increase in the yield of many crops such as rice and wheat (Abrol and Bhumbra, 1975; and Gazia *et al.*, 1996), wheat and berseem (Hussain *et al.*, 1988), bean and barley (Ghowail *et al.* 1978) and wheat and broad bean (Dora, 1996).

On the other hand, the availability of macro and micro-nutrients (e.g. N, P, K, Fe or Zn ...etc) in the soil is affected by organic manure application (Mahmoud, 1994; Nadia *et al.*, 2000; El-Gala *et al.*, 1998; Gazia, 2001; Rechcigl, 1995; and Swarup, 1982). Moreover, it is stated that organic manure application leads to an increase in the yield of many crops such as rice (Thakur *et al.*, 1995), maize (Hamoud, 1992), berseem, sorgum and barley, (Ghazy, 1994), barley (Abdel Karim, 1989), and (El-Sherief, 1997) and wheat (El-Koumey, 1998). Consequently, organic matter application increased macro and micro-nutrients uptake (e.g. N (Kaloosh *et al.*, 1989), P (Nafady *et al.*, 1993) and K, Fe, Cu and Zn (El-Koumey, 1998).

The current study was carried out to study the effect of some amelioration processes on macro and microelements availability, elemental uptake and yield productivity of salt affected soil at North Delta.

MATERIALS AND METHODS

A field experiment was conducted at Sakha Agricultural Research Station Farm, Kafr El-Sheikh, Egypt for two successive seasons; summer of 2000, where sorgum was cultivated and winter of 2000/2001, where barley was cultivated to study the effect of some amelioration processes on some soil properties, some elements availability, and productivity of salt affected

soil at North Delta. Area of 2100 m² of deteriorated bared soil was chosen to implement these processes. A split-split plot design with four replicates. Plot area was 24m². The main plots were occupied by phosphogypsum (PG) treatment with three levels namely: PG₀, PG₁ and PG₂ to represent zero, 50% and 100% of correspondence gypsum requirements (10 ton/fed, 80% purity). The sub plots were occupied by subsoiling treatment namely: without subsoiling (S₀) and with subsoiling (S₁) (with 2-m spacing and 50cm depth). Sub-sub plots were assigned to farmyard manure (FYM) treatment namely: FYM₀ and FYM₁ to represent without and with 20m³ FYM /fed application respectively. Soil texture was clayey with 46.92% clay and 28.94% silt in average. Chemical properties of the experimental soil and its content of macro and micronutrients are given in Tables' (1and 2). Chemical composition of FYM is given in Table (3). Chemical analyses were done according to Jackson, 1973. Micronutrients were determined according to Cotteine *et al.* (1982a). Statistical analyses were done according to Cochran and Cox, (1960).

Table (1): Some chemical properties of the experimental soil.

Soil depth cm	PH 1:2.5S.W. sus.	EC dS/m (soil paste)	Soluble ions (meq/l)							O.M. %	CaC ₃ %	CEC meq/100g soil	SAR	ESP
			Cations				Anions							
			Ca	Mg	Na	K	Cl	HCO ₃	SO ₄					
0-15	8.87	17.3	22.8	21.6	152	3.7	153.8	2.1	44.2	1.4	3.6	33.8	32.4	31.7
15-30	8.85	15.2	20.2	14.9	116	1.3	116.3	2.0	34.1	1.2	3.2	41.6	27.7	28.4

Table (2): Macro and micronutrients content in soil (ppm).

Soil depth cm	N		P		K		Fe		Mn		Zn		Cu	
	Total	Available	Total	Available	Total	Available	Total	Available	Total	Available	Total	Available	Total	Available
0-15	160.4	8.9	25.3	1.3	8240	864	3214	10.5	621	2.5	52	0.8	51	.95
15-30	153.0	8.1	19.5	0.9	7560	845	3977	7.0	447	1.8	84	0.4	45	0.8

Table (3) chemical composition of FYM.

Properties		Values
PH	(1:2.5)	6.30
O.M.	%	33.02
O.C.	%	19.12
C:N	ratio	13.2
Macronutrients		
Total N	%	1.45
Total P	%	1.60
Total K	%	2.00
Micronutrients		Total Available
Fe	ppm	6608 158.1
Mn	ppm	3561.3 519.9
Zn	ppm	33.37 5.55
Cu	ppm	11.60 1.00

RESULTS AND DISCUSION

A - Effect of a melioration processes on the availability of macro and micronutrients.

1 - Nitrogen (N):

Application of PG and/or FYM have a significant effect on increasing the available N content in the sequent layers of soil profile especially in the

surface layer as shown in Table (4). But subsoiling has no significant effect on N availability. This result may be due to the improvement in physical and biological conditions that led to activation of microorganisms, consequently enhanced organic nitrogen mineralization and nitrogen fixation. This result is in agreement with Mohammed *et al.*, (1993), Nadia *et al.*, (2000) and Shams El-Din *et al.* (2000).

Table (4): Effect of amelioration processes on some macronutrients availability in soils.

Treatment	N ppm		P ppm		K ppm	
	0.0 – 15 Cm	15 – 30 cm	0.0 – 15 cm	15 – 30 cm	0.0 – 15 cm	15 – 30 Cm
Phosphogypsum(PG):						
PG ₀	15.662	15.516	2.71	1.68	1418.9	1003.4
PG ₁	17.908	15.780	2.91	2.02	1751.6	1269.3
PG ₂	21.204	16.650	2.66	1.78	1889.9	1342.3
F-test	*	ns	ns	ns	**	Ns
LSD.05	4.446	-	-	-	234.25	-
0.01	-	-	-	-	354.90	-
Subsoiling (S):						
S ₀	15.931	16.18	1.19	0.75	1510.6	1203.1
S ₁	16.585	15.77	4.33	2.91	1862.9	1207.0
F-test	ns	ns	**	**	**	Ns
Farmyard manure (FYM):						
FYM ₀	17.285	14.53	2.35	1.51	1662.9	1188.5
FYM ₁	19.231	1744	3.166	2.147	1686.5	1221.6
F-test	*	*	**	*	*	Ns
Interactions:						
PGxS	*	ns	*	*	*	Ns
PGxFYM	*	ns	*	*	*	**
SxFYM	Ns	ns	*	*	ns	Ns
PGxSxFYM	ns	ns	ns	ns	ns	Ns

2 – Phosphorus (P):

Data in Table (4) show that the available P in the plots treated with both FYM and subsoiling was greater than those found in the control plots. This may be attributed to that the phosphorus released from the added organic manure and also to the increase in the biological activity (Dora, 1996 and Nadia *et al.* 2000). On the other hand, addition of PG especially PG₂ treatment decreased the available P. this may be due to an enhancement of P fixation (Chhabra *et al.* 1981). Similar results were obtained by Swarup, (1982) And Dora, (1996). In addition, available P decreased with the soil depth. This result may be attributed to decreasing of organic matter with the depth.

3 - Potassium (K):

Data in Table (4) show that PG application and subsoiling treatment as well as FYM application resulted in an increase in available K. The highest content of available K was recorded under the combination treatment of PG₂S₁FYM₁, while the lowest one was obtained by the control treatment (PG₀S₀FYM₀). Moreover, data show that the available K decreased with soil depth. These results are in agreement with those obtained by Mahmoud *et al.* (1996), Nadia *et al.* (2000) and Shams El-Din *et al.* (2000).

4 – Iron (Fe):

Data in Table (5) show that all amelioration treatments resulted in a significant increase in the available Fe at the surface layer, but the increase was not significant in the subsurface layer except with *FYM*, which was highly significant. This may be attributed to the beneficial effect of *FYM* on soil physical and chemical properties that make more suitable conditions for Fe availability. These results are in harmony with those obtained by Dora (1996) and Shams El-Din *et al.* (2000). Fe availability was distinctly affected by phosphogypsum application. This result may be due to the negative effect of phosphogypsum on the soil *pH*. This result is in agreement with that of Sharma and Yadov (1989) and Shams El-Din *et al.* (2000).

5 – Manganese (Mn):

Data in Table (5) show that all amelioration treatments resulted in a significant increase in the available Mn in the surface layer (0 – 15 cm) and subsurface layer (15 – 30 cm). *PG* and *FYM* caused a significant increase in available Mn. This may be due to the resultant decrease in soil *pH*. The highest values of available Mn were obtained with both the combined treatments of *PG₁S₁FYM₁* and *PG₂S₁FYM₁*. These results are in harmony with that obtained by Sharma and Yadav (1989) and Shams El-Din *et al.* (2000).

Table (5): Effect of amelioration processes on some micronutrients availability.

treatment	Fe (ppm)		Mn (ppm)		Zn (ppm)		Cu (ppm)	
	0 - 15	15 - 30	0 - 15	15 - 30	0 - 15	15 - 30	0 - 15	15 - 30
Phosphogypsum(PG):								
<i>PG₀</i>	12.926	11.513	5.878	4.443	0.694	0.387	0.951	0.612
<i>PG₁</i>	14.294	11.722	12.536	8.306	1.034	0.465	1.424	0.789
<i>PG₂</i>	15.469	12.231	14.571	9.959	1.507	0.578	1.634	0.945
F-test	**	Ns	**	**	**	*	**	**
LSD.05	0.776	-	2.182	1.971	0.334	0.124	0.167	0.102
0.01	1.172	-	3.305	2.986	0.475	-	0.209	0.145
Subsoiling (S):								
<i>S₀</i>	13.556	11.878	9.600	6.887	0.975	0.360	1.203	0.781
<i>S₁</i>	14.103	11.765	12.390	8.252	1.182	0.527	1.470	0.783
F-test	**	Ns	**	*	*	*	**	Ns
Farmyard manure (FYM):								
<i>FYM₀</i>	13.384	11.528	9.038	6.042	1.002	0.428	1.087	0.771
<i>FYM₁</i>	14.276	12.116	12.952	9.097	1.115	0.459	1.586	0.793
F-test	**	**	**	**	*	*	**	Ns
Interactions:								
<i>PGxS</i>	Ns	Ns	Ns	Ns	Ns	Ns	Ns	Ns
<i>PGxFYM</i>	*	Ns	Ns	Ns	Ns	Ns	**	Ns
<i>SxFYM</i>	Ns	Ns	Ns	Ns	Ns	Ns	*	Ns
<i>PGxSxFYM</i>	ns	ns	ns	ns	ns	ns	ns	Ns

6 – Zinc (Zn):

Data in Table (5) show that all the amelioration treatments resulted in a significant increase in the available Zn especially in the surface layer. The highest values of available Zn were achieved with both the combined

treatments of $PG_1S_1FYM_1$ and $PG_2S_1FYM_1$. Similar results were obtained by Abd El-Fattah *et al.* (1987) and Dora (1996).

7 - Copper (Cu):

Data in Table (5) show that all the amelioration treatments resulted in a significant increase in the available Cu in the surface layer (0 - 15 cm) and subsurface layer (15 - 30 cm). The highest values of available Cu were achieved with the combined treatments of $PG_1S_1FYM_1$ and $PG_2S_1FYM_1$. These results may be due to the improving effect of the the amelioration treatments on soil physical and chemical properties particularly pH and ESP. Similar results were obtained by Shams El-Din *et al.* (2000) and Abd El-Fattah *et al.* (1987).

It could be concluded that the tested amelioration treatments (PG, Subsoiling and FYM) have a positive effect on the availability of micronutrients (Fe, Mn, Zn and Cu). This effect diminishes with the soil depth.

B - Effect of amelioration processes on sorghum and barley yield:

Data in Table (6) show that the three amelioration processes have a highly significant effect on sorghum and barley yield. Phosphogypsum (PG) has a highly significant effect on the yield of both crops. These results may be attributed to that PG improved the soil properties such as porosity, ESP, pH and nutrients availability, which enhance plant growth.

Table (6): Effect of amelioration processes on sorghum and barley grain yield

Treatment	Sorghum (ton/fed.)	Barley (ton/fed.)
Phosphogypsum(PG):		
PG ₀	0.613	1.082
PG ₁	1.185	2.309
PG ₂	1.277	2.647
F-test	**	**
LSD 0.05	0.044	0.079
0.01	0.066	0.110
Subsoiling (S):		
S ₀	0.867	1.545
S ₁	1.183	2.481
F-test	**	**
Farmyard manure (FYM):		
FYM ₀	0.920	1.598
FYM ₁	1.130	2.428
F-test	**	**
Interactions:		
PGxS	**	**
PGxFYM	**	*
SxFYM	**	Ns
PGxSxFYM	**	Ns

Regarding subsoiling and FYM, it was noticed that subsoiling and FYM as well have a highly significant effect on both yields. Moreover, all possible combinations of interaction between the three treatments are highly

significant with respect to sorghum yield. But for barley, only the paired interaction between *PG* and the other two treatments (*Subsoiling* or *FYM*) were significant.

C-Effect of amelioration processes on elements Uptake by sorghum plants:

Data in Table (7) show the average uptake (g/plot) of N, P, K and Na by sorghum plants for two cuts. Data show that the uptake of the determined macronutrients was increased under the different amelioration processes. The highest increase occurred when the combination of the three amelioration processes were included, while the lowest increase was in the case of single process. It was noticeable that the opposite direction was observed with Na uptake, since the lowest values of Na uptake was occurred with the combination treatments of *PG₁S₁FYM₁* and *PG₂S₁FYM₁*, while the highest values were occurred with the control *PG₀SoFYM₀*. This may be due to the high level of salinity. These results are in agreement with those of Abdel-Rahman and Mikkelson (1986).

Regarding micronutrients, Table (8) shows Fe, Mn, Zn and Cu uptake. It was observed that the more combination the more increase in the uptake, i. e. the effect of each individual process on micronutrients uptake by sorghum plants was less than that of double or triple combination. These results are in agreement with those of Mohammed et al. (1993) and Dora (1996).

Table (7): Effect of amelioration processes on macronutrients (N, P, K) and Na uptake by sorghum.

Treatment	N Gr/plot	P Gr/plot	K Gr/plot	Na Gr/plot
Phosphogypsum(PG):				
PG ₀	30.428	7.264	35.241	7.693
PG ₁	65.722	20.796	85.199	9.825
PG ₂	74.245	24.015	89.644	10.199
F-test	**	**	**	**
LSD.05	7.467	2.802	10.797	1.033
0.01	11.265	4.239	16.261	1.505
Subsoiling (S):				
S ₀	45.857	13.937	56.951	9.431
S ₁	67.73	20.780	83.106	9.047
F-test	**	**	**	**
Farmyard manure (FYM):				
FYM ₀	45.073	14.005	61.447	9.064
FYM ₁	68.523	20.711	78.610	9.414
F-test	**	**	**	**
Interactions:				
PGxS	**	**	**	**
PGxFYM	**	*	*	*
SxFYM	Ns	Ns	*	*
PGxSxFYM	Ns	Ns	Ns	Ns

Table (8): Effect of amelioration processes on micronutrients (Fe, Mn, Zn and Cu) uptake by sorghum.

Treatment	Fe Mg/plot	Mn Mg/plot	Zn Mg/plot	Cu Mg/plot
Phosphogypsum (PG):				
PG ₀	610.59	78.244	72.191	6.058
PG ₁	1889.30	180.142	162.535	17.653
PG ₂	2125.76	200.336	182.623	19.944
F-test	**	**	**	**
LSD.05	315.36	19.280	28.822	1.729
0.01	477.67	28.870	43.543	2.602
Subsoiling (S):				
S ₀	1222.01	121.168	113.201	11.832
S ₁	1861.76	184.647	165.031	17.271
F-test	**	**	**	**
Farmyard manure (FYM):				
FYM ₀	1246.08	129.516	119.847	11.358
FYM ₁	1837.68	176.299	158.385	17.746
F-test	**	**	**	**
Interactions:				
PGxS	**	**	**	**
PGxFYM	**	**	**	**
SxFYM	**	Ns	Ns	Ns
PGxSxFYM	ns	*	ns	Ns

REFERENCES

- Abd El-Fattah, M.; K. Matter.; A.A. Khamis and M.M. Ghaly (1987). Effect of application gypsum under different water regimes on chemical properties of new reclaimed alkali soil. *Egypt. J. Soil.*; 27(3): 245 – 258.
- Abdel-Kharim, A. M. M. (1989). Role of organic matter on availability of some micronutrients in soils M. Sc. Thesis Fac. Agric. Zagazig University.
- Abdel-Rahman, A. M. and D. S. Mikkelson (1986). Chemical constituents of some rice varieties as influenced by different levels of some salinity. *J. Agric. Sci. Mansoura Univ.*, 11:463-467.
- Abrol, I. P. and D. R. Bhumbla (1975). Effect of application of different levels of gypsum on the yield of rice, wheat and barley grown in saline sodic soil. (C. F., El-Sabry 1986).
- Alcorido, I. S. and J. E. Rechcigl (1993). Phosphogypsum in agriculture: a review, in *Advances in Agronomy*, Vol. 49, Sparks, D. L., Ed., Academic Press, San Diego, 55
- Alva, A. K. and G. J., Gascho (1991) Differential leaching of cations and sulfate in gypsum amended soils, *Commun. Soil Sci. Plant Anal.*, 22, :1195.
- Chhabra, R.; I. P. Abrol and M. Y. Singh (1981). Dynamics of phosphorus during reclamation of sodic soils. *Soil Sci.* 132:319-324.

- Cochran, W. G. and G. M. Cox (1960). *Experimental Designs*, 2nd ed. John Wiley, New York, :293-316.
- Collings, R. K. (1980) Phosphogypsum in Canada, in *Phosphogypsum*, Borris, D. P. and Boody, P. W. Eds. Florida Institute of Phosphate research, Bartow, FL. :583.
- Cottenie, A.; M. Verso; L. Kiekens; G. Velghe and R. Camerlynck (1982a). *Chemical analysis of plant and soils*. Lab. Anal. And Agrochem. State Univ., Gent, Belgium, Chapter 2, 3 pp:14-54.
- Dora, A. M. (1996). Effect of some amendments on alkali soil properties and their relation to plant. M. Sc. Thesis, Fac. Of Agric., Minufia University.
- El-Fakharani, Y. M. (1997). Response of Broad beans to gypsum and poultry manure in a salt – affected Arid Environment *J. Agric. Sci. Mansoura Univ.*, 22:4023 – 4034.
- El-Gala, A. M.; M. A. Eid and H. G. Al-Shandoudy (1998). The effect of organic matter, sulphure and Fe application on the availability of certain nutrients in the soils of El-Dhahera area, Sultanat of Oman. *Arab. Univ. J. of Agric. Sci.*, 6(2):607-623.
- El-Koumei, B. Y.(1998). Influence of Zn, Cu and farmyard manure on wheat plants. *Zagazig J. Agric. Res.* Vol. 25 No. (4) 1998.
- El-Sabry, W. S. (1986). Reclamation processes and their effects on some properties and crops yield in deteriorated soils, at North Delta Ph. D. Thesis, Fac. Agric. Ain Shams.
- El-Sherief, M. A. B. (1997). Phosphorus nutrition of barley plant as influenced by Zinc, Manganese and organic matter application to saline soil. M. Sc. Thesis Fac. Agric. Menoufia University.
- Gazia, E. A. E. (2001) Effect of farmyard manure, gypsum and nitrogen on sugarbeet yield and NPK uptake. *Minufia J. Agric. Res.*, 26:1117-1128.
- Gazia, E. A. E.; S. A. Hassanein; H. A. Shams El-Din and M. S. M. Abo Soliman (1996). Effect of some amendments and mole drain systems on rice yield, water use efficiency and some soil properties. *J. Agric. Sci. Mansoura University*, 21:4701-4711.
- Genaidy, S. A.; H. A. El-Attar and M. A. Barakat (1989). Some major factors affecting rice yield in the soil of Egypt. *Egypt. J. Soil Sci.*, 29 (1):67-78.
- Ghazy, M. A. (1994). Effect of water regime, gypsum and sewage sludge increments on the improvement and productivity of saline-sodic soils. Ph. D. Thesis, Fac. Of Agric. Tanta University.
- Ghowail, S. I.; A. F. El-Ashtar; W. M. El-Ghamry and I. M. Antar (1978). Effect of added gypsum on ion uptake and chemical properties of some Egyptian soils. *Agric., Res. Rev.*, Cairo, 56 (5):41-51.
- Hamoud, H.S.M.(1992). Some factors affecting sugarbeet yield in some Egyptian soils. M.Sc. Thesis, Fac. Agric., Tanta University.
- Hussain, G. ; A. H. Hsanulh and D. G Westfall (1988). Reclamation of saline sodic-non-gypsiferous soils. *Pakistan: Arid Soil Res. And Rehabil* 2(1):29-47.
- Ismail, A. S.; A. A. Saker and A. A. Radwan (1991). Effect of leaching processes and application of some amendments on certain chemical properties and barley grown on an alluvial soil 2nd African Soil Sci. Soc., Conf. Pp. 144.

- Jackson, M. L. (1973). *Soil Chemical Analysis*. Constable and Company Ltd. London.
- Kaloosh, A.; M. Abou Bakr and E. El-Hadad (1989). Effect of addition organic materials from different sources to soil on CO₂ evaluation and nitrogen forms. *Com. Sci. Dev.*, 26:1-19.
- Mahmoud, H. M. M. (1994). Microbiological studies on the management urban wastes. Ph. D. Thesis, Fac. Of Agric. Moshtohor, Zagazig University, Egypt.
- Mahmoud, S. A.; T. H. Sheta and M. H. Gomaa (1996). Effect of applied gypsum and water regime on some chemical and nutritional properties of newly reclaimed saline alkali soils. *J. Agric. Sci. Mansoura University*, 21:2399-2407.
- Mohammed, S. S.; E., A. Abou Hussien and A. S. Ahmed (1993). Gypsum as amendment against soil alkalization in relation to tomato plants: 1- Effect on growth and chemical composition. *Egypt. J. of Appl.Sci.*, 8:73-84.
- Nadia, M. B.; M. E. A. Khalil and M. A. A. El-Emam (2000). Availability of N, P and K in sandy and clayey soils as affected by the addition of organic materials. *Egypt. J. Soil Sci.*, 40:265-283.
- Nafady, M. H.; H. G. Hassanien; M. I. Abu Zeid, and M. M. Ahmed (1993). Availability of phosphorus to Alfa-alfa and ryegrass in sandy calcareous soil as affected by adding organic manure or Nile sediments. *Assiut. J. of Agric. Sci.*, 24(1):59-71.
- Novikov, A. A.; P. V. Klassen and S. D. Evenchik (1990) The status and trends of phosphogypsum utilization in the USSR, in *Proc. 3rd Int. Symp. Phosphogypsum*, 2, Florida Institute Phosphate Research, Bartow, FL, :594.
- Rehcgil, Jack E. (1995). *Soil amendments and environmental Quality*. Lewis. Publishers New York, London.
- Shams El-Din, H. A.; E. A. E. Gazia; M. A. Abo El-Soud, and M. M. Ragab (2000). On-Farm Soil amelioration procedures of some salt affected soils at North Delta. 8th Conf. Agric. Dev. Res. Fac. Agric. Ain Shams Univ., Cairo, November 20-22, 2000. *Annals Agric. Sci. Sp. Issue1*, 269-288.
- Sharma, B. M. and J. S. P. Yodav (1989). Removal during leaching and availability of iron and manganese in pyrite and farmyard manure treated alkali soil. *Soil Sci.*, 147:17-22.
- Sonbol, H. A.; M. W. M. El-Agroudi and G. Labeeb (2001). Effect of nitrogen and gypsum on rice plant under saline sodic condition. 11-rice yield and nutrients uptake. *J. Agric. Sci. , Mansoura University* , 26:2465-2480.
- Swarup, A.(1982).Availability of Iron, Manganese, Zinc and Phosphorus in submerged sodic soils as affected by amendments during the growth period of rice crop, plant and soil, 66:37-43.
- Thakur, R. C. ; A. D. Bindra; R. D. Sood and M. Bhargeva (1995). Effect of fertilizer application and manuring on phisico-chemical properties of soil and grain yield in rice-wheat crop sequence. *Indian. J. Agron.*, 40(1):4-13.

Weast, R. C., (1981) Ed., Handbook of Chemistry and Physics. 61st ed., CRC press, Boca Raton, FL. B-72.

Youssef, N. N. (1992). Availability of some essential nutrients during submergence in gypsum treated saline sodic soil under different conditions. Zagazig J. Agric. Res. 19:2509-2522.

تأثير بعض عمليات تحسين الأراضي على صلاحية وامتصاص المذبات في
الأراضي المتأثرة بالأملاح
السيد عامر السيد جازية
معهد بحوث الأراضي والمياه والبيئة - مركز البحوث الزراعية

تنتشر الأراضي المتأثرة بالأملاح في مصر في منطقة شمال الدلتا، وان تحسين مثل هذه الأراضي عادة يكون باستخدام مصلحات كيميائية (مثل الجبس أو الجبس المفسفر) أو باستخدام الأسمدة العضوية، بالإضافة الى استخدام الحرث العميق تحت التربة. وقد نفذت هذه الدراسة على مدى موسمين زراعيين متتاليين (٢٠٠ و ٢٠٠١/٢٠٠٠) في أراضى بور ومتأثرة بالأملاح لدراسة تأثير عمليات التحسين (إضافة الجبس المفسفر - الحرث العميق تحت التربة - و/أو السماد البلدي) على صلاحية العناصر الكبرى والصغرى وامتصاصها بواسطة نبات الذرة الرفيعة (السورجم)، الى جانب تأثير عمليات التحسين على إنتاجية هذه الأراضي ومحصولي الذرة الرفيعة (السورجم) والشعير. وقد أوضحت الدراسة أن عمليات التحسين المستعملة (إضافة الجبس المفسفر - الحرث العميق تحت التربة - و/أو السماد البلدي) أدت الى زيادة صلاحية كل من العناصر الكبرى (النيتروجين - الفوسفور - البوتاسيوم) والصغرى (حديد - منجنيز - زنك - نحاس) خصوصا في الطبقة السطحية. بالإضافة الى ذلك، فإن لهذه العمليات تأثير عالي المعنوية على كل من محصولي الذرة الرفيعة (السورجم) والشعير. فضلا عن أن هذه العمليات قد أدت الى زيادة كمية العناصر الكبرى والصغرى الممتصة بواسطة نبات الذرة الرفيعة (السورجم) ما عدا عنصر الصوديوم الذي تناقص امتصاصه بسبب إضافة الجبس المفسفر والحرث العميق تحت التربة.